

Rate Constants for the Decay and Reactions of the Lowest Electronically Excited Singlet State of Molecular Oxygen in Solution. An Expanded and Revised Compilation

Francis Wilkinson

Department of Chemistry, University of Technology, Loughborough, Leicestershire, LE11 3TU, United Kingdom

W. Phillip Helman and Alberta B. Ross

Radiation Chemistry Data Center, Radiation Laboratory, University of Notre Dame, Notre Dame, Indiana 46556

Received 1 November 1994; revised manuscript received 13 February 1995

An expanded and revised compilation on the reactivity of singlet oxygen, the lowest electronically excited singlet state of molecular oxygen, ${}^1\text{O}_2^*({}^1\Delta_g)$, in fluid solution is presented, which supersedes the publication of Wilkinson and Brummer, J. Phys. Chem. Ref. Data **10**, 809 (1981). Rate constants for the chemical reaction and physical deactivation of singlet oxygen available through 1993 have been critically compiled. Solvent deactivation rates (k_d) are tabulated for 145 solvents or solvent mixtures and second-order rate constants for interaction of singlet oxygen with 1915 compounds are reported. ©1995 American Institute of Physics and American Chemical Society.

Key words: Chemical kinetics; data compilation; decay; oxidation; photochemistry; photosensitization; quenching; rate constants; singlet oxygen; solution.

Contents

1. Introduction.....	664
2. Sources of Singlet Oxygen in Solution.....	665
2.1. Direct Absorption by O_2 (${}^3\Sigma_g^-$).	665
2.2. Production Following Absorption into O_2 (${}^3\Sigma_g^-$): Organic Molecule Charge-Transfer Bands.....	665
2.3. Photosensitized Production.....	665
2.4. Sensitized Production by Pulsed Radiolysis....	666
2.5. Microwave Generation of Singlet Oxygen....	666
2.6. Chemical Production of Singlet Oxygen.....	667
3. Methods for the Direct Study of the Kinetics of the Reactions of Singlet Oxygen.....	667
3.1. Luminescence Decay—Time Resolved Infrared Luminescence.....	667
3.2. Time Dependent Thermal Lensing due to Singlet Oxygen Decay.....	667
3.3. Time Dependent Acceptor Disappearance....	667
3.3.1. Following Pulsed Excitation of a Sensitizer.....	667
3.3.2. Following Pulsed Excitation of Solutions Containing High Pressures of O_2	668
3.4. Kinetic Analysis to give the Decay of Singlet Oxygen by Monitoring the Decay of Triplet β -carotene Produced by Energy Transfer from Singlet Oxygen.....	668
4. Methods for Obtaining Kinetic Information from Studies using Continuous Irradiation i.e., Under Photostationary State Conditions.....	668
4.1. Luminescence Intensity Quenching.....	668
4.2. Monitoring Sensitized Photo-oxidations by Consumption of Reactants or Appearance of Products Under Steady-State Conditions.....	669
4.2.1. Sensitized Photo-oxygenation of a Single Substrate A.....	669
4.2.2. Photo-Auto-Oxygenation of a Single Substrate A.....	670
4.2.3. Sensitized Photo-oxygenations of a Substrate A in the Presence of a Second Substrate A' which also Reacts with Singlet Oxygen.....	670
4.2.4. Sensitized Photo-Oxygenations of a Substrate A' in the Presence of A=Q, a Physical Quencher of Singlet Oxygen.....	670
4.2.5. Sensitized Photo-Oxygenation of a Substrate A' in the Presence of A=Q, a Quencher of Both Singlet Oxygen and the Sensitizer Triplet.....	671
4.2.6. Separation of k_t and k_q	672

5. Kinetic Analysis for Oxygenation by Singlet Oxygen Generated by Chemical Reaction, by Microwave Discharge, or by Direct Laser Excitation.....	671
5.1. When Singlet Oxygen is Produced Chemically.....	671
5.2. When Singlet Oxygen is Produced by Microwave Discharge in the Gas Phase.....	672
5.3. When Singlet Oxygen is Produced by Direct CW Laser Excitation of Oxygen.....	672
6. First Order Rate Constants for the Decay of Singlet Oxygen in Various Solvents—Comments on Table 1.....	672
7. Second Order Rate Constants for Reactive and Physical Quenching of Singlet Oxygen by Various Substrates—Comments on Table 2–17....	673
7.1. Scope of the Compilation.....	673
7.2. Arrangement of Tables.....	673
7.2.1. Solvent	674
7.2.2. Rate Constants (k) and Beta (k_d/k) Values.....	674
7.2.3. Temperature.....	674
7.2.4. Method.....	674
7.2.5. Comment.....	674
7.2.6. References.....	674
8. Mechanism of Quenching.....	674
9. List of Abbreviations and Symbols.....	675
10. Acknowledgements.....	677
11. References to Text.....	678
12. References to Tables.....	917
13. Molecular Formula Index.....	935
14. Chemical Name Index.....	965
11. Rate constants for interaction of singlet oxygen with diazo compounds and dyes.....	840
12. Rate constants for interaction of singlet oxygen with metal complexes.....	858
13. Rate constants for interaction of singlet oxygen with sulfur, selenium and tellurium compounds...	875
14. Rate constants for interaction of singlet oxygen with some inorganic compounds.....	893
15. Rate constants for interaction of singlet oxygen with oximes, hydrazones, nitrones, nitroso compounds and <i>N</i> -oxides.....	898
16. Rate constants for interaction of singlet oxygen with some compounds of biological interest.....	904
17. Rate constants for interaction of singlet oxygen with miscellaneous substrates.....	911

1. Introduction

Reactions of singlet oxygen ${}^1\text{O}_2^*({}^1\Delta_g)$, are of much current interest because of their importance in many photooxidations of chemical and biological systems including reactions used in photo-chemotherapy. There is strong evidence for the involvement of singlet oxygen, a powerful oxidant, in many photosensitized oxidations, photodynamic inactivation of viruses and cells, in phototherapy for cancer, in photocarcinogenesis, in the photodegradation of dyes and polymers and in the dye sensitization of the photodegradation of polymers. [See Refs. 1–7 and references therein].

The ground electronic state of molecular oxygen, which has zero angular momentum about the internuclear axis and contains two unpaired p electrons, has the group theoretical symbol ${}^3\Sigma_g^-$. The two electronically excited singlet states which arise from this same electron configuration but with spin pairing of the two electrons are the ${}^1\Delta_g$ and the ${}^1\Sigma_g^+$ states which lie at 94 and 157 kJ mol⁻¹, respectively, above the ${}^3\Sigma_g^-$ ground state. The electronic transitions ${}^1\Delta_g \leftarrow {}^3\Sigma_g^-$ and ${}^1\Sigma_g^+ \leftarrow {}^3\Sigma_g^-$ although highly forbidden are readily observed in absorption and emission in the upper atmosphere with zero-zero transitions at 1,269 and 762 nm and estimated radiative lifetimes of 64 min and 10 s, respectively.^{8,9} The measured lifetimes in the gas phase and in solution are very much shorter than this. In condensed media, the lifetime of ${}^1\text{O}_2^*({}^1\Sigma_g^+)$ is short, 130 ns in carbon tetrachloride, and until recently very little was known about its properties,¹⁰ and thus the term singlet oxygen is used throughout this review to refer to the ${}^1\Delta_g$ state.

In 1981 we published a critical comprehensive compilation which reported first order decay constants for 50 solvents and second order rate constants for deactivation and chemical reaction of singlet oxygen with 690 different compounds. This article updates and replaces our 1981 article.⁵ It uses identical equation numbers and symbols wherever possible as did our recent compilation,⁶ "Quantum Yields for the Photosensitized Formation of the Lowest Electronically Excited Singlet State of Molecular Oxygen in Solution." It covers the literature up to the end of 1993 and includes solvent deactivation rates (k_d) for 145 solvents and solvent mixtures and rate constants for chemical reaction and physical

List of Tables


1. Decay constants for singlet oxygen in various solvents.....	680
2. Rate constants for interaction of singlet oxygen with olefins.....	699
3. Rate constants for interaction of singlet oxygen with aromatic substrates.....	740
4. Rate constants for interaction of singlet oxygen with phenols and naphthols.....	755
5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds.....	779
6. Rate constants for interaction of singlet oxygen with pyrroles, oxazoles, pyridines and other heterocyclic nitrogen compounds.....	797
7. Rate constants for interaction of singlet oxygen with porphyrins, phthalocyanines and polypyroles.....	804
8. Rate constants for interaction of singlet oxygen with aliphatic and alicyclic amines.....	817
9. Rate constants for interaction of singlet oxygen with aromatic amines.....	829
10. Rate constants for interaction of singlet oxygen with amino acids, peptides and proteins.....	833

deactivation of singlet oxygen with over 1,900 different compounds.

2. Sources of Singlet Oxygen in Solution


2.1. Direct Absorption by O_2 ($^3\Sigma_g^-$)

Generation of $^1O_2^*$ ($^1\Delta_g$) by direct absorption has been accomplished by irradiation with Nd:YAG lasers of Freon solutions subjected to high pressures of O_2 , by Matheson and Lee¹¹⁻¹³ and also by Evans¹⁴ using the simultaneous transition $2 O_2$ ($^3\Sigma_g$, $^3\Sigma_g$) $\rightarrow 2 O_2^*$ ($^1\Delta_g$, $^1\Delta_g$). In addition Singh *et al.*¹⁵ have shown that it is also possible to produce $^1O_2^*$ ($^1\Delta_g$) by irradiation of aerated water at 600 nm via vibrationally excited water with a yield of $\sim 3 \times 10^{-5}$, i.e.,



2.2. Production Following Absorption into O_2 ($^3\Sigma_g^-$): Organic Molecule Charge-Transfer Bands

The perturbing effect of dissolved oxygen on the uv/vis absorption spectra of organic molecules is a well known phenomenon first studied in detail by Evans¹⁶ and subsequently discussed by Mulliken¹⁷ and others.¹⁸ The additional absorption bands observed include enhanced absorption of the lowest energy transition in the organic molecule, M, which corresponds to the $S_0 \rightarrow T_1$ transition of the organic molecule in intimate contact with an oxygen molecule. More intense absorption is also often observed at shorter wavelengths which is attributed to CT transitions within $^3(M..O_2)$ contact complexes, viz,



This effect is easily observed by comparing the absorption spectra of liquid aromatic hydrocarbons such as benzene,

toluene, or *o*-, *m*-, or *p*-xylenes saturated with oxygen and in the absence of dissolved oxygen. The oxygen induced red shift observed, which is reversible when the oxygen is removed, has been assigned to charge transfer absorption. This assignment is supported by the broad structureless appearance of the bands. Scurlock and Ogilby^{19,20} have demonstrated that irradiation of O_2 ($^3\Sigma_g^-$):organic molecule charge-transfer bands produces singlet oxygen with relatively high yields. Picosecond laser flash photolysis studies have been carried out in attempts to understand the mechanism involved.^{21,22}

2.3. Photosensitized Production

The method most frequently used for producing singlet oxygen in the laboratory is photo-sensitization (see Scheme 1). Both continuous irradiation and pulsed excitation studies have made much use of this method of production which also frequently occurs in nature. In this mechanism (Scheme 1) the lowest excited singlet and triplet states of the sensitizers are represented as S_1 , and T_1 , respectively, and the ground electronic state by S_0 . Any substrate which deactivates singlet oxygen by physical or chemical quenching is represented in Scheme 1 by M. However, when quenching is known to be mainly by chemical reaction we shall use A to represent a reactive substrate and if a second reactive substrate is employed A' will be used. When the molecule is known to be a physical quencher with negligible reaction, Q will be used in place of M. The singlet oxygen sensitizer, even when the sensitizer quenches singlet oxygen, will be represented by S. Thus M is a general symbol in Scheme 1 which may be replaced by A, A', Q, or S or any combination of these. In the tables we have used A throughout for all substrates irrespective of whether they were known to have little measured chemical reactivity.

Scheme 1

	Rate Constants	Quantum Yields and Fractional Probabilities
1 $S_0 + h\nu \rightarrow S_1$	rate = I_a	
2 $S_1 \rightarrow S_0 + h\nu_F$	k_F	$\phi_F = \frac{k_F}{k_{SD}}$
3 $S_1 \rightarrow S_0$	k_{ic}	
4 $S_1 \rightarrow T_1$	k_{isc}	$\phi_T = \frac{k_{isc}}{k_{SD}}$
5 $S_1 + O_2 \rightarrow T_1 + ^1O_2^*$	$k_{SA}^{O_2}$	$f_\Delta^S = k_{SA}^{O_2}/k_{SQ}^{O_2}$
6 $S_1 + O_2 \rightarrow T_1 + O_2$	$k_{isc}^{O_2}$	$f_T^{O_2} = (k_{SA}^{O_2} + k_{isc}^{O_2})/k_{SQ}^{O_2}$
7 $S_1 O_2 \rightarrow S_0 + ^3O_2$	$k_{sd}^{O_2}$	$P_S^{O_2} = k_{SQ}^{O_2}[O_2]/(k_{SD} + k_{SQ}^{O_2}[O_2])$
8 $S_1 + O_2 \rightarrow \text{products}$	$k_{sr}^{O_2}$	$1 - P_S^{O_2} = 1/(1 + k_{SQ}^{O_2}\tau_S^0[O_2])$
		$\phi_T^{O_2} = \phi_T(1 - P_S^{O_2}) + f_T^{O_2}P_S^{O_2}$
9 $T_1 \rightarrow S_0 + h\nu_P$	k_{TP}	
10 $T_1 \rightarrow S_0$	k_{TD}	$k_{TD} = (\tau_T^0)^{-1}$
11 $T_1 + O_2 \rightarrow S_0 + ^1O_2^*$	$k_{TA}^{O_2}$	$f_\Delta^T = k_{TA}^{O_2}/k_{TQ}^{O_2}$
12 $T_1 + O_2 \rightarrow S_0 + ^3O_2$	$k_{Td}^{O_2}$	
13 $T_1 + O_2 \rightarrow \text{products}$	$k_{Tr}^{O_2}$	$P_T^{O_2} = k_{TQ}^{O_2}[O_2]/(k_{TD} + k_{TQ}^{O_2}[O_2])$
14 $^1O_2^* \rightarrow ^3O_2 + h\nu_P$	$k_{\Delta P}$	$f_P^\Delta = k_{\Delta P}/(k_{\Delta P} + k_{\Delta d})$
15 $^1O_2^* \rightarrow ^3O_2$	$k_{\Delta d}$	
16 $^1O_2^* + M \rightarrow \text{products}$	k_r^M	$f_r^M = k_r^M/(k_r^M + k_q^M)$
17 $^1O_2^* + M \rightarrow ^1M + ^3O_2$	k_q^M	

f_Δ^S = fraction of S_1 quenched by O_2 which gives $^1O_2^*$

$f_T^{O_2}$ = fraction of S_1 quenched by O_2 which gives T_1

f_Δ^T = fraction of T_1 quenched by O_2 which gives $^1O_2^*$

$P_S^{O_2}$ = proportion of S_1 quenched by O_2

$P_T^{O_2}$ = proportion of T_1 quenched by O_2

The quantum yield of singlet oxygen production, ϕ_Δ , via sensitization is given by the sum of contributions due to oxygen quenching of S_1 and T_1 , i.e.,

$$\phi_\Delta = \phi_\Delta(S) + \phi_\Delta(T) = P_S^{O_2}f_\Delta^S + \phi_T^{O_2}P_T^{O_2}f_\Delta^T.$$

In the presence of oxygen quenching of S_1 and T_1

$$\phi_\Delta = P_S^{O_2}f_\Delta^S + \phi_T(1 - P_S^{O_2})P_T^{O_2}f_\Delta^T + P_S^{O_2}f_T^{O_2}P_T^{O_2}f_\Delta^T,$$

which can be written as

$$\phi_\Delta = \phi_T P_T^{O_2}f_\Delta^T + P_S^{O_2}\{f_\Delta^S + P_T^{O_2}f_\Delta^T(f_T^{O_2} - \phi_T)\}. \quad (1)$$

Methods for determining ϕ_Δ were reviewed by us⁶ and 1,600 quantum yield determinations were tabulated for 750 different chemical species.

2.4. Sensitized Production by Pulsed Radiolysis

When a high energy electron beam is passed through liquid benzene, for example, a high yield of excited states can be formed. Since the lifetimes of singlet and triplet states of

benzene are only a few nanoseconds in neat benzene, quenching by dissolved oxygen does not produce much singlet oxygen. However, in the presence of a triplet energy acceptor, e.g., 10^{-2} mol L⁻¹ naphthalene, energy transfer from triplet benzene occurs to give triplet naphthalene which is quenched by oxygen producing singlet oxygen. This enables the properties of singlet oxygen to be probed using the technique of pulsed radiolysis.²³

2.5. Microwave Generation of Singlet Oxygen

When a microwave discharge is passed through oxygen gas in a gaseous flow system, singlet molecular oxygen ($^1\Delta_g$ and $^1\Sigma_g^+$), oxygen atoms, and ozone are produced. The latter two oxidizing species can be removed by reaction with mercury vapor and the $^1O_2^*$ ($^1\Delta_g$) is rapidly quenched, probably to give $^1O_2^*$ ($^1\Delta_g$) when the emerging gases are bubbled through solutions containing oxidizable substrates. Often the flow is bubbled simultaneously through two cells, one containing the substrate only and the other containing substrate

and potential singlet oxygen quencher.²⁴ This makes it easier to allow for variations in the concentration of dissolved singlet oxygen produced in these bubbling experiments.

2.6. Chemical Production of Singlet Oxygen

Among the reactions used to form singlet oxygen are (a) reaction of hydrogen peroxide with hypochlorite or hypobromite,²⁵ (b) reaction of potassium superoxide with water, and (c) thermal decomposition of aryl peroxides or of the ozonide of triphenyl phosphite.²⁴ Experiments are usually carried out so that a fixed amount of singlet oxygen is produced in the presence of variable amounts of reactive substrate A and/or A' with or without added quencher, Q.

3. Methods for the Direct Study of the Kinetics of the Reactions of Singlet Oxygen

3.1. Luminescence Decay—Time Resolved Infrared Luminescence

Although the quantum yield of phosphorescence of singlet oxygen is very low in most solvents, the development of germanium photodiodes with high gain, coupled with wide-band amplifiers which can give overall time responses of less than a microsecond, means that the lifetime of singlet oxygen in most solvents can be measured accurately by monitoring infrared luminescence at ~ 1270 nm.^{26,27} If the logarithm of the phosphorescence intensity is plotted as a function of time then the slope gives the decay constant

$$k_D = k_\Delta + \Sigma k_M [M], \quad (2)$$

where M represents any molecule including the sensitizer which quenches singlet oxygen, physically or chemically. If k_D is measured as a function of the concentration of a particular quencher M, keeping all other quencher concentrations constant, the bimolecular rate constant for quenching of singlet oxygen by M, k_M can be obtained.^{28,29} Measurement of k_D as a function of sensitizer concentration allows k_S to be evaluated. When a decay constant for singlet oxygen is not extrapolated to zero sensitizer concentration we shall use $k_d = (k_\Delta + k_S[S])$ for the decay constant. It is often assumed that under many experimental conditions $k_d = k_\Delta$, however this is not always the case.

3.2. Time Dependent Thermal Lensing due to Singlet Oxygen Decay

The absorption of even part of the energy in a laser pulse gives rise to local temperature changes in gases or liquids. This leads to changes in density and refractive index which causes the system to act as a diverging lens.³⁰ Time resolved thermal lensing (TL) due to release of energy by decaying excited states can be used to measure lifetimes of singlet oxygen in the range 0.1 to 100 μs . The probe source is a continuous laser beam which is deflected by the thermal lens. The signal gives the relative magnitudes of heat contributions (U) for fast and slow non-radiative processes relative to

the acoustic transit time. The time dependence of the slow process gives rate information. The TL signal at time t is defined as

$$U(t) = \frac{V_0 - V(t)}{V_0}, \quad (3)$$

where $V(t)$ is the time dependent voltage generated by the detector. Following the sensitized production of ${}^1\text{O}_2^*$, the thermal buildup $\Delta U = U_{\text{total}} - U_{\text{fast}}$ decreases exponentially with the decay constant k_D given by Eq. (2).

3.3. Time Dependent Acceptor Disappearance

3.3.1. Following Pulsed Excitation of a Sensitizer

According to the mechanism given in Scheme 1, in this case with M=A, since we are discussing a reactive substrate, it follows that

$$-\frac{d[A]}{dt} = k_r^A [A] [{}^1\text{O}_2^*].$$

Following pulsed excitation in aerated solutions most singlet and triplet states have decayed within 1 μs and then

$$-\frac{d[{}^1\text{O}_2^*]}{dt} = (k_d + k_A [A]) [{}^1\text{O}_2^*]$$

and thus

$$-d[A] = k_r^A [A] [{}^1\text{O}_2^*]_{t=0} \left[\exp \left(- \left(k_d t + \int_0^t k_A [A] dt \right) \right) \right]. \quad (4)$$

Equation (4) has been treated slightly differently as follows:

(a) Merkel and Kearns^{31,32} assumed that [A] on the right hand side of Eq. (4) can be treated as constant (i.e., as $[A]_{av}$ since typically [A] only varies by $\sim 10\%$). Taking $k_D = k_d + k_A [A]_{av}$, integration of Eq. (4) gives,

$$[A]_{t=0} - [A]_\infty = \frac{k_r^A [{}^1\text{O}_2^*]_{t=0} [A]_{av}}{k_D} e^{-k_D t}. \quad (5)$$

Since the change in concentration of A is proportional to ΔA , the change in absorbance by the oxidizable acceptor at some convenient wavelength, $[A]_{t=0} - [A]_\infty = \Delta A / \epsilon l$, where ϵ is the extinction coefficient and l the analyzing path-length. It follows from Eq. (5) that

$$\ln \Delta A = -k_D t + \text{const}, \quad (6)$$

and a plot of $-\ln \Delta A$ vs t should have a slope equal to $k_D = k_d + k_A [A]_{av}$. Thus by varying $[A]_{av}$ values k_d and k_A can be determined.

(b) Adams and Wilkinson³³ and Young *et al.*³⁴ replaced [A] by $[A]_{av}$ only in the exponential term in Eq. (4) and then integrated Eq. (4) to give

$$\ln \left[\frac{[A]}{[A]_0} \right] = \frac{(k_r^A [{}^1\text{O}_2^*]_{t=0})}{k_D} [e^{-k_D t} - 1] \quad (7)$$

or

$$\ln \left[\frac{[A]}{[A]_\infty} \right] = \frac{(k_r^A [{}^1\text{O}_2^*]_{t=0})}{k_D} e^{-k_D t}. \quad (8)$$

Experimental results were fitted to Eqs. (7) or (8) to evaluate k_D and thus values of k_d and k_A were obtained.

Comparison of the values obtained from the same data for k_D using Eqs. (5) or (7) and (8) give agreement to within $\pm 5\%$ and do not differ, within an experimental error of about $\pm 10\%$ when experiments are repeated many times, from computer treatments which do not make any assumptions about [A] being relatively constant. If k_A is partly due to physical quenching and partly due to reaction, the value of k_A obtained will be the total rate constant for quenching due to both processes.


In the presence of several possible singlet oxygen quenchers, i.e., Scheme 1 with M=A, S, and Q, we have

$$k_D = k_\Delta + k_S[S] + k_Q[Q] + k_A[A]. \quad (9)$$

Measurement of k_D as a function of [Q] keeping [S] and [A] constant allows values of k_Q to be obtained. N.B. If k_Q is only partly due to physical quenching and partly due to reaction, the value of k_Q obtained is the total rate constant for quenching due to both processes.

3.3.2. Following Pulsed Excitation of Solutions Containing High Pressures of O₂

A variation on this method has been developed by Matheson *et al.*^{11,35} in which singlet oxygen is directly generated by absorption of the output at 1065 nm of a pulsed Nd-glass laser by oxygen dissolved under pressure (up to 130 atm) in 1,1,2-trichlorotrifluoroethane (Freon 113). The disappearance of the singlet oxygen acceptor 1,3-diphenylisobenzofuran (DPBF) was monitored. Because of the high concentration of oxygen present, quenching by ground state oxygen, i.e., due to the reaction,



contributes substantially to singlet oxygen decay, and




$$k_D = k_d + k_A[A] + k_q^{O_2}[\text{O}_2]. \quad (10)$$

Under these conditions $k_q^{O_2}[\text{O}_2] \gg k_d$, therefore only values of k_A and $k_q^{O_2}$ can be obtained by measuring k_D as a function of [A] or [O₂] respectively. Recent measurements suggest that k_A is pressure dependent.³⁶

3.4. Kinetic Analysis to give the Decay of Singlet Oxygen by Monitoring the Decay of Triplet β-carotene Produced by Energy Transfer from Singlet Oxygen

Farmilo and Wilkinson³⁷ have developed a method for measuring singlet oxygen decay which monitors absorption by triplet β-carotene, $^3\text{C}^*$, formed by energy transfer from singlet oxygen in aerated solutions containing a sensitizer and a low concentration of β-carotene. Consider the mechanisms given in Scheme 1 in the presence of β-carotene, C,

i.e., a mechanism involving steps 1–17 with M=Q together with steps 18–21 given below:



The differential rate equations which can be written for $d[T_1]/dt$, $d[^1\text{O}_2^*]/dt$, and $d[^3\text{C}^*]/dt$ can be solved without making the steady-state approximation and this gives the concentration of $[^3\text{C}^*]$ as

$$[^3\text{C}^*] = V[e^{-k_T t} - e^{-k_C t}] + W[e^{-k_D t} - e^{-k_C t}],$$

where $k_T = k_{TQ}^{O_2}[\text{O}_2] + k_{TC}[C]$; $k_C = k_{CO}[\text{O}_2] + k_{dC}$, and $k_D = k_d + k_{\Delta C}[C] + k_Q[Q]$. In aerated solutions the values of $k_{TQ}^{O_2}[\text{O}_2]$ and $k_{CO}[\text{O}_2]$ are such that after $\sim 1 \mu\text{s}$, $\exp(-k_T t)$ and $\exp(-k_C t)$ become negligibly small and thus

$$[^3\text{C}^*] = We^{-k_D t}, \quad (11)$$

where $W = [T_1]_{t=0} k_{\Delta C}[C] k_{TQ}^{O_2}[\text{O}_2] / [(k_T - k_D)(k_C - k_D)]$. W is a constant provided [O₂], [C], and [Q] are constant. It is usually possible to arrange for these to be present in excess so that $[\text{O}_2] \gg [^1\text{O}_2^*]$, $[C] \gg [^3\text{C}^*]$ and any consumption of Q or O₂ must also be negligibly small. It is important to bear these conditions in mind especially for work which involves, for example, focussed, high-energy laser pulses. However, when these conditions are met it follows from Eq. (11) that the decay of triplet β-carotene after $\sim 1 \mu\text{s}$ becomes first order with a decay constant k_D equal to that of its precursor, singlet oxygen as confirmed by Farmilo and Wilkinson.³⁷ Thus the decay of absorption at 520 nm due to $^3\text{C}^*$ under these conditions mirrors the singlet oxygen decay yielding values of k_D and hence values of k_d , $k_{\Delta C}$, and k_Q .

4. Methods for Obtaining Kinetic Information from Studies using Continuous Irradiation, i.e., Under Photostationary State Conditions

4.1. Luminescence Intensity Quenching

The phosphorescence intensity due to emission of singlet oxygen at 1270 nm, I_Δ , observed when a sensitizer is continuously irradiated depends on the steady-state concentration of singlet oxygen present in a solution. When a photostationary state of singlet oxygen is produced its rate of production $I_\Delta \phi_\Delta$ equals the rate of decay $k_d [^1\text{O}_2^*]_{ss}$ and thus

$$[^1\text{O}_2^*]_{ss} = \phi_\Delta \frac{I_\Delta}{k_d}.$$

In the presence of a singlet oxygen quencher M the steady-state singlet oxygen concentration $[{}^1\text{O}_2^*]_{\text{ss}}^{\text{M}}$ is reduced, i.e.,

$$[{}^1\text{O}_2^*]_{\text{ss}}^{\text{M}} = \frac{\phi_{\Delta} I_a}{k_d + (k_r^{\text{M}} + k_q^{\text{M}})[\text{M}]}$$

Krasnovsky³⁸ and others have shown that a Stern-Volmer relationship

$$\frac{L_{\Delta}^0}{L_{\Delta}} = 1 + \frac{(k_r^{\text{M}} + k_q^{\text{M}})[\text{M}]}{k_d} \quad (12)$$

holds for quenching of singlet oxygen phosphorescence by physical and chemical quenchers; from the slopes of plots of L_{Δ}^0/L_{Δ} vs [M] values of the total quenching constant ($k_r^{\text{M}} + k_q^{\text{M}}$) can be obtained provided the singlet oxygen decay constant in the absence of the quencher M is known.

Krasnovsky³⁸ pioneered the use of phosphorescence intensity measurements from singlet oxygen in solution. He and other workers have used this method to derive quenching rate constants using the Stern-Volmer equation. In addition, by assuming that $k_{\Delta P}$ is independent of solvent, Turro³⁹ has given estimates of lifetimes relative to a standard, in this case benzene, for which a value of $k_d = 4 \times 10^4 \text{ s}^{-1}$ was used.⁵ Measurements were made of the decrease in fluorescence intensity of a CCl_4 solution upon addition of other solvents and it was found that the Stern-Volmer relationship was observed, i.e.,

$$\frac{I_0}{I} = \kappa x + 1, \quad \text{where } \kappa = \frac{\tau_{\text{CCl}_4}}{\tau_{\text{solvent}}},$$

where x is the volume fraction. The question of the extent to which the radiative probability for phosphorescence from singlet oxygen varies with solvent is debatable.⁴⁰

4.2. Monitoring Sensitized Photo-oxidations by Consumption of Reactants or Appearance of Products Under Steady-State Conditions

Two major classes of photosensitized oxygenations have been designated as Type I and Type II. In the former, the excited sensitizer interacts directly with the substrate resulting, for example, in either H-atom transfer or electron transfer. The radicals so produced from the sensitizer react in the presence of oxygen to regenerate the sensitizer, while radicals produced from the substrate, for example, initiate free radical chain reactions, as observed in auto-oxidations. Type II reactions involve the direct interaction of the excited sensitizer and oxygen with energy transfer to give singlet oxygen, which reacts with various substrates.⁴¹

4.2.1. Sensitized Photo-oxygenation of a Single Substrate A

Many photosensitized reactions have been carried out under conditions such that no quenching by A of the sensitizer singlet or triplet states occurs, in which case the mechanism is as given in Scheme 1, with M=A. For continuous irradiation,

$$-\frac{d[A]}{dt} = -\frac{d[\text{O}_2]}{dt} = \frac{d[P]}{dt} = k_r^{\text{A}} [{}^1\text{O}_2^*]_{\text{ss}}[A], \quad (13)$$

and the steady-state approximation can be applied to give


$$\phi_{\text{ox}} = \frac{r_{\text{ox}}}{I_a} = \frac{\phi_{\Delta} k_r^{\text{A}} [A]}{k_d + k_A [A]}, \quad (14)$$

where the rate of oxygenation (r_{ox}) may be followed by determining the rate of appearance of some product (Pa) and/or the rate of consumption of either the substrate A (Ac) and/or of oxygen (Oc). All three have been used, and this is indicated in the tables by an entry in the methods column. N.B. It follows from Eq. (14) that when $k_d \gg k_A [A]$ or $k_A [A] \gg k_d$ the rate of oxygenation will be first or zero order with respect to A, respectively. Equation (14) can be rearranged to give

$$r_{\text{ox}}^{-1} = (I_a \phi_{\Delta} f_r^{\text{A}})^{-1} \left[1 + \frac{k_d}{k_A} \frac{1}{[A]} \right], \quad (15)$$

where $f_r^{\text{A}} = k_r^{\text{A}}/k_A$ is the fraction of reactive quenching of singlet oxygen by A.

Most reactive substrates, A, react with singlet oxygen with unit stoichiometry as implied in Eqs. (13) to (15) which apply when each time a molecule of ${}^1\text{O}_2^*$ reacts it leads to the consumption of one molecule of A and one of O_2 and produces one molecule of the product P. If there are stoichiometry factors α different from unity, the overall reaction is given by the equation



then

$$r_{\text{ox}} = -\frac{1}{\alpha_A} \frac{d[A]}{dt} = -\frac{1}{\alpha_{\text{O}_2}} \frac{d[\text{O}_2]}{dt} = \frac{1}{\alpha_P} \frac{d[P]}{dt},$$

and Eqs. (14) and (15) still apply. If, however, the rate of the reaction is monitored by observing the rate of consumption of a reactive substrate such as thiourea where $\alpha=2$ and this rate is not divided by $\alpha=2$, then αf_r^{A} appears in Eqs. (14) and (15). This needs to be borne in mind whenever unit stoichiometry does not apply.

According to Eq. (15) linear plots of r_{ox}^{-1} or ϕ_{ox}^{-1} vs $[A]^{-1}$ should give

$$\frac{\text{slope}}{\text{intercept}} = \frac{k_d}{k_A} = \beta_A,$$



where $\beta_A = k_d/k_A$ represents the concentration at which the decay of singlet oxygen in the solvent alone (steps 14 and 15) equals the decay due to quenching by A (steps 16 and 17), i.e., it is the half-quenching concentration. [N.B. Eq. (15) only predicts a linear relationship if there is (i) constant light intensity, (ii) constant absorption by the sensitizer and with no absorption by A although this could be allowed for, and (iii) a constant oxygen concentration in the solution]. Tables 2 to 17 list hundreds of β values along with the values of k_d used to derive values of k_A from these β values.

4.2.2 Photo-Auto-Oxygenation of a Single Substrate A

When a substrate acts as its own sensitizer yielding singlet oxygen one can observe Type II photo-auto-oxygenations. Good examples are rubrene (Rub) and 9,10-dimethylanthracene (DMA) and for such cases this is indicated in the *Comments* column of the tables by S=A. Since the self-sensitizer is consumed in the photo-auto-oxygenation this usually leads to a change in I_a which must be allowed for. Alternatively, a totally absorbing solution can be used and then the rate of appearance of some product or the rate of consumption of oxygen can be measured and substituted directly into Eq. (15) to obtain values of β since under these conditions I_a is effectively constant.

4.2.3. Sensitized Photo-oxygenations of a Substrate A in the Presence of a Second Substrate A' which also Reacts with Singlet Oxygen

Consider in addition to steps 1 to 17 (Scheme 1) with M=A the further steps 22 and 23



In the presence of a second substrate [A'] it follows that

$$-\frac{d[\text{A}']} {dt} = \frac{I_a \phi_\Delta k_r^{A'} [\text{A}']} {k_d + k_\Delta [\text{A}] + k_{A'} [\text{A}']},$$

which when $k_{A'}[\text{A}'](k_d + k_\Delta[\text{A}])$ becomes

$$-\frac{d \ln[\text{A}']} {dt} = \frac{I_a \phi_\Delta k_r^{A'}} {k_d + k_\Delta [\text{A}]} = \text{Slope.}$$

In the absence of A under identical conditions, the slope of the first order plot for the disappearance of A'

$$-\frac{d \ln[\text{A}']} {dt} = \frac{I_a \phi_\Delta k_r^{A'}} {k_d} = \text{Slope}_0,$$

thus

$$\frac{\text{Slope}_0} {\text{Slope}} = 1 + \frac{k_A} {k_d} [\text{A}] = 1 + \frac{[\text{A}]} {\beta_A}. \quad (16)$$

Young and co-workers^{42,43} and others^{44,45} have used this Stern-Volmer equation to obtain β values using low concentrations of the highly reactive fluorescent substrate 1,3-diphenylisobenzofuran (DPBF) as A'. This substrate is more reactive than most so that it decays in a first order manner during continuous irradiation in photosensitized oxidation experiments even in the presence of other substrates. The low concentrations of DPBF are often monitored by following the decrease in its fluorescence. Note that when $k_q^{A'} \gg k_r^A$ (i.e., if A were a physical quencher of singlet oxygen) Eq. (16) applies even for much less reactive substrates than DPBF.

One can also use measurements of the relative rates of disappearance of two substrates in the same solution. Application of the steady-state approximation to steps 1 to 17 (Scheme 1) together with 22 and 23 gives

$$\frac{d[\text{AO}_2]/dt} {d[\text{A}'\text{O}_2]/dt} = \frac{-d[\text{A}]/dt} {-d[\text{A}']/dt} = \frac{k_r^A[\text{A}]} {k_r^{A'}[\text{A}']}, \quad (17)$$

which upon integration gives

$$\frac{\ln ([\text{A}]_0/[\text{A}])} {\ln ([\text{A}']_0/[\text{A}'])} = \frac{k_r^A} {k_r^{A'}}.$$

Equation (17) and its integrated forms have been used to determine values of $\beta_r = k_d/k_r^A$. Equation (17) has also been used to compare rates of oxidation of two substrates A and A' separately irradiated under identical conditions in the same solvent and to compare the rates for the same substrate in different solvents.⁴²

Alternatively, $r_{\text{ox}}^{A'}(\text{A})$ and $r_{\text{ox}}^0(\text{A})$, the rates of oxygenation of A in the presence and absence of A' respectively, can be evaluated and these will be related by the equation,

$$\left(\frac{r_{\text{ox}}^0(\text{A})} {r_{\text{ox}}^{A'}(\text{A})} \right)_{t=0} = 1 + \frac{[\text{A}']_0/\beta_{A'}} {1 + [\text{A}]_0/\beta_A}, \quad (18)$$

provided A' does not absorb any exciting light, quench or react with S₁ or T₁. If initial rates are measured keeping [A]₀ constant and varying [A']₀ then Eq. (18) allows values of $\beta_{A'}$ to be obtained. If $\beta_A \gg [\text{A}]_0$ the determination of $\beta_{A'}$ values is simplified.

4.2.4. Sensitized Photo-Oxygenations of a Substrate A' in the Presence of A=Q, a Physical Quencher of Singlet Oxygen

Consider Scheme 1 with M equal to both A' and Q, i.e., assuming for the moment that there is no chemical reaction with singlet oxygen by the quencher Q and that it does not absorb the exciting light. Application of the steady-state approximation then predicts

$$(r_{\text{ox}}^Q)^{-1} = (I_a \phi_\Delta k_r^{A'})^{-1} \left[1 + [\text{A}']^{-1} \left(\beta_{A'} + \frac{k_Q[\text{Q}]} {k_{A'}} \right) \right], \quad (19)$$

i.e., a linear relationship between $(r_{\text{ox}}^Q)^{-1}$ and $[\text{A}']^{-1}$ with

$$\frac{\text{slope}} {\text{intercept}} - \left(\beta_{A'} + \frac{k_Q[\text{Q}]} {k_{A'}} \right). \quad (20)$$

In the absence of quencher, slope/intercept = $\beta_{A'}$ [see also Eq. (15)]. Thus

$$\frac{(\text{slope}/\text{intercept})_Q} {(\text{slope}/\text{intercept})_0} = 1 + \frac{k_Q[\text{Q}]} {k_d} = 1 + \beta_Q^{-1}[\text{Q}]. \quad (21)$$

In the absence of quenching of excited singlet and triplet states of the sensitizer the intercepts of $(r_{\text{ox}}^Q)^{-1}$ vs $[\text{A}']^{-1}$ plots are identical and it follows from Eqs. (19) and/or (21) that

$$\frac{(\text{slope})_Q} {(\text{slope})_0} - 1 + \beta_Q^{-1}[\text{Q}]. \quad (22)$$

In addition, from Eq. (19) the ratio of the rates of oxygenation in the absence and presence of Q is given by

$$\left(\frac{r_{\text{ox}}^0(A')}{r_{\text{ox}}^0(A')} \right)_{t=0} = 1 + \frac{k_Q[Q]}{k_d + k_{A'}[A']_0} = 1 + \frac{[Q]/\beta_Q}{1 + [A']_0/\beta_{A'}} \quad (23)$$

cf. Eq. (18). Alternatively if one assumes that [A] does not change appreciably (i.e., for low fractional conversions) Eq. (19) can be integrated to give the concentration of $A'\text{O}_2$ produced after time t , which equals the change in concentration of A' , $\Delta[A']$, and


$$\begin{aligned} \frac{[A']_0 - [A']_0}{[\text{A}'\text{O}_2]} &= \frac{[A']_0 - A^0}{\Delta[A']} = \frac{A^0}{\Delta A} \\ &= (I_a t \phi_\Delta f_r^A)^{-1} \left([A']_0 + \frac{k_d}{k_{A'}} + \frac{k_Q[Q]}{k_{A'}} \right), \end{aligned} \quad (24)$$

where A^0 and ΔA represent the initial absorbance and the change in the absorbance by the reactant A' . Plots of $A^0/\Delta A$ vs $[Q]$ have been shown to be linear and from Eq. (24) it follows that for such plots

$$\frac{\text{slope}}{\text{intercept}} = (k_d + k_{A'}[A']_0). \quad (25)$$

4.2.5. Sensitized Photo-Oxygenation of a Substrate A' in the Presence of $A=Q$, a Quencher of Both Singlet Oxygen and the Sensitizer Triplet

The mechanism is that given by Scheme 1 with $M=A$ and Q , together with the reaction 24.



Application of the steady-state approximation gives

$$\begin{aligned} (r_{\text{ox}}^Q)^{-1} &= (I_a \phi_\Delta f_r^A)^{-1} \left(\frac{k_{TQ}^Q[Q]}{k_{TQ}^Q[\text{O}_2]} + 1 \right) \\ &\times \left[1 + [A']^{-1} \left(\beta_{A'} + \frac{k_Q[Q]}{k_{A'}} \right) \right]. \end{aligned} \quad (26)$$

Note that for a plot of $(r_{\text{ox}}^Q)^{-1}$ vs $[A']^{-1}$ the slope/intercept is still given by Eq. (20) since the extra term in Eq. (26) affects both the intercept and slope equally. The presence or absence of this change in intercept for $(r_{\text{ox}}^Q)^{-1}$ vs $[A']^{-1}$ plots can be used as a diagnostic test for the presence of steps such as 24 which reduce the yield of singlet oxygen produced. Another test is the occurrence of a dependence on the concentration of oxygen in solution since reaction 24 competes with oxygen quenching and thus the yield of singlet oxygen becomes dependent on the pressure of oxygen above the solution. (e.g., see Refs. 45, 46).

4.2.6. Separation of k_r and k_q

Apart from Eq. (17) application of all of the equations given so far only allows values of $k_A = k_r^A + k_q^A$ or $k_Q = k_r^Q + k_q^Q$ to be obtained. Methods which have been used to separate k_r and k_q values usually involve the direct measurement

of the quantum yields of oxygenation, ϕ_{ox} , for example at high concentrations of A such that $k_A[A] \gg k_d$, in which case it follows from Eq. (14) that

$$(\phi_{\text{ox}})_{[A] \rightarrow \infty} = \phi_\Delta f_r^A. \quad (27)$$

Alternatively at low values of [A] when $k_d \gg k_A[A]$ Eq. (14) gives

$$\phi_{\text{ox}} = \frac{\phi_\Delta k_r^A[A]}{k_d}, \quad (28)$$

and measurement of ϕ_{ox} together with a knowledge of ϕ_Δ , and k_d or β_A allows values of k_r^A or f_r^A to be determined.

The limiting yield of oxygenation of a very reactive acceptor e.g., α -terpinene or 2,5-dimethylfuran (2,5-DMF) for which f_r^A is close to unity (i.e., $k_r^A \gg k_q^A$) may be used as a reference substrate to give values of ϕ_Δ and whence k_r^A for other additives.^{47,48}

Gollnick and Gricsbck⁴⁹ have, for example, measured the rates of oxygen consumption of various substrates relative to that of 2,5-DMF at concentrations greater than $4 \times 10^{-5} \text{ L}^{-1}$. In this method $k_r^{\text{DMF}} \gg k_q^{\text{DMF}}$, $r_{\text{ox}}^{\text{DMF}} = I_a \phi_\Delta$, and

$$\frac{r_{\text{ox}}^{\text{DMF}}}{r_{\text{ox}}^A} = \frac{(k_r^A + k_q^A)[A] + k_S[S] + k_\Delta}{k_r^A[A]},$$

or

$$\frac{r_{\text{ox}}^{\text{DMF}}}{r_{\text{ox}}^A} = \left(1 + \frac{k_q^A}{k_r^A} \right) + \left(\frac{k_\Delta}{k_r^A} + \frac{k_S[S]}{k_r^A} \right) \frac{1}{[A]}, \quad (29)$$

Plots of $r_{\text{ox}}^{\text{DMF}}/r_{\text{ox}}^A$ versus $[A]^{-1}$ are linear and the value of k_q^A/k_r^A can be determined from the intercept. Variations in the slope of such plots with the sensitizer concentration allow values of k_S/k_r^A and k_Δ/k_r^A to be determined. If k_Δ is known then k_r^A , k_S , and k_q^A can be determined.⁴⁹

5. Kinetic Analysis for Oxygenation by Singlet Oxygen Generated by Chemical Reaction, by Microwave Discharge, or by Direct Laser Excitation

5.1. When Singlet Oxygen is Produced Chemically

Oxygenation reactions arising from singlet oxygen produced chemically have been studied in the presence and absence of singlet oxygen quenchers. In the absence of quencher the decrease in A is given by the relative probabilities of steps 14, 15, and 16, and 17 with $M=A$

$$-\frac{d[A]}{dt} = \frac{d[{}^1\text{O}_2^*]}{dt} \left[\frac{k_r^A[A]}{k_d + k_A[A]} \right], \quad (30)$$

which integrates to give

$$\beta_A \ln \left[\frac{[A]_0}{[A]_\infty} \right] + [A]_0 - [A]_\infty = f_r^A [{}^1\text{O}_2^*]_\infty,$$

where $[{}^1\text{O}_2^*]_\infty$ is the total amount of singlet oxygen generated chemically. For small fractional conversions

$$\ln\left(\frac{[A]_0}{[A]_\infty}\right) = \left(\frac{[A]_0 - [A]_\infty}{[A]_\infty}\right)$$

and substitution gives

$$\frac{1}{\Delta[A]} = (f_r^A [{}^1O_2^*]_\infty)^{-1} \left[\frac{\beta_A}{[A]_\infty} + 1 \right], \quad (31)$$

where $\Delta[A] = [A]_0 - [A]_\infty$. Thus a plot of $\Delta[A]^{-1}$ vs $[A]^{-1}$ for small conversions has slope/intercept = β_A . In the presence of a quencher of chemically produced singlet oxygen and a substrate A'

$$\left(\beta_{A'} + \frac{k_Q[Q]}{k_{A'}} \right) \left(\ln\left(\frac{[A']_0}{[A']_\infty}\right) + \Delta[A'] \right) = f_r^{A'} [{}^1O_2^*]_\infty, \quad (32)$$

so that if k_d and $k_{A'}$ are known, Eq. (32) allows k_Q to be evaluated from values of $[A']_0$ and $[A']_\infty$ in the presence of $[Q]$. With the decomposition of the ozonide $(C_6H_5O)_3P...O_3$ in the presence of pyridine, Mendenhall⁵⁰ has shown that there is quantitative generation of ${}^1O_2^*$ so that $[{}^1O_2^*]_\infty$ can be replaced by the initial concentration of $(C_6H_5O)_3PO_3$, i.e., $[{}^1O_2^*]_\infty = [(C_6H_5O)_3PO_3]_0$ and $f_r^{A'}$ is often assumed to be unity when A' is rubrene.⁵⁰

5.2. When Singlet Oxygen is Produced by Microwave Discharge in the Gas Phase

Equations (31) to (32) also apply to the disappearance of a substrate as a result of reaction of singlet oxygen generated in microwave experiments (as well as for singlet oxygen produced via photosensitization). Taking Eq. (32) for relative measurements with the same initial concentration of substrate $[A'_0]$ in the presence and absence of a quencher gives

$$k_Q = \frac{k_{A'}([A']_\infty^Q - [A']_\infty^0) + k_d \ln\left(\frac{[A']_\infty^Q}{[A']_\infty^0}\right)}{[Q] \ln\left(\frac{[A']_0}{[A']_\infty^0}\right)}, \quad (33)$$

where $[A']_\infty^Q$ and $[A']_\infty^0$ represent the final values after exposure to microwave generated singlet oxygen in the presence and absence of quencher, respectively. Equation (33) has also been used following chemical and photochemical production of ${}^1O_2^*$ in the absence and presence of a quencher Q (or acceptor A).

5.3. When Singlet Oxygen is Produced by Direct CW Laser Excitation of Oxygen


Evans and Tucker¹⁴ and Matheson and co-workers^{11-13,35,51} have used laser excitation of highly concentrated oxygen solutions in high pressure cells, with gaseous oxygen up to 130 atm, to give directly the ${}^1O_2^*(^1\Delta_g)$ state by absorption at 1065 nm from a CW Nd YAG laser and ${}^1O_2^*(^1\Delta_g)$ directly from the "double simultaneous transition" using He-Ne laser excitation at 632.8 nm. Evans and Tucker¹⁴ find the quantum yields of photo-oxygenation of 9,10-dimethylanthracene and tetraphenylcyclopentadiene are twice as high for absorption of a photon in the "double simultaneous transition" as for the direct excitation of a single

${}^1\Delta_g$ state. Modification of Eq. (15) to account for the direct excitation by replacing ϕ_Δ by $n=1$ or 2 for excitation at 1065 nm or 633 nm, respectively, gives

$$\frac{1}{\phi_{ox}} = \frac{1}{n} \left[\frac{k_A}{k_r^A} + \frac{k_d}{k_r^A [A]} \right] \quad (34)$$

and plots of ϕ_{ox}^{-1} vs $[A]^{-1}$ can be used to give f_r^A and β_A values.

Matheson *et al.*^{11-13,35,51,52} have obtained experimental pseudo first order rate constants for the disappearance of various substrates while directly producing ${}^1O_2^*(^1\Delta_g)$ with a CW Nd YAG laser. Since the absorption involves the process



Eq. (14) has been modified by them to give

$$\frac{d[A]}{dt} = \frac{\sigma E [O_2]^2 k_r^A [A]}{k_d + k_A [A] + k_{O_2} [O_2]}, \quad (35)$$

where E is the laser intensity and σ the absorption cross section for double photon absorption. At high oxygen concentration where $k_{O_2} [O_2] \gg (k_d + k_A [A])$ the observed first order rate constant k_1 is given by

$$k_1 = \frac{\sigma E [O_2] k_r^A}{k_{O_2}}. \quad (36)$$

Thus from a knowledge of σ , E , and $[O_2]$ values of k_r^A relative to k_{O_2} can be obtained. In the presence of a singlet oxygen quencher at relatively high concentrations and a substrate A'

$$k_1 = \frac{\sigma E [O_2]^2 k_r^A}{k_{O_2} [O_2] + k_Q [Q]} \quad (37)$$

and a plot of $([O_2]/k_1)$ vs $([Q]/[O_2])$ will have a slope/intercept = k_Q/k_{O_2} .

Values obtained using this method have often been substantially lower than those given by other workers. According to Matheson^{51,52} this was due, partially at least, to an artifact with the over depletion of the acceptor concentration in the laser beam cross section so that the observed chemical reaction rate may be affected by diffusion of unreactive acceptor into the depleted region. Thus values for k_r^A determined in this way published prior to 1977 may be low by up to an order of magnitude and have been omitted from this compilation.

6. First Order Rate Constants for the Decay of Singlet Oxygen in Various Solvents—Comments on Table 1

Table 1 contains decay rates of singlet oxygen in 145 solvents and solvent mixtures. Many of the data have been reported as lifetimes (τ), the reciprocal of the decay rate, which have been tabulated. The decay rate (k_d) column contains reported values as well as values calculated from lifetimes. The values in both columns are rounded to two significant figures. When the data were reported with one significant

figure, or when the quoted error exceeded 20%, only one significant figure is given. The *Method*, *Comment*, and *Reference* columns (see Secs. 7.2.4–7.2.6) provide details of the measurements. Only data obtained by time-resolved methods have been included except for solvents where no such measurements have been reported but where estimates have been made from steady-state measurements.

The data for each solvent are listed in reverse chronological order, so that the most recent entries appear first. It is apparent from Table 1 that k_d values tend to decrease with year of publication especially for solvents with values $<10^2 \text{ s}^{-1}$. For such solvents the presence of impurities including water and high concentrations of sensitizers can give inaccurate values. It is likely that the lower the k_d values the higher the accuracy. Usually there is good agreement between different authors for k_d values $>10^3 \text{ s}^{-1}$.

The lifetime of singlet oxygen shows an enormous solvent dependence, including a large isotope effect, and variations over almost five orders of magnitude from a few microseconds in water to hundreds of milliseconds in carbon tetrachloride. The reason for this is that the nonradiative deactivation is essentially a bimolecular energy transfer reaction between $\text{O}_2^*(^1\Delta_g)$ and the solvent molecules which thereby become vibrationally excited. The first interpretation by Merkel and Kearns³² that the electronic vibrational energy transfer was a dipole resonance transfer has been replaced by an exchange transfer mechanism following important contributions by Schuster *et al.*,^{27,53} Rodgers,⁵⁴ and Schmidt *et al.*^{55–58} The bimolecular rate constant for solvent deactivation k_{sol} can be obtained from

$$k_D = k_0 + k_{\text{sol}}[\text{sol}],$$

where k_0 is an intrinsic molecular rate constant probably equal to $k_{\Delta P}$, the rate constant for radiative decay, and $[\text{sol}]$ represents the concentration of the solvent in mol L⁻¹ calculated from the density and the molecular weight of the solvent; k_{sol} can be shown to depend on the availability of high energy X–Y, especially C–H and O–H, vibrations in the molecule. Thus several workers have shown that

$$k_{\text{sol}} = \sum_{XY} N_{XY} k_{XY},$$

where N_{XY} is the number of times a particular X–Y bond occurs in the molecule. In this way values for k_{XY} have been calculated, which can be used to calculate k_{sol} values for solvents for which values of k_d have not been experimentally determined. Since $k_{\Delta P}$ varies from 0.1 to 3 s⁻¹, k_0 is often negligible compared with $k_{\text{sol}}[\text{sol}]$, but for solvents containing no high energy vibrations values of $k_{\Delta P}$ have been used by Schmidt to obtain k_{XY} values from k_D , for example in CCl_4 and CS_2 which show the lowest values for k_D .

We made several attempts to calculate k_{XY} values from the k_d values in Table 1 which includes solvents with C–H, C–D, C=O, O–H, O–D, C≡N, C–F, C–Cl, C–Br, C–I, C=S, and S=O bonds. We used the formula

$$k_d / [\text{sol}] = \sum_{XY} N_{XY} k_{XY}$$

as the basis for a least squares fitting of the measured values of k_d to obtain a set of values for k_{XY} . The procedure resulted in negative values for some parameters and statistical uncertainties for many that were larger than the value of the parameter. Restricting the data set, for example by eliminating solvents containing halogens, deuterium and sulfur and measured values which deviated significantly from the average, did not eliminate this problem. Separation of C–H bonds into classes (CH_3 , CH_2 , CH, cycloalkyl, aromatic) and perhaps other X–Y bonds as well, and the selection of standard k_d values appear to be crucial to establishing a set of k_{XY} values.

7. Second-order Rate Constants for Reactive and Physical Quenching of Singlet Oxygen by Various Substrates—Comments on Tables 2–17

7.1. Scope of the Compilation

Second-order rate constants for quenching of singlet oxygen are given in 16 tables containing 4683 separate entries for 1915 substrates. The data have been extracted from the literature published through 1993. The papers have been selected from the bibliographic database of the Radiation Chemistry Data Center and from citations in those papers. Data reported by the same author(s) which have been clearly superseded have been omitted. For some well-studied systems many independent measurements have been made; all reported values have been included allowing the user to make comparisons based on method, similarity of values, etc. Data from dissertations and conference papers have been included only if no other measurements for the substrate in that solvent have been reported. The authors would welcome having our attention drawn to any data which have been missed.

7.2. Arrangement of Tables

Each table includes related groups of compounds, e.g., olefins, aromatic hydrocarbons, phenols, etc. In some cases choices were made based on functionality, for example tyrosine and its peptides are in Table 4 with phenols instead of Table 10 (amino acids and peptides). Benzopyranols (such as tocopherol) are in Table 4 (phenols) instead of Table 5 (furans and pyrans). Cysteine and methionine and their derivatives are in Table 11 (sulfur compounds), not Table 10 (amino acids).

Within each table the various substrates are arranged alphabetically by name, given an entry number which includes the table number (2.1, 2.2, etc.) and within each entry the arrangement is by solvent. Inverted names have been used as headings whenever possible so that substituted anthracenes, porphines, phenols, etc. are grouped. Systematic names have been used in most cases, otherwise, common names have been given. An index of chemical names and synonyms and a molecular formula index follow the tables (Secs. 13 and 14) and refer to the entry numbers in Tables 2–17.

7.2.1. Solvent

Data in each entry are grouped by solvent and listed in reverse chronological order of the year of publication. The *Solvent* column gives the solvent or mixture of solvents with the proportions (volume:volume). Micellar systems are labelled (mic), and the surfactant present is given in the *Comment*. When water or D₂O is the solvent the pH or pD is given, if reported. If the reaction was studied over a range of pH values and the *k* derived for a particular ionic form of the substrate no pH is given in the *Solvent* column and the pH range is given in the *Comment* column.

7.2.2. Rate Constants (*k*) and Beta (*k_d/k*) Values

The rate constants in the second column are for total quenching by chemical reaction and/or physical quenching, $k = k_r + k_q$, except where separate reactive quenching (*k_r*) and physical quenching (*k_q*) rate constants have been measured. It may be known from other studies that a particular substrate interacts with singlet oxygen totally by chemical reaction ($f_r^A = 1$) or totally by physical quenching; an example of the latter is azide ion. We have, however, only provided the labels *k_r* and *k_q* when a method was used which made the distinction. Values of β which have been reported are listed in the third column. When the ratio is *k_d/k_r* the label β_r is used.

Values of *k* and β have been rounded to two significant figures unless they were given to only one significant figure by the authors. The authors' error limits have been omitted; when the reported error is greater than 20% the values have been given to one significant figure. Upper or lower limits have been given to only one significant figure.

7.2.3. Temperature

Temperatures in Kelvin are listed whenever they have been reported. When the information has not been reported it is assumed that the temperature is ambient. If the study has been carried out over a range of temperatures the range is given in the *Comment* column.

7.2.4. Method

The *Method* column describes the method by symbols (see Sec. 9) representing the excitation technique, the measurement technique, and the equation numbers given in the previous sections which have been used for analyzing the data. For example, PL/Ld-2 represents the pulsed-laser luminescence-decay technique and refers to Eq. (2). If two measurement techniques or equations were used the symbols are combined. For example, CP/Oc,Ac-17 indicates that excitation was by continuous photolysis, oxygen consumption (Oc) and substrate consumption (Ac) were measured, and the rate constant was derived using Eq. (17).

7.2.5. Comment

Some details of the determination are given in the *Comment* column. Abbreviations and symbols are listed in Sec. 9 for chemical species which have been used as photosensitizer

(S) or reference substrate (A'). The source of the ${}^1\text{O}_2^*$ is given when it was generated in a chemical reaction. Values for *k_d* which have been used by the author to calculate *k* from β (*k_d/k*) are given. The reader is referred to Table 1 to compare these values with the various measured values of *k_d* for a particular solvent. In some cases values of $\beta_{A'}$ for the reference substrate have been used to calculate β (or to calculate *k* also using a value of *k_d*) and those values are given in the *Comment*. Data are available for some substrates only as ratios with a reference substrate (e.g., $k_r^A/k_r^{A'}$). When a value for the rate constant for the reference substrate has been used by the author(s) to calculate *k*, the reference value which was used is given in the *Comment* column. When rate constants have not been calculated by the authors the ratios are given in the *Comment* column and no value appears in the *k* column; the ratios have been rounded to two significant figures. Since ratios of rate constants refer equally to both substrates, the reference substrates (A') have been included in the Chemical Name Index (Sec. 14) with the entry number(s) in which ratios of rate constants with another substrate are present or can be derived. For example, Acetone azine appears in Table 15 (entry 15.3) but rate constants for another substrate relative to *k_r* for acetone azine are found in entry 2.40. There is no entry in the tables for 9-Acetyl-2,3,4,9-tetrahydrocarbazole but relative data are found in entries 6.2, 6.3, 6.10, 6.14, 6.18, and 6.19.


Other data are included when they have been reported: activation energy (*E_a*), activation volume (ΔV^\ddagger), enthalpy of activation (ΔH^\ddagger), entropy of activation (ΔS^\ddagger), along with the temperature or pressure ranges of the study. For some entries where product appearance (Pa) was measured the products have been given.

7.2.6. References


The rate constants have been compiled from more than 700 references which are identified in the *Reference* column by codes from the Radiation Chemistry Data Center Bibliographic Database. The first two characters of the reference code represents the year of publication. The complete references to Tables 1–17 are listed in Sec. 12 which follows the tables.

8. Mechanism of Quenching

Two recent reviews^{1,2} discuss bimolecular processes of singlet molecular oxygen in detail. Most results can be explained in terms of exciplex formation followed by competing reactions involving energy transfer (ET), charge transfer (CT) or chemical reaction as shown.




It is well established that quenching of singlet oxygen arises from chemical reaction and/or physical quenching or a combination of both. Physical quenching due to the spin allowed electronic energy transfer process



has been demonstrated to occur by direct observation of the triplet state of the quencher, e.g., in the case of various carotenes.³⁷ This type of quenching is diffusion controlled when the triplet energy of the quencher lies below 94 kJ mol⁻¹ and has been proposed as the mechanism of quenching to account for the high efficiency of quenching of singlet oxygen by certain dyes and several coordination complexes.^{59,60} Quenching of singlet oxygen by electronic energy transfer is the reverse of reaction 11 which is likely to be a limiting factor for finding sensitizers absorbing in the red for use in photodynamic therapy.⁶¹ Physical quenching of singlet oxygen as discussed in Sec. 6, also arises as a result of electronic to vibrational energy transfer, where the pseudo-first order decay constant in a particular solvent depends on the availability of high energy vibrations for example O—H and C—H vibrations in the solvent molecule. This kind of process is obviously possible for all substrates but is often much less efficient than physical quenching resulting from charge transfer interactions.

Physical quenching resulting from favorable charge transfer interactions was first demonstrated with various amines.^{62,63} This type of quenching may be represented as follows:



If the charge transfer complexes shown above are able to dissociate to give separated ions this constitutes chemical quenching, although the production of O₂[−] (superoxide) has only been demonstrated for highly polar solvents such as water and with compounds with exceptionally low ionization potentials such as *N,N,N',N'*-tetramethyl-*p*-phenylenediamine.⁶⁴ The temperature dependence of the rate constants observed⁶⁵ for singlet oxygen quenching by strychnine and DABCO in toluene has established complex formation during the quenching process. The efficiency of quenching by amines increases when the ionization potential of the amine decreases, which is also the case for quenching by phenols and sulfides which give similar Hammett plots.^{66,67} Quenching by amines, phenols and, sulfides shows varying amounts of chemical reaction.

Chemical reactions of singlet oxygen have been discussed in detail in a number of reviews.^{68–72} Alkenes react to form hydroperoxides, 1,2-dioxetanes, and endoperoxides, which are often produced in the case of other substrates, for example in the case of aromatic hydrocarbons, furans, etc. Recent research^{73,74} has revealed that intermediates exist along the reaction coordinate and there has been much speculation concerning the nature of these intermediates. Since it is known that tryptophan is one of the main targets when O₂^{*}(¹Δ_g) reacts with protein^{75,76} there has been considerable

interest in studying rate constants for reaction of singlet oxygen with tryptophan in a variety of solvents. As with many other amino acids⁷⁷ reactive quenching only accounts for part of the total quenching of singlet oxygen observed. It has been shown⁷⁸ that singlet oxygen causes single-strand breaks in DNA and that guanine is the nucleoside base from which oxidative products are derived. As with all substrates it is important to establish that particular products arise as a result of reactions of singlet oxygen. It is hoped that this compilation will help with establishing the importance of singlet oxygen reactions in the photodegradation of dyes, pigments, polymers, etc, as well as in harmful and/or beneficial photo-oxidations in biological systems.

9. List of Abbreviations and Symbols

Excitation Methods

CL	continuous laser photolysis
CP	continuous photolysis
CR	chemical reaction
FP	flash photolysis
MD	microwave discharge
MP	modulated photolysis
PL	pulsed laser photolysis
PR	pulsed radiolysis

Measurement Methods

Ac	substrate consumption under steady-state conditions
A'c	reference substrate consumption under steady-state conditions
Ad	time resolved substrate disappearance
A'd	time resolved reference substrate disappearance
βCd	triplet beta-carotene decay as a probe for singlet oxygen decay
Ld	luminescence decay
LI	luminescence intensity under steady-state conditions
Oc	oxygen consumption under steady-state conditions
Pa	product appearance under steady-state conditions
P'a	product appearance (from reference substrate) under steady-state conditions
Pb	time-resolved product buildup
P'b	time-resolved product buildup (from reference substrate)
Tb	time-resolved thermal buildup

Other Symbols

A	primary substrate
A'	reference substrate
A''	second reference substrate
β	beta value including both β _q and β _r
β _q	beta value for physical quenching of ¹ O ₂ [*]
β _r	beta value for chemical reaction
E _a	activation energy
ΔH [‡]	enthalpy of activation

ΔS^\ddagger	entropy of activation	CQ	Camphoroquinone
ΔV^\ddagger	activation volume	CTAB	Hexadecyltrimethylammonium bromide
f_r^A	fraction of ${}^1O_2^*$ quenched by A which reacts	CTAC	Hexadecyltrimethylammonium chloride
k	rate constant including both k_q and k_r	CuTCPc	Copper(II) tetracarboxyphthalocyanine
k_A	rate constant for primary substrate, including both k_q and k_r	DABCO	1,4-Diazabicyclo[2.2.2]octane
$k_{A'}$	rate constant for reference substrate, including both k_q and k_r	DAP	Dodecylammonium propionate
k_d	first-order decay rate of singlet oxygen in solvent	DBrA	9,10-Dibromoanthracene
k_Δ	first-order decay rate of singlet oxygen for $[S] \rightarrow 0$	DCA	9,10-Dicyanoanthracene
k_Q	rate constant for reference substrate which is a physical quencher	DCIA	9,10-Dichloroanthracene
k_q	rate constant for physical quenching of ${}^1O_2^*$	DDAB	Didodecyldimethylammonium bromide
k_r	rate constant for chemical reaction of ${}^1O_2^*$	DDM	Diazodiphenylmethane [1,1'-(Diazomethylene) bisbenzene]
k_s	rate constant for reaction of ${}^1O_2^*$ with sensitizer	DLPC	Dilauroyl phosphatidylcholine
mic	micelles	DMA	9,10-Dimethylanthracene
P	product of primary substrate	DMAA	<i>N,N</i> -Dimethylacetamide
P'	product of reference substrate	DMAO ₂	9,10-Dimethylanthracene endoperoxide
ϕ_Δ	quantum yield of singlet oxygen formation	DMBA	9,10-Dimethyl-1,2-benzanthracene
ϕ_{isc}	quantum yield of intersystem crossing	DMDPA	1,4-Dimethoxy-9,10-diphenylanthracene
ϕ_{ox}	quantum yield of oxidation of substrate	2,5-DMF	2,5-Dimethylfuran
Q	reference substrate which is a physical quencher	DMF	<i>N,N</i> -Dimethylformamide
r_{ox}	rate of oxygenation	DMHD	2,5-Dimethyl-2,4-hexadiene
S	sensitizer	DMNO ₂	1,4-Dimethylnaphthalene 1,4-endoperoxide
ves	vesicles	DNT	1,8-Dinaphthalene thiophene (Diacenaphtho[1,2- <i>b</i> :1',2'- <i>d</i>]thiophene)

Sensitizers, Reference Substrates, and Medium Components

Ac	Acridine	DODAC	Dioctadecyltrimethylammonium chloride
2-ACN	2-Acetonaphthone	DPA	9,10-Diphenylanthracene
ADC	Anthra[1,9- <i>bc</i> :4,10- <i>b'c'</i>]dichromene	DPB	1,4-Diphenyl-1,3-butadiene
ADPA	9,10-Anthracenedipropionate ion	DPBF	1,3-Diphenylisobenzofuran
AES	Anthracene-9,10-bis(ethanesulfate ion)	DPF	2,5-Diphenylfuran
AlCl(tspc)	Chloroaluminum(III) sulfophthalocyanine	DPPC	Dipalmitoyl phosphatidylcholine
An	Anthracene	DTAC	Dodecyltrimethylammonium chloride
AnS	Anthracenesulfonate	Eos	Eosin (Tetrabromofluorescein)
AOT	Di(2-ethylhexyl) sulfosuccinate	Ery	Erythrosin (Tetraiodofluorescein)
azine	2-Pivaloyl-4-phenyl-6-diethylamino-8-methyl- quinoxal-3-one	ES	Ergosterol
Azo dye 10	<i>N</i> -[3-(Aminosulfonyl)phenyl]-3-[(3-cyano- 5-hydroxy-1-phenylpyrazol-4-yl)azo]-4- methoxybenzenesulfonamide	EtOH	Ethanol
BA	1,2-Benzanthracene	FFA	Furfuryl alcohol
BChl	Bacteriochlorophyll	Fl	Fluorene
BDX	Benzo[1,2,3- <i>k</i> l:4,5,6- <i>k'l</i> ']dixanthene	Fl ²⁻	Fluorescein
BHMF	2,5-Bis(hydroxymethyl)furan	FIN ₂	9-Diazofluorene
BHT	2,6-Di- <i>tert</i> -butyl-4-methylphenol	FMN	Flavin mononucleotide (Riboflavin-5'-phosphate)
Biph	Biphenyl	GaCl(tspc)	Chlorogallium(III) sulfophthalocyanine
BP	Benzophenone	GV	Gilvocarcin V
BR	Bilirubin	HA	Hypocrellin A
Brij 35	Polyoxyethylene(23) dodecyl ether	HCD	Heterocoerdianthrone
BuOH	Butanol	3-HF	3-Hydroxyflavone
BXP	Benoxaprofen	His	Histidine
Car	all- <i>trans</i> -Carotene	HP	Hematoporphyrin
CHD	Cyclohexadiene	HPD	Hematoporphyrin derivative
Chl	Chlorophyll	HYP	Hypericin
Chr	Chrysene	Im	Imidazole
Cor	Coronene	InH	Indole
		MB	Methylene Blue
		2M2B	2-Methyl-2-butene
		MC 540	Merocyanine 540
		MCH	Methylcyclohexane
		MDH	Mesodiphenylhelianthrene
		MeOH	Methanol

Met	Methionine	RF	Riboflavin
MNPO ₂	4-Methyl-1-naphthalenepropionate endoperoxide	RF(OAc) ₄	Riboflavin tetraacetate
8-MOP	8-Methoxypsoralen	Rub	Rubrene (5,6,11,12-Tetraphenylnaphthacene)
MP	Mesoporphyrin	Rubi	Rubicene (Benz[<i>a</i>]indenol[1,2,3- <i>hi</i>]aceanthrylene)
2M2P	2-Methyl-2-pentene	SDS	Sodium dodecylsulfate
MPDME	Mesoporphyrin dimethyl ester	SiNC	Bis(<i>tri-n</i> -hexylsiloxy)-(2,3-naphthalocyaninato) silicon
NAZ	Nitrazepam	TAN	2,2,6,6-Tetramethylpiperidone-1-oxyl
NDPO ₂	1,4-Naphthalenedipropionate endoperoxide	TEMP	2,2,6,6-Tetramethylpiperidine
NMTA	<i>N</i> -Methylthioacridone	TEMPO	2,2,6,6-Tetramethylpiperidin-1-oxyl
Np	Naphthalene	TEMPOH	2,2,6,6-Tetramethylpiperidin-1-ol
NT2	Octaethylporphyrin ethyl ester	TEMP-4-OH	2,2,6,6-Tetramethylpiperidin-4-ol
<i>t</i> -octane	2,2,4-Trimethylpentane	Tetr	Tetracene (Naphthacene)
OMAPB	2-Methylacetophenone biradical	Th	Thionine
PBA	1-Pyrenebutyrate ion	THF	Tetrahydrofuran
PBN	Phenyl- <i>N</i> - <i>tert</i> -butylnitronite	TMDT	2,6,10-Trimethyl-2,6,10-dodecatriene
PEO	Polyethylene oxide	TME	2,3-Dimethyl-2-butene (Tetramethylethylene)
Per	Perylene	T(m-HOP)P	Tetra(3-hydroxyphenyl)porphyrin
Ph	Pheophytin	TMS	α,β,β -Trimethylstyrene
Phen	Phenanthrene	TPBC	<i>trans</i> -Tetraphenylbacteriochlorin
PHO	Phenalen-1-one	TPP	Tetraphenylporphyrin
Poly-RB	Polymer supported Rose Bengal	H ₂ TPPS ⁴⁻	Tetra(4-sulfonatophenyl)porphyrin
PP	Protoporphyrin	Triton X-100	Polyoxyethylene(10)4-(1,1,3,3-tetramethylbutyl)phenyl ether
PPDME	Protoporphyrin dimethyl ester	TTMPP	Tetra(3,4,5-trimethoxyphenyl)porphyrin
PrOH	Propanol	TTT	Terthiophene
PrPor	2,7,12,17-Tetrapropylporphycene	UP	Uroporphyrin I
Pso	Psoralen	VD ₃	Vitamin D ₃
Pt(DHBA)(DPA)	Platinum(II) (3,4-dihydroxybenzoate) (2,2'-dipyridylamine)	Zn(pc)	Zinc(II) phthalocyanine
Pt(phen)(BCAT)	Platinum(II) (1,10-phenanthroline) (<i>tert</i> -butylcatechol)	Zn(pc)(py) ₂	Zinc(II) phthalocyaninebis(pyridine)
Pt(phen)(CAT)	Platinum(II) (1,10-phenanthroline) (catechol)	Zn(tspc)	Zinc(II) sulfophthalocyanine
Pt(phen)(DMT)	Platinum(II) (1,10-phenanthroline) (3,4-dimercaptotoluene)	ZnTPP	Zinc(II) tetraphenylporphyrin
PTSA	Pyrenetetrasulfonate ion		
Pur	Purpurin		
py	Pyridine		
Py	Pyrene		
Pz	Phenazine		
RB	Rose Bengal (Tetrachlorotetraiodofluorescein)		
RBCE	Rose Bengal complexed with dicyclohexyl-18-crown-6		
Ret	<i>all-trans</i> -Retinal		

10. ACKNOWLEDGMENTS

The Radiation Laboratory at the University of Notre Dame is operated under Contract DE-AC02-76ER0038 with the Department of Energy. The Radiation Chemistry Data Center is supported jointly by the Office of Basic Energy Sciences of the Department of Energy and the National Institute of Standards and Technology, Standard Reference Data. This is Radiation Laboratory Document No. NDRL-3772.

11. References to Text

- ¹Lissi, E.A.; Encinas, M.V.; Lemp, E.; Rubio, M.A., *Chem. Rev.* **93**: 699-723 (1993) [93Z073]
- ²Gorman, A.A., *Adv. Photochem.* **17**: 217-74 (1992) [92Z144]
- ³Dougherty, T.J., *Adv. Photochem.* **17**: 275-311 (1992) [92Z145]
- ⁴Foote, C.S., *Free Radicals in Biology. Volume 2.*, W.A. Pryor (ed.), Academic Press, New York, 1976, p.85-133 [763101]
- ⁵Wilkinson, F.; Brummer, J.G., *J. Phys. Chem. Ref. Data* **10**: 809-999 (1981) [81Z251]
- ⁶Wilkinson, F.; Helman, W.P.; Ross, A.B., *J. Phys. Chem. Ref. Data* **22**: 113-262 (1993) [93Z042]
- ⁷Gorman, A.A.; Rodgers, M.A.J., *CRC Handbook of Organic Photochemistry, Vol. II*, J.C. Scaiano (ed.), CRC Press, Inc., Boca Raton, FL, 1989, p.229-47 [89Z221]
- ⁸Childs, W.H.J.; Mecke, R., *Z. Phys.* **68**: 344-61 (1931) [319004]
- ⁹Wallace, L.; Hunten, D.M., *J. Geophys. Res.* **73**: 4813-34 (1968) [68M108]
- ¹⁰Bodesheim, M.; Schuetz, M.; Schmidt, R., *Chem. Phys. Lett.* **221**: 7-14 (1994) [94E113]
- ¹¹Matheson, I.B.C.; Lee, J.; Yamanashi, B.S.; Wolbarsht, M.L., *J. Am. Chem. Soc.* **96**: 3343-8 (1974) [747102]
- ¹²Matheson, I.B.C.; Lee, J., *J. Am. Chem. Soc.* **94**: 3310-3 (1972) [72F513]
- ¹³Matheson, I.B.C.; Lee, J., *Photochem. Photobiol.* **29**: 879-81 (1979) [79A112]
- ¹⁴Evans, D.F.; Tucker, J.N., *J. Chem. Soc., Faraday Trans. 2* **72**: 1661-6 (1976) [767417]
- ¹⁵Singh, A.; Koroll, G.W.; Antonsen, S.A., *J. Photochem.* **25**: 99-104 (1984) [84F190]
- ¹⁶Evans, D.F., *J. Chem. Soc.* : 1351-7 (1957) [57B004]
- ¹⁷Tsubomura, H.; Mulliken, R.S., *J. Am. Chem. Soc.* **82**: 5966 (1960) [60B009]
- ¹⁸Birks, J.B., *Photophysics of Aromatic Molecules*, Wiley-Interscience, New York, 1970, 704p. [706229]
- ¹⁹Scurlock, R.D.; Ogilby, P.R., *J. Phys. Chem.* **93**: 5493-500 (1989) [89E309]
- ²⁰Kristiansen, M.; Scurlock, R.D.; Iu, K.-K.; Ogilby, P.R., *J. Phys. Chem.* **95**: 5190-7 (1991) [91E297]
- ²¹Logunov, S.L.; Rodgers, M.A.J., *J. Phys. Chem.* **97**: 5643-8 (1993) [93E220]
- ²²McGarvey, D.J.; Wilkinson, F.; Worrall, D.R.; Hobley, J.; Shaikh, W., *Chem. Phys. Lett.* **202**: 528-34 (1993) [93A026]
- ²³Gorman, A.A.; Lovering, G.; Rodgers, M.A.J., *J. Am. Chem. Soc.* **100**: 4527-32 (1978) [78E263]
- ²⁴Carlsson, D.J.; Suprunchuk, T.; Wiles, D.M., *Can. J. Chem.* **52**: 3728-37 (1974) [747341]
- ²⁵Foote, C.S., *Acc. Chem. Res.* **1**: 104-10 (1968) [68F282]
- ²⁶Byteva, I.M.; Gurinovich, G.P., *J. Lumin.* **21**: 17-20 (1979) [79E669]
- ²⁷Hurst, J.R.; McDonald, J.D.; Schuster, G.B., *J. Am. Chem. Soc.* **104**: 2065-7 (1982) [82E104]
- ²⁸Ogilby, P.R.; Foote, C.S., *J. Am. Chem. Soc.* **104**: 2069-70 (1982) [82E106]
- ²⁹Gorman, A.A.; Hamblett, I.; Land, E.J., *J. Am. Chem. Soc.* **111**: 1876-7 (1989) [89E113]
- ³⁰Rossbroich, G.; Garcia, N.A.; Braslavsky, S.E., *J. Photochem.* **31**: 37-48 (1985) [85E591]
- ³¹Merkel, P.B.; Kearns, D.R., *Chem. Phys. Lett.* **12**: 120-22 (1971) [719325]
- ³²Merkel, P.B.; Kearns, D.R., *J. Am. Chem. Soc.* **94**: 7244-53 (1972) [72Z260]
- ³³Adams, D.R.; Wilkinson, F., *J. Chem. Soc., Faraday Trans. 2* **68**: 586-93 (1972) [72T126]
- ³⁴Young, R.H.; Brewer, D.; Keller, R.A., *J. Am. Chem. Soc.* **95**: 375-9 (1973) [73T014]
- ³⁵Matheson, I.B.C.; Lightner, D.A., *Photochem. Photobiol.* **29**: 933-5 (1979) [79A113]
- ³⁶Okamoto, M., *J. Phys. Chem.* **96**: 245-8 (1992) [92A014]
- ³⁷Parmilo, A.; Wilkinson, F., *Photochem. Photobiol.* **18**: 447-50 (1973) [73T438]
- ³⁸Krasnovsky, A.A., Jr., *Photochem. Photobiol.* **29**: 29-36 (1979) [79A010]
- ³⁹Jenny, T.A.; Turro, N.J., *Tetrahedron Lett.* **23**: 2923-6 (1982) [82E329]
- ⁴⁰Losev, A.P.; Nichiporovich, I.N.; Byteva, I.M.; Drozdov, N.N.; Al Joghami, I.F., *Chem. Phys. Lett.* **181**: 45-50 (1991) [91E288]
- ⁴¹Gollnick, K., *Adv. Chem. Ser.* **77**: 78-101 (1968) [68F289]
- ⁴²Young, R.H.; Wehrly, K.; Martin, R.L., *J. Am. Chem. Soc.* **93**: 5774-9 (1971) [717398]
- ⁴³Young, R.H.; Martin, R.L.; Chinh, N.; Mallon, C.; Kayser, R.H., *Can. J. Chem.* **50**: 932-8 (1972) [72F510]
- ⁴⁴Thomas, M.J.; Foote, C.S., *Photochem. Photobiol.* **27**: 683-93 (1978) [78A171]
- ⁴⁵Greenstock, C.L.; Wiebe, R.H., *Photochem. Photobiol.* **28**: 863-7 (1978) [78N003]
- ⁴⁶Foote, C.S.; Chang, Y.C.; Denny, R.W., *J. Am. Chem. Soc.* **92**: 5216-8 (1970) [707188]
- ⁴⁷Schenck, G.O.; Collnick, K.; Neumüller, O.A., *Ann. Chem.* **603**: 46-59 (1957) [57F008]
- ⁴⁸Tanielian, C.; Chaineaux, J., *J. Photochem.* **9**: 19-32 (1978) [78A344]
- ⁴⁹Gollnick, K.; Griesbeck, A., *Tetrahedron* **40**: 3235-50 (1984) [84F335]
- ⁵⁰Carlsson, D.J.; Mendenhall, G.D.; Suprunchuk, T.; Wiles, D.M., *J. Am. Chem. Soc.* **94**: 8960-2 (1972) [72Z319]
- ⁵¹Matheson, I.B.C.; Toledo, M.M., *Photochem. Photobiol.* **25**: 243-8 (1977) [77Z129]
- ⁵²Matheson, I.B.C., *Photochem. Photobiol.* **29**: 875-8 (1979) [79A111]
- ⁵³Hurst, J.R.; Schuster, G.B., *J. Am. Chem. Soc.* **105**: 5756-60 (1983) [83E398]
- ⁵⁴Rodgers, M.A.J., *J. Am. Chem. Soc.* **105**: 6201-5 (1983) [83A223]
- ⁵⁵Schmidt, R.; Brauer, H.-D., *J. Am. Chem. Soc.* **109**: 6976-81 (1987) [87E658]
- ⁵⁶Schmidt, R., *J. Am. Chem. Soc.* **111**: 6983-7 (1989) [89E388]
- ⁵⁷Schmidt, R.; Afshari, E., *Ber. Bunsenges. Phys. Chem.* **96**: 788-94 (1992) [92E256]
- ⁵⁸Schmidt, R., *Ber. Bunsenges. Phys. Chem.* **96**: 794-9 (1992) [92Y093]
- ⁵⁹Herkstroeter, W.G., *J. Am. Chem. Soc.* **97**: 4161-7 (1975) [75Z247]
- ⁶⁰Singh, R.J.; Feix, J.B.; Kalyanaraman, B., *Photochem. Photobiol.* **55**: 483-9 (1992) [92A123]
- ⁶¹Rihter, B.D.; Kenney, M.E.; Ford, W.E.; Rodgers, M.A.J., *J. Am. Chem. Soc.* **112**: 8064-70 (1990) [90E508]
- ⁶²Young, R.H.; Martin, R.L., *J. Am. Chem. Soc.* **94**: 5183-5 (1972) [72F514]
- ⁶³Monroe, B.M., *J. Phys. Chem.* **81**: 1861-4 (1977) [77Z486]
- ⁶⁴Manring, L.E.; Foote, C.S., *J. Phys. Chem.* **86**: 1257-9 (1982) [82A080]
- ⁶⁵Gorman, A.A.; Hamblett, I.; Lambert, C.; Spencer, B.; Standen, M.C., *J. Am. Chem. Soc.* **110**: 8053-9 (1988) [88A427]
- ⁶⁶Young, R.H.; Brewer, D.; Kayser, R.; Martin, R.; Feriozi, D.; Keller, R.A., *Can. J. Chem.* **52**: 2889-93 (1974) [74Z245]
- ⁶⁷Monroe, B.M., *Photochem. Photobiol.* **29**: 761-4 (1979) [79A085]
- ⁶⁸Gollnick, K.; Kuhn, H.J., *Singlet Oxygen*, H.H. Wasserman and R.W. Murray (eds.), Academic Press, New York, N.Y., 1979, Vol. 40 of Organic Chemistry, p.287-427 [79Z111]
- ⁶⁹Frimer, A.A., *Singlet Oxygen. Vol. II*, CRC Press, Inc., Boca Raton, FL, 1985, 284p. [85Z260]
- ⁷⁰Frimer, A.A., *Singlet Oxygen. Vol. III*, CRC Press, Inc., Boca Raton, FL, 1985, 269p. [85Z261]
- ⁷¹Clennan, E.L., *Adv. Oxygenated Processes* **1**: 85-122 (1988) [88Z421]
- ⁷²Clennan, E.L., *Tetrahedron* **47**: 1343-82 (1991) [91Z295]
- ⁷³Hurst, J.R.; Schuster, G.B., *J. Am. Chem. Soc.* **104**: 6854-6 (1982), 105: 4501 (1983) [82A342]

⁷⁴Gorman, A.A.; Gould, I.R.; Hamblett, I., *J. Am. Chem. Soc.* **104**: 7098-104 (1982) [82A349]

⁷⁵Edwards, A.M.; Silva, E., *Radiat. Environ. Biophys.* **24**: 141-8 (1985) [85F114].

⁷⁶Reddi, E.; Lambert, C.R.; Jori, G.; Rodgers, M.A.J., *Photochem. Photobiol.* **45**: 345-51 (1987) [87A064]

⁷⁷Miskoski, S.; Garcia, N.A., *Photochem. Photobiol.* **57**: 447-52 (1993) [93R059]

⁷⁸Sies, H.; Menck, C.F.M., *Mutat. Res.* **275**: 367-75 (1992) [92R287]

TABLE 1. Decay constants for singlet oxygen in various solvents.

No.	Solvent	k_d (s^{-1})	τ ($1/k_d$) (s)	Method	Comment	Ref.
1.1 Acetone						
		2.0×10^4	5.1×10^{-5}	PL/Ld-2	S = MB, TPP or PHO.	92E256
		1.9×10^4	5.4×10^{-5}	PL/Ld-2	S = HVD and PPDME.	92E274
		2.6×10^4	3.8×10^{-5}	PL/Ld-2	S = RB.	89E324
		2.3×10^4	4.3×10^{-5}	PL/Ld-2	S = TPP.	89E324
		2.4×10^4	4.1×10^{-5}	PL/Ld-2	S = MB.	89E700
		2.2×10^4	4.6×10^{-5}	PL/Ld-2	S = RB.	89E700
		2.1×10^4	4.8×10^{-5}	PL/Ld-2	S = HP.	89E700
		2.0×10^4		PL/Ld-2	S = Ac.	87E466
		1.8×10^4	5.5×10^{-5}	PL/Ld-2	S = An.	87E668
		2.1×10^4	4.7×10^{-5}	PL/Ld-2	S = An.	87E959
		2.5×10^4	4.0×10^{-5}	PL/Tb-3	S = An.	85E591
		2.1×10^4	4.7×10^{-5}	PL/Tb-3	S = ZnTPP.	85E591
		2.0×10^4	5.0×10^{-5}	PL/Ld-2	S = 2-ACN.	84E066
		4.0×10^4	2.5×10^{-5}	PL/Tb-3	S = Ery.	83A050
		3.3×10^4	3.0×10^{-5}	PL/Tb-3	S = TPP.	83A050
		2.0×10^4	5.0×10^{-5}	PL/Ld-2	S = HPDME.	83A223
		2.1×10^4	4.7×10^{-5}	PL/Ld-2	S = MPDME; no change in k at $T = 192$ to 295 K.	83E235 82E106
		2.0×10^4	5.1×10^{-5}	PL/Ld-2	S = Sulfo- and carboxyanthracenes.	83E844
		2.2×10^4	4.6×10^{-5}	PL/Ld-2	S = MPDME.	83F196
		2.8×10^4	3.6×10^{-5}	PL/Ad-8	S = An; A = DPBF.	82A349
		2.6×10^4	3.9×10^{-5}	PL/Ld-2	S = MB.	82E104
		2.5×10^4	4.0×10^{-5}	PL/Ld-2	S = RB.	82E104
		2.0×10^4	5.1×10^{-5}	PL/Ld-2	S = TPP.	82E104
		1.5×10^4	6.5×10^{-5}	PL/Ld-2	S = HP.	82E105
		2.6×10^4	3.9×10^{-5}	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329
		1.8×10^4	5.5×10^{-5}	PL/Ad-5	S = 2-ACN; A = DPBF; cor. for k_S .	81A287
		2.0×10^4	5.1×10^{-5}	PL/Ld-2	S = PdMPDME.	81E398 83E818 79E846
		2.0×10^4	5.0×10^{-5}	PL/Ld-2	S = TPP.	81E631
		2.4×10^4	4.2×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	76F903
		4×10^4	3×10^{-5}	PL/Ad-5	S = MB, A = DPBF.	727260
1.2 Acetone-<i>d</i>₆						
		1.0×10^3	9.9×10^{-4}	PL/Ld-2	S = MB, TPP or PHO.	92E256
		1.6×10^3	6.4×10^{-4}	PL/Ld-2	S = RB; varied [S] and extrapolated to [S] = 0.	89E324
		8.2×10^3	1.2×10^{-4}	PL/Tb-3	S = An.	85E591
		1.7×10^3	5.8×10^{-4}	PL/Ld-2	S = HPDME.	83A223
		1.2×10^3	8.6×10^{-4}	PL/Ld-2	S = MB, RB or TPP.	83E398
		2.6×10^3	3.9×10^{-4}	PL/Ld-2	S = PdMPDME.	83E818
		1.2×10^3	8.4×10^{-4}	PL/Ld-2	S = HP.	82E105
		1.4×10^3	6.9×10^{-4}	PL/Ld-2	S = MPDME.	82E106 83E235
		1.3×10^3	7.7×10^{-4}	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329
		1×10^3	8×10^{-4}	PL/Ad-5	S = 2-ACN; A = Rub; cor. for k_S .	81A287

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	$\tau (1/k_d)$ (s)	Method	Comment	Ref.
1.3 Acetonitrile						
		1.4×10^4	7.1×10^{-5}	PL/Ld-2	S = DCA.	93A244
		1.3×10^4	7.7×10^{-5}	PL/Ld-2	S = Ac.	93A244
		1.4×10^4		PL/Ld-2	S = Chr; rate decreased linearly with magnetic field strength, by 4-7% at 21kG.	93E183
		1.6×10^4	6.0×10^{-5}	PL/Ld-2	S = Ru(bpy) ₃ ²⁺ .	92A095
		1.3×10^4	7.7×10^{-5}	PL/Ld-2	S = MB, TPP or PHO.	92E256
		1.3×10^4	8.0×10^{-5}	PL/Ld-2	S = 1-MeNp.	91E297
		1.3×10^4		PL/Ld-2	S = Ac.	89A099
		1.3×10^4	7.5×10^{-5}	PL/Ld-2	S = OMAPB; extrapolated to zero laser dose.	89A241
		1.2×10^4	8.3×10^{-5}	PL/Ld-2	S = RB or H ₂ TPPS ⁴⁻ .	89A322
		3.0×10^4	3.3×10^{-5}	PL/Tb-3	S = (C ₆ H ₅) ₂ CO.	89E232
		1.8×10^4	5.5×10^{-5}	PL/Ld-2	S = RB.	89E324
		1.7×10^4	6.0×10^{-5}	PL/Tb-3	S = Ac.	88Z155
		1.5×10^4	6.7×10^{-5}	PL/Ld-2	S = Np, Biph, or Fl.	87E234
		1.2×10^4		PL/Ld-2	S = Ac.	87E466
		1.8×10^4	5.5×10^{-5}	PL/Ld-2	S = An.	87E668
		1.2×10^4	8.7×10^{-5}	PL/Ld-2	S = RB; from high intensity photolysis; M = ¹ O ₂ *	87F333
		1.9×10^4	5.4×10^{-5}	PL/Ld-2	¹ O ₂ * from 9,10-diphenylanthracene endoperoxide.	86F337
		2.1×10^4	4.8×10^{-5}	PL/Tb-3	S = An.	85E591
		1.8×10^4	5.6×10^{-5}	PL/Ld-2	S = 2-ACN.	84E066
		2.8×10^4	3.6×10^{-5}	PL/Tb-3	S = Ery.	83A050
		1.7×10^4	5.8×10^{-5}	PL/Ld-2	S = HPDME.	83A223
		1.6×10^4	6.4×10^{-5}	PL/Ld-2	S = MPDME.	83F196
		2.9×10^4	3.5×10^{-5}	PL/Ad-8	S = An; A = DPBF.	82A349
		1.6×10^4	6.1×10^{-5}	PL/Ld-2	S = RB.	82E104
		1.5×10^4	6.8×10^{-5}	PL/Ld-2	S = MB.	82E104
		1.8×10^4	5.4×10^{-5}	PL/Ld-2	S = MPDME.	82E106
		1.7×10^4	5.8×10^{-5}	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329
		1.1×10^4	9.2×10^{-5}	PL/Ad-5	S = 2-ACN; A = DPBF; cor. for k_S .	81A287
		1.8×10^4	5.7×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	76F903
		3×10^4	3×10^{-5}	PL/Ad-5	S = MB, A = DPBF.	727260
1.4 Acetonitrile/Water (50:50)						
		6.7×10^4	1.5×10^{-5}	PL/Ld-2	S = 9-Anthracenemethanol.	93F069
		4.5×10^4	2.2×10^{-5}	PL/Ld-2	S = RB.	93R059
1.5 Acetonitrile-<i>d</i>₃						
		9.0×10^2	1.1×10^{-3}	PL/Ld-2	S = Chr.	93E183
		8.2×10^2	1.2×10^{-3}	PL/Ld-2	S = RB or H ₂ TPPS ⁴⁻	89A322
		6.7×10^2	1.5×10^{-3}	PL/Ld-2	S = Pz.	89E597
		1.7×10^3	6.0×10^{-4}	PL/Ld-2	S = RB; from high intensity photolysis; M = ¹ O ₂ *	87F333
		9.3×10^3	1.1×10^{-4}	PL/Tb-3	S = An.	85E591
		1.6×10^3	6.2×10^{-4}	PL/Ld-2	S = HPDME.	83A223

TABLE I. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s ⁻¹)	t (1/k _d) (s)	Method	Comment	Ref.
1.5 Acetonitrile-<i>d</i>₃ — Continued						
		2.3×10^3	4.4×10^{-4}	PL/Ld-2	S = MB, RB or TPP.	83E398
		1.7×10^3	6.0×10^{-4}	PL/Ld-2	S = MPDME.	82E106
						83E235
		1.1×10^3	9.5×10^{-4}	CP/LI-12	S = Benz[<i>de</i>]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4$ s ⁻¹ in benzene.	82E329
		8×10^2	1×10^{-3}	PL/Ad-5	S = 2-ACN; A = Rub; cor. for k_S .	81A287
1.6 Acetophenone						
		3.8×10^4		PL/Ad-5	S = MB; A = DPBF.	83A371
						83A006
1.7 Benzene						
		3.2×10^4	3.1×10^{-5}	PL/Ld-2	S = TPP; $\Delta V \ddagger = -8.3$ cm ³ mol ⁻¹ ; studied from 0.1 MPa to 250 MPa.	93A017
		3.3×10^4	3.0×10^{-5}	PL/Ld-2	S = Ac.	93A244
		3.3×10^4	3.0×10^{-5}	PL/Ld-2	S = Chr.	93E183
		3.3×10^4	3.0×10^{-5}	PL/Ld-2	S = MB, TPP or PHO.	92E256
		3.3×10^4	3.0×10^{-5}	PL/Ld-2	S = TPP.	92E555
		3.4×10^4		PL/Ld-2	S = TPP.	92F251
		3.2×10^4	3.1×10^{-5}	PL/Ld-2	S = 1-MeNp.	91E297
		3.3×10^4	3.0×10^{-5}	PL/Ld-2	S = Np, Pz and BP.	90A328
		3.2×10^4	3.1×10^{-5}	PL/Ld-2	S = Por.	90E374
		3.2×10^4	3.1×10^{-5}	PL/Ld-2	S = An.	90E400
		7.4×10^4	1.4×10^{-5}	PL/Ld-2	S = CHD.	89A235
		3.3×10^4	3.0×10^{-5}	PL/Ld-2	S = TPP.	89A322
						93U041
		3.9×10^4	2.6×10^{-5}	PL/Ld-2	S = Ac.	89E324
		3.2×10^4	3.1×10^{-5}	PL/Ld-2	S = TPP.	89E388
		3.1×10^4	3.2×10^{-5}	PL/Ld-2	S = TPP.	89E700
		$\sim 3.4 \times 10^4$	$\sim 2.9 \times 10^{-5}$	PL/Ld-2	S = (C ₆ H ₅) ₂ CO.	88E452
		3.2×10^4	3.1×10^{-5}	PL/Ld-2	S = TPP.	88F503
		3.3×10^4	3.0×10^{-5}	PL/Tb-3	S = (C ₆ H ₅) ₂ CO.	88Z155
		3.0×10^4	3.3×10^{-5}	PL/Ld-2	S = TPP.	87E055
		3.1×10^4	3.2×10^{-5}	PL/Ld-2	S = Np, Bp, or Fl.	87E234
		3.3×10^4		PL/Ld-2	S = Ac.	87E466
		3.3×10^4	3.0×10^{-5}	PL/Ld-2	S = An.	87E668
		3.6×10^4	2.8×10^{-5}	PL/Ld-2	S = An.	87E959
		2.9×10^4	3.5×10^{-5}	PL/Ld-2	S = SiNC; reversible energy transfer with sensitizer proposed.	87R032
		4.1×10^4		PL/Ad-8	S = PP; A = DPBF.	85A124
		3.4×10^4	2.9×10^{-5}	PL/Tb-3	S = ZnTPP.	85E591
		3.3×10^4	3.0×10^{-5}	PL/Tb-3	S = An.	85E591
		3.2×10^4	3.1×10^{-5}	PL/Ld-2	S = 2-ACN.	84E066
		4.0×10^4		PL/Ad-5	S = 2-ACN; A = DPBF.	84F005
		4.3×10^4	2.3×10^{-5}	PL/Tb-3	S = TPP.	83A050
		3.2×10^4	3.1×10^{-5}	PL/Ld-2	S = HPDME.	83A223
		3.1×10^4	3.2×10^{-5}	PL/Ld-2	S = MB, RB or TPP.	83E398
		3.4×10^4	2.9×10^{-5}	PL/Ld-2	S = MPDME.	83F196
		3.7×10^4	2.7×10^{-5}	PL/Ld-2	S = MPDME.	82E106
						83E235

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	$\tau(1/k_d)$ (s)	Method	Comment	Ref.
1.7 Benzene — Continued						
		3.5×10^4	2.8×10^{-5}	PL/Ad-5	S = 2-ACN; A = DPBF; cor. for k_S .	81A287
		3.3×10^4	3.0×10^{-5}	PL/Ld-2	S = PdMPDME.	81E398 83E818 79E846
		4.0×10^4		PL/Ld-2	S = An.	80A143
		3.7×10^4	2.7×10^{-5}	PR/Ad-5	S = Np, A = DPBF.	78E263
		4.1×10^4		PL/ β Cd-11	S = An, Q = Car.	78F276
		3.7×10^4	2.7×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	76F903
		4.1×10^4		PL/ β Cd-11	S = An, Q = Car.	737438
		3.9×10^4		PL/Ad-8	S = An, A = DPBF.	737438
		4×10^4	2×10^{-5}	PL/Ad-5	S = MB, A = DPBF.	727260
1.8 Benzene (mic)						
		5.0×10^4	2.0×10^{-5}	PL/Tb-3	S = Ery; reverse micelles contg. 0.04 mol L ⁻¹ DAP and 0.1 mol L ⁻¹ water.	83A050
1.9 Benzene/Methanol (80:20)						
		4×10^4	2.6×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	737014
1.10 Benzene-d₆						
		1.7×10^3	5.9×10^{-4}	PL/Ld-2	S = Chr.	93E183
		1.3×10^3	7.8×10^{-4}	PL/Ld-2	S = C ₆₀ or C ₇₀ .	93E301
		1.5×10^3	6.8×10^{-4}	PL/Ld-2	S = MB, TPP or PHO.	92E256
		1.5×10^3	6.5×10^{-4}	PL/Ld-2	S = Np, Pz and BP.	90A328
		1.5×10^3	6.8×10^{-4}	PL/Ld-2	S = An.	90E400
		1.5×10^3		PL/Ld-2	S = TPP.	89A331
		1.6×10^3	6.2×10^{-4}	PR/Ld-2	S = 2-ACN.	89E113
		1.3×10^3	7.9×10^{-4}	PL/Ld-2	S = TPP.	89E388
		1.3×10^3	7.7×10^{-4}	PL/Ld-2	S = Pz.	89E597
		1.6×10^3	6.3×10^{-4}	PL/Ld-2	S = Pur.	89R044
		2.6×10^3	3.8×10^{-4}	PL/Tb-3	S = An.	85E591
		1.6×10^3	6.3×10^{-4}	PL/Ld-2	S = HPDME.	83A223
		1.4×10^3	7.0×10^{-4}	PL/Ld-2	S = MB, RB or TPP.	83E398
		3.2×10^3	3.1×10^{-4}	PL/Ld-2	S = PdMPDME.	83E818
		1.6×10^3	6.2×10^{-4}	PL/Ld-2	S = Cercosporin.	83R123
		1.8×10^3	5.5×10^{-4}	PL/Ld-2	S = MPDME.	82E106 83E235
		1.5×10^3	6.8×10^{-4}	CP/LI-12	S = Renz[de]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329
		1×10^3	1×10^{-3}	PL/Ad-5	S = 2-ACN; A = DPBF; cor. for k_S .	81A287
		2×10^3	7×10^{-4}	PL/Ad-5	S = 3,6-Bis(dibutylamino)phenothiazinium bromide; A = Rub.	81A287
		2.8×10^4	3.6×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	76F902
1.11 Benzene-d₆/Benzene (95:5)						
		4.0×10^3	2.5×10^{-4}	PL/Ld-2	S = TPP.	89A322 93U041
1.12 Benzonitrile						
		2.5×10^4		PL/Ld-2	S = Ac.	87E466
		3.1×10^4		PL/Ad-5	S = A = Rub; .	83F075

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	$\tau(1/k_d)$ (s)	Method	Comment	Ref.
1.13 Bromobenzene						
		2.4×10^4	4.2×10^{-5}	PL/Ld-2	S = TPP.	89E388
		2.0×10^4		PL/Ld-2	S = Rub.	87E466
		2.0×10^4	5.0×10^{-5}	PL/Ld-2	S = MB, RB or TPP.	83E398
1.14 Bromobenzene/Methanol (80:20)						
		4.3×10^4	2.3×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	737014
1.15 Bromobenzene-<i>d</i>₅						
		7.4×10^2	1.4×10^{-3}	PL/Ld-2	S = TPP.	89E388
		1.2×10^3	8.1×10^{-4}	PL/Ld-2	S = MB, RB or TPP.	83E398
1.16 Bromoform						
		5.9×10^3	1.7×10^{-4}	PL/Ld-2	S = PdMPDME.	81E398 83E818
		2×10^4	5×10^{-5}	PL/Ld-2	S = PdMPDME.	79E846
1.17 Bromopentafluorobenzene						
		46	0.022	MP/Ld-2	S = TPP.	89E388
1.18 1-Bromopropane						
		1×10^5	1×10^{-5}	PL/βCd-11	S = An, Q = Car.	76F902
		1×10^5	8×10^{-6}	PL/Ad-8	S = MB, A = DPBF.	76F902
1.19 1-Butanol						
		5.7×10^4	1.8×10^{-5}	PL/Ld-2	S = An.	87E668
		5.7×10^4	1.8×10^{-5}	PL/Ld-2	S = HPDME.	83A223
		5.2×10^4	1.9×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	737014
1.20 2-Butanol						
		6.2×10^4	1.6×10^{-5}	PL/Ld-2	S = RB.	89E324
		5.1×10^4	2.0×10^{-5}	PL/Ld-2	S = HPDME.	83A223
1.21 2-Butoxyethanol						
		$\sim 3.8 \times 10^5$		MD/Ac-33	A = Rub, Q = DABCO. Assumed $k_Q = 3.4 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$ and $k_A = 7.0 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$; T = 273K.	727319
1.22 Butylbenzene						
		4.0×10^4	2.5×10^{-5}	PL/Ld-2	S = O ₂ :solvent CT state.	90E220
1.23 Carbon disulfide						
		22	0.045	MP/Ld-2	S = MB, TPP or PHO; $k_d = 13 \text{ s}^{-1}$ for [O ₂] ≤ $10^{-5} \text{ mol L}^{-1}$.	92E256
		34	0.029	MP/Ld-2	S = TPP.	89A400
		29	0.034	MP/Ld-2	S = TPP.	87E658
		6.6×10^2	1.5×10^{-3}	FP/Ld-2	S = Per.	82A322
		5×10^3	2×10^{-4}	PL/Ad-5	S = MB, A = DPBF, k_d cor. for 1% MeOH content.	727260
1.24 Carbon disulfide/Methanol (98:2)						
		4.0×10^3	2.5×10^{-4}	FP/Ad-5	S = MB, A = DPBF.	737334
1.25 Carbon tetrachloride						
		17	0.059	MP/Ld-2	S = MB, TPP or PHO; $k_d = 7.8 \text{ s}^{-1}$ for [O ₂] ≤ $10^{-5} \text{ mol L}^{-1}$.	92E256
		1.3×10^3	7.7×10^{-4}	PL/Ld-2	S = TPP.	89A322 93U041

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	$\tau(1/k_d) \cdot$ (s)	Method	Comment	Ref.
1.25 Carbon tetrachloride — Continued						
		36	0.028	FP/Ld-2	S = PdMP; very dilute soln.	88E018
		36	0.028	PL/Ld-2	S = TPP.	88F503
		36	0.028	PL/Ld-2	S = Ph a.	88R193
		11	0.087	MP/Ld-2	S = TPP.	87E658
		1.0×10^3	1.0×10^{-3}	PL/Ld-2	S = An.	87E668
		36	0.028	PL/Ld-2	S = TPP.	86F316
		1.1×10^3	9.0×10^{-4}	PL/Ld-2	S = TPP.	82E104
		32	0.031	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329
		38	0.026	PL/Ld-2	S = PdMPDME.	81E398 79E846
		38	0.026	PL/Ld-2	S = TPP.	81E631
		38	0.026	PL/Ld-2	S = PdMPDME or MPDME; decay at 1272 or 1588 nm.	80E558
		36	0.028	PL/Ld-2	S = Ret.	79F463
1.26 Carbon tetrachloride/Methanol (98:2)						
		3.3×10^3		PL/Ad-5	S = MB; A = DPBF.	83A006
		3.2×10^3	3.1×10^{-4}	FP/Ad-5	S = MB, A = DPBF.	78E238
1.27 Chlorobenzene						
		2.3×10^4	4.3×10^{-5}	PL/Ld-2	S = TPP.	89E388
		2.3×10^4		PL/Ld-2	S = Ac.	87E466
		2.2×10^4	4.5×10^{-5}	PL/Ld-2	S = An.	87E668
		2.2×10^4		PL/Ad-5	S = MB; A = DPBF.	83A371 83A006
		2.0×10^4	5.1×10^{-5}	PL/Ld-2	S = MB, RB or TPP.	83E398
		2.4×10^4	4.2×10^{-5}	PL/Ld-2	S = PdMPDME.	81E398 83E818
1.28 Chlorobenzene-d₅						
		8.3×10^2	1.2×10^{-3}	PL/Ld-2	S = MB, RB or TPP.	83E398
1.29 Chloroform						
		4.4×10^3	2.3×10^{-4}	PL/Ld-2	S = MB, TPP or PHO.	92E256
		6.0×10^3	1.7×10^{-4}	PL/Ld-2	S = HVD and PPDME.	92E274
		4.0×10^3	2.5×10^{-4}	PL/Ld-2	S = TPP.	90E731 92R076
		3.8×10^3	2.6×10^{-4}	PL/Ld-2	S = TPP.	89E700
		5.0×10^3	2.0×10^{-4}	PL/Tb-3	S = Pz.	88Z155
		4.0×10^3	2.5×10^{-4}	PL/Ld-2	S = An.	87E668
		1.6×10^4	6.2×10^{-5}	PL/Tb-3	S = TPP.	83A050
		4.0×10^3	2.5×10^{-4}	PL/Ld-2	S = MB, RB or TPP.	83E398
		6.3×10^3	1.6×10^{-4}	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329
		2.4×10^3	4.1×10^{-4}	PL/Ad-5	S = 2-ACN; A = DPBF; cor. for k_S .	81A287
		4.0×10^3	2.5×10^{-4}	PL/Ld-2	S = PdMPDME.	81E398 83E818 79E846
		4.2×10^3	2.4×10^{-4}	PL/Ld-2	S = PdMPDME or MPDME; decay at 1588 nm.	80E558

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	$\tau(1/k_d)$ (s)	Method	Comment	Ref.
1.29 Chloroform — Continued						
		9.0×10^3		PL/Ad-5	S = MB, A = DPBF.	78F061
		4×10^3	2×10^{-4}	PL/Ad-8	S = MB, A = DPBF.	76F903
		2×10^4	6×10^{-5}	PL/Ad-5	S = MB, A = DPBF. k_d decreases by 50% on lowering the temperature from 298 to 223K.	757088
1.30 Chloroform/Ethanol (50:50)						
		3.0×10^4	3.3×10^{-5}	PL/Ld-2	S = Zn(pc)(py) ₂ .	92E654
1.31 Chloroform-d						
		1.4×10^2	7.0×10^{-3}	MP/Ld-2	S = MB, TPP or PHO; $k_d = 1.1 \times 10^2 s^{-1}$ for $[O_2] \leq 10^{-5} \text{ mol L}^{-1}$.	92E256
		1.1×10^2	9.4×10^{-3}	MP/Ld-2	S = TPP.	87E658
		1.6×10^3	6.4×10^{-4}	PL/Ld-2	S = MB, RB or TPP.	83E398
		2.8×10^2	3.6×10^{-3}	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329
		2.8×10^2	3.6×10^{-3}	PL/Ld-2	S = Benz[de]anthracen-7-one, phenalenone or fluorenone.	82E329
		1×10^3	9×10^{-4}	PL/Ad-5	S = 2-ACN; A = Rub; cor. for k_S .	81A287
		1.2×10^3	8.4×10^{-4}	PL/Ld-2	S = PdMPDME.	81E398 83E818 79E846
		3×10^3	3×10^{-4}	PL/Ad-5	S = MB, A = DPBF.	757088
1.32 Chloropentfluorobenzene						
		41	0.024	MP/Ld-2	S = TPP.	89E388
1.33 Chlorotrifluoromethane						
		1.0×10^3	1.0×10^{-3}	PL/Ad-5	S = MB, A = DPBF, k_d cor. for 1-2% MeOH content.	757088
1.34 Cyclohexane						
		4.3×10^4	2.3×10^{-5}	PL/Ld-2	S = MB, TPP or PHO.	92E256
		4.3×10^4	2.3×10^{-5}	PL/Ld-2	S = 1-MeNp.	91E297
		4.2×10^4	2.4×10^{-5}	PL/Ld-2	S = 2-ACN.	90NU78
		4.3×10^4	2.4×10^{-5}	PL/Ld-2	S = Np, Bp, or Fl.	87E234
		5.2×10^4		PL/Ld-2	S = Ac.	87E466
		5.0×10^4		PL/Ad-5	S = 2-ACN; A = DPBF.	84F005
		5.0×10^4	2.0×10^{-5}	PL/Tb-3	S = TPP.	83A050
		4.3×10^4	2.3×10^{-5}	PL/Ld-2	S = HPDME.	83A223
		5.0×10^4	2.0×10^{-5}	PL/Ad-8	S = An; A = DPBF.	82A349
		5.9×10^4	1.7×10^{-5}	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329
		4.3×10^4	2.3×10^{-5}	PL/Ld-2	S = PdMPDME.	81E398
		5.9×10^4	1.7×10^{-5}	PL/Ad-5	S = MB, A = DPBF, k_d cor. for 2% MeOH content.	727260
1.35 Cyclohexane (mic)						
		5.9×10^4	1.7×10^{-5}	PL/Tb-3	S = Ery; reverse micelles contg. 0.04 mol L^{-1} DAP and 0.1 mol L^{-1} water.	83A050
1.36 Cyclohexane-d₁₂						
		2.2×10^3	4.5×10^{-4}	PL/Ld-2	S = Chr; from high intensity photolysis; M = $^1O_2^*$.	87F333

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	$\tau(1/k_d)$ (s)	Method	Comment	Ref.
1.36 Cyclohexane-<i>d</i>₁₂ — Continued						
		3.1×10^3	3.2×10^{-4}	CP/LI-12	$S = \text{Benz}[de]\text{anthracen-7-one, phenalenone or fluorenone; rel. to } k_d = 4.0 \times 10^4 \text{ s}^{-1} \text{ in benzene.}$	82E329
1.37 Cyclohexanol						
		6.6×10^4	1.5×10^{-5}	PL/Ld-2	$S = \text{HPDME}; T = 308\text{K.}$	83A223
		6.3×10^4	1.6×10^{-5}	PL/Ad-8	$S = \text{MB}, A = \text{DPBF}.$	76F903
1.38 Decane						
		3.6×10^4	2.8×10^{-5}	PL/Ld-2	$S = \text{TPP}.$	83A223
1.39 Decyl alcohol						
		5.6×10^4	1.8×10^{-5}	PL/Ld-2	$S = \text{HPDME}.$	83A223
1.40 Deuterium oxide						
		1.8×10^4	5.5×10^{-5}	PL/Ld-2	$S = \text{Ru(bpy)}_3^{2+}; \text{pH} = 6.8.$	92A095
		1.5×10^4	6.8×10^{-5}	PL/Ld-2	$S = \text{MB}, \text{TPP or PHO}.$	92E256
		1.8×10^4	5.5×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^{4-}, \text{CAPcS}, \text{SnTPPSCl}_2^{4-} \text{ or ZnPCs}^{3-}; \text{Soln. contg. } 2 \times 10^{-3} \text{ mol L}^{-1} \text{ phosphate buffer and 1% NaCl wt/wt.}$	90A022
		1.7×10^4	6.0×10^{-5}	PL/Ld-2	$S = 4\text{-Thiouridine}.$	90E312
		1.4×10^4	7.0×10^{-5}	PL/Ld-2	$S = \text{RB or H}_2\text{TPPS}^{4-}; \text{pD} = 3.$	89A322 93U041
		1.6×10^4	6.4×10^{-5}	PL/Ld-2	$S = \text{RB, TPP or H}_2\text{TPPS}^{4-}; 1.0 \text{ mol L}^{-1} \text{ NaOD.}$	89A322 93U041
		1.5×10^4	6.8×10^{-5}	PL/Ld-2	$S = \text{MB}.$	89E388
		1.6×10^4	6.1×10^{-5}	PL/Ld-2	$S = \text{Chlorin e}_6; \text{pD} = 8.1.$	89E505
		1.5×10^4	6.5×10^{-5}	PL/Ld-2	$S = \text{HP}.$	89E700
		1.4×10^4	7.0×10^{-5}	PL/Ld-2	$S = \text{MB}.$	89E700
		1.9×10^4	5.2×10^{-5}	PL/Ld-2	$S = \text{AlCl(tspe); pH} = 7.$	89R092
		1.7×10^4	6.0×10^{-5}	PL/Ld-2	$S = \text{Ru(bpy)}_3^{2+}; \text{pH} = 7; \text{O}_2 \text{ saturated.}$	88A105
		1.7×10^4	5.8×10^{-5}	PL/Ld-2	$S = 2\text{-AnS}; \text{pH} = 7.$	88A398
		1.5×10^4	6.7×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^{4-}.$	88F503
		1.5×10^4	6.8×10^{-5}	PL/Ld-2	$S = \text{UP}.$	87E234
		1.7×10^4	6.0×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^{4-}.$	87E941
		1.5×10^4	6.7×10^{-5}	PL/Ld-2	$S = \text{RF}; \text{pD} = 7.$	87F290
		1.7×10^4	5.9×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^{4-}; \text{pH} = 7.$	86A198
		1.6×10^4	6.3×10^{-5}	PL/Ld-2	${}^1\text{O}_2^* \text{ from NDPO}_2.$	86A264
		1.6×10^4	6.3×10^{-5}	PL/Ld-2	$S = \text{C}_6\text{H}_5\text{CO}_2^-.$	86A264
		1.5×10^4	6.7×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^{4-}.$	86F316
		1.5×10^4	6.8×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TMpyP}^{4+}.$	84E296
		1.8×10^4	5.5×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^{4-}.$	83A223
		2.3×10^4	4.4×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^{4-}.$	83E756
		2.3×10^4	4.4×10^{-5}	PL/Ld-2	$S = \text{PdTPPS}^{4-}.$	83E818
		1.5×10^4	6.7×10^{-5}	PL/Ld-2	$S = \text{Sulfo- and carboxyanthracenes.}$	83E844
		1.9×10^4	5.4×10^{-5}	PL/Ad-5	$S = \text{PBA}; A = \text{Crocetin}; \text{pD} = 8.4.$	82A204
		1.7×10^4	5.8×10^{-5}	PL/Ld-2	$S = \text{MB}.$	82E104
		1.5×10^4	6.8×10^{-5}	PL/Ld-2	$S = \text{RB or Th}.$	82E106 83E235
		-2.2×10^4	-4.5×10^{-5}	PL/Ld-2	$S = \text{HP}.$	81E704
		1.9×10^4		PL/Ad-5	$S = \text{MB}; A = \text{ADPA}.$	80A205

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	τ ($1/k_d$) (s)	Method	Comment	Ref.
1.41 Deuterium oxide (mic)						
		1.6×10^4	6.1×10^{-5}	PL/Ld-2	$S = \text{Chlorin e}_6$; $pD = 8.1$; Soln. cont. 10^{-3} mol L $^{-1}$ Triton X-100.	89E505
		2.1×10^4	4.9×10^{-5}	PL/Ad-5	$S = \text{HYP}$; $A = \text{DPBF}$; BRIJ 35 micelles.	88N343
		2.8×10^4	3.6×10^{-5}	PL/Ld-2	$S = \text{Ppb a}$; Triton X-100 micelles.	88R193
		2.3×10^4	4.3×10^{-5}	PL/Ld-2	$S = \text{UP}$; 0.5 mol L $^{-1}$ SDS.	87E234
		1.8×10^3	5.7×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^4$; 10^{-2} mol L $^{-1}$ SDS.	83N084
		1.8×10^3	5.6×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^4$; 10^{-2} mol L $^{-1}$ CTAB.	83N084
		1.9×10^3	5.2×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^4$; 0.1 mol L $^{-1}$ Brij.	83N084
		2.9×10^3	3.5×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^4$; 10^{-2} mol L $^{-1}$ DDAB.	83N084
		1.6×10^3	6.2×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^4$; 10^{-5} mol L $^{-1}$ HSA.	83N084
		3.1×10^3	3.2×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^4$; 20 mg/mL Triton X-100.	83N084
		2.2×10^4	4.6×10^{-5}	PL/Ad-5	$S = 2\text{-ACN}$; $A = \text{DPBF}$; vesicles (4.0×10^{-2} mol L $^{-1}$ DDAB).	82N027
		2.3×10^4		PL/Ad-5	$S = \text{MB}$ or MB-tetrol; $A = \text{DPBF}$; vesicles (4.0×10^{-2} mol L $^{-1}$ DDAB).	82N027
		3.8×10^4	2.6×10^{-5}	PL/Ad-5	$S = 2\text{-ACN}$; $A = \text{DPBF}$; 0.1 mol L $^{-1}$ Brij 35.	79N041
		4.7×10^4	2.2×10^{-5}	PL/Ad-5	$S = 2\text{-ACN}$; $A = \text{DPBF}$; 0.1 mol L $^{-1}$ Igepal CO-660.	79N041
		4.3×10^4	2.4×10^{-5}	PL/Ad-5	$S = \text{MB}$; $A = \text{DPBF}$; 0.1 mol L $^{-1}$ Igepal CO-660.	79N041
		2.1×10^4	4.8×10^{-5}	PL/Ad-5	$S = 2\text{-ACN}$; $A = \text{DPBF}$; 0.1 mol L $^{-1}$ sodium laurate.	79N041
		1.9×10^4	5.4×10^{-5}	PL/Ad-5	$S = \text{MB}$; $A = \text{DPBF}$; 0.1 mol L $^{-1}$ SDS.	79N041
		1.9×10^4	5.4×10^{-5}	PL/Ad-5	$S = 2\text{-ACN}$; $A = \text{DPBF}$; 0.1 mol L $^{-1}$ SDS.	79N041
		1.9×10^4	5.4×10^{-5}	PL/Ad-5	$S = \text{MB}$; $A = \text{DPBF}$; 0.1 mol L $^{-1}$ CTAB.	79N041
		1.7×10^4	5.7×10^{-5}	PL/Ad-5	$S = 2\text{-ACN}$; $A = \text{DPBF}$; 0.1 mol L $^{-1}$ CTAB.	79N041
		4.2×10^4	2.4×10^{-5}	PL/Ad-5	$S = 2\text{-ACN}$; $A = \text{DPBF}$; 0.1 mol L $^{-1}$ Igepal CO-630.	79N041
		2.8×10^4	3.6×10^{-5}	PL/Ad-8	$S = \text{MB}$, $A = \text{DPBF}$; 0.1 mol L $^{-1}$ CTAB.	78E143
		3.0×10^4	3.3×10^{-5}	PL/Ad-8	$S = \text{MB}$; $A = \text{DPBF}$; $pD = 7.1$; 0.1 mol L $^{-1}$ SDS.	78E143
		3.3×10^4		PL/Ad-5	$S = 2\text{-ACN}$, $A = \text{DPBF}$; 0.1 mol L $^{-1}$ SDS.	78E144
1.42 Deuterium oxide/Ethanol (95:5)						
		$\sim 2 \times 10^4$	$\sim 6 \times 10^{-5}$	PL/Ld-2	$S = \text{Ppb a}$.	88R193
1.43 Deuterium oxide/Ethanol (70:30)						
		4.0×10^4	2.5×10^{-5}	PL/Ld-2	$S = \text{HP}$.	86F316
1.44 Deuterium oxide/Ethanol (67:33)						
		3.8×10^4	2.6×10^{-5}	PL/Ld-2	$S = \text{H}_2\text{TPPS}^4$.	88F503
1.45 Deuterium oxide/Methanol (50:50)						
		9.1×10^4	1.1×10^{-5}	PL/Ad-5	$S = \text{MB}$, $A = \text{DPBF}$.	727027
1.46 1,2-Dibromotetrafluoroethane						
		32	0.031	MP/Ld-2	$S = \text{MB}$, TPP or PHO; $k_d = 9.0 \text{ s}^{-1}$ for $[\text{O}_2] \leq 10^{-5} \text{ mol L}^{-1}$.	92E256
		20	0.050	MP/Ld-2	$S = \text{TPP}$.	89E388
1.47 1,1-Dichloroethane						
		1.4×10^4	7.0×10^{-5}	PL/Ld-2	$S = \text{PdMPDME}$; Paper says ethylene dichloride.	81E398
		1.5×10^4		PL/Ad-5	$S = \text{MB}$, $A = \text{DPBF}$.	78F061

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	τ ($1/k_d$) (s)	Method	Comment	Ref.
1.48 1,2-Dichloroethane						
		1.6×10^4	6.3×10^{-5}	PL/Ld-2	S = HPDME.	83A223
		1.6×10^4	6.3×10^{-5}	PL/Ld-2	S = PdMPDME; Paper says dichloroethane.	81E398 79E846
1.49 2,2-Dichloroethanol						
		2×10^4	5×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	76F903
1.50 Dichloromethane						
		1.1×10^4	9.5×10^{-5}	PL/Ld-2	S = C ₆₀ or C ₇₀ .	93E301
		1.0×10^4	9.9×10^{-5}	PL/Ld-2	S = MB, TPP or PHO.	92E256
		1.3×10^4	8.0×10^{-5}	PL/Ld-2	S = HVD and PPDME.	92E274
		1.0×10^4	1.0×10^{-4}	PL/Tb-3	S = Pz.	88Z155
		1.2×10^4	8.2×10^{-5}	PL/Ld-2	S = An.	87E668
		1.9×10^4	5.4×10^{-5}	PL/Tb-3	S = TPP.	83A050
		1.2×10^4	8.3×10^{-5}	PL/Ld-2	S = HPDME.	83A223
		1.8×10^4	5.5×10^{-5}	PL/Ld-2	S = MPDME.	83F196
		1.0×10^4	1.0×10^{-4}	PL/Ld-2	S = TPP.	82E104
		1.7×10^4	5.9×10^{-5}	CP/LI-12	S = Benz[<i>de</i>]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329
		1.1×10^4	9.1×10^{-5}	PL/Ld-2	S = PdMPDME.	81E398 79E970
		1.6×10^4		PL/Ad-5	S = MB, A = DPBF.	78F061
		7.1×10^3	1.4×10^{-4}	PL/Ad-8	S = MB, A = DPBF.	76F903
1.51 Dichloromethane (mic)						
		1.9×10^4	5.2×10^{-5}	PL/Tb-3	S = Ery; reverse micelles contg. 0.04 mol L ⁻¹ DAP and 0.1 mol L ⁻¹ water.	83A050
1.52 Dichloromethane-d₂						
		8.3×10^3	1.2×10^{-4}	CP/LI-12	S = Benz[<i>de</i>]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329
1.53 Diethyl ether						
		3.1×10^4	3.2×10^{-5}	PL/Ld-2	S = Ph a.	88R193
		2.7×10^4	3.7×10^{-5}	PL/Ld-2	S = An.	87E668
		3.3×10^4	3.0×10^{-5}	PL/Ld-2	S = HPDME.	83A223
		2.9×10^4	3.4×10^{-5}	PL/Ld-2	S = PdMPDME.	81E398 79E846
1.54 1,3-Difluorobenzene						
		1.6×10^4	6.2×10^{-5}	PL/Ld-2	S = MB, RB or TPP.	83E398
1.55 N,N-Dimethylacetamide						
		6.8×10^4	1.5×10^{-5}	PL/Ld-2	S = HVD and PPDME.	92E274
1.56 N,N-Dimethylacetamide-d₆						
		9.5×10^3	1.0×10^{-4}	PL/Ld-2	S = HVD and PPDME.	92E274
1.57 Dimethyl adipate						
		3.1×10^4	3.2×10^{-5}	PL/Ld-2	S = Pz.	89E597
1.58 1,1-Dimethylethyl methyl ether						
		3.4×10^4	2.9×10^{-5}	PL/Ld-2	S = TPP.	92A386
		2.9×10^4	3.5×10^{-5}	PL/Ld-2	S = HPDME.	83A223

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	$\tau(1/k_d)$ (s)	Method	Comment	Ref.
1.59 <i>N,N</i>-Dimethylformamide						
		7.1×10^4	1.4×10^{-5}	PL/Ld-2	S = Chr.	93E183
		5.3×10^4	1.9×10^{-5}	PL/Ld-2	S = HVD and PPDME.	92E274
		4.0×10^4	2.5×10^{-5}	PL/Ld-2	S = An.	87E668
1.60 Dimethyl sulfoxide						
		$\sim 5.2 \times 10^4$		CP/Ac-16	S = RB, A = DPBF, $A' = 2M2P$. Measured $\beta_{A'}$ = 0.055 mol L ⁻¹ , assumed $k_{A'}$ in MeOH and DMSO are the same, used $\beta_{A'} = 0.15$ and $k_d = 1.4 \times 10^5$ s ⁻¹ in MeOH.	766072
		$\sim 3.3 \times 10^4$	$\sim 3.0 \times 10^{-5}$		Method not reported, may be direct method given in ref. [719325].	74F643
1.61 1,4-Dioxane						
		4.8×10^4	2.1×10^{-5}	PL/Tb-3	S = TPP.	83A050
		3.7×10^4	2.7×10^{-5}	PL/Ld-2	S = HPDME.	83A223
		4.0×10^4	2.5×10^{-5}	PL/Ld-2	S = TPP.	82E104
		3×10^4	3×10^{-5}	PL/Ld-2	S = PdMPDME.	79E846
1.62 Dodecane						
		4.0×10^4	2.5×10^{-5}	PL/Ld-2	S = TPP.	83A223
1.63 Epibromohydrin						
		2×10^4	5×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	76F903
1.64 Ethanol						
		7.9×10^4	1.3×10^{-5}	PL/Ld-2	S = BPhe a.	93R131
		7.1×10^4	1.4×10^{-5}	PL/Ld-2	S = HP.	89E700
		6.3×10^4	1.6×10^{-5}	PL/Ld-2	S = RB.	89E700
		7.7×10^4	1.3×10^{-5}	PL/Ld-2	S = MB.	89E700
		7.1×10^4	1.4×10^{-5}	PL/Ld-2	S = Zn(pc)(py) ₂ .	88A284
		6.7×10^4	1.5×10^{-5}	PL/Ld-2	S = H ₂ TPPS ⁴ .	88F503
		7.4×10^4	1.4×10^{-5}	PL/Tb-3	S = HP.	87E054
		7.2×10^4	1.4×10^{-5}	PL/Ld-2	S = An.	87E668
		8.3×10^4		PL/Ad,Pb-5	S = An, A = MDH.	87F541
		8.3×10^4	1.2×10^{-5}	PL/Tb-3	S = An.	85E591
		7.7×10^4	1.3×10^{-5}	PL/Ld-2	S = 2-ACN.	84E066
		9.1×10^4	1.1×10^{-5}	PL/Tb-3	S = Ery.	83A050
		6.5×10^4	1.5×10^{-5}	PL/Ld-2	S = HPDME.	83A223
		1.0×10^5	9.6×10^{-6}	PL/Ld-2	S = PdMPDME.	81E398 83E818 79E846
		-9.1×10^4	$\sim 1.1 \times 10^{-5}$	PL/Ld-2	S = HP.	81E704
		1.0×10^5	1.0×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	76F902
		5.3×10^4	1.9×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	76F903
		8×10^4	1×10^{-5}	PL/Ad-5	S = MB, A = DPBF.	727260
1.65 Ethanol/Water (95:5)						
		2×10^5	5×10^{-6}	PL/ β Cd-11	S = An, Q = Car.	76F902
1.66 Ethanol-d						
		3.2×10^4	3.1×10^{-5}	PL/Ld-2	S = Zn(pc)(py) ₂ .	88A284
		3.3×10^4	3.0×10^{-5}	PL/Ld-2	S = HP.	87E054
		3.4×10^4	2.9×10^{-5}	PL/Tb-3	S = HP.	87E054

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	$\tau(1/k_d)$ (s)	Method	Comment	Ref.
1.67	Ethanol- <i>d</i> ₆					
		5×10^3	2×10^{-4}	PL/Tb-3	S = HP.	87E054
		4.3×10^3	2.3×10^{-4}	PL/Ld-2	S = PdMPDME.	81E398 83E818
1.68	Ethyl acetate					
		2.2×10^4	4.5×10^{-5}	PL/Ld-2	S = HVD and PPDME.	92E274
		2×10^4	5×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	76F903
1.69	Ethylbenzene					
		3.8×10^4	2.6×10^{-5}	PL/Ld-2	S = O ₂ ; solvent CT state.	90E220
		3.8×10^4	2.6×10^{-5}	PL/Ld-2	S = An; 5% decrease in lifetime as T increased 298 to 353K.	89E597
		4.0×10^4		PL/Ad-5	S = MB; A = DPBF.	83A371 83A006
1.70	Ethylene glycol					
		1.3×10^5	7.5×10^{-6}	PL/Ld-2	S = HPDME.	83A223
1.71	Ethylene glycol/Methanol (50:50)					
		1.2×10^5		PL/Ad-8	S = MB or RB, A = DPBF, $k_D = 1.6 \times 10^5 s^{-1}$ when [DPBF] = $1.9 \times 10^{-5} \text{ mol L}^{-1}$ and $\beta = 6.3 \times 10^{-5} \text{ mol L}^{-1}$.	737014
1.72	Fluorobenzene					
		2.3×10^4	4.3×10^{-5}	PL/Ld-2	S = TPP.	89E388
		2.2×10^4		PL/Ld-2	S = Ac.	87E466
		2.0×10^4	4.9×10^{-5}	PL/Ld-2	S = MB, RB or TPP.	83E398
1.73	Fluorobenzene- <i>d</i> ₅					
		9.1×10^2	1.1×10^{-3}	PL/Ld-2	S = MB, RB or TPP.	83E398
1.74	2-Fluoroethanol					
		6×10^4	2×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	76F903
1.75	Formamide					
		1.5×10^5	6.7×10^{-6}	PL/Ld-2	S = An.	87E668
1.76	Furan					
		8.3×10^4	1.2×10^{-5}	PL/Ld-2	S = PdMPDME.	80E549
1.77	Heptane					
		3.4×10^4		PL/Ld-2	S = 3-HF.	89E365
		3.4×10^4	2.9×10^{-5}	PL/Ld-2	S = An.	87E668
		3.4×10^4	3.0×10^{-5}	PL/Ld-2	S = TPP.	83A223
		3.6×10^4	2.8×10^{-5}	PL/Ld-2	S = PdMPDME.	81E398
1.78	Heptane/Water (mic)					
		3.4×10^4		PL/Ld-2	S = RB; reverse micelles contg. AOT.	83N171
1.79	1-Heptanol					
		5.5×10^4	1.8×10^{-5}	PL/Ld-2	S = HPDME.	83A223
1.80	Hexadecane					
		4.1×10^4	2.4×10^{-5}	PL/Ld-2	S = TPP.	83A223
1.81	Hexafluorobenzene					
		2×10^2	5×10^{-3}	PL/Ld-2	S = C ₆₀ or C ₇₀ .	93E301

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	$\tau(1/k_d)$ (s)	Method	Comment	Ref.
1.81 Hexafluorobenzene — Continued						
		48	0.021	MP/Ld-2	$S = MB, TPP \text{ or PHO}; k_d = 33 s^{-1} \text{ for } [O_2] \leq 10^{-5} \text{ mol L}^{-1}$.	92E256
		40	0.025	MP/Ld-2	$S = TPP$.	87E658
		2.9×10^2	3.4×10^{-3}	PL/Ld-2	$S = Chr$; from high intensity photolysis; $M = ^1O_2^*$.	87F333
		2.6×10^2	3.9×10^{-3}	PL/Ld-2	$S = MB, RB \text{ or TPP}$.	83E398
		1.1×10^2	9.4×10^{-3}	CP/LI-12	$S = Benz[de]anthracen-7-one, phenalenone \text{ or fluorenone}; \text{rel. to } k_d = 4.0 \times 10^4 s^{-1} \text{ in benzene}$.	82E329
		2×10^3	6×10^{-4}	PL/Ad-5	$S = MB, A = DPBF, k_d \text{ cor. for 1-2% MeOH content}$.	757088
1.82 1,1,1,3,3-Hexafluoro-2-propanol						
		1.9×10^4	5.2×10^{-5}	PL/Ld-2	$S = RB$.	89E324
1.83 Hexane						
		3.2×10^4	3.1×10^{-5}	PL/Ld-2	$S = An$.	87E668
		3.2×10^4	3.1×10^{-5}	PL/Ld-2	$S = TPP$.	83A223
		3.3×10^4	3.0×10^{-5}	PL/Ld-2	$S = PdMPDME$.	81E398
1.84 1-Hexanol						
		5.6×10^4	1.8×10^{-5}	PL/Ld-2	$S = HPDME$.	83A223
1.85 Iodobenzene						
		2.5×10^4	3.9×10^{-5}	PL/Ld-2	$S = TPP$.	89E388
		4.2×10^4		PL/Ld-2	$S = Rub$.	87E466
		2.9×10^4	3.5×10^{-5}	PL/Ld-2	$S = MB, RB \text{ or TPP}$.	83E398
1.86 Iodobenzene-d₅						
		3.6×10^3	2.8×10^{-4}	PL/Ld-2	$S = MB, RB \text{ or TPP}$.	83E398
1.87 Iodoethane						
		2.9×10^4	3.4×10^{-5}	PL/Ld-2	$S = PdMPDME$.	81E398 79E846
1.88 Iodomethane						
		2.7×10^4	3.7×10^{-5}	PL/Ld-2	$S = PdMPDME$.	81E398 79E846
1.89 Iodopentafluorobenzene						
		66	0.015	MP/Ld-2	$S = TPP$.	89E388
1.90 3-Iidotoluene						
		3.4×10^4	2.9×10^{-5}	PL/Ld-2	$S = PdMPDME$.	81E398 83E818
1.91 Mesitylene						
		6.4×10^4	1.6×10^{-5}	PL/Ld-2	$S = TPP; \Delta V_{\ddagger}^{\ddagger} = -20.7 \text{ cm}^3 \text{ mol}^{-1}$; studied from 0.1 MPa to 250 MPa.	93A017
		6.7×10^4	1.5×10^{-5}	PL/Ld-2	$S = An$.	90E400
1.92 Methanol						
		1.1×10^5	9.5×10^{-6}	PL/Ld-2	$S = MB, TPP \text{ or PHO}$.	92E256
		8.9×10^4	1.1×10^{-5}	PL/Ld-2	$S = T(m-HOP)P$.	90R164
		9.8×10^4	1.0×10^{-5}	PL/Ld-2	$S = RB \text{ or H}_2\text{TPPS}^{4-}$.	89A322 93U041
		1.1×10^5	9.0×10^{-6}	PL/Ld-2	$S = MB \text{ and RF}$.	88A165

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	τ ($1/k_d$) (s)	Method	Comment	Ref.
1.92 Methanol — Continued						
		1.0×10^5	9.7×10^{-6}	PL/Ld-2	$S = C_6H_5CO_2^-$.	86A264
		1.0×10^5	1.0×10^{-5}	PL/Ld-2	$^1O_2^*$ from NDPO ₂ .	86A264
		1.0×10^5		PL/Ad-5	$S = 2\text{-ACN}; A = DPBF$.	84F005
		1.1×10^5	9.0×10^{-6}	PL/Tb-3	$S = \text{Ery}$.	83A050
		9.6×10^4	1.0×10^{-5}	PL/Ld-2	$S = \text{HPDME}$.	83A223
		1.0×10^5	1.0×10^{-5}	PL/Ad-8	$S = \text{An}; A = DPBF$.	82A349
		$\sim 1 \times 10^5$	$\sim 7 \times 10^{-6}$	PL/Ld-2	$S = \text{HP}$.	81E704
		8.3×10^4	1.2×10^{-5}	PL/Tb-3	$S = CH_3COCOCH_3; A = 2,5\text{-DMF}$.	80E606
		1.1×10^5		PL/Ad-5	$S = \text{MB}, A = DPBF$.	78F061
		1×10^5	9×10^{-6}	PL/Ad-8	$S = \text{MB}, A = DPBF$.	76F902
		9.7×10^4	1.0×10^{-5}	PL/Ad-8	$S = \text{RB}, A = DPBF$.	737014
		8.8×10^4	1.1×10^{-5}	PL/Ad-8	$S = \text{MB}, A = DPBF$.	737014
		1×10^5	7×10^{-6}	PL/Ad-5	$S = \text{MB}, A = DPBF$.	719325
1.93 Methanol/Water (50:50)						
		3×10^5		PL/Ad-8	$S = \text{MB or RB}, A = DPBF, k_D = 3.7 \times 10^5 s^{-1}$ when $[DPBF] = 1.7 \times 10^{-5} \text{ mol L}^{-1}$ and $\beta = 5.5 \times 10^{-5} \text{ mol L}^{-1}$.	737014
		2.9×10^5	3.5×10^{-6}	PL/Ad-5	$S = \text{MB}, A = DPBF$.	727027
1.94 Methanol-d						
		4.1×10^3	2.5×10^{-4}	PL/Ld-2	$S = \text{RB or H}_2\text{TPPS}^{4-}$.	89A322 93U041
		3.4×10^4	2.9×10^{-5}	PL/Ld-2	$S = \text{MB and RF}$.	88A165
		3.6×10^4	2.8×10^{-5}	PL/Ld-2	$S = \text{SnPPCl}_2$.	88R194
		3.1×10^4	3.2×10^{-5}	PL/Tb-3	$S = \text{HP}$.	88Z155
		2.5×10^4	4.0×10^{-5}	PL/Tb-3	$S = \text{Ery}$.	83A050
		4.1×10^4	2.5×10^{-5}	PL/Ld-2	$S = \text{MPDME}$.	83F196
		2.7×10^4	3.7×10^{-5}	PL/Ld-2	$S = \text{MB}$.	82E104
1.95 Methanol-d₄						
		3.7×10^3	2.7×10^{-4}	PL/Ld-2	$S = \text{RB}$.	93A326
		4.2×10^3	2.4×10^{-4}	PL/Ld-2	$S = \text{RB}$.	92F063
		3.8×10^3		PL/Ld-2	$S = H_2\text{TPPS}^{4-}$.	89A331
		4.4×10^3	2.3×10^{-4}	PL/Ld-2	$S = \text{MB and RF}$.	88A165
		4.4×10^3	2.3×10^{-4}	PL/Ld-2	$S = \text{HPDME}$.	83A223
1.96 2-Methoxyethanol						
		9.3×10^4	1.0×10^{-5}	PL/Ad-5	$A = DPBF; S = \text{MB}$.	87E690
1.97 Methyl benzoate						
		3×10^4	4×10^{-5}	PL/Ad-8	$S = \text{MB}, A = DPBF$.	76F903
1.98 3-Methyl-1-butanol						
		5.4×10^4	1.8×10^{-5}	PL/Ld-2	$S = \text{An}$.	87E668
1.99 3-Methylpentane						
		3.1×10^4	3.2×10^{-5}	PL/Ld-2	$S = \text{HPDME}$.	83A223
1.100 2-Methyl-1-propanol						
		4.7×10^4	2.1×10^{-5}	PL/Ld-2	$S = \text{An}$.	87E668
		4.7×10^4	2.1×10^{-5}	PL/Ld-2	$S = \text{HPDME}$.	83A223

TABLE I. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	τ ($1/k_d$) (s)	Method	Comment	Ref.
1.101 2-Methyl-2-propanol						
		3.2×10^4	3.1×10^{-5}	PL/Ld-2	S = HPDME.	83A223
		3.0×10^4	3.4×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	737014
1.102 Methyl propionate						
		2.7×10^4	3.7×10^{-5}	PL/Ld-2	S = Chr.	93E183
		2.6×10^4	3.8×10^{-5}	PL/Ld-2	S = An; 5% decrease in lifetime as T increased 298 to 353K.	89E597
1.103 Nonane						
		4.2×10^4	2.4×10^{-5}	PL/Ld-2	S = TPP.	83A223
1.104 1-Nonanol						
		5.4×10^4	1.9×10^{-5}	PL/Ld-2	S = HPDME.	83A223
1.105 1-Octanol						
		5.4×10^4	1.9×10^{-5}	PL/Ld-2	S = HPDME.	83A223
1.106 Oxygen						
		2.3×10^4		PL/Ld-2	k_d in $^{18}\text{O}_2 = 50 \text{ s}^{-1}$. $T = 77\text{K}$.	84E289
1.107 Pentachloroethane						
		2.1×10^3	4.8×10^{-4}	PL/Ld-2	S = PdMPDME.	81E398
		2.2×10^3	4.5×10^{-4}	PL/Ld-2	S = PdMPDME or MPDME; decay at 1588 nm.	80E558
1.108 Pentfluorobenzene						
		3.2×10^3	3.2×10^{-4}	PL/Ld-2	S = MB, RB or TPP.	83E398
1.109 Pentane						
		2.9×10^4	3.5×10^{-5}	PL/Ld-2	S = MB, TPP or PHO.	92E256
		2.9×10^4	3.5×10^{-5}	PL/Ld-2	S = TPP.	83A223
		4.5×10^4	2.2×10^{-5}	PL/Ld-2	S = PdMPDME.	81E398
1.110 1-Pentanol						
		5.6×10^4	1.8×10^{-5}	PL/Ld-2	S = HPDME.	83A223
1.111 Perfluorodecalin						
		17	0.059	MP/Ld-2	S = MB, TPP or PHO; $k_d = 3.2 \text{ s}^{-1}$ for $[\text{O}_2] \leq 10^{-5} \text{ mol L}^{-1}$.	92E256
1.112 Perfluoroethyl ether						
		9.1×10^2	1.1×10^{-3}	CP/LI-12	S = Benz[<i>de</i>]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 \text{ s}^{-1}$ in benzene.	82E329
1.113 Perfluorohexane						
		15	0.068	MP/Ld-2	S = MB, TPP or PHO; $k_d = 4.7 \text{ s}^{-1}$ for $[\text{O}_2] \leq 10^{-5} \text{ mol L}^{-1}$.	92E256
		20		PL/Ld-2	S = Phen; from delayed fluorescence of phenanthrene attributed to singlet(oxygen)-triplet(Phen) annihilation; $T = 200\text{K}$.	88E775
		10	0.096	MP/Ld-2	S = TPP.	87E658
1.114 Perfluorohexyl iodide						
		40	0.025	MP/Ld-2	S = MB, TPP or PHO; $k_d = 19 \text{ s}^{-1}$ for $[\text{O}_2] \leq 10^{-5} \text{ mol L}^{-1}$.	92E256
		27	0.036	MP/Ld-2	S = TPP.	89E388

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	$\tau(1/k_d)$ (s)	Method	Comment	Ref.
1.115 1-Phenylethanol						
		4.6×10^4		PL/Ad-5	S = MB; A = DPBF.	83A371 83A006
1.116 1-Propanol						
		5.7×10^4	1.8×10^{-5}	PL/Ld-2	S = An.	87E668
		6.1×10^4	1.6×10^{-5}	PL/Ld-2	S = HPDME.	83A223
1.117 2-Propanol						
		4.5×10^4	2.2×10^{-5}	PL/Ld-2	S = RB.	89E324
		4.9×10^4		PL/Ld-2	S = Ac.	87E466
		4.0×10^4	2.5×10^{-5}	PL/Ld-2	S = An.	87E668
		4.5×10^4	2.2×10^{-5}	PL/Ld-2	S = HPDME.	83A223
1.118 Propylene carbonate						
		3.3×10^4	3.0×10^{-5}	PL/Ad-5	A = DPBF; S = MB.	87E690
1.119 Pyridine						
		1.8×10^5	5.7×10^{-6}	PL/Ld-2	S = RB or H_2TPPS^{4-} .	89A322 93U041
		6.5×10^4	1.5×10^{-5}	PL/Ld-2	S = Chlorin e ₆ .	89E505
		7.7×10^4	1.3×10^{-5}	PL/Ld-2	S = An.	87E959
		7.1×10^4	1.4×10^{-5}	PL/Tb-3	S = TPP.	83A050
		5.9×10^4	1.7×10^{-5}	PL/Tb-3	S = Ery.	83A050
		6.3×10^4	1.6×10^{-5}	PL/Ld-2	S = PdMPDME.	81E398 79E846
		6×10^4	2×10^{-5}	PL/Ad-8	S = RB, A = DPBF.	76F903
1.120 1,1,2,2-Tetrachloro-1,2-difluoroethane						
		24	0.042	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329
1.121 1,1,2,2-Tetrachloroethane						
		8.3×10^3		PL/Ad-5	S = MB, A = DPBF.	78F061
1.122 Tetrachloroethylene						
		8.3×10^2	1.2×10^{-3}	PL/Ld-2	S = PdMPDME.	81E398
		7.7×10^2	1.3×10^{-3}	PL/Ld-2	S = PdMPDME or MPDME; decay at 1588 nm.	80E558
1.123 Tetradecane						
		4.0×10^4	2.5×10^{-5}	PL/Ld-2	S = TPP.	83A223
1.124 1,2,4,5-Tetrafluorobenzene						
		6.5×10^3	1.6×10^{-4}	PL/Ld-2	S = MB, RB or TPP.	83E398
1.125 Tetrahydrofuran						
		4.8×10^4		PL/Ld-2	S = RB.	87E466
		4.3×10^4	2.3×10^{-5}	PL/Ld-2	S = An.	87E668
		5.4×10^4	1.9×10^{-5}	PL/Tb-3	S = An.	85E591
		5.0×10^4	2.0×10^{-5}	PL/Ld-2	S = TPP.	82E104
		3.3×10^4	3.0×10^{-5}	PL/Ld-2	S = PdMPDME.	81E398
		4.3×10^4	2.3×10^{-5}	PL/Ad-8	S = RB, A = DPBF.	76F903
1.126 Toluene						
		3.4×10^4	2.9×10^{-5}	PL/Ld-2	S = TPP; $\Delta V^\ddagger = -10.0 \text{ cm}^3 \text{ mol}^{-1}$; studied from 0.1 MPa to 250 MPa.	93A017

TABLE I. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	$\tau(1/k_d)$ (s)	Method	Comment	Ref.
1.126 Toluene — Continued						
		3.5×10^4	2.9×10^{-5}	PL/Ld-2	S = Ac.	93A017
		3.4×10^4	2.9×10^{-5}	PL/Ld-2	S = O ₂ ; solvent CT state.	90E220
		3.6×10^4	2.8×10^{-5}	PL/Ld-2	S = An.	90E400
		3.4×10^4	2.9×10^{-5}	PL/Ld-2	S = 2-ACN; $\Delta H^\ddagger = -1.7$ kJ mol ⁻¹ ; studied at 203-333 K.	88A427
		3.5×10^4		PL/Ld-2	S = Ac.	87E466
		3.4×10^4	2.9×10^{-5}	PL/Ld-2	S = An.	87E668
		4.0×10^4	2.5×10^{-5}	PL/Ld-2	S = 2-ACN.	84E066
		4.0×10^4	2.5×10^{-5}	PL/ β Cd-11	S = An.	82A349
		3.8×10^4	2.6×10^{-5}	PL/Ad-8	S = An; A = DPBF.	82A349
		3.2×10^4		PL/Ad-5	S = A = Rub; .	82E072
		3.7×10^4	2.7×10^{-5}	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329
		3.4×10^4	2.9×10^{-5}	PL/Ld-2	S = PdMPDME.	81E398 83E818 79E846
		4.3×10^4	2.3×10^{-5}	CP/Ac-14	S = TPP; A = TME; Used $k_A = 4 \times 10^7 L mol^{-1} s^{-1}$.	80C002
		4.0×10^4		PR/Ad-5	S = Np; A = DPBF; $E_a = -2$ kJ mol ⁻¹ ; studied at 222 to 333K.	79A106 78E263
		5×10^4	2×10^{-5}	CP/Ac-16	S = A = DMA; compared with $\beta = 1.2 \times 10^{-3} mol L^{-1}$ in benzene where $\tau = 24 \mu s$.	79F148
1.127 Toluene-d₈						
		3.5×10^3	2.8×10^{-4}	PL/Ld-2	S = Pz.	89E597
		3.2×10^3	3.1×10^{-4}	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329
		3.1×10^3	3.2×10^{-4}	PL/Ld-2	S = PdMPDME.	81E398 83E818 79E846
1.128 2,2,2-Trichloroethanol						
		2×10^4	5×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	76F903
1.129 Trichloroethylene						
		4.0×10^3	2.5×10^{-4}	PL/Ld-2	S = MB, TPP or PHO.	92E256
		4.5×10^3	2.2×10^{-4}	PL/Ld-2	S = PdMPDME.	81E398
		4.8×10^3	2.1×10^{-4}	PL/Ld-2	S = PdMPDME or MPDME; decay at 1588 nm.	80E558
1.130 Trichlorofluoromethane						
		9.1	0.11	MP/Ld-2	S = TPP.	89E388
		42	0.024	MP/Ld-2	S = TPP.	87E658
1.131 1,1,2-Trichlorotrifluoroethane						
		14	0.072	MP/Ld-2	S = MB, TPP or PHO; $k_d = 7.5 s^{-1}$ for $[O_2] \leq 10^{-5} mol L^{-1}$.	92E256
		10	0.099	MP/Ld-2	S = TPP.	87E658
		51	0.020	FP/Ld-2	S = Per.	82A322
		24	0.042	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; rel. to $k_d = 4.0 \times 10^4 s^{-1}$ in benzene.	82E329

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	$\tau(1/k_d)$ (s)	Method	Comment	Ref.
1.132	1,1,2-Trichlorotrifluoroethane/Ethanol (99:1)	7.1×10^2	1.4×10^{-3}	PL/Ld-2	S = HP.	81E704
1.133	1,1,2-Trichlorotrifluoroethane/Ethanol (90:10)	6.7×10^3	1.5×10^{-4}	PL/Ld-2	S = HP.	81E704
1.134	1,3,5-Trifluorobenzene	1.0×10^4	9.9×10^{-5}	PL/Ld-2	S = MB, RB or TPP.	83E398
1.135	2,2,2-Trifluoroethanol	4.1×10^4	2.4×10^{-5}	PL/Ld-2	S = RB.	89E324
		3.1×10^4		PL/Ld-2	S = RB; not extrapolated to $[S] = 0$.	87E466
		2×10^4	4×10^{-5}	PL/Ad-8	S = MB, A = DPBF.	76F903
1.136	α,α,α -Trifluorotoluene	1.6×10^4		PL/Ld-2	S = Ac.	87E466
1.137	2,4,4-Trimethyl-1-pentanol	4.1×10^4	2.4×10^{-5}	PL/Ld-2	S = HPDME.	83A223
1.138	2,2,4-Trimethylpentane	2.7×10^4	3.8×10^{-5}	PL/Ld-2	S = HPDME.	83A223
		2.7×10^4		PL/Ld-2	S = RB; reverse micelles contg. AOT.	83N171
1.139	n-Undecane	4.0×10^4	2.5×10^{-5}	PL/Ld-2	S = TPP.	83A223
1.140	Water	2.4×10^5	4.2×10^{-6}	PL/Ld-2	S = MB or PHO.	92E256
		3.2×10^5	3.1×10^{-6}	PL/Ld-2	S = H_2TPPS^{4-} ; Nonmonotonic change in lifetime over 283-368 K; $\tau_A = 4.8 \mu s$ at 368 K.	91A478
		2.4×10^5	4.1×10^{-6}	PL/Ld-2	S = RB or H_2TPPS^{4-} ; 1.0 mol L ⁻¹ NaOH.	89A322
		2.4×10^5	4.2×10^{-6}	PL/Ld-2	S = MB.	89E388
		3.2×10^5	3.1×10^{-6}	PL/Ld-2	S = H_2TPPS^{4-} ; pH = 8; oxygen pressure 1-15 atm.	89E585
		2.5×10^5	4.0×10^{-6}	PL/Ld-2	S = H_2TPPS^{4-} ; pH = 7.	86A198
		2.4×10^5	4.1×10^{-6}	PL/Ld-2	S = $C_6H_5CO_2^-$.	86A264
		3.1×10^5	3.2×10^{-6}	PL/Ld-2	S = H_2TMyP^{4+} .	84E296
		2.4×10^5	4.2×10^{-6}	PL/Ld-2	S = H_2TPPS^{4-} .	83A223
		2.9×10^5	3.5×10^{-6}	PL/Ld-2	S = PdTTPS ⁴⁻ .	83E818
		3.2×10^5	3.1×10^{-6}	PL/Ld-2	S = Sulfo- and carboxyanthracenes.	83E844
1.141	Water (mic)	2.2×10^5	4.6×10^{-6}	PL/Ld-2	S = H_2TPPS^{4-} ; 1% Triton X-100.	91A478
		3.4×10^5	2.9×10^{-6}	PL/Ac-5	S = MB; A = DPBF; BRIJ 35 micelles.	88N343
		5.0×10^4		PL/Ld-2	S = RB; 0-1 mol L ⁻¹ SDS.	83N171
		4.0×10^4		PL/Ld-2	S = RB; 0-0.5 mol L ⁻¹ CTAB.	83N171
		2.5×10^5	4.0×10^{-6}	PL/Ad-5	S = 2-ACN; A = DPBF; 0.1 mol L ⁻¹ CTAB; extrapolated from D ₂ O/H ₂ O mixtures to 100% water.	79N041
		2.9×10^5	3.5×10^{-6}	PL/Ad-5	S = 2-ACN; A = DPBF; 0.1 mol L ⁻¹ Igepal CO-630; extrapolated from D ₂ O/H ₂ O mixtures to 100% water.	79N041
		3.3×10^5		PL/Ad-5	S = 2-ACN, A = DPBF; 0.1 mol L ⁻¹ SDS.	78E144

TABLE 1. Decay constants for singlet oxygen in various solvents. — Continued

No.	Solvent	k_d (s^{-1})	τ ($1/k_d$) (s)	Method	Comment	Ref.
1.142 Water/Methanol (80:20)						
		2.0×10^5	4.9×10^{-6}	PL/Ld-2	${}^1\text{O}_2^*$ from NDPO ₂ .	86A264
1.143 <i>m</i>-Xylene						
		4.0×10^4	2.5×10^{-5}	PL/Ld-2	$S = \text{TPP}; \Delta V \ddagger = -13.0 \text{ cm}^3 \text{ mol}^{-1}$; studied from 0.1 MPa to 250 MPa.	93A017
1.144 <i>o</i>-Xylene						
		4.4×10^4	2.3×10^{-5}	PL/Ld-2	$S = \text{TPP}; \Delta V \ddagger = -14.2 \text{ cm}^3 \text{ mol}^{-1}$; studied from 0.1 MPa to 250 MPa.	93A017
		5.3×10^4	1.9×10^{-5}	PL/Ld-2	$S = \text{An.}$	90E400
1.145 <i>p</i>-Xylene						
		5.0×10^4	2.0×10^{-5}	PL/Ld-2	$S = \text{TPP}; \Delta V \ddagger = -19.2 \text{ cm}^3 \text{ mol}^{-1}$; studied from 0.1 MPa to 250 MPa.	93A017

TABLE 2. Rate constants for interaction of singlet oxygen with olefins.

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_q/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.1 Acrolein (Propenal)							
	MeOH	1.6×10^3	89	263	CP/Oc-15	S = MB; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	737479
2.2 Adamantane, 2-(hydroxymethylene)-, acetate							
	CD ₃ COCD ₃	3.4×10^4			PL/Ld-2	S = MB.	83A026
	CD ₃ OD	2.1×10^4			PL/Ld-2	S = MB.	83A026
2.3 Benzvalene							
	C ₆ H ₆	1.1 (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. $1 + k_q/k_r = 1$.	84F065	
	CCl ₄	0.059 (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. $1 + k_q/k_r = 1$.	84F065	
	CH ₂ Cl ₂	4.9×10^{-3} (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. $1 + k_q/k_r = 1$.	84F065	
	CH ₃ CN	4.0×10^{-3} (β_r)	286	CP/Oc-29	S = RB; A' = 2,5-DMF; meas. $1 + k_q/k_r = 1$.	84F065	
	CH ₃ COCH ₃	7.2×10^{-3} (β_r)	286	CP/Oc-29	S = RB; A' = 2,5-DMF; meas. $1 + k_q/k_r = 1$.	84F065	
	CHCl ₃	0.020 (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. $1 + k_q/k_r = 1$.	84F065	
	MeOH	0.019 (β_r)	286	CP/Oc-29	S = RB; A' = 2,5-DMF; meas. $1 + k_q/k_r = 1$.	84F065	
2.4 Bicyclo[2.2.1]hepta-2,5-diene (Norbornadiene)							
	CHCl ₃	1.7×10^4			CR/A'c-33	A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ${}^1\text{O}_2^*$ from DMNO ₂ .	81E003
2.5 Bicyclo[2.2.1]hepta-2,5-diene, 7,7-dimethyl-2-(trimethylsiloxy)- [Norbornadiene, 7,7-dimethyl-2-(trimethylsiloxy)-]							
	CDCl ₃			CP/Ac,A'c-17	S = TPP; A' = 7,7-Dimethyl-2-(trimethylsiloxy)norborn-2-ene; meas. $k_q/k_r A' = 4.2$.	78F290	
2.6 Bicyclo[2.2.1]hepta-2,5-diene, 2-methyl- (Norbornadiene, 2-methyl-)							
	CH ₃ CN		273	CP/Ac,A'c-17	S = MB; A' = 5-Methylene-2-norbornene; meas. $k_q/k_r A' = 34$.	78F149	
2.7 Bicyclo[2.2.1]hepta-2,5-diene, 2-(trimethylsiloxy)-							
	CDCl ₃			CP/Ac,A'c-17	S = TPP; A' = 7,7-Dimethyl-2-(trimethylsiloxy)norborn-2-ene; meas. $k_q/k_r A' = 0.17$.	78F290	
2.8 Bicyclo[2.2.1]heptane, 2,3-bis(methylene)- (Norbornane, 2,3-dimethylene-)							
	CD ₃ COCD ₃	3.6×10^5		195	CP/A'c-16	S = RB; A' = DPBF; k_d not given.	87A368
2.9 Bicyclo[2.2.1]heptane, 7,7-dimethyl-2-methylene-							
	CH ₃ CN		273	CP/Ac,A'c-17	S = MB; A' = 2,7,7-Trimethylnorborn-2-ene; meas. $k_q/k_r A' = 3.4$.	78F149	
	CH ₃ CN		273	CP/Ac,A'c-17	S = MB; A' = 2-Methylenenorbornane; meas. $k_q/k_r A' = 0.26$.	74F647	
2.10 Bicyclo[2.2.1]heptane, 2,3-dioxymethyl-7-tricyclo[3.3.1.1^{3,7}]decylidene- (exo,exo)							
	C ₆ H ₅ Cl/ 2-PrOH (90:10)	1.8×10^6		273	CR/LI-12	used $k_d = 4.1 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from hydrotrioxide, e.g. (CH ₃) ₂ C(OH)OOOH.	90M125
2.11 Bicyclo[2.2.1]heptane, 2-methylene- (Norbornane, 2-methylene-)							
	CH ₃ CN		273	CP/Ac,A'c-17	S = MB; A' = 5-Methylene-2-norbornene; meas. $k_q/k_r A' = 4.3$.	78F149	
	CH ₃ CN		273	CP/Ac,A'c-17	S = MB; A' = 2,7,7-Trimethylnorborn-2-ene; meas. $k_q/k_r A' = 13$.	78F149	
	CH ₃ CN		?		S = MB; A' = c-C ₆ H ₉ CH ₃ ; meas. $k_A/k_{A'} = 0.065$.	73F664	
2.12 Bicyclo[2.2.1]heptane, 2,3-(2¹-oxatrimethylene)-7-tricyclo[3.3.1.1^{3,7}]decylidene- (exo,exo)							
	C ₆ H ₅ Cl/ 2-PrOH (90:10)	1.2×10^6		273	CR/LI-12	used $k_d = 4.1 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from hydrotrioxide, e.g. (CH ₃) ₂ C(OH)OOOH.	90M125

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.13	Bicyclo[2.2.1]heptane-2- <i>d</i> , 3-methylene-, <i>endo</i> -						
	CH ₃ CN				CP/Ac,A'c-17	S = MB; A' = 2-Methylenenorbornane; meas. $k_t/k_{t'}^{A'} = 1.0$.	74F647
2.14	Bicyclo[2.2.1]heptane-2- <i>d</i> , 3-methylene-, <i>exo</i> -				CP/Ac,A'c-17	S = MB; A' = 2-Methylenenorbornane; meas. $k_t/k_{t'}^{A'} = 1.1$.	74F647
2.15	Bicyclo[2.2.1]heptane-2,3-dicarboxylic acid, 7-tricyclo[3.3.1.1 ^{3,7}]decylidene-, anhydride (<i>endo, endo</i>)						
	CHCl ₃	2.8×10^4		295	CL/LI-12	S = An; used $k_d = 4.9 \times 10^3$ s ⁻¹ .	90F473
2.16	Bicyclo[2.2.1]heptane-2,3-dicarboxylic acid, 7-tricyclo[3.3.1.1 ^{3,7}]decylidene-, anhydride (<i>exo, exo</i>)						
	CHCl ₃	1.0×10^4		295	CL/LI-12	S = An; used $k_d = 4.9 \times 10^3$ s ⁻¹ .	90F473
2.17	Bicyclo[2.2.1]heptane-2,3-dicarboxylic acid, 7-tricyclo[3.3.1.1 ^{3,7}]decylidene-, dimethyl ester (<i>endo, endo</i>)						
	CHCl ₃	3.1×10^4		295	CL/LI-12	S = An; used $k_d = 4.9 \times 10^3$ s ⁻¹ .	90F473
2.18	Bicyclo[2.2.1]heptane-2,3-dicarboxylic acid, 7-tricyclo[3.3.1.1 ^{3,7}]decylidene-, dimethyl ester (<i>exo, exo</i>)						
	CHCl ₃	1.5×10^4		295	CL/LI-12	S = An; used $k_d = 4.9 \times 10^3$ s ⁻¹ .	90F473
2.19	Bicyclo[2.2.1]hept-2-ene, 5,6-bis(methylene)- [Norborn-2-ene, 5,6-bis(methylene)-]						
	CH ₂ Cl ₂			253	CP/Ac-17	S = TPP; A' = 2,3-Dimethylcyclo[2.2.1]heptane; meas. $k_t/k_{t'}^{A'} = 0.12$.	82F450
2.20	Bicyclo[2.2.1]hept-2-ene, 2-methoxy- (Norborn-2-ene, 2-methoxy-)						
	c-C ₆ H ₁₂	$<1.0 \times 10^4$			PL/A'd-8	S = An; A' = DPBF.	82A349
	C ₆ H ₅ CH ₃	1.1×10^4			PL/A'd-8	S = An; A' = DPBF.	82A349
	CH ₃ CN	6.2×10^5			PL/A'd-8	S = An; A' = DPBF; $\Delta H^\ddagger = -1.3$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -141$ J K ⁻¹ mol ⁻¹ ; studied at 250-273 K.	82A349
	CH ₃ COCH ₃	2.5×10^5			PL/A'd-8	S = An; A' = DPBF.	82A349
	MeOH	6.2×10^5			PL/A'd-8	S = An; A' = DPBF; $\Delta H^\ddagger = -0.8$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -130$ J K ⁻¹ mol ⁻¹ ; studied at 250-273 K.	82A349
2.21	Bicyclo[2.2.1]hept-2-ene, 2-methyl- (Norborn-2-ene, 2-methyl-)						
	CH ₃ CN			273	CP/Ac,A'c-17	S = MB; A' = 5-Methylene-2-norbornene; meas. $k_t/k_{t'}^{A'} = 14$.	78F149
	CH ₃ CN				CP/Ac,A'c-17	S = MB; A' = 2-Methylenenorbornane; meas. $k_t/k_{t'}^{A'} = 3.1$.	74F647
	CH ₃ CN				?	S = MB; A' = c-C ₆ H ₉ CH ₃ ; meas. $k_A/k_{A'} = 0.14$.	73F664
2.22	Bicyclo[2.2.1]hept-2-ene, 5-methylene-						
	CH ₃ CN			273	CP/Ac,A'c-17	S = MB; A' = c-C ₅ H ₇ CH ₃ ; meas. $k_t/k_{t'}^{A'} = 2.0 \times 10^{-3}$.	78F149
2.23	Bicyclo[2.2.1]hept-2-ene, 2,7,7-trimethyl-						
	CH ₃ CN			273	CP/Ac,A'c-17	S = MB; A' = 2-Methylnorborn-2-ene; meas. $k_t/k_{t'}^{A'} = 0.025$.	78F149
	CH ₃ CN				CP/Ac,A'c-17	S = MB; A' = 2-Methylenenorbornane; meas. $k_t/k_{t'}^{A'} = 0.077$.	74F647
2.24	Bicyclo[2.2.1]hept-2-ene, 2-(trimethylsiloxy)-						
	CDCl ₃				CP/Ac,A'c-17	S = TPP; A' = 7,7-Dimethyl-2-(trimethylsiloxy)norborn-2-ene; meas. $k_t/k_{t'}^{A'} = 8.4$.	78F290
2.25	Bicyclo[3.1.1]hept-2-ene, 2-ethenyl-6,6-dimethyl- (Nopadiene)						
	MeOH	-2.0		293	CP/Oc-15	S = RB; $E_a = 16$ kJ mol ⁻¹ .	68F288

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.26	Bicyclo[4.1.0]hept-2-ene, 3,7,7-trimethyl- (Δ^2 -Carene)						
	MeOH	0.17		?		Method not reported.	68F289
2.27	Bicyclo[4.1.0]hept-2-ene, 4,7,7-trimethyl- (Δ^4 -Carene)						
	MeOH	0.21		?		Method not reported.	68F289
2.28	Bicyclo[4.1.0]hept-3-ene, 3,7,7-trimethyl- (Δ^3 -Carene)						
	MeOH	0.40		?		Method not reported.	68F289
2.29	Bicyclo[2.2.0]hexa-2,5-diene, hexamethyl-						
	CHCl ₃	1.1×10^7			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005
2.30	Bicyclo[3.1.0]hex-2-ene, 4-methyl-1-(1-methylethyl)-, (1 <i>R</i>)- (α -Thujene)						
	CD ₃ OD	1.8×10^6 (k_r) $<10^4$ (k_d)		295	CP/Oc-15,28	$S = \text{RB}$; used $k_d = 4.0 \times 10^3$ s ⁻¹ ; used $\phi_A(\text{RB}) = 0.76$.	91F332
2.31	Bicyclo[3.2.2]nonane, 6,7-bis(methylene)-						
	CD ₃ COCD ₃	1.7×10^6		195	CP/A'c-16	$S = \text{RB}$; $A' = \text{DPBF}$; k_d not given.	87A368
2.32	Bicyclo[4.2.0]octa-2,4-diene, 7,8-dibromo- (Cyclooctatetraene dibromide)						
	MeOH	-0.30		293	CP/Oc-15	$S = \text{RB}$; $E_a = 11$ kJ mol ⁻¹ .	68F288
2.33	Bicyclo[2.2.2]octane, 2,3-bis(methylene)-						
	CD ₃ COCD ₃	1.7×10^6		195	CP/A'c-16	$S = \text{RB}$; $A' = \text{DPBF}$; k_d not given.	87A368
2.34	Bicyclo[7.2.0]undec-4-ene, 4,11,11-trimethyl-8-methylene-, [<i>IR</i> -(<i>IR</i> [*] , <i>4E</i> , <i>9S</i> [*])]- (β -Caryophyllene)						
	MeOH/ C ₆ H ₆ (50:50)			293	CP/Ac,A'c-17	$S = \text{BP}$, triphenylene, quinoline, Np, Py, RB, MB; $A' = \gamma$ -Caryophyllene; meas. $k_r/k_r^{A'}$ = 5.1.	70F735
2.35	1,3-Butadiene, 1,4-di(4,4'-dimethoxyphenyl)-						
	CCl ₄	1.1×10^6 (k_r)			CP/Pa,P'a-17	$S = \text{TPP}$; $A' = \text{DPB}$; used $k_r^{A'} = 2.1 \times 10^5$ L mol ⁻¹ s ⁻¹ ; P = 1,4-Di(4,4'-dimethoxyphenyl)-1,3-butadiene endoperoxide.	90F292
2.36	1,3-Butadiene, 1,4-di(1,1-dimethylethoxy)-, (<i>E,E</i>)-						
	CH ₂ Cl ₂	8.5×10^7			CP/A'c-16	$S = \text{MPDME}$; $A' = \text{DPBF}$; used $k_d = 1.0 \times 10^4$ s ⁻¹ .	85A261 86A159
	CH ₃ CN	2.0×10^8			CP/A'c-16	$S = \text{MPDME}$; $A' = \text{DPBF}$; k_d not given.	86A159
	CH ₃ COCH ₃	9.1×10^7			CP/A'c-16	$S = \text{MPDME}$; $A' = \text{DPBF}$; used $k_d = 2.2 \times 10^4$ s ⁻¹ .	85A261 86A159
	CH ₃ COCH ₃	9.6×10^7 (k_r)	2.3×10^{-3} (β_r)		CP/Ac,A'c-17	$S = \text{MPDME}$; $A' = (\text{Z,Z})\text{-1,4-Di}(\text{tert}\text{-butoxy})\text{-1,3-butadiene}$; used $k_d = 2.2 \times 10^4$ s ⁻¹ ; β_r meas. rel. to $\beta_r(\text{TME}) = 1.2 \times 10^{-3}$ mol L ⁻¹ .	86A159
	THF	1.1×10^8			CP/A'c-16	$S = \text{MPDME}$; $A' = \text{DPBF}$; k_d not given.	86A159
2.37	1,3-Butadiene, 1,4-di(1,1-dimethylethoxy)-, (<i>E,Z</i>)-						
	CH ₂ Cl ₂	5.3×10^7			CP/A'c-16	$S = \text{MPDME}$; $A' = \text{DPBF}$; used $k_d = 1.0 \times 10^4$ s ⁻¹ .	85A261 86A159
	CH ₃ CN	1.1×10^8			CP/A'c-16	$S = \text{MPDME}$; $A' = \text{DPBF}$; k_d not given.	86A159
	CH ₃ COCH ₃	5.7×10^7			CP/A'c-16	$S = \text{MPDME}$; $A' = \text{DPBF}$; used $k_d = 2.2 \times 10^4$ s ⁻¹ .	85A261 86A159
	THF	5.5×10^7			CP/A'c-16	$S = \text{MPDME}$; $A' = \text{DPBF}$; k_d not given.	86A159
2.38	1,3-Butadiene, 1,4-di(1,1-dimethylethoxy)-, (<i>Z,Z</i>)-						
	CH ₂ Cl ₂	2.0×10^7			CP/A'c-16	$S = \text{MPDME}$; $A' = \text{DPBF}$; used $k_d = 1.0 \times 10^4$ s ⁻¹ .	85A261 86A159
	CH ₃ CN	3.8×10^7			CP/A'c-16	$S = \text{MPDME}$; $A' = \text{DPBF}$; k_d not given.	86A159

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.38 1,3-Butadiene, 1,4-di(1,1-dimethylethoxy)-, (Z,Z) — Continued							
	CH ₃ COCH ₃	1.5×10^7			CP/A'c-16	S = MPDME; A' = DPBF; used $k_d = 2.2 \times 10^4$ s ⁻¹ .	85A261
	CH ₃ COCH ₃	1.4×10^7 (k_t)	1.6×10^{-3} (β_r)		CP/Ac,A'c-17	S = MPDME; A' = TME; used $k_d = 2.2 \times 10^4$ s ⁻¹ , $\beta_r A' = 1.2 \times 10^{-3}$ mol L ⁻¹ .	85A261
	THF	1.9×10^7			CP/A'c-16	S = MPDME; A' = DPBF; k_d not given.	86A159
2.39 1,3-Butadiene, 2,3-dimethyl-							
	CHCl ₃	7.8×10^4			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
2.40 1,3-Butadiene, 1,4-diphenyl- (DPB)							
	CCl ₄	2.1×10^5	6.9×10^{-3}		CP/Pa-15	S = TPP; used $k_d = 1.4 \times 10^3$ s ⁻¹ ; P = 1,4-Diphenyl-1,3-butadiene endoperoxide.	90F292
	CCl ₄			273	CP/Pa-17	S = TPP; A' = Acetone azine; meas. $k_t/k_r A' = 3.5$.	79F278
	C ₆ F ₆			273	CP/Pa-17	S = TPP; A' = Acetone azine; meas. $k_t/k_r A' = 3.6$.	79F278
	CDCl ₃			253	CP/Pa-17	S = MB; A' = Acetone azine; meas. $k_t/k_r A' = 3.3$.	79F278
	CDCl ₃			253	CP/Pa-17	S = TPP; A' = Acetone azine; meas. $k_t/k_r A' = 4.5$.	79F278
	CDCl ₃			253	CP/Pa-17	S = Poly-RB; A' = Acetone azine; meas. $k_t/k_r A' = 3.6$.	79F278
	CH ₃ CN	6.0×10^5	0.056		CP/Pa-15	S = RB; used $k_d = 3.3 \times 10^4$ s ⁻¹ ; P = 1,4-Diphenyl-1,3-butadiene endoperoxide.	90F292
2.41 1,3-Butadiene, 2-ethyl-							
	CHCl ₃	7.2×10^4			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
2.42 1,3-Butadiene, 2-methyl- (Isoprene)							
	C ₆ H ₅ Cl/ 2-PrOH (90:10)	3×10^4		273	CR/LI-12	used $k_d = 4.1 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from hydrotrioxide, e.g. (CH ₃) ₂ C(OH)OOOH.	90M125
	CHCl ₃	3.7×10^4			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
2.43 1,3-Butadiene, 2-(phenylmethyl)-							
	C ₆ H ₆	7.3		277	CP/Oc-14	S = TPP.	81F582
	CCl ₄	0.82		277	CP/Oc-14	S = TPP.	81F582
	CH ₃ COCH ₂ CH ₃	5.7		277	CP/Oc-14	S = RB.	81F582
	CHCl ₃	3.4		277	CP/Oc-14	S = TPP.	81F582
	EtOH	11		277	CP/Oc-14	S = RB.	81F582
2.44 (E)-2-Butenal (Crotonaldehyde)							
	MeOH	2.2×10^3	65	263	CP/Oc-15	S = MB; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	737479
2.45 1-Butene, 2,3-dimethyl-							
	MeOH			288	CP/Ac,A'c-17	S = MB; A' = c-C ₆ H ₁₀ ; meas. $k_t/k_r A' = 1.7$.	65F028
2.46 2-Butene							
	MeOH		13	293	CP/Oc-15	S = RB; $E_a = 42$ kJ mol ⁻¹ .	68F288
2.47 2-Butene, (E)							
	CCl ₄ / MeOH (96:4)	2×10^3			PL/A'd-8	S = MB; A' = DPBF.	777162
	CS ₂	7.2×10^3		298	PL/Ld-2	S = TPP; $\Delta H^\ddagger = 1$ kJ mol ⁻¹ , $\Delta S^\ddagger = -176$ J K ⁻¹ mol ⁻¹ ; studied at 183-310 K.	85A167
2.48 2-Butene, (Z)							
	CCl ₃ F			193	CP/Ac-17	S = MPDME; A' = (E)-2-Butene; meas. $k_t/k_r A' = 18$.	90F111

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.48 2-Butene, (Z) — Continued							
	CCl ₄ /MeOH (96:4)	6×10^3			PL/A'd-8	S = MB; A' = DPBF.	77F162
	CS ₂	4.8×10^4		298	PL/Ld-2	S = TPP; $\Delta H^\ddagger = 7 \text{ kJ mol}^{-1}$; $\Delta S^\ddagger = -134 \text{ J K}^{-1}$ mol ⁻¹ ; studied at 183–310 K.	85A167
2.49 2-Butene, 2-cyclopropyl-, (E)							
	CH ₃ COCH ₃				CP/Ac,A'c-17	S = Eos; A' = (Dicyclopropylmethylidene)cyclobutane; meas. $k_q/k_{\tau}^{A'} = 0.53$.	78F430
2.50 2-Butene, 2-cyclopropyl-3-methyl-							
	CH ₃ CN			283	CP/Ac,A'c-17	S = MB; A' = TME; meas. $k_q/k_{\tau}^{A'} = 1.0$; C.k. with 1,2-dimethylcyclohexene.	79F646
	CH ₃ COCH ₃				CP/Ac,A'c-17	S = Eos; A' = (Dicyclopropylmethylidene)cyclobutane; meas. $k_q/k_{\tau}^{A'} = 0.76$.	78F430
2.51 2-Butene, 2,3-dimethyl- (TME)							
	1-BuOH	4.4×10^{-3}		296	CP/A'c-16	S = MB; A' = DPBF.	71F398
	tert-BuOH	2.9×10^{-3}		296	CP/A'c-16	S = MB; A' = DPBF.	71F398
	C ₅ H ₅ N	$\sim 9 \times 10^6$		286	?	S = Poly-RB; value from graph; method not reported.	81F534
	C ₅ H ₅ N		1.3×10^{-3}	285	CP/Pa-20	S = Th, MB, DMA; A' = DMA.	73F7202
	C ₅ H ₅ N			~283	CP/A'c-17	S = A' = DPBF; meas. $k_q/k_{\tau}^{A'} = 0.026$.	66F041
	c-C ₆ H ₁₂ (mic)	1×10^7			PL/Tb-3	S = Ery; reverse micelles contg. 0.04 mol L ⁻¹ DAP and 0.1 mol L ⁻¹ water.	83A050
	C ₆ H ₅ CH ₃	3.6×10^7			PL/Ld-2	S = 2-ACN; $\Delta H^\ddagger = -3 \text{ kJ mol}^{-1}$; $\Delta S^\ddagger = -126 \text{ J K}^{-1} \text{ mol}^{-1}$; $E_a = -1 \text{ kJ mol}^{-1}$; studied at 183–363 K; pre-exciplex-equilibrium limit activation parameters.	88A427
	C ₆ H ₅ CH ₃		1.1×10^{-3}		CP/Ac-14	S = TPP.	80C002
	C ₆ H ₅ CH ₃	4.2×10^7			PR/A'd-5	S = Np; A' = DPBF.	78E263
	C ₆ H ₅ Cl/ 2-PrOH (90:10)	2.6×10^7		273	CR/LI-12	used $k_d = 4.1 \times 10^4 \text{ s}^{-1}$; ${}^1\text{O}_2^*$ from hydrotrioxide, e.g. (CH ₃) ₂ C(OH)OOOH.	90M125
	C ₆ H ₆		1.3×10^{-3} (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. $1 + k_q/k_{\tau} = 1$.	84F065
	C ₆ H ₆	1.8×10^7	2.2×10^{-3}		CP/Oc-19	S = Chl (os); Q = Chl (os); used $k_d = 4.0 \times 10^4 \text{ s}^{-1}$; Chl (os) is a commercial mixture contg. a small % of chlorophyll.	81F321
	C ₆ H ₆	$\sim 8 \times 10^6$		286	?	S = Poly-RB; value from graph; method not reported.	81F534
	C ₆ H ₆		1.2×10^{-3}	298	CP/A'c-23	S = A' = Tet; used $\beta_{A'} = 1.7 \times 10^{-3} \text{ mol L}^{-1}$.	70F079
	C ₆ H ₆		7.1×10^{-4}	298	CP/A'c-23	S = A' = Rub; used $\beta_{A'} = 3.0 \times 10^{-4} \text{ mol L}^{-1}$.	70F079
	C ₆ H ₆		1.2×10^{-3}	298	CP/A'c-23	S = A' = DMBA; used $\beta_{A'} = 7.1 \times 10^{-4} \text{ mol L}^{-1}$.	70F079
	C ₆ H ₆		7.7×10^{-4}	298	CP/A'c-23	S = A' = DMA; used $\beta_{A'} = 3.0 \times 10^{-4} \text{ mol L}^{-1}$.	70F079
	CCl ₄		1.0×10^{-4} (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. $1 + k_q/k_{\tau} = 1$.	84F065
	CCl ₄	$\sim 5 \times 10^5$		286	?	S = Poly-RB; value from graph; method not reported.	81F534
	CH ₂ Cl ₂	5.2×10^7			PL/Ld-2	S = MoO ₅ .	93F368
	CH ₂ Cl ₂		4.0×10^{-4} (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. $1 + k_q/k_{\tau} = 1$.	84F065
	CH ₂ Cl ₂	4×10^7			PL/Tb-3	S = TPP.	83A050
	CH ₂ Cl ₂	$\sim 2 \times 10^6$		286	?	S = Poly-RB; value from graph; method not reported.	81F534
	CH ₃ CN	3.7×10^7		300	PL/P'b-5	S = MB; A' = FIN ₂ ; P' = 9-Fluorenone oxide.	85A206

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.51 2-Butene, 2,3-dimethyl- (TME) — Continued							
	CH ₃ CN	3.5×10^7		300	PL/P'b-5	S = MB; A' = DDM; P' = Benzophenone oxide.	84A339
	CH ₃ CN		$7.6 \times 10^{-4} (\beta_r)$	286	CP/Oc-29	S = RB; A' = 2,5-DMF; meas. $1 + k_q/k_r = 1$.	84F065
	CH ₃ CN	$\sim 7 \times 10^6$		286	?	S = Poly-RB; value from graph; method not reported.	81F534
	CH ₃ COCH ₃		$1.2 \times 10^{-3} (\beta_r)$	286	CP/Oc-29	S = RB; A' = 2,5-DMF; meas. $1 + k_q/k_r = 1$.	84F065
	CH ₃ COCH ₃	3×10^7			PL/Ld-2	S = MPDME.	83E235
	CH ₃ COCH ₃		6.3×10^{-4}	294	CP/A'c-16	S = MPDME; A' = DPBF.	83E235
	CH ₃ COCH ₃		7.7×10^{-4}	206			
	CH ₃ COCH ₃	$\sim 10^7$		286	?	S = Poly-RB; value from graph; method not reported.	81F534
	CH ₃ COCH ₃	5.4×10^7	7.2×10^{-4}	288	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	77F876
	CHCl ₃		$3.6 \times 10^{-4} (\beta_r)$	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. $1 + k_q/k_r = 1$.	84F065
	CHCl ₃	5.6×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81E003
	CHCl ₃	5.4×10^7			CR/A'c-33	A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ${}^1\text{O}_2^*$ from DMNO ₂ .	81E003
	CHCl ₃	$\sim 5 \times 10^6$		286	?	S = Poly-RB; value from graph; method not reported.	81F534
	CHCl ₃	5.8×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005
	CS ₂	2.2×10^7		298	PL/Ld-2	S = TPP; $\Delta H^\ddagger = 2$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -96$ J K ⁻¹ mol ⁻¹ ; studied at 183-310 K.	85A167
	CS ₂	$\sim 8 \times 10^5$		286	?	S = Poly-RB; value from graph; method not reported.	81F534
	EtOH		0.56		CP/Ac-15	S = MB; Reported values are suspect since r_{ox} depends on [O ₂] and β value was determined from nonlinear data plots.	767041
	EtOH	1.6×10^7			CP/Oc-23	S = RB; A' = Hexamethylenedithiocarbamate; used $k_d = 1.0 \times 10^4$ s ⁻¹ , $k_{A'} = 1.5 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k/(k_d + k_{A'}[A']) = 270$ at [A'] = 2.8×10^{-4} mol L ⁻¹ .	727116
	H ₂ O pH = 7.0	$1.1 \times 10^8 (k_r)$		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	87A180
	MeOH		$4.0 \times 10^{-3} (\beta_r)$	286	CP/Oc-29	S = RB; A' = 2,5-DMF; meas. $1 + k_q/k_r = 1$.	84F065
	MeOH	3×10^7			PL/Tb-3	S = Ery.	83A050
	MeOH	1.3×10^7		286	?	S = Poly-RB; value from graph; method not reported.	81F534
	MeOH		4.6×10^{-3}	296	CP/A'c-16	S = MB; A' = DPBF.	717398
	MeOII	4×10^7			PL/A'd-5	S = MB; A' = DPBF.	719325
	MeOH		6.2×10^{-3}	293	CP/Oc-15	S = RB; $E_a = 2.1$ kJ mol ⁻¹ .	68F288
	MeOH		3.0×10^{-3}		?	Method not reported.	68F289
	MeOH/ <i>tert</i> -BuOH (50:50)		2.3×10^{-3}		CP/A'c-16	S = RB; A' = DPBF.	717398
2.52 2-Butene, 2,3-dimethyl-<i>d</i>₆, (E)-	CD ₃ COCD ₃			263	CP/Pa-17	S = RB; A' = TME; meas. $k_r/k_r^{A'} = 0.71$.	79F155
2.53 2-Butene, 2,3-dimethyl-<i>d</i>₆, (Z)-	CD ₃ COCD ₃			263	CP/Pa-17	S = RB; A' = TME; meas. $k_r/k_r^{A'} = 0.91$.	79F155

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
1.34	2-Butene, 2,3-dimethyl-, (<i>d</i> ₁₂)- C ₆ H ₅ CH ₃	3.2 × 10 ⁷			PL/Ld-2	S = 2-ACN; ΔH‡ = -3 kJ mol ⁻¹ ; ΔS‡ = -126 J K ⁻¹ mol ⁻¹ ; E _a = -0.2 kJ mol ⁻¹ ; studied at 183-363 K; pre-exciplex-equilibrium limit activation parameters.	88A427
1.35	2-Butene, 2-methoxy-						
	c-C ₆ H ₁₂	2.5 × 10 ⁶			PL/A'd-8	S = An; A' = DPBF; ΔH‡ = 4 kJ mol ⁻¹ ; ΔS‡ = -103 J K ⁻¹ mol ⁻¹ ; studied at 290-340 K.	82A349
	C ₆ H ₅ CH ₃	1.2 × 10 ⁷			PL/A'd-8	S = An; A' = DPBF; ΔH‡ = -1 kJ mol ⁻¹ ; ΔS‡ = -115 J K ⁻¹ mol ⁻¹ ; studied at 250-340 K.	82A349
	CH ₃ CN	2.4 × 10 ⁷			PL/A'd-8	S = An; A' = DPBF; ΔH‡ = -3 kJ mol ⁻¹ ; ΔS‡ = -114 J K ⁻¹ mol ⁻¹ ; studied at 250-340 K.	82A349
	CH ₃ COCH ₃	1.1 × 10 ⁷			PL/A'd-8	S = An; A' = DPBF; ΔH‡ = -2.5 kJ mol ⁻¹ ; ΔS‡ = -119 J K ⁻¹ mol ⁻¹ ; studied at 240-320 K.	82A349
	MeOH	1.3 × 10 ⁷			PL/A'd-8	S = An; A' = DPBF; ΔH‡ = -2 kJ mol ⁻¹ ; ΔS‡ = -116 J K ⁻¹ mol ⁻¹ ; studied at 250-310 K.	82A349
1.36	2-Butene, 2-methyl- (2M2B)						
	C ₆ H ₆	0.035 (β_r)		286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. 1 + k _q /k _r = 1.	84F065
	CCl ₄	1.5 × 10 ⁻³ (β_r)		286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. 1 + k _q /k _r = 1.	84F065
	CD ₃ COCD ₃	1.3 × 10 ⁶		298	PL/Ld-2	S = TPP; ΔH‡ = 5 kJ mol ⁻¹ ; ΔS‡ = -113 J K ⁻¹ mol ⁻¹ ; studied at 183-310 K.	85A167
	CD ₃ OD	9.4 × 10 ⁵			CP/LI-12	S = RB; used k _d = 4.0 × 10 ³ s ⁻¹ .	92A386
	CH ₂ Cl ₂	5.8 × 10 ⁻³ (β_r)		286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. 1 + k _q /k _r = 1.	84F065
	(CH ₃) ₃ CCH ₂ OCH ₃	1.1 × 10 ⁶			CP/LI-12	S = TPP; used k _d = 3.5 × 10 ⁴ s ⁻¹ .	92A386
	CH ₃ CN	0.011 (β_r)		286	CP/Oc-29	S = RB; A' = 2,5-DMF; meas. 1 + k _q /k _r = 1.	84F065
	CH ₃ COCH ₃	0.020 (β_r)		286	CP/Oc-29	S = RB; A' = 2,5-DMF; meas. 1 + k _q /k _r = 1.	84F065
	CHCl ₃	7.2 × 10 ⁻³ (β_r)		286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. 1 + k _q /k _r = 1.	84F065
	CHCl ₃	1.5 × 10 ⁶			CP/A'c-33	S = A' = Rub; used k _d = 1.7 × 10 ⁴ s ⁻¹ , k _{A'} = 5.3 × 10 ⁷ L mol ⁻¹ s ⁻¹ .	81E003
	CHCl ₃	2.3 × 10 ⁶			CR/A'c-33	A' = Rub; used k _d = 1.7 × 10 ⁴ s ⁻¹ , k _{A'} = 5.3 × 10 ⁷ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from DMNO ₂ .	81E003
	CHCl ₃	2.3 × 10 ⁶			CP/A'c-33	S = A' = Rub; used k _d = 1.7 × 10 ⁴ s ⁻¹ , k _{A'} = 5.3 × 10 ⁷ L mol ⁻¹ s ⁻¹ .	78A005
	CS ₂	7.2 × 10 ⁵		298	PL/Ld-2	S = TPP; ΔH‡ = 3 kJ mol ⁻¹ ; ΔS‡ = -125 J K ⁻¹ mol ⁻¹ ; studied at 183-310 K.	85A167
	EtOH	1.6			CP/Ac-15	S = MB; Reported values are suspect since r _{ox} depends on [O ₂] and β value was determined from nonlinear data plots.	767041
	MeOII	0.096 (β_r)		286	CP/Oc-29	S = RB; A' = 2,5-DMF; meas. 1 + k _q /k _r = 1.	84F065
	MeOH	1.1 × 10 ⁶		286	?	S = Poly-RB; method not reported.	81F534
	MeOH	1.1 × 10 ⁶	0.10		?	used k _d = 1.1 × 10 ⁵ s ⁻¹ ; Unpublished data.	79F074
	MeOH	0.060		293	CP/Oc-15	S = FeCl ₄ ²⁻ ; E _a = 6.7 kJ mol ⁻¹ .	68F288
	MeOH	0.030		293	CP/Oc-15	S = DNT; E _a = 5.9 kJ mol ⁻¹ .	68F288
	MeOH	0.60		293	CP/Oc-15	S = RB; E _a = 6.7 kJ mol ⁻¹ .	68F288
	MeOH	0.055		293	CP/Oc-15	S = FeBr ₄ Cl ₄ ²⁻ ; E _a = 6.3 kJ mol ⁻¹ .	68F288
	MeOH	0.10		293	CP/Oc-15	S = MB; E _a = 9.2 kJ mol ⁻¹ .	68F288
	MeOH	0.055		?		Method not reported.	68F289
	MeOH/tert-BuOH (50:50)			CP/Ac,A'c-17	S = RB; A' = TME; meas. k _r /k _{r'} ^{A'} = 0.045.	68F292	
	MeOH/tert-BuOH (50:50)			CR/Ac,A'c-17	A' = TME; meas. k _r /k _{r'} ^{A'} = 0.043; ¹ O ₂ * from H ₂ O ₂ /Ca(OCl) ₂ .	68F292	

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.56	2-Butene, 2-methyl- (2M2B) — Continued						
	MeOH/ <i>tert</i> -BuOH (50:50)				CR/Ac,A'c-17	$A' = \text{TME}$; meas. $k_t/k_{t'}^{A'} = 0.067$; ${}^1\text{O}_2^*$ from $\text{H}_2\text{O}_2/\text{NaOCl}$.	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)	303			CP/Pa,P'a-17	$S = \text{RB}$; $A' = \text{TME}$; meas. $k_t/k_{t'}^{A'} = 0.024$.	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)	298			CR/Pa,P'a-17	$A' = \text{TME}$; meas. $k_t/k_{t'}^{A'} = 0.029$; ${}^1\text{O}_2^*$ from $\text{H}_2\text{O}_2/\text{NaOCl}$.	68F292
2.57	(E)-2-Butenoic acid (Crotonic acid)						
	C ₆ D ₆	3.3×10^3			PL/Ld-2	$S = \text{TPP}$.	89A331
2.58	2-Butenoic acid, 2-methyl-, (E)						
	C ₆ D ₆	6.8×10^3			PL/Ld-2	$S = \text{TPP}$.	89A331
2.59	3-Butenoic acid						
	C ₆ D ₆	4.0×10^3			PL/Ld-2	$S = \text{TPP}$.	89A331
2.60	2-Buten-1-ol, 3-methyl-						
	MeOH/ H ₂ O (95:5)		0.14		CP/Oc-15	$S = \text{RB}$; Soln. contg. 1% sodium acetate.	79F137
2.61	1-Buten-3-one, 1-(2,6,6-trimethyl-1-cyclohexen-1-yl)- (β-Ionone)						
	C ₆ H ₆	1.6×10^6			PL/A'd-5	$S = A$; $A' = \text{DPBF}$.	85E293
2.62	β-apo-8'-Carotenal						
	C ₆ H ₆	1.4×10^{10}			PL/βCd-11	$S = \text{An}$.	78F276
	CH ₂ Cl ₂	3.1×10^9		298	CP/P'a-19	$S = \text{Chl } a$; $A' = \text{Soybean oil}$; used $k_{A'} = 1.0 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 3.0 \times 10^4$; obs. peroxide formation.	91U180
2.63	α-Carotene						
	C ₆ H ₅ CH ₃	8.2×10^9			PL/Ld-2	$S = \text{Pz}$.	91E465
	CHCl ₃ / EtOH (50:50)	6×10^9		310	CR/LI-12	${}^1\text{O}_2^*$ from NDPO ₂ ; reported $k = 1.9 \times 10^{10}$ L mol ⁻¹ s ⁻¹ using $k_d = 1 \times 10^5$ s ⁻¹ , recalcd. using $k_d = 3.0 \times 10^4$ s ⁻¹ [92E654].	89R188 90E622
2.64	9-cis-β-Carotene						
	C ₆ H ₅ CH ₃	9×10^9			PL/Ld-2	$S = \text{Pz}$.	91E465
	C ₆ H ₆	1.1×10^{10}			PL/Ld-2	$S = \text{Pz}$.	91E465
2.65	15,15'-(Z)-β-Carotene						
	C ₆ H ₅ CH ₃	1.2×10^{10}			PL/Ld-2	$S = \text{Pz}$.	91E465
	C ₆ H ₆	1.1×10^{10}			PL/Ld-2	$S = \text{Pz}$.	91E465
2.66	β-Carotene						
	1-BuOH		5.5×10^{-6}		CP/A'c-16	$S = \text{RB}$; $A' = \text{DPF}$.	717398
	<i>tert</i> -BuOH		3.8×10^{-6}		CP/A'c-16	$S = \text{RB}$; $A' = \text{DPF}$.	717398
	C ₅ H ₅ N	6.5×10^9			CP/A'c-23	$S = A' = \text{Rub}$; used $k_d = 6.0 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	743112
	C ₆ D ₆	2.1×10^{10}			PL/Ld-2	$S = \text{Pur}$.	89R044
	n-C ₆ H ₁₄	1.4×10^{10}			PL/Ld-2	$S = \text{Pz}$.	91E465
	n-C ₆ H ₁₄	3.7×10^7 (k_t)			CP/Ac-?	$S = A' = \text{An}$; used $k_d = 6 \times 10^4$ s ⁻¹ ; calcd. using $k_A = 3 \times 10^{10}$ L mol ⁻¹ s ⁻¹ .	80F169
	C ₆ H ₅ Br	3.4×10^9		273	MD/A'c-33	$A' = \text{Rub}$; used $k_d = 1.3 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k/[(k_d/[A']) + k_{A'}] = 33.7$ at $[A'] = 1.5 \times 10^{-4}$ mol L ⁻¹ .	737333
	C ₆ H ₅ CH ₃	1.4×10^{10}			PL/Ld-2	$S = \text{Pz}$.	91E465

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
146 β-Carotene — Continued							
	C ₆ H ₅ CH ₃	1.1 × 10 ¹⁰			PL/Ld-2	S = 2-ACN; $\Delta H^\ddagger = 2$ (5) kJ mol ⁻¹ ; $\Delta S^\ddagger = -60$ (-50) J K ⁻¹ mol ⁻¹ ; $E_a = 4$ (7) kJ mol ⁻¹ ; studied at 183-363 K; pre-exciplex-equilibrium limit (and diffusion limit).	88A427
	C ₆ H ₅ CH ₃	1 × 10 ¹⁰			CP/A'c-20	S = A' = MDH; used $k_d = 4.0 \times 10^4$ s ⁻¹ .	83F406
	C ₆ H ₅ CH ₃	3.0 × 10 ¹⁰			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
	C ₆ H ₆	1.3 × 10 ¹⁰			PL/Ld-2	S = Pz.	91E465
	C ₆ H ₆	1.3 × 10 ¹⁰		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
	C ₆ H ₆	2.0 × 10 ⁶ (k_d)					
	C ₆ H ₆	1.2 × 10 ¹⁰ (k_d)					
	C ₆ H ₆	1.5 × 10 ¹⁰		303	CP/P'a-19	S = A' = Rub; k_d not given.	87A107
	C ₆ H ₆	1.2 × 10 ¹⁰			PL/ β Cd-11	S = An.	78F276
	C ₆ H ₆	1.3 × 10 ¹⁰		298	PL/A'd-8	S = An; A' = DPBF.	737438
	C ₆ H ₆	1.1 × 10 ¹⁰		298	PL/ β Cd-11	S = An.	737438
	C ₆ H ₆	2 × 10 ¹⁰			PL/A'd-5	S = MB; A' = DPBF; Solvent contained 2% MeOH.	727260
	C ₆ H ₆ /EtOH (67:33)	1.6 × 10 ¹⁰			CP/A'c-19	S = RB; A' = Chl a; used $k_d = 1 \times 10^5$ s ⁻¹ .	78F404
	C ₆ H ₆ /EtOH (89:11)	1.3 × 10 ¹⁰		295	CP/P'a-19	S = RB; A' = TEMP-4-OH; used $k_d = 3 \times 10^4$ s ⁻¹ ; P' = TEMPOL.	757445
	C ₆ H ₆ /MeOH (60:40)	2.3 × 10 ¹⁰	4.3 × 10 ⁻⁶		CP/A'c-22	S = MB; A' = Rub; used $k_d = 1.0 \times 10^5$ s ⁻¹ .	747042
	C ₆ H ₆ /MeOH (80:20)	9.7 × 10 ⁹	3.9 × 10 ⁻⁶		CP/A'c-16	S = RB; A' = DPF; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	717398
	C ₆ H ₆ /MeOH (80:20)			298	CP/P'a-20	S = MB; A' = 2M2P; meas. $k_A/k_{A'} = 1.5 \times 10^4$.	70F734
	CCl ₄	9.9 × 10 ⁹			PL/Ld-2	S = Pz.	91E465
	CCl ₄	1.0 × 10 ¹⁰		296	CR/LI-12	used $k_d = 1.7 \times 10^3$ s ⁻¹ ; ${}^1\text{O}_2^*$ from DMNO ₂ .	90F069
	CCl ₄	5.9 × 10 ⁹			CP/LI-12	S = PHO; used $k_d = 35$ s ⁻¹ .	88E479
	CCl ₄	3.8 × 10 ⁶ (k_d)			CP/Ac-14,27	S = TPP; A' = Tetr; k_t derived using $k_A = 7 \times 10^9$ L mol ⁻¹ s ⁻¹ .	83A336
	CCl ₄	7.0 × 10 ⁹			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	83A336 78E892
	CCl ₄	6.5 × 10 ⁹		295	PL/Ld-2	S = PdMPDME or MPDME; decay at 1588 nm; decay at 1272 nm gave 7×10^9 L mol ⁻¹ s ⁻¹ .	80E558
	CCl ₄	8.0 × 10 ⁹			MP/LI-12	S = Ret; used $k_d = 36$ s ⁻¹ .	79F463 78F700
	CCl ₄	7 × 10 ⁹			MP/LI-12	S = PP, Ph a, or BChl a; used $k_d = 35$ s ⁻¹ .	77E617 78E881 79A010
	CCl ₄ /CHCl ₃ (90:10)	6.2 × 10 ⁹			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k/(k_A[A'] + k_d) = 3.5 \times 10^6$ L mol ⁻¹ at [A'] = 5×10^{-6} mol L ⁻¹ .	74F645
	CD ₃ OD	1.4 × 10 ⁹			CP/LI-12	S = RB; used $k_d = 4.4 \times 10^3$ s ⁻¹ .	88E479
	CD ₃ OD	1.5 × 10 ⁹			CP/LI-12	S = PHO; used $k_d = 4.4 \times 10^3$ s ⁻¹ .	88E479
	CH ₂ Cl ₂	4.6 × 10 ⁹		298	CP/P'a-19	S = Chl a; A' = Soybean oil; used $k_{A'} = 1.0 \times 10^5$ L mol ⁻¹ s ⁻¹ ; obs. peroxide formation.	91U180
	CH ₂ Cl ₂	1.3 × 10 ¹⁰			PL/Tb-3	S = TPP.	83A050
	CH ₂ Cl ₂	1.3 × 10 ¹⁰		298	CP/A'c-23	S = A' = Rub; used $k_d = 8.0 \times 10^3$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.66 β-Carotene — Continued							
	CH ₂ Cl ₂	8.5×10^9			CP/A'c-23	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	732066
	CHCl ₃	5×10^9			CR/LI-12	used $k_d = 3.0 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from NDPO ₂ .	92R054
	CHCl ₃	1.1×10^{10}			PL/Ld-2	S = Pz.	91E465
	CHCl ₃	6.5×10^9			PL/Ld-2	S = MPDEE or PdMPDEE.	81E472
	CHCl ₃ / EtOH (50:50)	1.2×10^{10}			PL/Ld-2	S = Pz; Soln. cont. 1% H ₂ O.	91E465
	CHCl ₃ / EtOH (50:50)	4×10^9		310	CR/LI-12	$^1\text{O}_2^*$ from NDPO ₂ ; reported $k = 1.4 \times 10^{10}$ L mol ⁻¹ s ⁻¹ using $k_d = 1 \times 10^5$ s ⁻¹ , recalcd. using $k_d = 3.0 \times 10^4$ s ⁻¹ [92E654].	89R188 90E622
	CS ₂		1.5×10^{-7}		CP/P'a-20	S = TPP; A' = 2M2P.	727028
	ClCF ₂ CCl ₂ F	1.1×10^{10}		303	CR/P'a-32	A' = Rub; $^1\text{O}_2^*$ from DMNO ₂ ; $k_{A'}$ and k_d not given.	87A107
	ClCF ₂ CCl ₂ F/ CH ₂ Cl ₂ (96:4)	9.5×10^9			PL/Tb-3	S = TPP.	83A050
	EtOH/ H ₂ O (95:5)	2×10^9			CR/LI-12	$^1\text{O}_2^*$ from pyrogallol autoxidation by O ₂ /KOH; k_d not given.	78F605
	MeOH	9.3×10^8		297	CR/P'a-31	A' = TEMP; formn. of TEMPO monitored by esr; soln. cont. MeONa and CoCl ₂ ; $^1\text{O}_2^*$ from autoxidation of adrenaline; k_d not given.	92D227
	MeOH		6.1×10^{-6}		CP/A'c-16	S = RB.	717398
	MeOH/ H ₂ O (95:5)	6.3×10^9		310	CR/LI-12	used $k_d = 1.8 \times 10^5$ s ⁻¹ ; soln. cont. 0.05 mol L ⁻¹ MeONa and 5×10^{-4} mol L ⁻¹ CoCl ₂ ; $^1\text{O}_2^*$ from autoxidation of oxytetracycline.	92M228
2.67 β,β-Carotene, 3'-(acetoxy)-6',7'-didehydro-5,6-epoxy-5,5',6,6',7,8-hexahydro-3,3',5'-trihydroxy-8-oxo-, (3S,3'S,5R,5'R,6S,6'R)- (Fucoxanthin)							
	CCl ₄	2.6×10^5 (k_r)			CP/Ac-14,27	S = TPP; A' = Tetr; k_r derived using $k_A = 7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	83A336
	CCl ₄	7.0×10^8			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	83A336
2.68 ψ-Carotene, 3,4-didehydro-1,2,7',8'-tetrahydro-1-methoxy- (all-E)- (Spheroidene)							
	C ₆ D ₆	1.6×10^{10}			PL/Ld-2	S = Pur.	89R044
2.69 ψ-Carotene, 3,4-didehydro-1,2,7',8'-tetrahydro-1-methoxy-2-oxo- (all-E)- (Spheroidenone)							
	C ₆ D ₆	1.9×10^{10}			PL/Ld-2	S = Pur.	89R044
2.70 ψ-Carotene, 7,8-dihydro- (all-E)- (Neurosporene)							
	C ₆ D ₆	1.7×10^{10}			PL/Ld-2	S = Pur.	89R044
2.71 ψ-Carotene, 3,3',4,4'-tetradehydro-1,1',2,2'-tetrahydro-1,1'-dimethoxy- (all-E)- (Spirilloxanthin)							
	C ₆ D ₆	1.3×10^{10}			PL/Ld-2	S = Pur.	89R044
2.72 ψ-Carotene, 7,8,11,12-tetrahydro- (γ-Carotene)							
	C ₆ D ₆	3.5×10^9			PL/Ld-2	S = MPDME.	89R044
2.73 ψ,ψ-Carotene, 7,7',8,8',11,11,12-hexahydro-, 15-cis- (15-cis-Phytofluene)							
	CHCl ₃	$<1 \times 10^7$			CR/LI-12	used $k_d = 3.0 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from NDPO ₂ .	92R054
	C ₆ H ₆ / MeOH (60:40)	$\leq 1.0 \times 10^8$	$\geq 1.0 \times 10^{-3}$		CP/A'c-22	S = MB; A' = Rub; used $k_d = 1.0 \times 10^5$ s ⁻¹ .	747042
2.74 15(Z)-ψ-Carotene, 7,7',8,8',11,11',12,12'-octahydro- (15-cis-Phytoene)							
	CHCl ₃	$<1 \times 10^7$			CR/LI-12	used $k_d = 3.0 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from NDPO ₂ .	92R054
	C ₆ H ₆ / MeOH (60:40)	$\leq 1.9 \times 10^7$	$\geq 5.3 \times 10^{-3}$		CP/A'c-22	S = MB; A' = Rub; used $k_d = 1.0 \times 10^5$ s ⁻¹ .	747042

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.76 (E)-β,ϵ-Carotene, 3,3'-dihydroxy- (3R,3'R,6,6'R)- (Lutein)							
	C ₆ H ₆ / MeOH (60:40)	2.1×10^{10}	4.8×10^{-6}		CP/A'c-22	S = MB; A' = Rub; used $k_d = 1.0 \times 10^5$ s ⁻¹ .	747042
	CH ₂ Cl ₂	5.7×10^9	1.8×10^{-6}	293	CP/Oc-19	S = Chl a; A' = Soybean oil; used $k_d = 1.0 \times 10^4$ s ⁻¹ .	90R128
	CHCl ₃	1.6×10^{10}			PL/Ld-2	S = Pz; Contains ~5% zeaxanthin.	91E465
	CHCl ₃ / EtOH (50:50)	2×10^9		310	CR/LI-12	$^1\text{O}_2^*$ from NDPO ₂ ; reported $k = 8 \times 10^9$ L mol ⁻¹ s ⁻¹ using $k_d = 1 \times 10^5$ s ⁻¹ , recalcd. using $k_d = 3.0 \times 10^4$ s ⁻¹ [92E654].	89R188 90E622
3.77 γ-Carotene							
	CHCl ₃ / EtOH (50:50)	7×10^9		310	CR/LI-12	$^1\text{O}_2^*$ from NDPO ₂ ; reported $k = 2.5 \times 10^{10}$ L mol ⁻¹ s ⁻¹ using $k_d = 1 \times 10^5$ s ⁻¹ , recalcd. using $k_d = 3.0 \times 10^4$ s ⁻¹ [92E654].	89R188 90E622
	CHCl ₃	8.3×10^8			PL/Ld-2	S = Pz.	91E465
3.78 Carotene analog, C-30							
	C ₆ H ₆ / MeOH (80:20)				CP/Pa-20	S = MB; A' = 2M2P; meas. $k/k_{A'} = (57 \pm 86)$.	707188
3.79 β,β-Carotene-3,3'-diol, (3R,3'R)- (Zeaxanthin)							
	C ₆ H ₆	1.2×10^{10}			PL/Ld-2	S = Pz.	91E465
	C ₆ H ₆	2.8×10^9			PL/ β Cd-2	S = An.	80A143
	CCl ₄	7.0×10^9			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	83A336
	CCl ₄	3.4×10^6 (k_r)			CP/Ac-27	S = TPP; A' = Tetr; k_r derived using $k_A = 7 \times 10^9$ L mol ⁻¹ s ⁻¹ .	83A336
	CH ₂ Cl ₂	6.8×10^9	1.5×10^{-6}	293	CP/Oc-19	S = Chl a; A' = Soybean oil; used $k_d = 1.0 \times 10^4$ s ⁻¹ .	90R128
	CHCl ₃ / EtOH (50:50)	3×10^9		310	CR/LI-12	$^1\text{O}_2^*$ from NDPO ₂ ; reported $k = 1.0 \times 10^{10}$ L mol ⁻¹ s ⁻¹ using $k_d = 1 \times 10^5$ s ⁻¹ , recalcd. using $k_d = 3.0 \times 10^4$ s ⁻¹ [92E654].	89R188 90E622
3.80 β,β-Carotene-3,3'-diol, 5,6:5',6'-diepoxy-5,5',6,6'-tetrahydro- (all-E) (Violaxanthin)							
	C ₆ H ₆	1.6×10^{10}			PL/Ld-2	S = Pz.	91E465
3.81 β,β-Carotene-4,4'-diol, all-E- (Isozeaxanthin)							
	C ₆ H ₆ / MeOH (60:40)	2.9×10^{10}	3.4×10^{-6}		CP/A'c-22	S = MB; A' = Rub; used $k_d = 1.0 \times 10^5$ s ⁻¹ .	747042
	CH ₂ Cl ₂	7.4×10^9	1.3×10^{-6}	293	CP/Oc-19	S = Chl a; A' = Soybean oil; used $k_d = 1.0 \times 10^4$ s ⁻¹ .	90R128
3.82 β-Carotene-4,4'-dione, 3,3'-dihydroxy- (Astaxanthin)							
	C ₆ H ₆	1.4×10^{10}			PL/Ld-2	S = Pz.	91E465
	CH ₂ Cl ₂	9.8×10^9	1.0×10^{-6}	293	CP/Oc-19	S = Chl a; A' = Soybean oil; used $k_d = 1.0 \times 10^4$ s ⁻¹ .	90R128
	CHCl ₃ / EtOH (50:50)	7×10^9		310	CR/LI-12	$^1\text{O}_2^*$ from NDPO ₂ ; reported $k = 2.4 \times 10^{10}$ L mol ⁻¹ s ⁻¹ using $k_d = 1 \times 10^5$ s ⁻¹ , recalcd. using $k_d = 3.0 \times 10^4$ s ⁻¹ [92E654].	89R188 90E622
3.83 β,β-Carotene-4,4'-dione (Canthaxanthin)							
	C ₆ H ₅ CH ₃	1.3×10^{10}			PL/Ld-2	S = Pz.	91E465
	C ₆ H ₆	1.8×10^{10}			PL/ β Cd-2	S = An.	80A143
	C ₆ H ₆	1.4×10^{10}			PL/ β Cd-11	S = An.	78F276

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.83	β,β -Carotene-4,4'-dione (Canthaxanthin) — Continued						
	CCl ₄	1.0×10^{10}			CP/LI-12	S = PHO; used $k_d = 35$ s ⁻¹ .	88E479
	CD ₃ OD	1.2×10^{10}			CP/LI-12	S = PHO; used $k_d = 4.4 \times 10^3$ s ⁻¹ .	88E479
	CH ₂ Cl ₂	1.1×10^{10}			CP/P'a-19	S = Chl a; A' = Soybean oil; used $k_{A'} = 1.0 \times 10^5$ L mol ⁻¹ s ⁻¹ ; obs. peroxide formation.	91U180
	CHCl ₃ / EtOH (50:50)	6×10^9		310	CR/LI-12	$^1\text{O}_2^*$ from NDPO ₂ ; reported $k = 2.1 \times 10^{10}$ L mol ⁻¹ s ⁻¹ using $k_d = 1 \times 10^5$ s ⁻¹ , recalcd. using $k_d = 3.0 \times 10^4$ s ⁻¹ [92E654].	89R188
2.84	κ,κ -Carotene-6,6'-dione, 2,2'-dihydroxy- (2S,2'S,5R,5'R) (C ₄₀ -Epiisocapsorubin)						
	CHCl ₃	8×10^9			CR/LI-12	used $k_d = 3.0 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from NDPO ₂ .	92R054
2.85	β -apo-8-Carotenoic acid, ethyl ester						
	C ₆ H ₆	1.2×10^{10}			PL/ β Cd-11	S = An.	78F276
2.86	β -apo-8'-Carotenol						
	C ₆ H ₆ / MeOH (80:20)				CP/P'a-20	S = MB; A' = 2M2P; meas. $k_A/k_{A'} = -1.9 \times 10^3$.	707188
	C ₆ H ₆ / MeOH (80:20)	1.2×10^{10}		298	CP/P'a-20	S = MB; A' = 2M2P; used $k_d = 1.0 \times 10^5$ s ⁻¹ , $\beta_{A'} = 0.04$ mol L ⁻¹ .	70F734
2.87	β,β -Caroten-3-ol (Cryptoxanthin)						
	CHCl ₃ / EtOH (50:50)	2×10^9		310	CR/LI-12	$^1\text{O}_2^*$ from NDPO ₂ ; reported $k = 6 \times 10^9$ L mol ⁻¹ s ⁻¹ using $k_d = 1 \times 10^5$ s ⁻¹ , recalcd. using $k_d = 3.0 \times 10^4$ s ⁻¹ [92E654].	89R188 90E622
2.88	β,β -Caroten-4-one (Echinone)						
	C ₆ H ₆	1.1×10^{10}			PL/ β Cd-2	S = An.	80A143
2.89	3,5-Cholestadiene						
	CHCl ₃	7.4×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
2.90	4-Chosten-3-one, 4-methyl-						
		$>2 \times 10^2$			CP/Ac-18	S = Poly-RB or RB or MB; A' = Linalool; used $\beta_{A'} = 0.18$ mol L ⁻¹ ; Solvent is CH ₂ Cl ₂ or MeOH or C ₆ H ₆ .	80F111
2.91	(E)-Cinnamic acid						
	CD ₃ OD	3.4×10^3		293	PL/Ld-2	S = HP.	90F411
2.92	Coumarin, 3,4,5,6,7,8-hexahydro-						
	CD ₃ COCD ₃	1.0×10^6 (k_r) 5.5×10^5 (k_q)		296	CP/Ad,P'a-17	S = MB; A' = 2M2P; used $k_r^{A'} = 8.1 \times 10^5$ L mol ⁻¹ s ⁻¹ ; $\Delta H^\ddagger = 4.6$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -130$ J K ⁻¹ mol ⁻¹ .	86F219
	CD ₃ COCD ₃	1.1×10^6		296	PL/Ld-2	S = MB.	86F219
2.93	Crocetin						
	D ₂ O pD = 8.4	6×10^8		293	PL/Ad,A'd-5	S = PBA; A' = DPBF; used $k_{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	82A204
	D ₂ O pD = 8.4	2.5×10^9		293	PL/Ad-5	S = PBA.	82A204
	DMF	$<5 \times 10^6$ (k_r) $\sim 7 \times 10^9$ (k_q)		308	CR/Ad-?	used $k_d = 1.4 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	87A318
	DMSO	$<5 \times 10^6$ (k_r) $\sim 2 \times 10^9$ (k_q)		308	CR/Ad-?	used $k_d = 4.2 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	87A318
	H ₂ O pH = 7.8	2.5×10^8 (k_r) 5.5×10^9 (k_q)		308	CR/Ad-?	used $k_d = 4.2 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	87A318

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_q/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
1.94 Cyclobutane, (1-cyclopropylethylidene)-							
	CH ₃ COCH ₃			CP/Ac,A'c-17	S = Eos; A' = (Dicyclopropylmethyliidene)cyclobutane; meas. $k_t/k_t^{A'} = 0.70$.		78F430
1.95 Cyclobutane, (cyclopropylmethylene)-							
	CH ₃ COCH ₃			CP/Ac,A'c-17	S = Eos; A' = (Dicyclopropylmethylene)cyclobutane; meas. $k_t/k_t^{A'} = 0.59$.		78F430
1.96 Cyclobutane, (dicyclopropylmethyliidene)-							
	C ₂ H ₅ I			CP/Ac,A'c-17	S = TPP; A' = TMS; meas. $k_t/k_t^{A'} = 0.21$.		78F430
	C ₆ H ₆			CP/Ac,A'c-17	S = TPP; A' = TMS; meas. $k_t/k_t^{A'} = 0.25$.		78F430
	CH ₂ Cl ₂			CP/Ac,A'c-17	S = MB; A' = TMS; meas. $k_t/k_t^{A'} = 0.21$.		78F430
	CH ₃ COCH ₃			CP/Ac,A'c-17	S = Eos; A' = TMS; meas. $k_t/k_t^{A'} = 0.26$.		78F430
	CS ₂			CP/Ac,A'c-17	S = TPP; A' = TMS; meas. $k_t/k_t^{A'} = 0.22$.		78F430
	MeOH			CP/Ac,A'c-17	S = Eos; A' = TMS; meas. $k_t/k_t^{A'} = 0.21$.		78F430
1.97 Cyclobutene, 1-<i>tert</i>-butyl-							
	CH ₃ CN			273	CP/Ac,A'c-17	S = MB; A' = 1- <i>tert</i> -Butylcyclohexene; meas. $k_t/k_t^{A'} = 5.8$.	81F018
1.98 Cyclobutene, 1-methyl-							
	CH ₃ CN			273	CP/Ac,A'c-17	S = MB; A' = 1- <i>tert</i> -Butylcyclohexene; meas. $k_t/k_t^{A'} = 9.3$.	81F018
1.99 1,3-Cycloheptadiene							
	CHCl ₃	1.1×10^6		CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .		81F445
2.100 Cycloheptane, ethylidene-							
	CHCl ₃	4.8×10^6		CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .		81F445
2.101 Cycloheptane, methylene-							
	CH ₃ CN			273	CP/Ac,A'c-17	S = MB; A' = Methylenecyclohexane; meas. $k_t/k_t^{A'} = 62$.	81F018
2.102 Cycloheptene, 1-methyl-							
	CH ₃ CN			273	CP/Ac,A'c-17	S = MB; A' = 1- <i>tert</i> -Butylcyclohexene; meas. $k_t/k_t^{A'} = 17$.	81F018
	CHCl ₃	2.4×10^6		CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .		81F445
2.103 1,3-Cyclohexadiene							
	C ₆ H ₆	0.015 (β_r)		286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. $1 + k_q/k_t = 1$.	84F065
	CCl ₄	4.4×10^{-4} (β_r)		286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. $1 + k_q/k_t = 1$.	84F065
	CCl ₄	1.8×10^6			CP/Ac-14	S = TPP; used $k_d = 1.4 \times 10^3$ s ⁻¹ .	80C002
	CH ₂ Cl ₂	3.2×10^{-3} (β_r)		286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. $1 + k_q/k_t = 1$.	84F065
	CH ₃ CN	9.4×10^{-3} (β_r)		286	CP/Oc-29	S = RB; A' = 2,5-DMF; meas. $1 + k_q/k_t = 1$.	84F065
	CH ₃ COCH ₃	0.013 (β_r)		286	CP/Oc-29	S = RB; A' = 2,5-DMF; meas. $1 + k_q/k_t = 1$.	84F065
	CHCl ₃	4.3×10^{-3} (β_r)		286	CP/Oc-29	S = TPP; A' = 2,5-DMF; meas. $1 + k_q/k_t = 1$.	84F065
	CHCl ₃	7.1×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
	MeOH	0.037 (β_r)		286	CP/Oc-29	S = RB; A' = 2,5-DMF; meas. $1 + k_q/k_t = 1$.	84F065
	MeOH	0.025		293	CP/Oc-15	S = FlBr ₄ Cl ₄ ²⁻ ; $E_a = 5.4$ kJ mol ⁻¹ .	68F288

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.103 1,3-Cyclohexadiene — Continued							
	MeOH	0.011		293	CP/Oc-15	S = DNT; $E_a = 5.9 \text{ kJ mol}^{-1}$.	68F288
	MeOH	0.040		293	CP/Oc-15	S = FeCl ₄ ²⁻ ; $E_a = 5.9 \text{ kJ mol}^{-1}$.	68F288
	MeOH	0.073		293	CP/Oc-15	S = MB; $E_a = 5.0 \text{ kJ mol}^{-1}$.	68F288
	MeOH	0.045		293	CP/Oc-15	S = RB; $E_a = 5.0 \text{ kJ mol}^{-1}$.	68F288
	MeOH/ <i>tert</i> -BuOH (50:50)			CR/Ac,A'c-17	A' = TME; meas. $k_r/k_r^{A'} = 0.33$; $^1\text{O}_2^*$ from $\text{H}_2\text{O}_2/\text{Ca}(\text{OCl})_2$.	68F292	
	MeOH/ <i>tert</i> -BuOH (50:50)			CP/Ac,A'c-17	S = RB; A' = TME; meas. $k_r/k_r^{A'} = 0.077$.	68F292	
	2-PrOH	7.3×10^5		273	CR/LI-12	used $k_d = 4.1 \times 10^4 \text{ s}^{-1}$; $^1\text{O}_2^*$ from hydrotrioxide, e.g. $(\text{CH}_3)_2\text{C}(\text{OH})\text{OOOH}$.	90M125
2.104 1,3-Cyclohexadiene, 5,6-bis(1-methylethylidene)- (Tetramethyl-<i>o</i>-xylylene)							
	CH ₃ CN	3.2×10^8			PL/Ld-2	S = Pz.	92A325
2.105 1,3-Cyclohexadiene, 1-methyl-4-(1-methylethyl)- (α-Terpinene)							
	CHCl ₃	1×10^8			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	81F445
	MeOH	4.0×10^{-3}		293	CP/Oc-15	S = MB; $E_a = 1.7 \text{ kJ mol}^{-1}$; studied at 173-293 K.	68F288
	MeOH	1.3×10^{-3}		293	CP/Oc-15	S = Acridine Orange; $E_a = 1.7 \text{ kJ mol}^{-1}$.	68F288
	MeOH	3.3×10^{-3}		293	CP/Oc-15	S = FeBr ₄ Cl ₄ ²⁻ ; $E_a = 1.3 \text{ kJ mol}^{-1}$.	68F288
	MeOH	1.4×10^{-3}		293	CP/Oc-15	S = DNT; $E_a = 1.3 \text{ kJ mol}^{-1}$.	68F288
	MeOH	6.0×10^{-3}		293	CP/Oc-15	S = FeCl ₄ ²⁻ ; $E_a = 0.84 \text{ kJ mol}^{-1}$.	68F288
	MeOH	3.1×10^{-3}		293	CP/Oc-15	S = RB; $E_a = 1.7 \text{ kJ mol}^{-1}$.	68F288
	MeOH	0.010		288	CP/Oc-15	S = RB.	587004
2.106 1,3-Cyclohexadiene, 2-methyl-5-(1-methylethyl)-							
	CHCl ₃	3.2×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	81F445
	MeOH	0.010		293	CP/Oc-15	S = RB; $E_a = 4.2 \text{ kJ mol}^{-1}$.	68F288
2.107 1,4-Cyclohexadiene, 1-methyl-							
	CHCl ₃	2.0×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	81F445
2.108 2,4-Cyclohexadien-1-one, 6,6-dimethyl-							
	MeOH	~1.9		293	CP/Oc-15	S = RB; $E_a = 15 \text{ kJ mol}^{-1}$.	68F288
2.109 Cyclohexane, cyclohexylidene-							
	MeOH	0.030		293	CP/Oc-15	S = RB; $E_a = 5.4 \text{ kJ mol}^{-1}$.	68F288
	MeOH	0.056		?		Method not reported.	627005
2.110 Cyclohexane, ethylidene-							
	CHCl ₃	8.6×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	81F445 78A005
2.111 Cyclohexane, (hydroxymethylene)-, acetate							
	CD ₃ COCD ₃	7.5×10^4			PL/Ld-2	S = MB.	83A026
	CD ₃ OD	4.8×10^4			PL/Ld-2	S = MB.	83A026
2.112 Cyclohexane, methylene-							
	CD ₃ COCD ₃	6.1×10^3		298	PL/Ld-2	S = TPP; $\Delta H^\ddagger = -0.8 \text{ kJ mol}^{-1}$; $\Delta S^\ddagger = -176 \text{ J K}^{-1} \text{ mol}^{-1}$; studied at 183-310 K.	85A167
	CH ₃ CN	$<1 \times 10^3 (k_r)$?		S = MB; A' = c-C ₆ H ₅ CH ₃ .	73F664

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.112 Cyclohexane, methylene- — Continued							
	CS ₂	5.2×10^3		298	PL/Ld-2	S = TPP; $\Delta H^\ddagger = -0.4$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -176$ J K ⁻¹ mol ⁻¹ ; studied at 183-310 K.	85A167
2.113 Cyclohexanol, 2-(cyclohexylidene)-							
	MeOH	0.44		?		Method not reported.	627005
2.114 Cyclohexanone, 2-cyclohexylidene-							
		0.34		CP/Ac-18	S = Poly-RB or RB; A' = Linalool; used $\beta_{A'} = 0.18$ mol L ⁻¹ ; Solvent is CH ₂ Cl ₂ or MeOH.	80F111	
	MeOH	0.35		293	CP/Oc-15	S = RB; $E_a = 11.7$ kJ mol ⁻¹ .	68F288
	MeOH	0.37		?		Method not reported.	627005
2.115 Cyclohexanone, 5-methyl-2-(1-methylethylene)- (Pulegone)							
		0.12		CP/Ac-18	S = Poly-RB or RB or MB; A' = Linalool; used $\beta_{A'} = 0.18$ mol L ⁻¹ ; Solvent is CH ₂ Cl ₂ or MeOH or C ₆ H ₆ .	80F111	
	CHCl ₃	6.1×10^5		CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445	
2.116 Cyclohexene							
	C ₅ H ₅ N	$\sim 10^4$ (k_t)		286	?	S = Poly-RB; method not reported; value from graph.	81F534
	C ₆ H ₆	$\sim 8 \times 10^3$ (k_t)		286	?	S = Poly-RB; method not reported; value from graph.	81F534
	C ₆ H ₆ /MeOH (75:25)	2.0×10^4	3.1	298	CP/Pa-15	S = RB; used $k_d = 6.3 \times 10^4$ s ⁻¹ .	79A457 78F586
	CCl ₄	$\sim 5 \times 10^2$ (k_t)		286	?	S = Poly-RB; method not reported; value from graph.	81F534
	CH ₂ Cl ₂	$\sim 7 \times 10^3$ (k_t)		286	?	S = Poly-RB; method not reported; value from graph.	81F534
	CH ₃ CN	$\sim 10^4$ (k_t)		286	?	S = Poly-RB; method not reported; value from graph.	81F534
	CH ₃ COCH ₃	$\sim 2 \times 10^4$ (k_t)		286	?	S = Poly-RB; method not reported; value from graph.	81F534
	CHCl ₃	$\sim 9 \times 10^3$ (k_t)		286	?	S = Poly-RB; method not reported; value from graph.	81F534
	CS ₂	$\sim 10^3$ (k_t)		286	?	S = Poly-RB; method not reported; value from graph.	81F534
	MeOH	2.7×10^4 (k_t)		286	?	S = Poly-RB; method not reported; value from graph.	81F534
	MeOH			288	CP/Ac,A'c-17	S = MB; A' = CH ₃ CH ₂ CH ₂ CH=CHCH ₃ ; meas. $k_t/k_t^{A'} = 0.12$.	65F028
	MeOH	26		?		Method not reported.	68F289
	MeOH/tert-BuOH (50:50)			303	CP/Pa,P'a-17	S = RB; A' = (CH ₃) ₂ CHCH=CHCH ₃ ; meas. $k_t/k_t^{A'} = 0.19$.	68F292
2.117 Cyclohexene, 1,2-dimethyl-							
	CHCl ₃	3.0×10^7		CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005	
	MeOH	3.9×10^6		286	?	S = Poly-RB; method not reported.	81F534
	MeOH	1×10^7		PL/A'd-5	S = MB; A' = DPBF.	727260	
	MeOH			288	CP/Ac,A'c-17	S = MB; A' = TME; meas. $k_t/k_t^{A'} = 0.53$.	65F028
	MeOH	0.030		?		Method not reported.	68F289

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.118	Cyclohexene, 1,3-dimethyl-						
	CHCl ₃	5.1×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
2.119	Cyclohexene, 1,4-dimethyl-				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
2.120	Cyclohexene, 2,3-dimethyl-				288 CP/Ac,A'c-17	S = MB; A' = <i>c</i> -C ₆ H ₉ CH ₃ ; meas. $k_r/k_r^{A'} = 1.1$.	65F028
2.121	Cyclohexene, 4-(1-hydroxy-1-methylethyl)-1-methyl- (Terpinolene)				293 CP/Oc-15	S = RB; $E_a = 1.7$ kJ mol ⁻¹ .	68F288
2.122	Cyclohexene, 1-methyl						
	C ₆ H ₆	0.15		298	CP/A'c-23	S = A' = Rub; used $\beta_{A'} = 3.0 \times 10^{-4}$ mol L ⁻¹ .	70E079
	C ₆ H ₆	0.14		298	CP/A'c-23	S = A' = Tetr; used $\beta_{A'} = 1.7 \times 10^{-3}$ mol L ⁻¹ .	70E079
	C ₆ H ₆	0.13		298	CP/A'c-23	S = A' = DMA; used $\beta_{A'} = 3.0 \times 10^{-4}$ mol L ⁻¹ .	70E079
	C ₆ H ₆	0.12		298	CP/A'c-23	S = A' = DMBA; used $\beta_{A'} = 7.1 \times 10^{-4}$ mol L ⁻¹ .	70E079
	CH ₃ CN			273	CP/Ac,A'c-17	S = MB; A' = 1- <i>tert</i> -Butylcyclohexene; meas. $k_r/k_r^{A'} = 2.3$.	81F018
	CH ₃ COCH ₃	2.1×10^5 (k_r)		281	?	S = ?; A' = 1,4-Dioxene; used $k_{A'} = 2.2 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.96$.	70F733
	CHCl ₃	3.3×10^5			CR/A'c-33	A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from DMNO ₂ .	81E003
	CHCl ₃	3.4×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81E003
	CHCl ₃	3.6×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
	MeOH	2.0×10^5		286	?	S = Poly-RB; method not reported.	81F534
	MeOH			288	CP/Ac,A'c-17	S = MB; A' = <i>c</i> -C ₅ H ₇ CH ₃ ; meas. $k_r/k_r^{A'} = 0.12$.	65F028
	MeOH			288	CP/Ac,A'c-17	S = Ery; A' = <i>c</i> -C ₅ H ₇ CH ₃ ; meas. $k_r/k_r^{A'} = 0.11$.	65F028
	MeOH			288	CP/Ac,A'c-17	S = HP; A' = <i>c</i> -C ₅ H ₇ CH ₃ ; meas. $k_r/k_r^{A'} = 0.11$.	65F028
	MeOH			288	CP/Ac,A'c-17	S = RB; A' = <i>c</i> -C ₅ H ₇ CH ₃ ; meas. $k_r/k_r^{A'} = 0.087$.	65F028
	MeOH			288	CP/Ac,A'c-17	S = Eos; A' = <i>c</i> -C ₅ H ₇ CH ₃ ; meas. $k_r/k_r^{A'} = 0.093$.	65F028
	MeOH	1.2			?	Method not reported.	68F289
	MeOH/ <i>tert</i> -BuOH (50:50)				CR/Ac,A'c-17	A' = 2M2B; meas. $k_r/k_r^{A'} = 0.050$; ¹ O ₂ * from H ₂ O ₂ /Ca(OCl) ₂ .	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)			276	CR/Pa,P'a-17	A' = 2M2P; meas. $k_r/k_r^{A'} = 0.26$; ¹ O ₂ * from H ₂ O ₂ /NaOCl.	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)			303	CP/Pa,P'a-17	S = RB; A' = 2M2P; meas. $k_r/k_r^{A'} = 0.22$.	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)				CP/Ac,A'c-17	S = RB; A' = 2M2B; meas. $k_r/k_r^{A'} = 0.043$.	68F292
2.123	Cyclohexene, 4-methyl-						
	MeOH			288	CP/Ac,A'c-17	S = MB; A' = <i>c</i> -C ₆ H ₁₀ ; meas. $k_r/k_r^{A'} = 0.67$.	65F028
2.124	Cyclohexene, 1-methyl-4-(1-methylethyl)-, (R)- (Limonene)						
	CH ₃ COCH ₃	1.7×10^5			PL/Ld-2	S = RB.	92F225
	MeOH	1.7		293	CP/Oc-15	S = MB; $E_a = 8.4$ kJ mol ⁻¹ .	68F288
	MeOH	1.7			?	Method not reported.	627005
2.125	Cyclohexene, 1-methyl-4-(1-methylethyl)- (Carvomenthene)						
	MeOH	-1.0		293	CP/Oc-15	S = MB; $E_a = 13$ kJ mol ⁻¹ .	68F288

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.126	Cyclohexen-1-ol, 6,6-dimethyl-, acetate						
	CD ₃ COCD ₃	$8 \times 10^3 (k_r)$		296	CP/Pa-17	S = MB; A' = Adamantylidenemethyl acetate; used $k_r^{A'} = 2.7 \times 10^4$ L mol ⁻¹ s ⁻¹ .	86F219
2.127	2-Cyclohexen-1-one, 2-ethyl-3-methyl-						
		$>2 \times 10^2$			CP/Ac-18	S = Poly-RB or RB; A' = Linalool; used $\beta_{A'} = 0.18$ mol L ⁻¹ ; Solvent is CH ₂ Cl ₂ or MeOH.	80F111
2.128	2-Cyclohexen-1-one, 2-hydroxy-3-methyl-						
	MeOH	$\sim 4 \times 10^7 (k_r)$		273	CP/Pa,P'a-17	S = RB; A' = TME; $k_r^{A'}$ not given.	85F150
2.129	1,3-Cyclooctadiene						
	CHCl ₃	6.5×10^4			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
2.130	1,4-Cyclooctadiene, 6-hydroperoxy-						
	CHCl ₃	2.9×10^4	0.58		CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	89F349
2.131	1,4-Cyclooctadiene, 6-methoxy-						
	MeOH	7.2×10^5	0.14		CP/A'c-33	S = A' = Rub; used $k_d = 1.0 \times 10^5$ s ⁻¹ , $k_{A'} = 3.1 \times 10^7$ L mol ⁻¹ s ⁻¹ .	89F349
2.132	1,4-Cyclooctadiene, 6-(trimethylsiloxy)-						
	CHCl ₃	6.2×10^4	0.28		CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	89F349
2.133	1,5-Cyclooctadiene						
	CHCl ₃	3.3×10^4	0.53		CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	89F349
	MeOH	4.6×10^5		293	CP/Ac-14	S = MB; used $k_d = 2.0 \times 10^5$ s ⁻¹ .	89F358
	MeOH	3.1×10^5		293	CP/Ac-14	S = HP; used $k_d = 2.0 \times 10^5$ s ⁻¹ .	89F358
	MeOH	2.7×10^5		293	CP/Ac-14	S = RB; used $k_d = 2.0 \times 10^5$ s ⁻¹ .	89F358
	MeOH	4.9×10^5		293	CP/Ac-14	S = HA; used $k_d = 2.0 \times 10^5$ s ⁻¹ .	89F358
2.134	2,5-Cyclooctadien-1-ol						
	CHCl ₃	8.6×10^4	0.20		CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	89F349
	MeOH	1.9×10^5	0.54		CP/A'c-33	S = A' = Rub; used $k_d = 1.0 \times 10^5$ s ⁻¹ , $k_{A'} = 3.1 \times 10^7$ L mol ⁻¹ s ⁻¹ .	89F349
2.135	Cyclooctane, ethylenic-						
	CHCl ₃	4.9×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
2.136	1,3,5,7-Cyclooctatetraene						
	C ₆ H ₅ Cl/ 2-PrOH (90:10)	1.3×10^5		273	CR/LI-12	used $k_d = 4.1 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from hydrotrioxide, e.g. (CH ₃) ₂ C(OH)OOOH.	90M125
2.137	Cyclooctene						
	CHCl ₃	4.8×10^4			CR/A'c-33	A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from DMNO ₂ .	81E003
	EtOH				CP/Ac,A'c-17	S = MB; A' = 1,5-Cyclooctadiene; meas. $k_r/k_r^{A'} = 0.14$.	71F581
2.138	Cyclooctene, 1-methyl-						
	CH ₃ CN			273	CP/Ac,A'c-17	S = MB; A' = 1- <i>tert</i> -Butylcyclohexene; meas. $k_r/k_r^{A'} = 4.7$.	81F018

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.138	Cyclooctene, 1-methyl-	— Continued					
	CHCl ₃	5.3×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
2.139	Cyclooctyne				CR/A'c-33	A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; $^1\text{O}_2^*$ from DMNO ₂ .	81E003
2.140	Cyclopentadiene				PR/A'd-5	S = Np; A' = DPBF; $\Delta H^\ddagger = 0$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -108$ J K ⁻¹ mol ⁻¹ ; studied at 223-353 K.	79A106
	C ₆ H ₅ CH ₃	3.9×10^7			253 CP/Ac-17	S = TPP; A' = 2,3-Dimethylenecyclo[2.2.1]heptane; meas. $k_t/k_t^{A'}$ = >10.	82F450
	CH ₂ Cl ₂				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
	CHCl ₃	1×10^8			293 CP/Oc-15	S = RB; $E_a = 1.3$ kJ mol ⁻¹ .	68F288
	MeOH	4.4×10^{-3}			293 CP/Oc-15	S = MB; $E_a = 0.84$ kJ mol ⁻¹ .	68F288
	MeOH	4.1×10^{-3}			CR/Ac,A'c-17	A' = TME; meas. $k_t/k_t^{A'} = 0.7$; $^1\text{O}_2^*$ from H ₂ O ₂ /Ca(OCl) ₂ .	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)				CR/Ac,A'c-17	A' = TME; meas. $k_t/k_t^{A'} = 2.0$; $^1\text{O}_2^*$ from H ₂ O ₂ /NaOCl.	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)				CP/Ac,A'c-17	S = RB; A' = TME; meas. $k_t/k_t^{A'} = 1.2$.	68F292
2.141	1,3-Cyclopentadiene, 1,4-bis(4-chlorophenyl)-				CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	CH ₂ Cl ₂	6.5×10^7					
2.142	1,3-Cyclopentadiene, 1,4-bis(4-fluorophenyl)-				CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	CH ₂ Cl ₂	7.6×10^7					
2.143	1,3-Cyclopentadiene, 1,4-bis(4-methoxyphenyl)-				CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	CH ₂ Cl ₂	1.3×10^8					
2.144	1,3-Cyclopentadiene, 1,4-bis(4-methylphenyl)-				CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	CH ₂ Cl ₂	1.0×10^8					
2.145	1,3-Cyclopentadiene, 1-(4-chlorophenyl)-4-(4-methoxyphenyl)-				CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	CH ₂ Cl ₂	8.8×10^7					
2.146	1,3-Cyclopentadiene, 1-(4-chlorophenyl)-4-phenyl-				CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	CH ₂ Cl ₂	6.8×10^7					
2.147	1,3-Cyclopentadiene, 1,4-diphenyl-				CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	CH ₂ Cl ₂	8.7×10^7					
2.148	1,3-Cyclopentadiene, 1-(4-methoxyphenyl)-4-phenyl-				CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	CH ₂ Cl ₂	1.1×10^8					
2.149	1,3-Cyclopentadiene, 5-(1-methylethylidene)-(6,6-Dimethylfulvene)				CP/Oc-15	S = RB; $E_a = 1.3$ kJ mol ⁻¹ .	68F288
	MeOH	6.0×10^{-3}		293			

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.150	1,3-Cyclopentadiene, 5-(1-methylethylidene)-, endoperoxide						
	MeOH	~2.5		293	CP/Oc-15	$S = RB; E_a = 14 \text{ kJ mol}^{-1}$.	68F288
2.151	Cyclopentane, (dicyclopropylmethylidene)-						
	CH ₃ COCH ₃				CP/Ac,A'c-17	$S = Eos; A' =$ (Dicyclopropylmethylidene)cyclobutane; meas. $k_t/k_{t'}^{A'} = 0.90$.	78F430
2.152	Cyclopentane, ethylidene-						
	CHCl ₃	8.7×10^6			CP/A'c-33	$S = A' = Rub; \text{used } k_d = 1.7 \times 10^4 \text{ s}^{-1}, k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	81F445 78A005
2.153	Cyclopentane, methylene-						
	CH ₃ CN			273	CP/Ac,A'c-17	$S = MB; A' = Methylene cyclohexane; \text{meas.}$ $k_t/k_{t'}^{A'} = 1.5 \times 10^2$.	81F018
	CH ₃ CN			273	CP/Ac,A'c-17	$S = MB; A' = 5\text{-Methylene-2-norbornene}; \text{meas.}$ $k_t/k_{t'}^{A'} = 16$.	78F149
	CH ₃ CN			?		$S = MB; A' = c\text{-C}_6\text{H}_9\text{CH}_3; \text{meas. } k_A/k_{A'} = 0.23$.	73F664
2.154	Cyclopentanone, 2-cyclopentylidene-						
		0.06			CP/Ac-18	$S = \text{Poly-RB or RB; } A' = \text{Linalool; used } \beta_{A'} = 0.18 \text{ mol L}^{-1}; \text{Solvent is CH}_2\text{Cl}_2 \text{ or MeOH.}$	80F111
2.155	Cyclopentanone, 2-(1-methylethenyl)-						
		0.25			CP/Ac-18	$S = \text{Poly-RB or RB; } A' = \text{Linalool; used } \beta_{A'} = 0.18 \text{ mol L}^{-1}; \text{Solvent is CH}_2\text{Cl}_2 \text{ or MeOH.}$	80F111
2.156	Cyclopentene						
	CS ₂	5.8×10^4		298	PL/Ld-2	$S = TPP; \Delta H^\ddagger = 5 \text{ kJ mol}^{-1}; \Delta S^\ddagger = -134 \text{ J K}^{-1}$ $\text{mol}^{-1}; \text{studied at } 183\text{-}310 \text{ K.}$	85A167
	MeOH	2.2×10^5		286	?	$S = \text{Poly-RB; method not reported.}$	81F534
	MeOH			288	CP/Ac,A'c-17	$S = MB; A' = 1,6\text{-Dimethylcyclohexene}; \text{meas.}$ $k_t/k_{t'}^{A'} = 0.33$.	65F028
2.157	Cyclopentene, 1- <i>tert</i> -butyl-						
	CH ₃ CN			273	CP/Ac,A'c-17	$S = MB; A' = 1\text{-}tert\text{-Butylcyclohexene}; \text{meas.}$ $k_t/k_{t'}^{A'} = 14$.	81F018
2.158	Cyclopentene, 1,2-dimethyl-						
	MeOH	9.5×10^6		286	?	$S = \text{Poly-RB; method not reported.}$	81F534
2.159	Cyclopentene, 1,5-dimethyl-						
	CHCl ₃	1.5×10^6			CP/A'c-33	$S = A' = Rub; \text{used } k_d = 1.7 \times 10^4 \text{ s}^{-1}, k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	81F445
2.160	Cyclopentene, 1-methoxy-						
	c-C ₆ H ₁₂	2.4×10^6			PL/A'd-8	$S = An; A' = DPBF; \Delta H^\ddagger = -3 \text{ kJ mol}^{-1}; \Delta S^\ddagger = -134 \text{ J K}^{-1} \text{ mol}^{-1}; \text{studied at } 290\text{-}340 \text{ K.}$	82A349
	C ₆ H ₅ CH ₃	1.2×10^7			PL/A'd-8	$S = An; A' = DPBF; \Delta H^\ddagger = -4 \text{ kJ mol}^{-1}; \Delta S^\ddagger = -125 \text{ J K}^{-1} \text{ mol}^{-1}; \text{studied at } 250\text{-}340 \text{ K.}$	82A349
	CH ₃ CN	2.5×10^7			PL/A'd-8	$S = An; A' = DPBF; \Delta H^\ddagger = -2 \text{ kJ mol}^{-1}; \Delta S^\ddagger = -112 \text{ J K}^{-1} \text{ mol}^{-1}; \text{studied at } 250\text{-}340 \text{ K.}$	82A349
	CH ₃ COCH ₃	1.6×10^7			PL/A'd-8	$S = An; A' = DPBF; \Delta H^\ddagger = -3 \text{ kJ mol}^{-1}; \Delta S^\ddagger = -119 \text{ J K}^{-1} \text{ mol}^{-1}; \text{studied at } 240\text{-}320 \text{ K.}$	82A349
	MeOH	1.3×10^7			PL/A'd-8	$S = An; A' = DPBF; \Delta H^\ddagger = -1 \text{ kJ mol}^{-1}; \Delta S^\ddagger = -114 \text{ J K}^{-1} \text{ mol}^{-1}; \text{studied at } 250\text{-}310 \text{ K.}$	82A349
2.161	Cyclopentene, 1-methyl-						
	C ₆ H ₆	0.018		298	CP/A'c-23	$S = A' = DMA; \text{used } \beta_{A'} = 3.0 \times 10^{-4} \text{ mol L}^{-1}$.	706079

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.161 Cyclopentene, 1-methyl- — Continued							
	C ₆ H ₆	0.018		298	CP/A'c-23	S = A' = Rub; used $\beta_{A'} = 3.0 \times 10^{-4}$ mol L ⁻¹ .	70F079
	C ₆ H ₆	0.016		298	CP/A'c-23	S = A' = DMBA; used $\beta_{A'} = 7.1 \times 10^{-4}$ mol L ⁻¹ .	70F079
	C ₆ H ₆	0.018		298	CP/A'c-23	S = A' = Tetr; used $\beta_{A'} = 1.7 \times 10^{-3}$ mol L ⁻¹ .	70F079
	CH ₃ CN			273	CP/Ac,A'c-17	S = MB; A' = 1- <i>tert</i> -Butylcyclohexene; meas. $k_r/k_r^{A'} = 17$.	81F018
	CH ₃ CN		?			S = MB; A' = <i>c</i> -C ₆ H ₉ CH ₃ ; meas. $k_r/k_r^{A'} = 7.7$.	73F664
	CHCl ₃	2.7×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_r = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445 78A005
	MeOH	9.2×10^5		286	?	S = Poly-RB; method not reported.	81F534
	MeOH			288	CP/Ac,A'c-17	S = MB; A' = 1,2-Dimethylcyclohexene; meas. $k_r/k_r^{A'} = 0.13$.	65F028
	MeOH/ <i>tert</i> -BuOH (50:50)				CR/Ac,A'c-17	A' = 2M2B; meas. $k_r/k_r^{A'} = 0.63$; ¹ O ₂ * from H ₂ O ₂ /Ca(OCl) ₂ .	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)				CP/Ac,A'c-17	S = RB; A' = 2M2B; meas. $k_r/k_r^{A'} = 1.0$.	68F292
2.162 2-Cyclopenten-1-one, 3-methyl-2-pentyl-							
		$>2 \times 10^2$			CP/Ac-18	S = Poly-RB or RB; A' = Linalool; used $\beta_{A'} = 0.18$ mol L ⁻¹ ; Solvent is CH ₂ Cl ₂ or MeOH.	80F111
2.163 2-Cyclopenten-1-one, 2-acetoxy-3-methyl-							
	CD ₃ COCD ₃	2.9×10^4			PL/Ld-2	S = Ac.	89F079
2.164 2-Cyclopenten-1-one, 3-acetoxy-2-methyl-							
	CD ₃ COCD ₃	3.6×10^4			PL/Ld-2	S = Ac.	89F079
2.165 2-Cyclopenten-1-one, 2,3-dimethyl-							
	CD ₃ COCD ₃	7.8×10^4			PL/Ld-2	S = Ac.	89F079
2.166 2-Cyclopenten-1-one, 2-ethyl-3-methyl-							
	CD ₃ COCD ₃	1.8×10^5			PL/Ld-2	S = Ac.	89F079
2.167 2-Cyclopenten-1-one, 3-methoxy-2-methyl-							
	CD ₃ COCD ₃	7.5×10^4			PL/Ld-2	S = Ac.	89F079
2.168 Cyclopropane, (dicyclopropylmethylenide)-							
	CH ₃ CN			293	CP/Ac-17	S = RB; A' = TME; meas. $k_r/k_r^{A'} = 2.1 \times 10^{-3}$; c.k. with 1-methylcyclohexene.	79F119
	CH ₃ COCH ₃				CP/Ac,A'c-17	S = Eos; A' = (Dicyclopropylmethylenide)cyclobutane; meas. $k_r/k_r^{A'} = 1.0$.	78F430
2.169 Cyclopropane, 1,1',1''-(1-ethenyl-2-ylidene)tris-							
	CH ₃ COCH ₃				CP/Ac,A'c-17	S = Eos; A' = (Dicyclopropylmethylenide)cyclobutane; meas. $k_r/k_r^{A'} = 0.40$.	78F430
2.170 Cyclopropane, (1-methylethylidene)-							
	CH ₃ CN			293	CP/Ac-17	S = RB; A' = TME; meas. $k_r/k_r^{A'} = 5.6 \times 10^{-3}$; c.k. with 1-methylcyclohexene.	79F119
2.171 1,6-Decadiene, 2,6,9-trimethyl-, (E)-							
	EtOH/ 2-PrOH (50:50)	0.080 (β_r)		292	CP/Oc-14,27	S = RB; A' = 2,5-DMF.	78F464

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.172	1,6-Decadiene, 2,6,9-trimethyl-, (Z)-EtOH/ 2-PrOH (50:50)	0.11 (β_p)		292	CP/Oc-14,27	S = RB; A' = 2,5-DMF.	78F464
2.173	8,8'-Diapo- ψ -carotenedioic acid, bis(6-O- β -D-glucopyranosyl- β -D-glucopyranosyl) ester (Crocin) CHCl ₃ / EtOH (50:50)	3×10^8		310	CR/LI-12	${}^1\text{O}_2^*$ from NDPO ₂ ; reported $k = 1.1 \times 10^9$ L mol ⁻¹ s ⁻¹ using $k_d = 1 \times 10^5$ s ⁻¹ , recalcd. using $k_d = 3.0 \times 10^4$ s ⁻¹ [92E654].	89R188 90E622
	H ₂ O pH = 7.4	5.6×10^7 (k_p) 1.8×10^9 (k_q)		308	CR/Ac-?	used $k_d = 3.3 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	87A318
2.174	6,6'-Diapo- Ψ,Ψ -carotenedioic acid (<i>trans</i> -Norbixin) DMF	8×10^9		308	CR/A'c-16	A' = DPBF; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMF	8×10^5 (k_p) 3×10^9 (k_q)		308	CR/A'c-?	used $k_d = 1.4 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMSO	4×10^9		308	CR/A'c-16	A' = DPBF; used $k_d = 5.2 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMSO	9×10^5 (k_p) 3×10^9 (k_q)		308	CR/A'c-?	used $k_d = 5.2 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
2.175	6,6'-Diapo- Ψ,Ψ -carotenedioic acid, dimethyl ester (<i>trans</i> -Methylbixin) DMF	9×10^9		308	CR/A'c-16	A' = DPBF; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMF	2×10^6 (k_p) 6×10^9 (k_q)		308	CR/A'c-?	used $k_d = 1.4 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMSO	4×10^9		308	CR/A'c-16	A' = DPBF; used $k_d = 5.2 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMSO	8×10^5 (k_p) 4×10^9 (k_q)		308	CR/A'c-?	used $k_d = 5.2 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
2.176	6,6'-Diapo- Ψ,Ψ -carotenedioic acid, dimethyl ester, 9-cis- (Methylbixin) DMF	1.6×10^{10}		308	CR/A'c-16	A' = DPBF; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMF	2×10^6 (k_p) 3×10^{10} (k_q)		308	CR/A'c-?	used $k_d = 1.4 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMSO	2.2×10^{10}		308	CR/A'c-16	A' = DPBF; used $k_d = 5.2 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMSO	1×10^6 (k_p) 1×10^{10} (k_q)		308	CR/A'c-?	used $k_d = 5.2 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
2.177	6,6'-Diapo- Ψ,Ψ -carotenedioic acid, monomethyl ester, 9-cis- (Bixin) CHCl ₃	9.2×10^9		PL/Ld-2	S = Pz.		91E465
	CHCl ₃ / EtOH (50:50)	4×10^9		310	CR/LI-12	${}^1\text{O}_2^*$ from NDPO ₂ ; reported $k = 1.4 \times 10^{10}$ L mol ⁻¹ s ⁻¹ using $k_d = 1 \times 10^5$ s ⁻¹ , recalcd. using $k_d = 3.0 \times 10^4$ s ⁻¹ [92E654].	89R188 90E622
	DMF	2.0×10^{10}		308	CR/A'c-16	A' = DPBF; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMF	2×10^6 (k_p) 2×10^{10} (k_q)		308	CR/A'c-?	used $k_d = 1.4 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMSO	1.5×10^{10}		308	CR/A'c-16	A' = DPBF; used $k_d = 5.2 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMSO	1×10^6 (k_p) 2×10^{10} (k_q)		308	CR/A'c-?	used $k_d = 5.2 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437
	H ₂ O pH = 7.8	9×10^7 (k_p) 2×10^{10} (k_q)		308	CR/Ac-?	used $k_d = 2.4 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	90A437

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.177 6,6'-Diapo-Ψ,Ψ-carotenedioic acid, monomethyl ester, 9-cis- (Bixin) — Continued							
	H ₂ O (mic) pH = 7.8	7×10^9		308	CR/A'c-16	A' = DPBF; used $k_d = 2.4 \times 10^5$ s ⁻¹ ; 0.1 mol L ⁻¹ SDS; $^1\text{O}_2^*$ from MNPO ₂ .	90A437
	H ₂ O (mic) pH = 7.8	$7 \times 10^7 (k_r)$ $1 \times 10^{10} (k_q)$		308	CR/A'c-?	used $k_d = 2.4 \times 10^5$ s ⁻¹ ; 0.1 mol L ⁻¹ SDS; $^1\text{O}_2^*$ from MNPO ₂ .	90A437
2.178 8,8'-Diapocarotenedioic acid, mono(6-O-β-D-glucopyranosyl-β-D-glucopyranosyl) ester (emi-Crocin)							
	H ₂ O pH = 7.4	$1.3 \times 10^8 (k_r)$ $5.1 \times 10^9 (k_q)$		308	CR/Ac-?	used $k_d = 3.3 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	87A318
2.179 9-cis-6,6'-Diapo-Ψ,Ψ-carotenedioic acid (Norbixin)							
	DMF	1.7×10^{10}		308	CR/A'c-16	A' = DPBF; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMF	$2 \times 10^6 (k_r)$ $1 \times 10^{10} (k_q)$		308	CR/A'c-?	used $k_d = 1.4 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMSO	9×10^9		308	CR/A'c-16	A' = DPBF; used $k_d = 5.2 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	90A437
	DMSO	$2 \times 10^6 (k_r)$ $2 \times 10^{10} (k_q)$		308	CR/A'c-?	used $k_d = 5.2 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	90A437
	H ₂ O pH = 7.8	$2 \times 10^8 (k_r)$ $2 \times 10^{10} (k_q)$		308	CR/Ac-?	used $k_d = 2.4 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	90A437
	H ₂ O (mic) pH = 7.8	6×10^9		308	CR/Ac-16	A' = DPBF; used $k_d = 2.4 \times 10^5$ s ⁻¹ ; 0.1 mol L ⁻¹ SDS; $^1\text{O}_2^*$ from MNPO ₂ .	90A437
	H ₂ O (mic) pH = 7.8	$8 \times 10^7 (k_r)$ $9 \times 10^9 (k_q)$		308	CR/Ac-?	used $k_d = 2.4 \times 10^5$ s ⁻¹ ; 0.1 mol L ⁻¹ SDS; $^1\text{O}_2^*$ from MNPO ₂ .	90A437
2.180 4a,9a:9,10-Diethenoanthracene, 1,4,9,10-tetrahydro-5,8-dimethoxy-							
	CH ₂ Cl ₂	2.5×10^4			PR/A'd-8	S = An; A' = DPBF.	79A457
2.181 4a,9a:9,10-Diethenoanthracene, 1,4,9,10-tetrahydro-							
	CH ₂ Cl ₂	1.0×10^6			PR/A'd-8	S = An; A' = DPBF.	79A457
2.182 2,3-Dioxabicyclo[2.2.2]oct-5-ene (3,6-Endoperoxycyclohexene)							
	MeOH	1.5×10^2		293	CP/Oc-15	S = RB; $E_a = 26$ kJ mol ⁻¹ .	68F288
2.183 1,4-Dioxene							
	CH ₃ COCH ₃	2.2×10^5	0.18	288	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	77F876
	CH ₃ COCH ₃	3.6×10^5		281	?	S = ?; A' = Car; used $k_{A'} = 3 \times 10^{10}$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 1.2 \times 10^{-5}$.	70F733
2.184 1,4-Dioxene, 2-(4-cyanophenyl)-3-(4-dimethylaminophenyl)-							
	CD ₃ COCD ₃	$2.3 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007
2.185 1,4-Dioxene, 2,3-di(4-acetaminophenyl)-							
	CD ₃ COCD ₃ / CD ₃ SOCD ₃ (95:5)	$1.5 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007
2.186 1,4-Dioxene, 2,3-di(3-chlorophenyl)-							
	CH ₃ COCH ₃	$3.6 \times 10^6 (k_r)$			CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007
2.187 1,4-Dioxene, 2,3-di(4-chlorophenyl)-							
	CH ₃ CN	$1.9 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 1.1 \times 10^8$ L mol ⁻¹ s ⁻¹ .	80F007
	CH ₃ COCH ₃	$7.5 \times 10^6 (k_r)$			CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.	
188	1,4-Dioxene, 2,3-di(4-dimethylaminophenyl)-							
	CD ₃ COCD ₃	$3.7 \times 10^8 (k_r)$			CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007	
189	1,4-Dioxene, 2,3-di(4-hydroxyphenyl)-				CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007	
190	1,4-Dioxene, 2,3-di(3-methoxyphenyl)-				CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007	
191	1,4-Dioxene, 2,3-di(4-methoxyphenyl)-				CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 1.1 \times 10^8$ L mol ⁻¹ s ⁻¹ .	80F007	
	CH ₃ CN	$1.3 \times 10^8 (k_r)$			CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007	
	CH ₃ COCH ₃	$4.8 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007	
192	1,4-Dioxene, 2-(4-dimethylaminophenyl)-3-phenyl-				CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007	
	CD ₃ COCD ₃	$7.5 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007	
193	1,4-Dioxene, 2,3-di(4-methylphenyl)-				CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007	
	CH ₃ CN	$2.9 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007	
194	1,4-Dioxene, 2,3-diphenyl-				298	PL/Ld-2	S = DCA.	91A311
	CH ₃ CN	1.7×10^7			CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007	
	CH ₃ COCH ₃	$1.6 \times 10^7 (k_r)$			281	?	S = ?; A' = 1,4-Dioxene; used $k_{A'} = 2.2 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 58$.	70F733
195	1,4-Dioxene, 2-(4-methoxyphenyl)-3-phenyl-				CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007	
	CH ₃ COCH ₃	$2.7 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = TPP or RB; A' = TME; used $k_r^{A'} = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80F007	
196	1,4-Dioxene, 2,3,5,6-tetraphenyl-				CP/Pa-17	S = A; A' = TME; used $k_r^{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ ; P = Benzil.	79A241	
	C ₆ H ₆	$1.5 \times 10^7 (k_r)$			CP/A'c-16	S = MPDME; A' = DPBF; used $k_d = 1 \times 10^4$ s ⁻¹ .	87A035	
197	1,3-Dioxolane, 4,5-diethylidene-2,2-dimethyl-, (E,Z)				CP/A'c-16	S = MPDME; A' = DPBF; used $k_d = 1 \times 10^4$ s ⁻¹ .	87A035	
	CH ₂ Cl ₂	2.3×10^7			CP/A'c-16	S = MPDME; A' = DPBF; used $k_d = 1 \times 10^4$ s ⁻¹ .	87A035	
198	1,3-Dioxolane, 4,5-diethylidene-2,2-dimethyl-, (Z,Z)				CP/A'c-16	S = MPDME; A' = DPBF; used $k_d = 1 \times 10^4$ s ⁻¹ .	87A035	
199	1,3-Dioxole				288	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	77F876
200	Docosahexaenoic acid				MP/LI-12	S = TPP; used $k_d = 42$ s ⁻¹ .	83E425 83F151	
201	4,8-Dodecadiene, 4,8-dimethyl-				253	CP/Oc-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	78A344
	MeOH	$6.9 \times 10^5 (k_r)$						

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.202	4,8-Dodecadiene, 4,8-dimethyl-, (E,E)- MeOH	1.1×10^6 (k_r)		253	CP/Oc-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	78A344
2.203	4,8-Dodecadiene, hydroperoxy-4,8-dimethyl- MeOH	3.8×10^5 (k_r)		253	CP/Oc,Ac,Pa-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	78A344
2.204	(E,E)-8,10-Dodecadienol $C_6H_5Cl/2\text{-PrOH}$ (90:10)	5.7×10^5		273	CR/LI-12	used $k_d = 4.1 \times 10^4$ s ⁻¹ ; $^1O_2^*$ from hydrotrioxide, e.g. $(CH_3)_2C(OH)OOOH$.	90M125
2.205	2,4,6,8,10-Dodecapentaene-1,12-dione, 1,12-bis(3-hydroxy-1,2,2-trimethylcyclopentyl)-4,7-dimethyl- [1R-[1 α [2E,4E,6E,8E,10E,12(1R*,3R*)],3 α]- (C ₃₀ -Capsorubin) CHCl ₃	$<1 \times 10^7$			CR/LI-12	used $k_d = 3.0 \times 10^4$ s ⁻¹ ; $^1O_2^*$ from NDPO ₂ .	92R054
2.206	2,4,6,8,10-Dodecapentaene-1,12-dione, 1,12-bis(3-hydroxy-1,2,2-trimethylcyclopentyl)-4,7-dimethyl- [1R-[1 α [2E,4E,6E,8E,10E,12(1R*,3R*)],3 β]- (C ₃₀ -Episocapsorubin) CHCl ₃	$<1 \times 10^7$			CR/LI-12	used $k_d = 3.0 \times 10^4$ s ⁻¹ ; $^1O_2^*$ from NDPO ₂ .	92R054
2.207	2,4,6,8,10-Dodecapentaene-1,12-dione, 1,12-bis(4-hydroxy-1,2,2-trimethylcyclopentyl)-4,7-dimethyl- [1R-[1 α [2E,4E,6E,8E,10E,12(1R*,4R*)],4 β]- (C ₃₀ -Isocapsorubin) CHCl ₃	$<1 \times 10^7$			CR/LI-12	used $k_d = 3.0 \times 10^4$ s ⁻¹ ; $^1O_2^*$ from NDPO ₂ .	92R054
2.208	2,6,10-Dodecatriene, dihydroperoxy-2,6,10-trimethyl- MeOH	6×10^5 (k_r)		253	CP/Oc,Pa-17	S = RB; A' = TMDT; used $k_r^{A'} = 2.0 \times 10^6$ L mol ⁻¹ s ⁻¹ .	83A078
2.209	2,6,10-Dodecatriene, hydroperoxy-2,6,10-trimethyl- MeOH	1.2×10^6 (k_r)		253	CP/Oc,Pa-17	S = RB; A' = TMDT; used $k_r^{A'} = 2.0 \times 10^6$ L mol ⁻¹ s ⁻¹ .	83A078
2.210	2,6,10-Dodecatriene, 2,6,10-trimethyl- (TMDT) MeOH	2.0×10^6	0.055	253	CP/Oc,Pa-15	S = RB; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	83A078
2.211	4,6,8,10,12,14,16-Eicosahptaene-3,18-dione, 8,13-dimethyl-2,2,19,19-tetramethoxy- (<i>all</i> -E) (C ₂₂ -Polyene-tetrone-diacetal) CHCl ₃	-2×10^9			CR/LI-12	used $k_d = 3.0 \times 10^4$ s ⁻¹ ; $^1O_2^*$ from NDPO ₂ .	92R054
2.212	5,8,11,14-Eicosatetraenoic acid (Arachidonic acid) CCl ₄	1×10^5			MP/LI-12	S = TPP; used $k_d = 42$ s ⁻¹ .	83E425 83F151
2.213	5,8,11,14-Eicosatetraenoic acid, methyl ester (Methyl arachidonate) C ₅ H ₅ N	2.4×10^5 (k_r)			CP/Ac,A'c-17	S = PP; A' = Cholesterol; used $k_d = 6 \times 10^4$ s ⁻¹ ; meas. $k_r/k_r^{A'} = 3.5$, $\beta_{A'} = 0.89$ mol L ⁻¹ .	743115
2.214	5,8,11,14-Eicosatetraenoic acid, phenyl ester (Phenyl arachidonate) CD ₃ OD	1.2×10^5			PL/Ld-2	S = MB and RF.	88A165
2.215	Ethanol, 2-cyclododecylidene- MeOH/ H ₂ O (95:5)	0.34			CP/Oc-15	S = RB; Soln. contg. 1% sodium acetate.	79F137
2.216	Ethanol, 2-cyclohexylidene- MeOH/ H ₂ O (95:5)	0.43			CP/Oc-15	S = RB; Soln. contg. 1% sodium acetate.	79F137
2.217	Ethanol, 2-cyclooctylidene- MeOH/ H ₂ O (95:5)	0.06			CP/Oc-15	S = RB; Soln. contg. 1% sodium acetate.	79F137
2.218	Ethanol, 2-cyclopentylidene- MeOH/ H ₂ O (95:5)	0.02			CP/Oc-15	S = RB; Soln. contg. 1% sodium acetate.	79F137

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.219	Ethanol, 1-(2-methylene-3,5-cyclohexadienylidene)- (2-Methylacetophenone enol)						
	CH ₃ CN	$\sim 10^9$			PL/Ld	S = OMAPB; estd. from obs. lifetimes for different laser doses.	89A241
2.220	Ethanol, 1-tricyclo[3.3.1.1 ^{3,7}]decylidene-, acetate (1-Adamantylideneethyl acetate)						
	CD ₃ COCD ₃	4.5×10^4		296	PL/Ld-2	S = MB.	86F219
2.221	Ethene, 1,1-diethoxy-						
	C ₆ H ₆	2.6×10^5	0.15	288	CP/A'c-16	S = RBCE; A' = DPBF; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	77F876
	CH ₃ COCH ₃	$<1 \times 10^4$ (k_r)		279	CP/Ac,A'c-17	S = RB; A' = 1,4-Dioxene; No measurable reaction of A.	77F876
	CH ₃ COCH ₃	4.6×10^5	0.083	288	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	77F876
2.222	Ethene, 1,2-diethoxy-, (E)-						
	CH ₃ COCH ₃	2.6×10^7 (k_r)		279	CP/Ac,A'c-17	S = RB; A' = (Z)-1,2-Diethoxyethene; used $k_r^{A'} = 5.7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.46$.	77F876
	CH ₃ COCH ₃	4.7×10^7	8.1×10^{-4}	288	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	77F876
	CH ₃ COCH ₃	1.0×10^7 (k_r)		281	?	S = ?; A' = 1,4-Dioxene; used $k_{A'} = 2.2 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 46$.	70F733
2.223	Ethene, 1,2-diethoxy-, (Z)-						
	CH ₃ COCH ₃	4.4×10^7	8.6×10^{-4}	288	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	77F876
	CH ₃ COCH ₃	5.7×10^7 (k_r)		279	CP/Ac,A'c-17	S = RB; A' = TME; used $k_r^{A'} = 5.4 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 1.1$; Found $k_r \gg k_q$.	77F876
	CH ₃ COCH ₃	3.3×10^7 (k_r)		281	?	S = ?; A' = 1,4-Dioxene; used $k_{A'} = 2.2 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 1.5 \times 10^2$.	70F733
2.224	Ethene, ethoxy-						
	C ₆ H ₆	2.3×10^5	0.18	288	CP/A'c-16	S = RBCE; A' = DPBF; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	77F876
	CH ₃ COCH ₃	3.1×10^4 (k_r)		279	CP/Ac,A'c-17	S = RB; A' = 1,4-Dioxene; used $k_r^{A'} = 2.2 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.14$; Found $k_r \gg k_q$.	77F876
	CH ₃ COCH ₃	4.3×10^4	0.88	288	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	77F876
	CH ₃ COCH ₃	5.7×10^4 (k_r)		281	?	S = ?; A' = 1,4-Dioxene; used $k_{A'} = 2.2 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 0.26$.	70F733
2.225	Ethene, tetraethoxy-						
	C ₆ H ₆	7.4×10^7	5.7×10^{-4}	288	CP/A'c-16	S = RBCE; A' = DPBF; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	77F876
	CH ₃ CN	4.3×10^7	7.7×10^{-4}	288	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	77F876
	CH ₃ COCH ₃	4.6×10^7 (k_r)		279	CP/Ac,A'c-17	S = RB; A' = TME; used $k_r^{A'} = 5.4 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.85$; Found $k_r \gg k_q$.	77F876
	CH ₃ COCH ₃	4.5×10^7	8.5×10^{-4}	288	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	77F876
	CHCl ₃	7.6×10^7	2.2×10^{-4}	288	CP/A'c-16	S = MB; A' = DPBF; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; Solvent contained 1% EtOII.	77F876
2.226	Ethene, 1,1,2-triethoxy-						
	CH ₃ COCH ₃	1.2×10^8	3.1×10^{-4}	288	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	77F876
2.227	2-Furanone, 3-(1-methylethylidene)-		0.65		CP/Ac-18	S = Poly-RB or RB; A' = Linalool; used $\beta_{A'} = 0.18$ mol L ⁻¹ ; Solvent is CH ₂ Cl ₂ or MeOH.	80F111
2.228	1-Heptene						
	CCl ₄ / MeOH (96:4)	2×10^3			PL/A'd-8	S = MB; A' = DPBF.	77F162
2.229	(Z)-2-Heptene						
	MeOH	1.3×10^5		286	?	S = Poly-RB; method not reported.	81F534

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k ($\text{L mol}^{-1} \text{s}^{-1}$)	$\beta (k_d/k)$ (mol L^{-1})	T (K)	Method	Comment	Ref.
2.230	(all-E)-1,3,5,7,9,11,13,15,17,19,21,23,25-Hexacosatridecaene, 3,7,11,16,20,24-hexamethyl-1,26-bis(2,6,6-trimethyl-1-cyclohexyl- (Decapreno- β -carotene)						
	$\text{C}_6\text{H}_5\text{CH}_3$	2.1×10^{10}			PL/Ld-2	$S = Pz.$	91E465
	C_6H_6	2.0×10^{10}			PL/Ld-2	$S = Pz.$	91E465
	C_6H_6	2.2×10^9			PL/ β Cd-2	$S = An.$	80A143
2.231	2,4,6,8,10,12,14-Hexadecaheptaene-1,16-dione, 1,16-bis(3-hydroxy-1,2,2-trimethylcyclopentyl)-6,11-dimethyl- [1R-[1 α][2E,4E,6E,8E,10E,12E,14E,16(1R*,3S*)],3 β]- (C_{34} -Capsorubin)						
	CHCl_3	$\sim 2 \times 10^9$			CR/LI-12	used $k_d = 3.0 \times 10^4 \text{ s}^{-1}$, ${}^1\text{O}_2^*$ from NDPO ₂ .	92R054
2.232	2,4,6,8,10,12,14-Hexadecaheptaene-1,16-dione, 1,16-bis(4-hydroxy-1,2,2-trimethylcyclopentyl)-6,11-dimethyl- [1R-[1 α][2E,4E,6E,8E,10E,12E,14E,16(1R*,4S*)],4 β]- (C_{34} -Epiisocapsorubin)				CR/LI-12	used $k_d = 3.0 \times 10^4 \text{ s}^{-1}$, ${}^1\text{O}_2^*$ from NDPO ₂ .	92R054
2.233	4,6,8,10,12-Hexadecapentaene-3,14-dione, 6,11-dimethyl-2,2,15,15-tetramethoxy- (all-E) (C_{18} -Polyene-tetrone-diacetal)				CR/LI-12	used $k_d = 3.0 \times 10^4 \text{ s}^{-1}$, ${}^1\text{O}_2^*$ from NDPO ₂ .	92R054
	CHCl_3	$<1 \times 10^7$					
2.234	4,6,8,10,12-Hexadecapentaene-2,3,14,15-tetraone, 6,11-dimethyl- (all-E) (C_{18} -Polyene-tetrone)				CR/LI-12	used $k_d = 3.0 \times 10^4 \text{ s}^{-1}$, ${}^1\text{O}_2^*$ from NDPO ₂ .	92R054
	CHCl_3	$\sim 1 \times 10^7$					
2.235	2,6,10,14-Hexadecatetraene, 2,6,10,14-tetramethyl-						
	MeOH	2.2×10^6	0.05	253	CP/Oc,Pa-15	$S = RB$; used $k_d = 1.1 \times 10^5 \text{ s}^{-1}$.	83A078
2.236	2-Hexadecene, 1-ethoxy-3,7,11,15-tetramethyl- (Ethyl phytol ether)						
	EtOH		3.0 (β_r)		CP/Pa,P'a-17	$S = RB$; $A' = c\text{-C}_6\text{H}_9\text{CH}_3$; used $\beta_r^{A'} = 1.5 \times 10^{-3} \text{ mol L}^{-1}$.	86F091
2.237	2-Hexadecene, 1-methoxy-3,7,11,15-tetramethyl- (Methyl phytol ether)						
	EtOH		3.7 (β_r)		CP/Pa,P'a-17	$S = RB$; $A' = c\text{-C}_6\text{H}_9\text{CH}_3$; used $\beta_r^{A'} = 1.5 \times 10^{-3} \text{ mol L}^{-1}$.	86F091
2.238	2-Hexadecen-1-ol, 3,7,11,15-tetramethyl- (Phytol)						
	EtOH		1.3 (β_r)		CP/Pa,P'a-17	$S = RB$; $A' = c\text{-C}_6\text{H}_9\text{CH}_3$; used $\beta_r^{A'} = 1.5 \times 10^{-3} \text{ mol L}^{-1}$.	86F091
2.239	2-Hexadecen-1-ol, 3,7,11,15-tetramethyl-, acetate (Phytol acetate)						
	EtOH		7.8 (β_r)		CP/Pa,P'a-17	$S = RB$; $A' = c\text{-C}_6\text{H}_9\text{CH}_3$; used $\beta_r^{A'} = 1.5 \times 10^{-3} \text{ mol L}^{-1}$.	86F091
2.240	1,5-Hexadiene						
	CCl_4/MeOH (96:4)	2×10^3			PL/A'd-8	$S = MB$; $A' = DPBF$.	777162
2.241	(E,E)-2,4-Hexadiene						
	CCl_4/MeOH (96:4)	2×10^4			PL/A'd-8	$S = MB$; $A' = DPBF$.	777162
	CHCl_3	2.4×10^5			CP/A'c-33	$S = A' = Rub$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	81F445
2.242	(E,Z)-2,4-Hexadiene						
	CHCl_3	1.0×10^5			CP/A'c-33	$S = A' = Rub$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	81F445
2.243	2,4-Hexadiene, 2,5-dimethyl- (DMHD)						
	C_6H_6	$1.8 \times 10^5 (k_r)$ $1.8 \times 10^6 (k_q)$	0.24 (β_r)	286	CP/Oc-29	$S = TPP$; $A' = 2,5\text{-DMF}$; used $k_d = 4.2 \times 10^4 \text{ s}^{-1}$; meas. $k_q/k_r = 9.74$: for formation of products (2) 3,3-dimethyl-2-(2-methyl-1-propenyl)-1,2-dioxetane, (3) 2,5-dimethyl-3-hydroperoxy-1,4-hexadiene, (4) 1,2-dioxa-3,3,6,6-tetramethyl-4-cyclohexene, (5) <i>trans</i> -2,5-dimethyl-5-hydroperoxy-1,3-hexadiene, $k_r = 0.05$, 0.59, 0.34 and $0.81 \times 10^5 \text{ L mol}^{-1} \text{ s}^{-1}$, resp.	84F335

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_q/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2,4-Hexadiene, 2,5-dimethyl- (DMHD) — Continued							
	C ₆ H ₆	4×10^6			PL/Ld-2	S = MPDME.	83F195 83F196
	C ₆ H ₆	$<2.5 \times 10^5$ (k_r)		295	CP/Ac-17	S = RB; A' = 2M2P; used $k_r^{A'} = 7.2 \times 10^5$ L mol ⁻¹ s ⁻¹ ; for formation of 3-hydroperoxy-2,5-dimethyl-1,4-diene $k_r = 1.4 \times 10^5$ L mol ⁻¹ s ⁻¹ .	83F196
	CCl ₄	1.0×10^5 (k_r) 9.8×10^5 (k_q)	0.014 (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; used $k_d = 1.4 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r = 9.81$; for formation of products (2) 3,3-dimethyl-2-(2-methyl-1-propenyl)-1,2-dioxetane, (3) 2,5-dimethyl-3-hydroperoxy-1,4-hexadiene, (4) 1,2-dioxa-3,3,6,6-tetramethyl-4-cyclohexene, (5) <i>trans</i> -2,5-dimethyl-5-hydroperoxy-1,3-hexadiene, $k_r = 0.07, 0.25, 0.23$ and 0.42×10^5 L mol ⁻¹ s ⁻¹ , resp.	84F335
	CH ₂ Cl ₂	8.6×10^5 (k_r) 1.9×10^6 (k_q)	0.011 (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; used $k_d = 9.5 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r = 2.21$; for formation of products (2) 3,3-dimethyl-2-(2-methyl-1-propenyl)-1,2-dioxetane, (3) 2,5-dimethyl-3-hydroperoxy-1,4-hexadiene, (4) 1,2-dioxa-3,3,6,6-tetramethyl-4-cyclohexene, (5) <i>trans</i> -2,5-dimethyl-5-hydroperoxy-1,3-hexadiene, $k_r = 0.26, 5.5, 0.34$ and 2.5×10^5 L mol ⁻¹ s ⁻¹ , resp.	84F335
	CH ₂ Cl ₂	5×10^6			PL/Ld-2	S = MPDME.	83F195 83F196
	CH ₂ Cl ₂	1×10^6 (k_r)		295	CP/Ac-17	S = RB; A' = 2M2P; used $k_r^{A'} = 9.7 \times 10^5$ L mol ⁻¹ s ⁻¹ ; for formation of 3-hydroperoxy-2,5-dimethyl-1,4-diene $k_r = 1.3 \times 10^6$ L mol ⁻¹ s ⁻¹ .	83F196
	CH ₂ Cl ₂				CP/Pa,P'a-17	S = ?; A' = <i>c</i> -C ₆ H ₉ CH ₃ ; meas. $k_r/k_r^{A'} = 5.0$.	73F662
	CH ₃ CN	1.4×10^6 (k_r) 1.3×10^6 (k_q)	0.024 (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; used $k_d = 3.3 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r = 0.91$; for formation of products (2) 3,3-dimethyl-2-(2-methyl-1-propenyl)-1,2-dioxetane, (3) 2,5-dimethyl-3-hydroperoxy-1,4-hexadiene, (4) 1,2-dioxa-3,3,6,6-tetramethyl-4-cyclohexene, (5) <i>trans</i> -2,5-dimethyl-5-hydroperoxy-1,3-hexadiene, $k_r = 0.41, 12.3, 0.55$ and 0.55×10^5 L mol ⁻¹ s ⁻¹ , resp.	84F335
	CH ₃ CN	5.2×10^6	0.029		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.5 \times 10^5$ s ⁻¹ .	83F195
	CH ₃ CN	6.3×10^6			PL/Ld-2	S = MPDME.	83F195 83F196
	CH ₃ CN	1.6×10^6 (k_r)		295	CP/Ac-17	S = RB; A' = 2M2P; used $k_r^{A'} = 1.2 \times 10^6$ L mol ⁻¹ s ⁻¹ ; for formation of 3-hydroperoxy-2,5-dimethyl-1,4-diene $k_r = 1.7 \times 10^6$ L mol ⁻¹ s ⁻¹ .	83F196
	CH ₃ CN				CP/Pa,P'a-17	S = ?; A' = <i>c</i> -C ₆ H ₉ CH ₃ ; meas. $k_r/k_r^{A'} = 6.3$.	73F662
	CH ₃ COCH ₃	7.4×10^5 (k_r) 1.1×10^6 (k_q)	0.051 (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; used $k_d = 3.8 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r = 1.44$; for formation of products (2) 3,3-dimethyl-2-(2-methyl-1-propenyl)-1,2-dioxetane, (3) 2,5-dimethyl-3-hydroperoxy-1,4-hexadiene, (4) 1,2-dioxa-3,3,6,6-tetramethyl-4-cyclohexene, (5) <i>trans</i> -2,5-dimethyl-5-hydroperoxy-1,3-hexadiene, $k_r = 0.59, 6.2, 0.22$ and 0.37×10^5 L mol ⁻¹ s ⁻¹ , resp.	84F335
	CH ₃ COCH ₃	3.9×10^6		295	PL/Ld-2	S = MPDME.	83F195 83F196

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.243 2,4-Hexadiene, 2,5-dimethyl- (DMHD) — Continued							
	CH ₃ COCH ₃	7×10^5 (k_r)		295	CP/Ac-17	S = RB; A' = 2M2P; used $k_r^{A'} = 8.1 \times 10^5$ L mol ⁻¹ s ⁻¹ ; $\Delta H^\ddagger = -0.96$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -138$ J K ⁻¹ mol ⁻¹ ; studied at 195-295 K; for formation of 3-hydroperoxy-2,5-dimethyl-1,4-diene $k_r = 3.4, 2.5$ and 3.2×10^5 L mol ⁻¹ s ⁻¹ at 295, 241 and 195 K, resp. and $\Delta H^\ddagger = 1.7$ kJ mol ⁻¹ and $\Delta S^\ddagger = -146$ J K ⁻¹ mol ⁻¹ .	83F196
	CH ₃ COCH ₃				CP/Pa,P'a-17	S = ?; A' = <i>c</i> -C ₆ H ₉ CH ₃ ; meas. $k_r/k_r^{A'} = 3.2$.	73F662
	CH ₃ COCH ₃ /H ₂ O (75:25)				CP/Pa,P'a-17	S = ?; A' = <i>c</i> -C ₆ H ₉ CH ₃ ; meas. $k_r/k_r^{A'} = 13$.	73F662
	CHCl ₃	4.8×10^5 (k_r) 9.0×10^5 (k_q)	8.3×10^{-3} (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; used $k_d = 4.0 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r = 1.86$; for formation of products (2) 3,3-dimethyl-2-(2-methyl-1-propenyl)-1,2-dioxetane, (3) 2,5-dimethyl-3-hydroperoxy-1,4-hexadiene, (4) 1,2-dioxa-3,3,6,6-tetramethyl-4-cyclohexene, (5) <i>trans</i> -2,5-dimethyl-5-hydroperoxy-1,3-hexadiene, $k_r = 0.14, 2.98, 0.29$ and 1.45×10^5 L mol ⁻¹ s ⁻¹ , resp.	84F335
	CHCl ₃	5.3×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
	MeOD	2.6×10^6			PL/Ld-2	S = MPDME.	83F196
	MeOH	2.4×10^6 (k_r) 1.1×10^6 (k_q)	0.059 (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; meas. $k_q/k_r = 0.45$; for formation of products (2) 3,3-dimethyl-2-(2-methyl-1-propenyl)-1,2-dioxetane, (3) 2,5-dimethyl-3-hydroperoxy-1,4-hexadiene, (4) 1,2-dioxa-3,3,6,6-tetramethyl-4-cyclohexene, (5) <i>trans</i> -2,5-dimethyl-5-hydroperoxy-1,3-hexadiene, (6) <i>trans</i> -2,5-dimethyl-1-hydroperoxy-5-methoxy-3-hexene, $k_r = 12.8, 5.1, 0.96, 0.48$ and 4.8×10^5 L mol ⁻¹ s ⁻¹ , resp.	84F335
	MeOH	2.6×10^6	0.042		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	83F195
	MeOH	3×10^6 (k_r)		295	CP/Ac-17	S = RB; A' = 2M2P; used $k_r^{A'} = 2.0 \times 10^5$ L mol ⁻¹ s ⁻¹ ; $\Delta H^\ddagger = 2.7$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -115$ J K ⁻¹ mol ⁻¹ ; studied at 195-295 K; for formation of 3-hydroperoxy-2,5-dimethyl-1,4-diene $k_r = 2.9, 1.0$ and 0.28×10^5 L mol ⁻¹ s ⁻¹ at 295, 241 and 195 K, resp. and $\Delta H^\ddagger = 9.2$ kJ mol ⁻¹ and $\Delta S^\ddagger = -113$ J K ⁻¹ mol ⁻¹ .	83F196
	MeOH	2×10^6			PL/A'd-5	S = MB; A' = DPBF.	727260
	MeOH				CP/Pa,P'a-17	S = ?; A' = <i>c</i> -C ₆ H ₉ CH ₃ ; meas. $k_r/k_r^{A'} = 28$.	73F662
	MeOH/H ₂ O (88:13)				CP/Pa,P'a-17	S = ?; A' = <i>c</i> -C ₆ H ₉ CH ₃ ; meas. $k_r/k_r^{A'} = 29$.	73F662
2.244 2-Hexene							
	MeOH			288	CP/Ac,A'c-17	S = MB; A' = <i>c</i> -C ₅ H ₈ ; meas. $k_r/k_r^{A'} = 0.56$.	65F028
2.245 (E)-3-Hexene							
	CS ₂	7.7×10^3		298	PL/Ld-2	S = TPP; $\Delta H^\ddagger = 2$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -163$ J K ⁻¹ mol ⁻¹ ; studied at 183-310 K.	85A167
2.246 (Z)-3-Hexene							
	CHCl ₃	3.5×10^4 3.1×10^4 (k_r) 4×10^3 (k_q)	0.14		CP/Oc-14,27	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; obs. rate of oxygen consumption as a ratio with TME; calcn. involves $k_q(MB) = 6.4 \times 10^6$ L mol ⁻¹ s ⁻¹ .	88A026
	CS ₂	3.9×10^4		298	PL/Ld-2	S = TPP; $\Delta H^\ddagger = 8$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -130$ J K ⁻¹ mol ⁻¹ ; studied at 183-310 K.	85A167

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.247	Indole, 1-methyl-3-(β -methoxyvinyl)- (<i>E</i>)						
	CD ₂ Cl ₂			195	CP/Pa,P'a-17	S = TPP; A' = <i>E</i> -1-(Phenylsulfonyl)-3-(β -methoxyvinyl)indole; meas. $k_t/k_{t'A'} = 5$.	93F367
2.248	Indole, 1-methyl-3-(β -methoxyvinyl)- (<i>Z</i>)						
	CD ₂ Cl ₂			195	CP/Pa,P'a-17	S = TPP; A' = <i>E</i> -1-Methyl-3-(β -methoxyvinyl)indole; meas. $k_t/k_{t'A'} = 4.1$.	93F367
2.249	Indole, 1-(phenylsulfonyl)-3-(β -methoxyvinyl)- (<i>Z</i>)						
	CD ₂ Cl ₂			195	CP/Pa,P'a-17	S = TPP; A' = <i>E</i> -1-(Phenylsulfonyl)-3-(β -methoxyvinyl)indole; meas. $k_t/k_{t'A'} = 5.2$.	93F367
2.250	Lycopene, (<i>all-E</i>)-						
	<i>n</i> -C ₆ H ₁₄	1.9×10^{10}			PL/Ld-2	S = Pz.	91E465
	C ₆ H ₅ CH ₃	1.8×10^{10}			PL/Ld-2	S = Pz.	91E465
	C ₆ H ₆	1.7×10^{10}			PL/Ld-2	S = Pz.	91E465
	C ₆ H ₆ /MeOH (80:20)				CP/P'a-20	S = MB; A' = 2M2P; meas. $k_A/k_{A'} = 1.5 \times 10^4$.	707188
	CCl ₄	1.4×10^{10}			PL/Ld-2	S = Pz.	91E465
	CH ₂ Cl ₂	6.9×10^9	1.4×10^{-6}	293	CP/Oc-19	S = Chl a; A' = Soybean oil; used $k_d = 1.0 \times 10^4$ s ⁻¹ .	90R128
	CHCl ₃	9×10^9			CR/LI-12	used $k_d = 3.0 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	92R054
	CHCl ₃	1.9×10^{10}			PL/Ld-2	S = Pz.	91E465
	CHCl ₃ /EtOH (50:50)	1.8×10^{10}			PL/Ld-2	S = Pz; Soln. cont. 1% H ₂ O.	91E465
	CHCl ₃ /EtOH (50:50)	9×10^9		310	CR/LI-12	${}^1\text{O}_2^*$ from NDPO ₂ ; reported $k = 3.1 \times 10^{10}$ L mol ⁻¹ s ⁻¹ using $k_d = 1 \times 10^5$ s ⁻¹ , recalcd. using $k_d = 3.0 \times 10^4$ s ⁻¹ [92E654].	89R188 90E622
2.251	Lycopene, dihydroxy-						
	C ₆ H ₆	5.1×10^9			PL/ β Cd-11	S = An.	80A143
2.252	Methacrylic acid						
	C ₆ D ₆	4.7×10^3			PL/Ld-2	S = TPP.	89A331
2.253	2,5-Methano-1,3-dithiolo[4,5- <i>d</i>][1,3]dithiin, dihydro-2,3a,5-trimethyl-7-methylene-						
	CHCl ₃	1.2×10^5		295	CL/LI-12	S = An; used $k_d = 4.9 \times 10^3$ s ⁻¹ .	90F473
2.254	4,7-Methanoindene, 4,5,6,7-tetrahydro- (Isodicyclopentadiene)						
	CD ₃ COCD ₃	1.5×10^8		195	CP/A'c-16	S = RB; A' = DPBF; k_d not given.	87A368
2.255	Methanol, cyclohexylidene-, acetate						
	CD ₃ COCD ₃	7.5×10^4		298	PL/Ld-2	S = MB.	85A167
	CD ₃ OD	4.8×10^4		298	PL/Ld-2	S = MB.	85A167
2.256	Methanol, 1-tricyclo[3.3.1.1 ^{3,7}]decylidene-, acetate (Adamantylidenemethyl acetate)						
	CD ₃ COCD ₃	3.4×10^4		296	PL/Ld-2	S = MB.	86F219 85A167
	CD ₃ COCD ₃	2.7×10^4 (k_t)		296	CP/Pa-17	S = MB; A' = 2M2P; used $k_t/A' = 8.1 \times 10^5$ L mol ⁻¹ s ⁻¹ .	86F219
	CD ₃ OD	2.1×10^4		298	PL/Ld-2	S = MB.	85A167
2.257	1,4-Methanonaphthalene, 5,8-dimethoxy-1,2,3,4-tetrahydro-9-isopropylidene-						
	CH ₂ Cl ₂	4.3×10^6			PL/A'd-8	S = An; A' = DPBF.	81A437
2.258	1,4-Methanonaphthalene, 5,6,7,8-tetrachloro-1,2,3,4-tetrahydro-9-isopropylidene-						
	CH ₂ Cl ₂	9.6×10^4			PL/A'd-8	S = An; A' = DPBF.	81A437

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.259	1,4-Methanonaphthalene, 5,6,7,8-tetrafluoro-1,2,3,4-tetrahydro-9-isopropylidene-						
	CH ₂ Cl ₂	5.2×10^4			PL/A'd-8	S = An; A' = DPBF.	81A437
2.260	1,4-Methanonaphthalene, 1,2,3,4-tetrahydro-9-isopropylidene-						
	CH ₂ Cl ₂	1.4×10^5			PL/A'd-8	S = An; A' = DPBF.	81A437
2.261	Naphthalene, 1,2,3,4,5,6,7,8-octahydro- ($\Delta^{9,10}$ -Octalin)						
	CHCl ₃	5.0×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005
2.262	2,4,6,8-Nonatetraenoic acid, 9-(4-methoxy-2,3,6-trimethylphenyl)-3,7-dimethyl-, ethyl ester, (all-E)-						
	CCl ₄	2×10^7			CL/LI-12	S = TPP; used $k_d = 1 \times 10^3$ s ⁻¹ .	85F667
2.263	1-Nonene						
	MeOH			288	CP/Ac,A'c-17	S = MB; A' = <i>c</i> -C ₆ H ₁₀ ; meas. $k_t/k_t^{A'} = 0.1$.	65F028
2.264	9,12-Octadecadienoic acid (Linoleic acid)						
	C ₆ D ₆	7.3×10^4			PL/Ld-2	S = TPP.	89A331
	CCl ₄	4.2×10^4			MP/LI-12	S = TPP; used $k_d = 42$ s ⁻¹ .	83E425
	CD ₃ OD	7.9×10^4			PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	89A331
	CH ₃ CN	2.8×10^5		298	CP/Ac-14	S = RF(OAc) ₄ ; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	89F484
2.265	9,12-Octadecadienoic acid, methyl ester (Methyl linoleate)						
	C ₅ H ₅ N	1.3×10^5 (k_t)			CP/Ac,A'c-17	S = PP; A' = Cholesterol; used $k_d = 6 \times 10^4$ s ⁻¹ ; meas. $k_t/k_t^{A'} = 2.0$, $\beta_{A'} = 0.89$ mol L ⁻¹ .	743115
	CHCl ₃	3.8×10^4 (k_t) 9.5×10^3 (k_q)	0.12 (β_t)	293	CP/Oc-14,27	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; meas. $k_q/k_t = 0.26$; obs. $\phi_{ox}^{TME}/\phi_{ox}^{A'}$; cor. for k_q (MB).	94R058
	EtOH		0.36	293	CP/Pa-15	S = MB; P = Methyl linoleate monohydroperoxide.	777378
2.266	9,12-Octadecadienoic acid, phenyl ester (Phenyl linoleate)						
	MeOD	5.5×10^4			PL/Ld-2	S = MB and RF.	88A165
	MeOD	5.4×10^4		298	PL/Ld-2	S = Ery, MB, RF and HP.	87R138
2.267	9,12,15-Octadecatrienoic acid (Linolenic acid)						
	C ₆ D ₆	1.0×10^5			PL/Ld-2	S = TPP.	89A331
	CCl ₄	8×10^4			MP/LI-12	S = TPP; used $k_d = 42$ s ⁻¹ .	83E425
	CH ₃ CN	6.4×10^5		298	CP/Ac-14	S = RF(OAc) ₄ ; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	89F484
2.268	9,12,15-Octadecatrienoic acid, ethyl ester (all-Z)- (Ethyl linolenate)						
	CHCl ₃	5.3×10^4 (k_t) 1.1×10^4 (k_q)	0.082 (β_t)	293	CP/Oc-14,27	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; meas. $k_q/k_t = 0.51$; obs. $\phi_{ox}^{TME}/\phi_{ox}^{A'}$; cor. for k_q (MB).	94R058
2.269	9,12,15-Octadecatrienoic acid, methyl ester (Methyl linolenate)						
	C ₅ H ₅ N	1.6×10^5			CP/A'c-23	S = A' = Rub; used $k_d = 6.0 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ .	743115
	C ₅ H ₅ N	1.9×10^5 (k_t)			CP/Ac,A'c-17	S = PP; A' = Cholesterol; used $k_d = 6 \times 10^4$ s ⁻¹ ; meas. $k_t/k_t^{A'} = 2.9$, $\beta_{A'} = 0.89$ mol L ⁻¹ .	743115
	EtOH		0.27	293	CP/Pa-15	S = MB.	777378
2.270	9,12,15-Octadecatrienoic acid, phenyl ester (Phenyl linolenate)						
	CD ₃ OD	8.3×10^4			PL/Ld-2	S = MB and RF.	88A165
2.271	9-Octadecenoic acid (Oleic acid)						
	C ₆ D ₆	5.3×10^4			PL/Ld-2	S = TPP.	89A331

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.271	9-Octadecenoic acid (Oleic acid) — Continued						
	CCl ₄	1.7×10^4			MP/LI-12	S = TPP; used $k_d = 42$ s ⁻¹ .	83E425 83F151
	CH ₃ CN	1.6×10^5		298	CP/Ac-14	S = RF(OAc) ₄ ; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	89F484
2.272	9-Octadecenoic acid, methyl ester (Methyl oleate)						
	C ₅ H ₅ N	7.4×10^4 (k_r)			CP/Ac,A'c-17	S = PP; A' = Cholesterol; used $k_d = 6 \times 10^4$ s ⁻¹ ; meas. $k_r/k_{A'} = 1.1$, $\beta_{A'} = 0.89$ mol L ⁻¹ .	743115
	CHCl ₃	2.4×10^4 (k_r) 1.2×10^4 (k_q)	0.18 (β_r)	293	CP/Oc-14,27	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r = 0.20$; obs. $\phi_{ox}^{TME}/\phi_{ox}^A$; cor. for k_q (MB).	94R058
	EtOH		0.63	293	CP/Pa-15	S = MB; P = Methyl oleate monohydroperoxide.	777378
2.273	9-Octadecenoic acid, phenyl ester (Phenyl oleate)						
	MeOD	2.4×10^4			PL/Ld-2	S = MB and RF.	88A165
	MeOD	2.4×10^4		298	PL/Ld-2	S = Ery, MB, RF and HP.	87R138
2.274	2,5-Octadiene, 7-hydroperoxy-2,7-dimethyl-						
	MeOH	4.2×10^5 (k_r)	0.27 (β_r)	253	CP/Oc,Ac,Pa-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; $k(P_1) = 1.9 \times 10^5$, $k(P_2) = 2.3 \times 10^5$ L mol ⁻¹ s ⁻¹ ; $P_1 = 2,7$ -dimethyl-2,7-dihydroperoxy-3,5-octadiene, $P_2 = 2,7$ -dimethyl-2,6-dihydroperoxy-3,7-octadiene.	78A357
2.275	2,6-Octadiene, 2,6-dimethyl-, (E)						
	MeOH	1.9×10^6 (k_r)		253	CP/Oc-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	78A344
2.276	2,6-Octadiene, 2,6-dimethyl-, (Z)						
	MeOH	1.1×10^6 (k_r)		253	CP/Oc-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	78A344
2.277	2,6-Octadiene, 2,7-dimethyl-						
	MeOH	1.7×10^6 (k_r)	0.065 (β_r)	253	CP/Oc,Pa,Ac-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; $k(P_1) = 9.0 \times 10^5$, $k(P_2) = 8.1 \times 10^5$ L mol ⁻¹ s ⁻¹ ; $P_1 = 2,7$ -dimethyl-7-hydroperoxy-2,6-octadiene, $P_2 = 2,7$ -dimethyl-6-hydroperoxy-2,7-octadiene.	78A357
2.278	2,6-Octadiene, hydroperoxy-2,6-dimethyl-						
	MeOH	7.4×10^5 (k_r)		253	CP/Oc,Pa,Ac-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; Mixture of all monohydroperoxides obtained from oxygenation of <i>trans</i> -2,6-dimethyl-2,6-octadiene.	78A344
2.279	2,7-Octadiene, 6-hydroperoxy-2,7-dimethyl-						
	MeOH	6.0×10^5 (k_r)	0.19 (β_r)		CP/Oc,Pa,Ac-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; $P_1 = 2,7$ -dimethyl-2,6-dihydroperoxy-3,7-octadiene, $P_2 = 2,7$ -dimethyl-3,6-dihydroperoxy-1,7-octadiene, $k(P_1) = 1.9 \times 10^5$, $k(P_2) = 4.1 \times 10^5$ L mol ⁻¹ s ⁻¹ .	78A357
2.280	1,6-Octadien-3-ol, 3,7-dimethyl- (Linalool)						
	CH ₂ Cl ₂				CR/Ac,A'c-17	A' = 2,6-[$(CH_3)_3C_2$] ₂ C ₆ H ₃ OH; meas. $k_r/k_{A'} = 24$; $^1O_2^*$ from $(PhO_3)_3PO_3$.	72F520
	MeOH				CP/Ac,A'c-17	S = RB; A' = 2,6-[$(CH_3)_3C_2$] ₂ C ₆ H ₃ OH; meas. $k_r/k_{A'} = 14$.	72F520
2.281	2,6-Octadien-1-ol, 3,7-dimethyl-, (E) (Geraniol)						
	MeOH	5.9×10^5 (k_r)	0.19 (β_r)	253	CP/Oc-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	80A430

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.282	2,7-Octadien-1-ol, 3,7-dimethyl-, (E)- MeOH/ H ₂ O (95:5)	0.15			CP/Oc-15	S = RB; Soln. contg. 1% sodium acetate.	79F137
2.283	2,7-Octadien-1-ol, 3,7-dimethyl-, (Z)- MeOH/ H ₂ O (95:5)	0.21			CP/Oc-15	S = RB; Soln. contg. 1% sodium acetate.	79F137
2.284	2,7-Octadien-1-ol, 6-hydroperoxy-3,7-dimethyl- (Monohydroperoxygeraniol) MeOH	2.8×10^4 (k_r)		253	CP/Oc-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; mixture with 7-hydroperoxy-3,7-dimethyl-2,5-octadien-1-ol.	80A430
2.285	2,4,6-Octatriene, 2,6-dimethyl- (Alloocimine) MeOH	0.070		293	CP/Oc-15	S = RB; $E_a = 8.4$ kJ mol ⁻¹ .	68F288
2.286	3-Octene, 5-hydroperoxy-4-methyl- MeOH	2.5×10^4 (k_r)		253	CP/Oc-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	80A430
2.287	(E)-4-Octene CH ₃ CN	2.5×10^4 1×10^4 (k_r) 2×10^4 (k_q)			CP/Oc-14,27	S = MB; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r = 1.5$; obs. rate of oxygen consumption as a ratio with TME; cor for $k_S[S] = 0.28$.	88A026
	CHCl ₃	2.2×10^4 4×10^3 (k_r) 2×10^4 (k_q)			CP/Oc-14,27	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r = 4.5$; obs. rate of oxygen consumption as a ratio with TME; cor for $k_S[S] = 0.2$.	88A026
2.288	(Z)-4-Octene CHCl ₃	2.7×10^4 2.2×10^4 (k_r) 5×10^3 (k_q)	0.20 (β_r)		CP/Oc-14,27	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r = 0.22$; obs. rate of oxygen consumption as a ratio with TME; calcn. involves $k_q(MB) = 6.4 \times 10^6$ L mol ⁻¹ s ⁻¹ .	88A026 89A254
2.289	4-Octene, 4-methyl- MeOH	0.27 (β_r)		253	CP/Oc-14,27	S = RB; A' = α -Terpinene; 46% trans, 54% cis.	76F909
	MeOH	0.29 (β_r)		253	CP/Oc-14,27	S = RB; A' = α -Terpinene; 25% trans, 75% cis.	76F909
2.290	4-Octene, 4-methyl-, (E)- MeOH	5.6×10^5 (k_r)	0.20 (β_r)	253	CP/Oc-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	79F074
2.291	4-Octene, 4-methyl-, (Z)- CHCl ₃	1.6×10^5 1.6×10^5 (k_r) $<8 \times 10^3$ (k_q)	0.027		CP/Oc-14/27	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r = <0.05$; obs. rate of oxygen consumption as a ratio with TME; calcn. involves $k_q(MB) = 6.4 \times 10^6$ L mol ⁻¹ s ⁻¹ .	85F148 89A254
	MeOH	3.3×10^5 (k_r)	0.34 (β_r)	253	CP/Oc-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	79F074
2.292	2-Octen-1-ol, 3,7-dimethyl-, (E)- MeOH/ H ₂ O (95:5)	0.14			CP/Oc-15	S = RB; Soln. contg. 1% sodium acetate.	79F137
2.293	2-Octen-1-ol, 3,7-dimethyl-, (Z)- MeOH/ H ₂ O (95:5)	0.23			CP/Oc-15	S = RB; Soln. contg. 1% sodium acetate.	79F137
2.294	6-Octen-1-ol, 3,7-dimethyl- (β -Citronellol) 1-BuOH	0.12		293	CP/Pa-15	S = Eos.	627005
	1-BuOH	0.11		293	CP/Pa-15	S = RB.	627005
	CD ₃ OD	7.8×10^5		295	CL/LI-12	S = RB; used $k_d = 4.0 \times 10^3$ s ⁻¹ .	91F332

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_q/k_r) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.294 6-Octen-1-ol, 3,7-dimethyl- (β-Citronellol) — Continued							
	CD ₃ OD	7.3×10^5 (k_r) 1×10^4 (k_q)		295	CP/Oc-15,28	S = RB; used $k_d = 4.0 \times 10^3$ s ⁻¹ ; used $\phi_{\Delta}(RB) = 0.76$.	91F332
	MeOH	0.16		293	CP/Pa-15	S = Ery.	627005
	MeOH	0.15		293	CP/Pa-15	S = Eos.	627005
	MeOH	0.16		293	CP/Pa-15	S = RB.	627005
	MeOH/ H ₂ O (70:30)	0.060		293	CP/Pa-15	S = RB or Eos.	627005
2.295 7-Oxabicyclo[2.2.1]heptane, 2,3-bis(methylene)-							
	CH ₂ Cl ₂			253	CP/Ac-17	S = TPP; A' = 2,3-Dimethylenebicyclo[2.2.1]heptane; meas. $k_r/k_r^{A'} = 0.07$.	82F450
2.296 7-Oxabicyclo[2.2.1]heptane, 2,3,5,6-tetrakis(methylene)-							
	CH ₂ Cl ₂			253	CP/Ac-17	S = TPP; A' = 2,3-Dimethylenebicyclo[2.2.1]heptane; meas. $k_r/k_r^{A'} = 0.02$.	82F450
2.297 7-Oxabicyclo[2.2.1]hept-2-ene, 5,6-bis(methylene)-							
	CH ₂ Cl ₂			253	CP/Ac-17	S = TPP; A' = 2,3-Dimethylenebicyclo[2.2.1]heptane; meas. $k_r/k_r^{A'} = <1 \times 10^{-3}$.	82F450
2.298 P-422 (8 conjugated bonds)							
	C ₆ H ₆ / MeOH (60:40)	1.2×10^{10}	8.3×10^{-6}		CP/A'c-22	S = MB; A' = Rub; used $k_d = 1.0 \times 10^5$ s ⁻¹ .	747042
2.299 P-438 (9 conjugated bonds)							
	C ₆ H ₆ / MeOH (60:40)	3.1×10^{10}	3.2×10^{-6}		CP/A'c-22	S = MB; A' = Rub; used $k_d = 1.0 \times 10^5$ s ⁻¹ .	747042
2.300 1,3-Pentadiene, 4-methyl-							
	CHCl ₃	1.6×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
2.301 1,3-Pentadiene, 1-phenyl-, (E,E)-							
	C ₆ H ₆	1.5		277	CP/Oc-14	S = TPP.	81F582
	CCl ₄	0.24		277	CP/Oc-14	S = TPP.	81F582
	CH ₃ COCH ₂ CH ₃	1.2		277	CP/Oc-14	S = RB.	81F582
	CHCl ₃	0.33		277	CP/Oc-14	S = TPP.	81F582
	EtOH	2.9		277	CP/Oc-14	S = RB.	81F582
2.302 1-Pentene							
	CCl ₄ / MeOH (96:4)	1×10^3			PL/A'd-8	S = MB; A' = DPBF.	777162
2.303 1-Pentene, 2,3,4-trimethyl-							
	CH ₃ CN			283	CP/Ac-17	S = MB; A' = TME; meas. $k_r/k_r^{A'} = 0.88$; C.k. with 1,2-dimethylcyclohexene.	79F646
2.304 (E)-2-Pentene							
	CCl ₄ / MeOH (96:4)	2×10^3			PL/A'd-8	S = MB; A' = DPBF.	777162
	CHCl ₃	2.1×10^4 7×10^3 (k_r) 1×10^4 (k_q)			CP/Oc-14,27	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r = 2.3$; obs. r_{ox} as a ratio with TME; cor. for $k_S[S] = 0.2$.	88A026

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.305 (Z)-2-Pentene							
	CCl ₄	4.1×10^4 $3.8 \times 10^4 (k_r)$ $3 \times 10^3 (k_q)$			CP/Oc-14,27	S = TPP; used $k_d = 1.1 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r = 0.08$; obs. r_{ox} as a ratio with TME; cor. for $k_S[S] = 1.4$.	88A026
	CCl ₄ / MeOH (96:4)	4×10^3			PL/A'd-8	S = MB; A' = DPBF.	77F162
	CHCl ₃	3.8×10^4 $4 \times 10^3 (k_q)$			CP/Oc-14,27	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; obs. r_{ox} as a ratio with TME; cor. for $k_S[S] = 0.2$.	88A026
2.306 2-Pentene, 2,4-dimethyl-							
	CHCl ₃	4.3×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005
	MeOH	1.3			?	Method not reported.	68F289
2.307 2-Pentene, 2-methyl- (2M2P)							
	tert-BuOH	0.081		298	CP/A'c-16	S = RB; A' = DPBF.	717398
	tert-BuOH	0.051		303	CP/Ac-14	S = RB.	68F292
	C ₂ H ₅ I	0.1		298	CP/Pa-15	S = ZnTPP.	717356
	C ₅ H ₅ N	0.043			CP/Oc-15	S = RB.	757166
	C ₅ H ₅ N	0.050		298	CP/Pa-15	S = RB.	717356
	c-C ₆ H ₁₁ OH	0.070		298	CP/Pa-15	S = RB.	717356
	1,3-C ₆ H ₄ (OCH ₃) ₂	0.2		298	CP/Pa-15	S = ZnTPP.	717356
	C ₆ H ₅ Br	0.05		298	CP/Pa-15	S = ZnTPP.	717356
	C ₆ H ₅ OCH ₃	0.13		298	CP/Pa-15	S = ZnTPP.	717356
	C ₆ H ₆	0.049			CP/Oc-14	S = TPP, Chl a, Chl b, Phe b, ZnTPP, MgTPP, PPDME.	88R136
	C ₆ H ₆	8.7×10^5	0.046		CP/Oc-14	S = RB; used $k_d = 4.0 \times 10^4$ s ⁻¹ .	84F191
	C ₆ H ₆	7.2×10^5		295	PL/Ld-2	S = MPDME; $k_r \gg k_q$.	83F196
	C ₆ H ₆	9.2×10^5	0.044		CP/Oc-19	S = Chl (os); Q = Chl (os); used $k_d = 4.0 \times 10^4$ s ⁻¹ ; Chl (os) is a commercial mixture contg. a small % of chlorophyll.	81F321
	C ₆ H ₆				CP/Oc-15	S = azine.	757166
	C ₆ H ₆	0.53			CP/Pa-15	S = ZnTPP.	717356
	C ₆ H ₆	0.10		298	CP/Pa-15	S = ZnTPP.	71F580
	C ₆ H ₆	0.2			CP/Pa-15	S = MB.	717356
	C ₆ H ₆ / MeOH (80:20)	0.050			CP/Pa-15	S = MB.	717356
	CH ₂ Cl ₂	9.7×10^5		295	PL/Ld-2	S = MPDME; $k_r \gg k_q$.	83F196
	CH ₃ CN	1.2×10^6	0.014		CP/Oc-14	S = RB; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	84F191
	CH ₃ CN	1.2×10^6		295	PL/Ld-2	S = MPDME; $k_r \gg k_q$.	83F196
	CH ₃ CN		0.018		CP/Oc-15	S = RB.	757166
	CH ₃ CN		0.014		CP/Pa-15	S = RB.	71F580
	CH ₃ CO ₂ CH ₃	0.04		298	CP/Pa-15	S = RB.	717356
	CH ₃ COCH ₃	8.1×10^5		295	PL/Ld-2	S = MPDME; $k_r \gg k_q$.	83F196
	CH ₃ COCH ₃	$4.0 \times 10^5 (k_r)$ $2.8 \times 10^5 (k_p)$		241	CP/Pa-17	S = RB; $\Delta H^\ddagger = 3$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -122$ J K ⁻¹ mol ⁻¹ ; P = 50:50 mixture of 2-methyl-3-penten-2-ol and 2-methyl-1-penten-3-ol; rel. to value at 295K from luminescence decay; $k_q \ll k_r$.	83F196
	CH ₃ COCH ₃			195			
	CHCl ₃	0.080		298	CP/Pa-15	S = RB.	717356
	CHCl ₃	7.6×10^5			PL/Ld-2	S = TPP.	92F103
	CHCl ₃	5.1×10^5	8.5×10^{-3}		CP/Oc-15	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; cor. for k_S .	85F148
	CHCl ₃	1.9×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.307 2-Pentene, 2-methyl- (2M2P) — Continued						
CS ₂	0.080		273	CP/Pa-15	S = ZnTPP.	717356
DMSO	0.055			CP/A'c-16	S = RB; A' = DPBF.	766072
DMSO	0.070		298	CP/Pa-15	S = RB.	717356
MeOH	6.9×10^5	0.16		CP/Oc-14	S = RB; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	84F191
MeOH	4.0×10^5 (k_r)		241	CP/Pa-17	S = RB; $\Delta H^\ddagger = 3.9 \pm 3.5$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -121 \pm 15$ J K ⁻¹ mol ⁻¹ ; rel. to $k_A = 6.7 \times 10^5$ L mol ⁻¹ s ⁻¹	83F196
	2.0×10^5 (k_r)		195		at 298 K; $k_q \ll k_r$.	
MeOH	6.2×10^5	0.18		CP/Oc-14	S = RB; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	81F321
MeOH	0.13		298	CP/Pa-15	S = Chl b.	717356
MeOH	0.2		298	CP/Pa-15	S = Chl a.	717356
MeOH	0.16		298	CP/Pa-15	S = RB.	717356
MeOH	0.15		298	CP/A'c-16	S = RB; A' = DPBF.	717398
MeOH	0.2			CP/Pa-15	S = ZnTPP.	71F580
MeOH	0.13		293	CP/Oc-15	S = RB; $E_a = 8.4$ kJ mol ⁻¹ .	68F288
MeOH	0.18				Method not reported.	68F289
MeOH/ <i>tert</i> -BuOH (50:50)	0.12			CP/A'c-16	S = RB; A' = DPBF.	717398
MeOH/ <i>tert</i> -BuOH (50:50)			303	CP/Pa,P'a-17	S = RB; A' = 2M2B; meas. $k_r/k_r^{A'} = 0.76$.	68F292
MeOH/ <i>tert</i> -BuOH (50:50)			298	CR/Pa,P'a-17	A' = 2M2B; meas. $k_r/k_r^{A'} = 0.78$; ${}^1\text{O}_2^*$ from $\text{H}_2\text{O}_2/\text{NaOCl}$.	68F292
MeOH/ <i>tert</i> -BuOH (50:50)	0.2		298	CR/Ac-14	${}^1\text{O}_2^*$ from $\text{H}_2\text{O}_2/\text{NaOCl}$.	68F292
MeOH/ <i>tert</i> -BuOH (50:50)	0.13		298	CP/Ac-14	S = RB.	68F292
2.308 2-Pentene, 3-methyl-						
C ₆ H ₅ Cl	1.0×10^6	0.016	298	CP/Ac-15	S = TPP; used $k_d = 1.6 \times 10^4$ s ⁻¹ ; <i>cis,trans</i> mixt.	747341
2.309 2-Pentene, 3-methyl-, (E)-						
CHCl ₃	2×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005
EtOH/ 2-PrOH (50:50)	0.060 (β_r)		292	CP/Oc-14,27	S = RB; A' = 2,5-DMF.	78F464
MeOH	9.3×10^5 (k_r)	0.12 (β_r)	253	CP/Oc-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	79F074
MeOH/ <i>tert</i> -BuOH (50:50)				CP/Ac,A'c-17	S = RB; A' = 2M2B; meas. $k_r/k_r^{A'} = 0.83$.	68F292
MeOH/ <i>tert</i> -BuOH (50:50)				CR/Ac,A'c-17	A' = <i>c</i> -C ₆ H ₉ CH ₃ ; meas. $k_r/k_r^{A'} = 7$; ${}^1\text{O}_2^*$ from $\text{H}_2\text{O}_2/\text{Ca}(\text{OCl})_2$.	68F292
MeOH/ <i>tert</i> -BuOH (50:50)				CR/Ac,A'c-17	A' = 2M2B; meas. $k_r/k_r^{A'} = 0.71$; ${}^1\text{O}_2^*$ from $\text{H}_2\text{O}_2/\text{Ca}(\text{OCl})_2$.	68F292
MeOH/ <i>tert</i> -BuOH (50:50)				CP/Ac,A'c-17	S = RB; A' = <i>c</i> -C ₆ H ₉ CH ₃ ; meas. $k_r/k_r^{A'} = 22$.	68F292
2.310 2-Pentene, 3-methyl-, (Z)-						
CHCl ₃	2.9×10^5	0.015		CP/Oc-15	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; cor. for quenching by MB.	85F148
CHCl ₃	2.0×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005
EtOH/ 2-PrOH (50:50)	0.050 (β_r)		292	CP/Oc-14,27	S = RB; A' = 2,5-DMF.	78F464

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.310 2-Pentene, 3-methyl-, (Z)- — Continued							
	MeOH	$5.8 \times 10^5 (k_r)$	0.19 (β_r)	253	CP/Oc-14,27	$S = RB; A' = \alpha\text{-Terpinene}$; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	79F074
	MeOH/ <i>tert</i> -BuOH (50:50)				CR/Ac,A'c-17	$A' = 2M2B$; meas. $k_r/k_r^{A'} = 0.40$; ${}^1O_2^*$ from $H_2O_2/Ca(OCl)_2$.	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)				CR/Ac,A'c-17	$A' = TME$; meas. $k_r/k_r^{A'} = 0.040$; ${}^1O_2^*$ from $H_2O_2/NaOCl$.	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)				CP/Ac,A'c-17	$S = RB; A' = 2M2B$; meas. $k_r/k_r^{A'} = 0.55$.	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)				CP/Ac,A'c-17	$S = RB; A' = TME$; meas. $k_r/k_r^{A'} = 0.028$.	68F292
2.311 2-Pentene, 4-methyl-, (E)-							
	MeOH/ <i>tert</i> -BuOH (50:50)			276	CR/Pa,P'a-17	$A' = 2M2P$; meas. $k_r/k_r^{A'} = 2.0 \times 10^{-3}$; ${}^1O_2^*$ from $H_2O_2/NaOCl$.	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)			303	CP/Pa,P'a-17	$S = RB; A' = 2M2P$; meas. $k_r/k_r^{A'} = 2.5 \times 10^{-3}$.	68F292
2.312 2-Pentene, 4-methyl-, (Z)-							
	MeOH/ <i>tert</i> -BuOH (50:50)	8.0			CP/A'c-16	$S = RB; A' = DPBF$.	717398
	MeOH/ <i>tert</i> -BuOH (50:50)			276	CR/Pa,P'a-17	$A' = 2M2P$; meas. $k_r/k_r^{A'} = 0.011$; ${}^1O_2^*$ from $H_2O_2/NaOCl$.	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)			303	CP/Pa,P'a-17	$S = RB; A' = 2M2P$; meas. $k_r/k_r^{A'} = 0.014$.	68F292
2.313 2-Pentene, 2,3,4-trimethyl-							
	CHCl ₃	3.9×10^7			CP/A'c-33	$S = A' = Rub$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005
2.314 2-Pentene, 2,4,4-trimethyl-							
	MeOH		4.2		?	Method not reported.	68F289
2.315 4-Pentenoic acid							
	C ₆ D ₆	4.9×10^3			PL/Ld-2	$S = TPP$.	89A331
2.316 2-Penten-1-ol, 3,4-dimethyl-, (E)-							
	MeOH/ H ₂ O (95:5)	0.14			CP/Oc-15	$S = RB$; Soln. contg. 1% sodium acetate.	79F137
2.317 2-Penten-1-ol, 3,4-dimethyl-, (Z)-							
	MeOH/ H ₂ O (95:5)	0.23			CP/Oc-15	$S = RB$; Soln. contg. 1% sodium acetate.	79F137
2.318 2-Penten-1-ol, 3-ethyl-							
	MeOH/ H ₂ O (95:5)	0.25			CP/Oc-15	$S = RB$; Soln. contg. 1% sodium acetate.	79F137
2.319 2-Penten-1-ol, 3-methyl-, (E)-							
	MeOH/ H ₂ O (95:5)	0.13			CP/Oc-15	$S = RB$; Soln. contg. 1% sodium acetate.	79F137
2.320 2-Penten-1-ol, 3-methyl-, (Z)-							
	MeOH/ H ₂ O (95:5)	0.19			CP/Oc-15	$S = RB$; Soln. contg. 1% sodium acetate.	79F137
2.321 2-Penten-1-ol, 3,4,4-trimethyl-, (E)-							
	MeOH/ H ₂ O (95:5)	0.09			CP/Oc-15	$S = RB$; Soln. contg. 1% sodium acetate.	79F137
2.322 2-Penten-1-ol, 3,4,4-trimethyl-, (Z)-							
	MeOH/ H ₂ O (95:5)	0.95			CP/Oc-15	$S = RB$; Soln. contg. 1% sodium acetate.	79F137
2.323 2-Penten-4-ol, 2-methyl-							
	MeOH		8.4 (β_r)	253	CP/Oc-14,27	$S = RB; A' = \alpha\text{-Terpinene}$.	76F909

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.324	3-Penten-2-one, 3,4-dimethyl-			28	CP/Ac-18	S = Poly-RB or RB; A' = Linalool; used $\beta_{A'} = 0.18$ mol L ⁻¹ ; Solvent is CH ₂ Cl ₂ or MeOH.	80F111
2.325	1-Phenanthrenecarboxylic acid, 1,2,3,4,4a,4b,5,6,10,10a-decahydro-1,4a-dimethyl-7-(1-methylethyl)-, [1R-(1 α ,4 α ,4 β ,4 β α ,10 α α)]- (Abietic acid)	CHCl ₃	3.8×10^6		CP/A' ^c -33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
2.326	α -Pinene						
	CH ₃ CN				?	S = MB; A' = <i>c</i> -C ₆ H ₉ CH ₃ ; meas. $k_q/k_{A'} = 0.25$.	73F664
	CH ₃ COCH ₃	4.3×10^4			PL/Ld-2	S = RB.	92F225
	MeOH	7.0		293	CP/Oc-15	S = MB; $E_a = 20$ kJ mol ⁻¹ .	68F288
	MeOH	5.0		293	CP/Oc-15	S = RB; $E_a = 19$ kJ mol ⁻¹ .	68F288
	MeOH	2.8		293	CP/Oc-15	S = DNT; $E_a = 21$ kJ mol ⁻¹ .	68F288
	MeOH	0.50		293	CP/Oc-15	S = FeCl ₄ ²⁻ ; $E_a = 17$ kJ mol ⁻¹ .	68F288
	MeOH	3.8		293	CP/Oc-15	S = FeBr ₄ Cl ₄ ²⁻ ; $E_a = 17$ kJ mol ⁻¹ .	68F288
	MeOH	8.2			?	Method not reported.	68F289
2.327	β -Pinene						
	CH ₃ CN				CP/Ac,A' ^c -17	S = MB; A' = <i>c</i> -C ₆ H ₉ CH ₃ ; meas. $k_q/k_{A'} = 0.19$.	73F664
	MeOH	-1.0		293	CP/Oc-15	S = RB; $E_a = 21$ kJ mol ⁻¹ .	68F288
2.328	(E)-Piperylene ((E)-1,3-Pentadiene)	CHCl ₃	2.4×10^4		CP/A' ^c -33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
2.329	(Z)-Piperylene ((Z)-1,3-Pentadiene)	CHCl ₃	9×10^3		CP/A' ^c -33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
2.330	Polybutadiene						
	CH ₂ Cl ₂	3.4×10^5			PL/Ld-2	S = An.	86E809 86E884
	CHCl ₃	2.3×10^4 1.7×10^4 (k_q) 6×10^3 (k_q)	0.26		CP/Oc-14,27	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; meas. $k_q/k_{A'} = 0.35$; obs. rate of oxygen consumption as a ratio with TME or 2,5-DMF; calcn. involves $k_q(MB) = 6.4 \times 10^6$ L mol ⁻¹ s ⁻¹ .	89A254
2.331	Polyisoprene						
	CH ₂ Cl ₂	5.5×10^5			PL/Ld-2	S = An.	86E809
	CHCl ₃	1.2×10^5 1.2×10^5 (k_q) $<6 \times 10^3$ (k_q)	0.035		CP/Oc-14,27	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; obs. rate of oxygen consumption as a ratio with TME or 2,5-DMF; calcn. involves $k_q(MB) = 6.4 \times 10^6$ L mol ⁻¹ s ⁻¹ .	89A254 85F148
2.332	Propene, 1,1-bis(cyclobutyl)-2-methyl-	CH ₃ CN		283	CP/Ac-17	S = MB; A' = TME; meas. $k_q/k_{A'} = 0.40$; C.k. with 1,2-dimethylcyclohexene.	79F646
2.333	Propene, 1,1-bis(cyclopropyl)-	CH ₃ COCH ₃			CP/Ac,A' ^c -17	S = Eos; A' = (Dicyclopropylmethyldiene)cyclobutane; meas. $k_q/k_{A'} = 0.50$.	78F430
2.334	Propene, 1,1-bis(cyclopropyl)-2-methyl-	CH ₃ CN		283	CP/Ac-17	S = MB; A' = TME; meas. $k_q/k_{A'} = 0.33$; C.k. with 1,2-dimethylcyclohexene.	79F646

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.334	Propene, 1,1-bis(cyclopropyl)-2-methyl-	— Continued					
	CH ₃ COCH ₃				CP/Ac,A'c-17	S = Eos; A' = (Dicyclopropylmethyldiene)cyclobutane; meas. $k_r/k_r^{A'} = 0.65$; meas. $k_H/k_D = 1$ for 2-(methyl- <i>d</i> ₃) isomer.	78F430
2.335	Propene, 1-cyclopropyl-2-methyl-	CH ₃ COCH ₃			CP/Ac,A'c-17	S = Eos; A' = (Dicyclopropylmethyldiene)cyclobutane; meas. $k_r/k_r^{A'} = 0.66$.	78F430
2.336	Propene, 1-ethoxy-2-methyl-	c-C ₆ H ₁₂	1.6×10^7		PL/A'd-8	S = An; A' = DPBF; $\Delta H^\ddagger = -2$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -120$ J K ⁻¹ mol ⁻¹ ; studied at 290-340 K.	82A349
	C ₆ H ₅ CH ₃	3.1×10^7			PL/A'd-8	S = An; A' = DPBF; $\Delta H^\ddagger = 0.8$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -98$ J K ⁻¹ mol ⁻¹ ; studied at 250-340 K.	82A349
	CH ₃ CN	3.9×10^7			PL/A'd-8	S = An; A' = DPBF; $\Delta H^\ddagger = -1$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -101$ J K ⁻¹ mol ⁻¹ ; studied at 250-340 K.	82A349
	CH ₃ COCH ₃	4.2×10^7			PL/A'd-8	S = An; A' = DPBF; $\Delta H^\ddagger = -2.5$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -108$ J K ⁻¹ mol ⁻¹ ; studied at 240-320 K.	82A349
	MeOH	2.3×10^7			PL/A'd-8	S = An; A' = DPBF; $\Delta H^\ddagger = 0.8$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -96$ J K ⁻¹ mol ⁻¹ ; studied at 250-310 K.	82A349
2.337	Propene, 2-methyl-	CS ₂	4.0×10^3	298	PL/Ld-2	S = TPP; $\Delta H^\ddagger = 0$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -180$ J K ⁻¹ mol ⁻¹ ; studied at 183-310 K.	85A167
	MeOH	1.6		293	CP/Oc-15	S = RB; $E_a = 24$ kJ mol ⁻¹ .	68F288
2.338	Propene, 2-methyl-1-(phenylsulfinyl)-	C ₆ H ₆	$>1 \times 10^2$ (β_r)		CP/Ac-17	S = TPP; A' = Linalool; used $\beta_r^{A'} = 0.18$ mol L ⁻¹ .	89F400
2.339	11-(Z)-Retinal	CCl ₄	10^6		MP/LI-12	S = PP; used $k_d = 36$ s ⁻¹ .	79F463
2.340	13-(Z)-Retinal	CCl ₄	3.6×10^5		MP/LI-12	S = PP; used $k_d = 36$ s ⁻¹ .	79F463 78F700
2.341	(all-E)-Retinal	CCl ₄	3.7×10^6		MP/LI-12	S = PP; used $k_d = 36$ s ⁻¹ .	79F463 78F700 83A336
	CCl ₄	7.5×10^5 (k_r) 3.7×10^6 (k_q)			CP/Ac-27	S = PP; A' = Tetr; meas. $k_q/k_r = 0.2 \pm 0.05$; used $\phi_A(S)$ and $k_A = 3.7 \times 10^6$ L mol ⁻¹ s ⁻¹ .	79F463 78F700 83A336
	CH ₃ CN	6.5×10^7			PL/A'd-8	S = MB; A' = DPBF.	83F256
	HCONHCH ₃	1.7×10^8			PL/A'd-8	S = MB; A' = DPBF.	83F256
2.342	(all-E)-Retinoic acid	CHCl ₃ /EtOH (50:50)		310	CR/LI-12	${}^1\text{O}_2^*$ from NDPO ₂ ; no quenching.	89R188
2.343	(all-E)-Retinol	c-C ₆ H ₁₂	$\leq 2 \times 10^6$		PL/A'd-8	S = Rub; A' = DPBF.	83F256
	n-C ₆ H ₁₄	$\leq 1.5 \times 10^7$			PL/A'd-8	S = Rub; A' = DPBF.	83F256
	C ₆ H ₆	4×10^6		296	PL/A'd-5	S = A; A' = DPBF.	85E190
	C ₆ H ₆	$\leq 1.2 \times 10^7$			PL/A'd-8	S = Rub; A' = DPBF.	83F256

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
1.343 (<i>all-E</i>)-Retinol — Continued							
	C ₆ H ₆ / MeOH (80:20)				CP/P'a-20	S = MB; A' = 2M2P; meas. $k_A/k_{A'} \leq 9$.	707188
	CH ₃ CN	2.6×10^7			PL/A'd-8	S = MB; A' = DPBF.	83F256
	CH ₃ COCH ₃	1×10^7			PL/A'd-8	S = MB; A' = DPBF.	83F256
	HCONHCH ₃	8.5×10^7			PL/A'd-8	S = MB; A' = DPBF.	83F256
	MeOH	6.0×10^6		296	PL/A'd-5	S = A; A' = DPBF.	85E190
	2-PrOH	1.7×10^7			PL/A'd-8	S = MB; A' = DPBF.	83F256
1.344 Retinyl acetate							
	CCl ₄	7×10^5			MP/LI-12	S = PP; used $k_d = 36$ s ⁻¹ .	79F463 78F700
1.345 Spirocyclopropane[4,7]methanoindene, 4,5,6,7-tetrahydro-							
	CD ₃ COCD ₃	1.3×10^8		195	CP/A'c-16	S = RB; A' = DPBF; k_d not given.	87A368
1.346 Squalene							
	EtOH	4.2×10^6		298	CP/Ac-14	S = TTT; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; $E_a = 11.2$ kJ mol ⁻¹ ; studied at 271-326 K.	93F327
	MeOH	3.6×10^6	0.031	253	CP/Oc-15	S = RB; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	83A078
1.347 4,6,8,10,12,14,16,18,20-Tetracosanonaene-3,22-dione, 6,10,15,19-tetramethyl-2,2,23,23-tetramethoxy- (<i>all-E</i>) (C₂₈-Polyene-tetrone-diacetate)							
	CHCl ₃	9×10^9			CR/LI-12	used $k_d = 3.0 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	92R054
1.348 4,6,8,10,12,14,16,18,20-Tetracosanonaene-2,3,22,23-tetraone, 6,10,15,19-tetramethyl- (<i>all-E</i>) (C₂₈-Polyene-tetrone)							
	CHCl ₃	1.6×10^{10}			CR/LI-12	used $k_d = 3.0 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	92R054
1.349 (Z)-9-Tricosene							
	CHCl ₃	2.4×10^4 $1 \times 10^4 (k_r)$ $1 \times 10^4 (k_q)$	0.37	293	CP/Oc-14,27	S = MB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r = 1.0$; obs. rate of oxygen consumption as a ratio with TME; calcn. involves $k_q(\text{MB}) = 6.4 \times 10^6$ L mol ⁻¹ s ⁻¹ .	88A026 83A394
1.350 Tricyclo[3.3.1.1^{3,7}]decane, 2-(3-chlorophenoxyethylene)- [2-(3-Chlorophenoxyethylene)adamantane]							
	C ₆ H ₆			283	CP/Pa-17	S = TPP; A' = 2-(Phenoxyethylene)adamantane; meas. $k_r/k_r^{A'} = 0.1$.	84F474
	CH ₂ Cl ₂			283	CP/Pa-17	S = MB; A' = 2-(Phenoxyethylene)adamantane; meas. $k_r/k_r^{A'} = 0.4$.	84F474
1.351 Tricyclo[3.3.1.1^{3,7}]decane, 2-(4-chlorophenoxyethylene)- [2-(4-Chlorophenoxyethylene)adamantane]							
	C ₆ H ₆			283	CP/Pa-17	S = TPP; A' = 2-(Phenoxyethylene)adamantane; meas. $k_r/k_r^{A'} = 0.2$.	84F474
	CH ₂ Cl ₂			283	CP/Pa-17	S = MB; A' = 2-(Phenoxyethylene)adamantane; meas. $k_r/k_r^{A'} = 0.6$.	84F474
1.352 Tricyclo[3.3.1.1^{3,7}]decane, 2-(4-methoxyphenoxyethylene)- [2-(4-Methoxyphenoxyethylene)adamantane]							
	C ₆ H ₆			283	CP/Pa-17	S = TPP; A' = 2-(Phenoxyethylene)adamantane; meas. $k_r/k_r^{A'} = 5$.	84F474
	CH ₂ Cl ₂			283	CP/Pa-17	S = MB; A' = 2-(Phenoxyethylene)adamantane; meas. $k_r/k_r^{A'} = 2$.	84F474

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.352 Tricyclo[3.3.1.1^{3,7}]decane, 2-(4-methoxyphenoxyethylene)- [2-(4-Methoxyphenoxymethyldene)adamantane] — Continued							
	CH ₃ CN			283	CP/Pa-17	S = MB; A' = 2-(Phenoxyethylene)adamantane; meas. $k_t/k_t^{A'}$ = 2.	84F474
	CH ₃ CO ₂ C ₂ H ₅			283	CP/Pa-17	S = TPP; A' = 2-(Phenoxyethylene)adamantane; meas. $k_t/k_t^{A'}$ = 6.	84F474
	CH ₃ COCH ₃			283	CP/Pa-17	S = MB; A' = 2-(Phenoxyethylene)adamantane; meas. $k_t/k_t^{A'}$ = 2.	84F474
	EtOH			283	CP/Pa-17	S = MB; A' = 2-(Phenoxyethylene)adamantane; meas. $k_t/k_t^{A'}$ = 2.	84F474
	MeOH			283	CP/Pa-17	S = MB; A' = 2-(Phenoxyethylene)adamantane; meas. $k_t/k_t^{A'}$ = 2.	84F474
	1-PrOH			283	CP/Pa-17	S = MB; A' = 2-(Phenoxyethylene)adamantane; meas. $k_t/k_t^{A'}$ = 2.	84F474
	THF			283	CP/Pa-17	S = TPP; A' = 2-(Phenoxyethylene)adamantane; meas. $k_t/k_t^{A'}$ = 6.	84F474
2.353 Tricyclo[3.3.1.1^{3,7}]decane, (1-methylethylene)- (2-Isopropylideneadamantane)							
	C ₆ H ₆				CP/Pa-17	S = TPP; A' = 1,1-Di- <i>tert</i> -butyl-2-methoxyethene; meas. $k_t/k_t^{A'}$ = 7.	82F102
	CH ₂ Cl ₂				CP/Pa-17	S = MB; A' = 1,1-Di- <i>tert</i> -butyl-2-methoxyethene; meas. $k_t/k_t^{A'}$ = 3.6.	82F102
	CH ₃ CN				CP/Pa-17	S = MB; A' = 1,1-Di- <i>tert</i> -butyl-2-methoxyethene; meas. $k_t/k_t^{A'}$ = 3.	82F102
	CH ₃ COCH ₃				CP/Pa-17	S = MB; A' = 1,1-Di- <i>tert</i> -butyl-2-methoxyethene; meas. $k_t/k_t^{A'}$ = 7.	82F102
	EtOH				CP/Pa-17	S = MB; A' = 1,1-Di- <i>tert</i> -butyl-2-methoxyethene; meas. $k_t/k_t^{A'}$ = 2.5.	82F102
	MeOH				CP/Pa-17	S = MB; A' = 1,1-Di- <i>tert</i> -butyl-2-methoxyethene; meas. $k_t/k_t^{A'}$ = 1.9.	82F102
	2-PrOH				CP/Pa-17	S = MB; A' = 1,1-Di- <i>tert</i> -butyl-2-methoxyethene; meas. $k_t/k_t^{A'}$ = 2.5.	82F102
2.354 Tricyclo[3.3.1.1^{3,7}]decane, 2-(4-methylphenoxyethylene)- [2-(4-Methylphenoxymethyldene)adamantane]							
	C ₆ H ₆			283	CP/Pa-17	S = TPP; A' = 2-(Phenoxyethylene)adamantane; meas. $k_t/k_t^{A'}$ = 2.	84F474
	CH ₂ Cl ₂			283	CP/Pa-17	S = MB; A' = 2-(Phenoxyethylene)adamantane; meas. $k_t/k_t^{A'}$ = 1.5.	84F474
2.355 Tricyclo[3.3.1.1^{3,7}]decane, (2-methylpropylidene)- [2-Isobutylideneadamantane]							
	C ₆ H ₆				CP/Pa-17	S = TPP; A' = 1,1-Di- <i>tert</i> -butyl-2-methoxyethene; meas. $k_t/k_t^{A'}$ = 2.	82F102
	CH ₂ Cl ₂				CP/Pa-17	S = MB; A' = 1,1-Di- <i>tert</i> -butyl-2-methoxyethene; meas. $k_t/k_t^{A'}$ = 1.	82F102
	CH ₃ CN				CP/Pa-17	S = MB; A' = 1,1-Di- <i>tert</i> -butyl-2-methoxyethene; meas. $k_t/k_t^{A'}$ = 0.7.	82F102
	CH ₃ COCH ₃				CP/Pa-17	S = MB; A' = 1,1-Di- <i>tert</i> -butyl-2-methoxyethene; meas. $k_t/k_t^{A'}$ = 1.2.	82F102

TABLE 2. Rate constants for interaction of singlet oxygen with olefins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
2.355 Tricyclo[3.3.1.1 ^{3,7}]decane, (2-methylpropylidene)- [2-Isobutylideneadamantane] — Continued							
	EtOH				CP/Pa-17	S = MB; A' = 1,1-Di- <i>tert</i> -butyl-2-methoxyethene; meas. $k_t/k_t^{A'} = 0.53$.	82F102
2.356 Tricyclo[3.3.1.1 ^{3,7}]decane, tricyclo[3.3.1.1 ^{3,7}]decylidene- (Adamantylideneadamantane)							
	C ₆ H ₅ Cl/ 2-PrOH (90:10)	2.2×10^5		273	CR/LI-12	used $k_d = 4.1 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from hydrotrioxide, e.g. (CH ₃) ₂ C(OH)OOOH.	90M125
	CHCl ₃	4.7×10^5		295	CL/LI-12	S = An; used $k_d = 4.9 \times 10^3$ s ⁻¹ .	90F473
	CHCl ₃	9.0×10^5			CR/A'c-33	A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from DMNO ₂ .	81E003
	CHCl ₃	1.0×10^6	0.016	288	CP/A'c-16	S = MB; A' = DPBF; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; Solvent contained 1% EtOH.	77F876
2.357 8,9,11-Trioxatetracyclo[4.2.2.1.0 ^{5,6}]undec-5-ene, 2,3-bis(methylene)-							
	CH ₂ Cl ₂			253	CP/Ac-17	S = TPP; A' = 2,3-Dimethylenebicyclo[2.2.1]heptane; meas. $k_t/k_t^{A'} = <1 \times 10^{-3}$.	82F450
2.358 (<i>all-E</i>)-1,3,5,7,9,11,13,15,17,19,21,23,25,27,29,31,33-Tritriacontaheptadecaene, 3,7,11,15,20,24,28,32-octamethyl-1,34-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)- (Dodecapreno- β -carotene)							
	C ₆ H ₅ CH ₃	2.9×10^{10}			PL/Ld-2	S = Pz.	91E465
	C ₆ H ₆	2.3×10^{10}			PL/Ld-2	S = Pz.	91E465
2.359 1,6-Undecadiene, 2,6-dimethyl-, (<i>E</i>)-							
	EtOH/ 2-PrOH (50:50)		0.10 (β_r)	292	CP/Oc-14,27	S = RB; A' = 2,5-DMF.	78F464
2.360 1,6-Undecadiene, 2,6-dimethyl-, (<i>Z</i>)-							
	EtOH/ 2-PrOH (50:50)		0.13 (β_r)	292	CP/Oc-14,27	S = RB; A' = 2,5-DMF.	78F464
2.361 Vitamin D ₃							
	C ₆ H ₆	4.0×10^5			PL/Ld-2	S = A.	90E030

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates.

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.1 2'-Acetonaphthone (2-ACN)							
	CDCl ₃	2×10^6			PL/A'd-5	S = A; A' = Rub.	81A287
3.2 Acetophenone							
	CCl ₄ / MeOH (98:2)	2.7×10^3		293	PL/A'd-8	S = MB; A' = DPBF.	83A371
3.3 Anthracene							
	n-C ₆ H ₁₄	3×10^5			CP/A'c-?	S = A; Q = Car; used $k_d = 6 \times 10^4$ s ⁻¹ , $k_Q = 3 \times 10^{10}$ L mol ⁻¹ s ⁻¹ .	80F169
	C ₆ H ₅ Br		0.12		CP/Ac-15	S = A.	59F003
	C ₆ H ₆	1.6×10^5	0.27	298	CP/A'c-20	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $\beta_{A'} = 1.0 \times 10^{-3}$ mol L ⁻¹ .	747312
	C ₆ H ₆		0.047		CP/Ac-15	S = DNT.	727028
	C ₆ H ₆		0.57		CP/Ac-15	S = A.	59F003
	C ₆ H ₆		0.42		CP/Pa-15	S = A; studied at 263-317 K.	55F004
	CCl ₄	3.5×10^5			CP/Ac-15	S = A; used $k_d = 1.4 \times 10^3$ s ⁻¹ .	80F718
	CHCl ₃		4.0×10^{-3}		CP/Oc-15	S = A.	537004
	CHCl ₃	5.4×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005
	CHCl ₃		0.036		CP/Pa-15	S = A; studied at 263-317 K.	55F004
	CHCl ₃		0.014		CP/Oc-15	S = A.	537004
	CS ₂		2.6×10^{-3}		CP/Ac-15	S = DNT.	727028
	CS ₂		3.6×10^{-3}		CP/Oc-15	S = A.	537004
	CS ₂ /C ₆ H ₆ (99:1)		0.015		CP/Oc-15	S = A; solvent in mole %.	537004
	CS ₂ /C ₆ H ₆ (96:4)		0.027		CP/Ac-15	S = DNT; solvent in mole %.	727028
	CS ₂ /C ₆ H ₆ (95:5)		0.038		CP/Oc-15	S = A; solvent in mole %.	537004
	CS ₂ /C ₆ H ₆ (93:7)		0.052		CP/Ac-15	S = DNT; solvent in mole %.	727028
	CS ₂ /C ₆ H ₆ (90:10)		0.062		CP/Oc-15	S = A; solvent in mole %.	537004
	CS ₂ /C ₆ H ₆ (75:25)		0.094		CP/Ac-15	S = TPP; solvent in mole %.	727028
	CS ₂ /C ₆ H ₆ (50:50)		0.077		CP/Ac-15	S = TPP; solvent in mole %.	727028
3.4 Anthracene, 1-chloro-							
	CCl ₄		4.1×10^{-3}		CP/Oc-15	S = A.	537004
	CHCl ₃		0.045		CP/Oc-15	S = A.	537004
	CS ₂		4.9×10^{-3}		CP/Oc-15	S = A.	537004
3.5 Anthracene, 9-chloro-							
	CCl ₄		3.2×10^{-3}		CP/Oc-15	S = A.	537004
	CHCl ₃		0.024		CP/Oc-15	S = A.	537004
	CS ₂		3.0×10^{-3}		CP/Oc-15	S = A.	537004
3.6 Anthracene, 1-chloro-9,10-diphenyl-							
	CHCl ₃	2×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005
3.7 Anthracene, 9,10-dichloro- (DCIA)							
	CCl ₄		0.010		CP/Oc-15	S = A.	537004
	CHCl ₃		0.060		CP/Oc-15	S = A.	537004
	CS ₂		0.017		CP/Oc-15	S = A.	537004
3.8 Anthracene, 9,10-dimethoxy-							
	C ₆ H ₆	1.4×10^7 (k_r)		298	CP/Ac,A'c-17	S = A' = Rub; used $k_r^{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.33$.	747312

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
J,9 Anthracene, 1,4-dimethoxy-9,10-diphenyl- (DMDPA)							
	(C ₂ H ₅) ₂ O		2 × 10 ⁻⁴		CP/Ac-15	S = A.	84F197
	C ₆ H ₅ CH ₃	1.3 × 10 ⁸			PL/Ld-2	S = A.	93F207
	CH ₃ CO ₂ C ₂ H ₅		9.3 × 10 ⁻⁵		CP/Ac-15	S = A.	84F197
	CHCl ₃	3.2 × 10 ⁸			CP/Ac-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005
J,10 Anthracene, 9,10-dimethyl- (DMA)							
	C ₅ H ₅ N		1.0 × 10 ⁻³	285	CP/Pa-20	S = MB; A' = TME.	737202
	C ₅ H ₅ N		1.4 × 10 ⁻³	285	CP/Pa-20	S = A; A' = TME.	737202
	C ₅ H ₅ N		1.4 × 10 ⁻³	285	CP/Pa-20	S = ThCl; A' = TME.	737202
	C ₅ H ₅ N				CP/Ac-17	S = MB; A' = TME; meas. $k_t/k_t^{A'} = 1.0$.	66F041
	c-C ₆ H ₁₂	1.2 × 10 ⁷	5 × 10 ⁻³	297	CP/Ac-15	S = A; used $k_d = 5.9 \times 10^4$ s ⁻¹ .	747207
	C ₆ H ₅ Br				CP/Ac-17	S = MB; A' = TME; meas. $k_t/k_t^{A'} = 1.0$.	66F041
	C ₆ H ₅ CH ₃		2 × 10 ⁻³		CP/Ac-15	S = A.	84F197
	C ₆ H ₅ CH ₃		1.5 × 10 ⁻³	298	CP/Ac-16	S = A; A' = Leucomalachite Green.	79F148
	C ₆ H ₆	2.5 × 10 ⁷		293	CP/Ac-15	S = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	85N293
	C ₆ H ₆		1.2 × 10 ⁻³	298	CP/Ac-?	S = A.	767422
	C ₆ H ₆	2.1 × 10 ⁷	2 × 10 ⁻³	298	CP/Ac-15	S = A; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	747312
	C ₆ H ₆		3.0 × 10 ⁻⁴	298	CP/Ac-15	S = A, Rub, Tetr.	706079
	C ₆ H ₆		3.0 × 10 ⁻⁴	298	CP/Ac-?	S = Per and An.	69F388
	C ₆ H ₆ / MeOH (60:40)	4.8 × 10 ⁷			CP/Ac-33	S = A' = Rub; k_d and $k_{A'}$ not given; 6 × 10 ⁻³ mol L ⁻¹ KOH.	88F650
	CCl ₄	7.0 × 10 ⁶ (k_t)		297	CP/Ac,A'c-17	S = A' = Rub; used $k_t^{A'} = 1.4 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 0.5$.	747207
	CH ₃ CN	1.7 × 10 ⁸	2 × 10 ⁻⁴		CP/Ac-15	S = MB; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	777113
	CHCl ₃	2.9 × 10 ⁷			PL/Ld-2	S = MB.	93A050
	CHCl ₃	3.2 × 10 ⁷	5.2 × 10 ⁻⁴		CP/Pa-14	S = 1,5-Diamino-4,8-dihydroxyanthraquinone; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; P = 9,10-Dimethylanthracene endoperoxide.	80E446
	CHCl ₃	9.3 × 10 ⁷	2 × 10 ⁻⁴		CP/Ac-15	S = MB; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	777113
	CHCl ₃	2.7 × 10 ⁷ (k_t)		297	CP/Ac,A'c-17	S = A' = Rub; used $k_t^{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 0.64$.	747207
	CS ₂	1.6 × 10 ⁷		295	MP/Ld-2	S = TPP.	89A400
	CS ₂		2 × 10 ⁻⁶	295	CP/Ac-14	S = A.	89A400
	CS ₂	1.3 × 10 ⁷ (k_t)		297	CP/Ac,A'c-17	S = A' = Rub; used $k_t^{A'} = 2.5 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 0.5$.	747207
	ClCF ₂ CCl ₂ F		1 × 10 ⁻⁶	295	CP/Ac-14	S = A; [DMA] = (4-11) × 10 ⁻⁵ mol L ⁻¹ .	89A400
	ClCF ₂ CCl ₂ F		2 × 10 ⁻⁵	295	CP/Ac-14	S = A; [DMA] = (3-9) × 10 ⁻⁶ mol L ⁻¹ .	89A400
	D ₂ O (mic)	7.4 × 10 ⁸	6.8 × 10 ⁻⁵	298	CP/Ac-15	S = MB; used $k_d = 5.0 \times 10^4$ s ⁻¹ ; A solubilized in DTAC micelles.	78F061
	D ₂ O (ves)	3.1 × 10 ⁸			CP/Ac-14	S = RB; used $k_d = 1.8 \times 10^4$ s ⁻¹ ; used ϕ_{isc} (RB) = 0.76; 0.05% egg yolk lecithin.	86N104
	EtOH	4.6 × 10 ⁷		293	CP/Ac-15	S = MB; used $k_d = 8.3 \times 10^4$ s ⁻¹ .	85N293
	EtOH	3.9 × 10 ⁷	2.1 × 10 ⁻³	298	CP/Ac-15	S = MB; used $k_d = 8.3 \times 10^4$ s ⁻¹ .	78F061
	EtOH	4.4 × 10 ⁷	1.9 × 10 ⁻³	298	CP/Ac-15	S = Eos; used $k_d = 8.3 \times 10^4$ s ⁻¹ .	78F061
	EtOH	1.2 × 10 ⁷	7.2 × 10 ⁻³	297	CP/Ac-15	S = A; used $k_d = 8.3 \times 10^4$ s ⁻¹ .	747207
	EtOH/H ₂ O (50:50)	2.0 × 10 ⁸		293	CP/Ac-15	S = MB; used $k_d = 2 \times 10^5$ s ⁻¹ .	85N293
	H ₂ O (mic)	5 × 10 ⁷		293	CP/Ac-15	S = MB; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; 0.015 mol L ⁻¹ CTAB.	85N293

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.10 Anthracene, 9,10-dimethyl- (DMA) — Continued							
	H ₂ O (mic)	5.4 × 10 ⁷		293	CP/Ac-15	S = MB; used $k_d = 2.4 \times 10^5$ s ⁻¹ ; 0.1 mol L ⁻¹ SDS.	85N293
	H ₂ O (mic)	7.5 × 10 ⁸	6.6 × 10 ⁻⁴	298	CP/Ac-15	S = MB; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; A solubilized in DTAC micelles.	78F061
	H ₂ O (mic)	9.1 × 10 ⁸	5.5 × 10 ⁻⁴	298	CP/Ac-15	S = Eos; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; S and A solubilized in DTAC micelles.	78F061
	H ₂ O (ves)	3.2 × 10 ⁸			CP/Ac-14	S = RB; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; used ϕ_{isc} (RB) = 0.76; 0.05% egg yolk lecithin.	86N104
	MeOH	6.9 × 10 ⁷		293	CP/Ac-15	S = MB; used $k_d = 1 \times 10^5$ s ⁻¹ .	85N293
	MeOH	4.8 × 10 ⁷	3 × 10 ⁻³		CP/Ac-15	S = MB; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	77F113
	MeOH	2.4 × 10 ⁷	8.0 × 10 ⁻³	297	CP/Ac-15	S = A; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	747207
	MeOH/H ₂ O (80:20)	2.0 × 10 ⁸		293	CP/Ac-15	S = MB; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	85N293
3.11 Anthracene, 9,10-diphenyl- (DPA)							
	C ₆ H ₅ N				CP/Ac-17	S = A; A' = TME; meas. $k_f/k_t^{A'} = 0.18$.	66F041
	C ₆ H ₅ CH ₃	7.8 × 10 ⁵			PL/Ld-2	S = A.	93F207
	C ₆ H ₅ CH ₃		0.031	293	CP/Ac-16	S = A' = Rub.	83F335
	C ₆ H ₆	1.2 × 10 ⁶	0.04	298	CP/Ac-20	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $\beta_{A'} = 1.0 \times 10^{-3}$ mol L ⁻¹ .	747312
	C ₆ H ₆		0.040	298	CP/Ac-15	S = A.	727196
	C ₆ H ₆		0.060		CP/Ac-20	S = ZnTPP; A' = (C ₂ H ₅) ₂ S.	71F580
	C ₆ H ₆		0.045		CP/Ac-15	S = A.	59F003
	C ₆ H ₆		0.057		CP/Ac-15	S = A.	55F004
	CCl ₄	6 × 10 ⁵			CP/Ac-14	S = TPP; used $k_d = 1.4 \times 10^3$ s ⁻¹ .	80C002
	CCl ₄		6.0 × 10 ⁻⁴		CP/Oc-15	S = A.	537004
	CH ₂ Cl ₂	4.2 × 10 ⁶ (k_p) 8 × 10 ⁶ (k_q)	1 × 10 ⁻³		CR/Ac-31	used $k_d = 1.2 \times 10^4$ s ⁻¹ , ${}^1\text{O}_2^*$ from triphenyl phosphite ozonide; $f_r^{A'} = 0.3$.	85M406
	CHCl ₃	3.0 × 10 ⁶			CP/Ac-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005
	CHCl ₃		4.0 × 10 ⁻³		CP/Oc-15	S = A.	537004
	CS ₂		5 × 10 ⁻⁵	295	CP/Ac-14	S = A.	89A400
	CS ₂	8.5 × 10 ⁵		295	MP/Ld-2	S = TPP.	89A400
	CS ₂		3.2 × 10 ⁻³		CP/Oc-15	S = A.	537004
	ClCF ₂ CCl ₂ F		1 × 10 ⁻⁴	295	CP/Ac-14	S = A.	89A400
3.12 Anthracene, 9-methoxy-							
	C ₆ H ₆	2.5 × 10 ⁶ (k_p)		298	CP/Ac,A'c-17	S = A' = Tetr; used $k_f^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_f/k_t^{A'} = 0.21$.	747312
3.13 Anthracene, 9-methyl-							
	C ₆ H ₆	2.3 × 10 ⁶		293	CP/Ac-15	S = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	85N293
	C ₆ H ₆	3.2 × 10 ⁶	0.01	298	CP/Ac-20	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $\beta_{A'} = 1.0 \times 10^{-3}$ mol L ⁻¹ .	747312
	CCl ₄		3.3 × 10 ⁻³		CP/Oc-15	S = A.	537004
	CHCl ₃	8.1 × 10 ⁶			CP/Ac-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78A005
	CHCl ₃		1.2 × 10 ⁻³		CP/Oc-15	S = A.	537004
	EtOH	2 × 10 ⁶		293	CP/Ac-15	S = MB; used $k_d = 8.3 \times 10^4$ s ⁻¹ .	85N293
	EtOH/H ₂ O (50:50)	1.7 × 10 ⁷		293	CP/Ac-15	S = MB; used $k_d = 2 \times 10^5$ s ⁻¹ .	85N293
	H ₂ O	3.8 × 10 ⁷		293	CP/Ac-15	S = MB; 0.068 mol L ⁻¹ PEO; k_d not given.	85N293

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.13 Anthracene, 9-methyl- — Continued							
	H ₂ O	9.6×10^7		293	CP/Ac-15	S = MB; used $k_d = 3 \times 10^5$ s ⁻¹ .	85N293
	H ₂ O (mic)	1.7×10^7		293	CP/Ac-15	S = MB; used $k_d = 2.4 \times 10^5$ s ⁻¹ ; 0.1 mol L ⁻¹ SDS.	85N293
	H ₂ O (mic)	1.4×10^7		293	CP/Ac-15	S = MB; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; 0.015 mol L ⁻¹ CTAB.	85N293
	MeOH	2×10^6		293	CP/Ac-15	S = MB; used $k_d = 1 \times 10^5$ s ⁻¹ .	85N293
3.14 Anthracene, 9-phenyl-							
	C ₆ H ₆	4.2×10^5	0.10	298	CP/A'c-20	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $\beta_{A'} = 1.0 \times 10^{-3}$ mol L ⁻¹ .	747312
	CCl ₄		7.4×10^{-4}		CP/Oc-15	S = A.	537004
	CHCl ₃		7.2×10^{-3}		CP/Oc-15	S = A.	537004
	CS ₂		8.4×10^{-4}		CP/Oc-15	S = A.	537004
3.15 9,10-Anthracenediethanesulfonate ion (AES)							
	D ₂ O/ H ₂ O (80:20) pH = 7.4		1.1×10^{-3}	310	CR/Pa-31	${}^1\text{O}_2^*$ from NDPO ₂ .	89M097
	H ₂ O pH = 7.4		7.7×10^{-3}	310	CR/Pa-31	${}^1\text{O}_2^*$ from NDPO ₂ .	89M097
	H ₂ O pH = 7.4		8×10^{-3}	310	CR/Pa-31	${}^1\text{O}_2^*$ from NaOCl/H ₂ O ₂ .	89M097
	H ₂ O	4.7×10^7 (k_p)			CR/Ac,Pa-31	${}^1\text{O}_2^*$ from H ₂ O ₂ /OCl ⁻ ; counter ion Na ⁺ ; meas. $k_q/k_t = 0.35$; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	85A176
	H ₂ O/ D ₂ O (62:38) pH = 7.4		3×10^{-3}	310	CR/Pa-31	${}^1\text{O}_2^*$ from NaOCl/H ₂ O ₂ .	89M097
3.16 9,10-Anthracenedipropionate ion (ADPA)							
	D ₂ O	8.2×10^7			PL/Ad-5	S = MB.	80A205
	H ₂ O pH = 6.8	1×10^8		295	CP/Ac-14	S = Ru(bpy) ₃ ²⁺ ; used $k_d = 2.5 \times 10^4$ s ⁻¹ .	92A095
3.17 1,5-Anthracenedisulfonate ion							
	H ₂ O	7.0×10^6	0.070	303	CP/Ac-15	S = A; used $k_d = 5.0 \times 10^5$ s ⁻¹ .	78A275
	H ₂ O	$\leq 10^7$		301	CP/P'a-23	S = A; A' = KI; Q = NaN ₃ ; P' = I ₃ ⁻ . No measurable effect.	777074
3.18 1-Anthracenesulfonate ion							
	H ₂ O	5.4×10^7	9.2×10^{-3}	303	CP/Ac-15	S = A; used $k_d = 5.0 \times 10^5$ s ⁻¹ .	78A275
	H ₂ O	5.0×10^8		301	CP/P'a-23	S = A; A' = KI; Q = NaN ₃ ; used $k_d = 5.0 \times 10^5$ s ⁻¹ , $k_Q = 2.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; P' = I ₃ ⁻ .	777074
3.19 2-Anthracenesulfonate ion							
	H ₂ O	4.5×10^7	0.011	303	CP/Ac-15	S = A; used $k_d = 5.0 \times 10^5$ s ⁻¹ .	78A275
	H ₂ O	3.0×10^8		301	CP/P'a-23	S = A; A' = KI; Q = NaN ₃ ; used $k_d = 5.0 \times 10^5$ s ⁻¹ , $k_Q = 2.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; P' = I ₃ ⁻ .	777074
3.20 Anthra[1,9-<i>bc</i>:4,10-<i>b'c'</i>]dichromene							
	C ₆ H ₅ CH ₃	5×10^8	1×10^{-4}		CP/Ac-15	S = A; used $k_d = 5 \times 10^4$ s ⁻¹ .	84F197 82F068
	CS ₂	3.4×10^8		295	MP/Ld-2	S = TPP.	89A400
3.21 9-Anthroate ion							
	H ₂ O	2.0×10^7			PL/P'a-16	S = Ery; A' = TMPD; used $k_d = 3.3 \times 10^5$ s ⁻¹ ; P' = [TMPD] ⁺ .	82A080

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.22	Benzamide, 4-(bromomethyl)-N-(diphenylmethyl)- CH ₂ Cl ₂	<10 ⁵			PL/Ld-2	S = MPDME.	91E134
3.23	Benz[a]anthracene C ₆ H ₆	4.8 × 10 ⁴ (k_t)		298	CP/Ac,A'c-17	S = A' = An; used $k_t^{A'} = 1.6 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 0.30$.	74F312
3.24	Benz[a]anthracene, 9,10-dimethyl- C ₆ H ₆	1.6 × 10 ⁻³		298	CP/Ac-?	S = A.	76F422
	C ₆ H ₆	1.4 × 10 ⁷ (k_t)		298	CP/Ac,A'c-17	S = A' = Rub; used $k_t^{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 0.33$.	74F312
	C ₆ H ₆	7.1 × 10 ⁻⁴		298	CP/Ac-15	S = A or Tetr.	70F079
	C ₆ H ₆	8.3 × 10 ⁻⁴		298	CP/Ac-15	S = Rub.	70F079
	C ₆ H ₆	7.4 × 10 ⁻⁴		298	CP/Ac-?	S = Per and An.	69F388
	H ₂ O (mic)	2.5 × 10 ⁹	4.0 × 10 ⁻⁴		CP/Ac-15	S = MB; used $k_d = 1 \times 10^6$ s ⁻¹ ; A solubilized in CTAB micelles.	78N003
3.25	Benzanthrone EtOH	9.6 × 10 ⁸			CP/Ac-14	S = A; used $k_d = 6.7 \times 10^4$ s ⁻¹ .	92D048
3.26	Benzene C ₆ D ₆	2.7 × 10 ³			PL/Ld-2	S = TPP.	83E398
	CCl ₄	3.9 × 10 ³		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329
	CCl ₄	2 × 10 ³			PL/A'd-8	S = MB; A' = DPBF.	76F903
	CCl ₄ / MeOH (96:4)	5 × 10 ²			PL/A'd-8	S = MB; A' = DPBF.	77F162
3.27	Benzene-d ₆ CCl ₄	1.3 × 10 ²		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329
3.28	Benzene, 1,5-bis(1,1-dimethylethyl)-2,4-dimethoxy- MeOH				CP/Pa,P'a-17	S = RB; A' = TME; meas. $k_t/k_t^{A'} = 2.3 \times 10^{-3}$.	70F454
	MeOH				CR/Pa,P'a-17	A' = TME; meas. $k_t/k_t^{A'} = 3.0 \times 10^{-3}$; ${}^1\text{O}_2^*$ from H ₂ O ₂ /NaOCl.	70F454
3.29	Benzene, 1,3-bis(1,1-dimethylethyl)-2-methoxy-5-methyl- MeOH	5.9 × 10 ⁵	0.20		CP/A'c-16	S = MB; A' = DPF; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	78A171
3.30	Benzene, bromo- CCl ₄	2 × 10 ³			PL/A'd-8	S = MB; A' = DPBF.	76F903
3.31	Benzene, chloro- CCl ₄ / MeOH (98:2)	2.1 × 10 ³		293	PL/A'd-8	S = MB; A' = DPBF.	83A371
3.32	Benzene, 2-chloro-1,4-dimethoxy- CH ₃ CN	4.7 × 10 ⁵			PL/Ld-2	S = RB.	91A341
	dioxane	<1 × 10 ⁴			PL/Ld-2	S = ZnTPP.	91A341
3.33	Benzene, 5-chloro-1,3-dimethoxy- CH ₃ CN	5.6 × 10 ⁵			PL/Ld-2	S = RB.	91A341
	dioxane	6.1 × 10 ⁵			PL/Ld-2	S = ZnTPP.	91A341
3.34	Benzene, 1-chloro-2-methoxy- MeOH	<10 ⁴			CP/Ac-23	S = A' = Rub; used $k_d = 1.4 \times 10^5$ s ⁻¹ , $k_{A'} = 5 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	90F518

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.15	Benzene, 1,3-dichloro-						
	C ₆ H ₆ / MeOH (60:40)	<10 ⁴			CP/A'c-33	S = A' = Rub; k_d and $k_{A'}$ not given; 6 × 10 ⁻³ mol L ⁻¹ KOH.	88F650
3.16	Benzene, 1,4-dichloro-						
	C ₆ H ₆ / MeOH (60:40)	<10 ⁴			CP/A'c-33	S = A' = Rub; k_d and $k_{A'}$ not given; 6 × 10 ⁻³ mol L ⁻¹ KOH.	88F650
3.17	Benzene, 2,6-dichloro-1-methoxy-						
	C ₆ H ₆ / MeOH (60:40)	<10 ⁴			CP/A'c-33	S = A' = Rub; k_d and $k_{A'}$ not given; 6 × 10 ⁻³ mol L ⁻¹ KOH.	88F650
3.18	Benzene, 1,2-dimethoxy-						
	MeOH	≤5 × 10 ⁶ (k_r)		293	CP/Ac,A'c-17	S = RB and MB; A' = 1,2,4,5-C ₆ H ₂ (CH ₃ O) ₄ ; No measurable effect.	72A020
3.19	Benzene, 1,3-dimethoxy-						
	MeOH	≤5 × 10 ⁶ (k_r)		293	CP/Ac,A'c-17	S = MB and RB; A' = 1,2,4,5-C ₆ H ₂ (CH ₃ O) ₄ ; No measurable effect.	72A020
3.20	Benzene, 1,4-dimethoxy-						
	MeOH	≤5 × 10 ⁶ (k_r)		293	CP/Ac,A'c-17	S = MB and RB; A' = 1,2,4,5-C ₆ H ₂ (CH ₃ O) ₄ ; No measurable effect.	72A020
3.21	Benzene, ethyl-						
	CCl ₄	1.3 × 10 ⁵		296	CR/LI-12	used $k_d = 1.7 \times 10^3$ s ⁻¹ ; ¹ O ₂ * from DMNO ₂ .	90F069
	CCl ₄ / MeOH (98:2)	3.6 × 10 ³		293	PL/A'd-8	S = MB; A' = DPBF.	83A371
	CCl ₄ / MeOH (96:4)	5 × 10 ²			PL/A'd-8	S = MB; A' = DPBF.	777162
3.22	Benzene, hexamethoxy-						
	MeOH			293	CP/Ac,A'c-17	S = MB and RB; A' = 1,2,4,5-C ₆ H ₂ (CH ₃ O) ₄ ; meas. $k_r/k_r^{A'} = 0.031$.	72A020
3.23	Benzene, methoxy-						
	C ₆ H ₆ / MeOH (60:40)	<10 ⁴			CP/A'c-33	S = A' = Rub; k_d and $k_{A'}$ not given; 6 × 10 ⁻³ mol L ⁻¹ KOH.	88F650
	CCl ₄	2.0 × 10 ⁵		296	CR/LI-12	used $k_d = 1.7 \times 10^3$ s ⁻¹ ; ¹ O ₂ * from DMNO ₂ .	90F069
	MeOH	≤5 × 10 ⁶ (k_r)		293	CP/Ac,A'c-17	S = RB and MB; A' = 1,2,4,5-C ₆ H ₂ (CH ₃ O) ₄ ; No measurable effect.	72A020
3.24	Benzene, 1-methoxy-2-nitro-						
	MeOH/ C ₆ H ₆ (60:40)	<10 ³			CP/Ac-23	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 5 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	90F518
3.25	Benzene, (3-methyl-2-butenyl)-						
	MeOH	0.15		293	CP/Oc-15	S = RB; $E_a = 9.6$ kJ mol ⁻¹ .	68F288
	MeOH	0.13			?	Method not given.	627005
3.26	Benzene, 1,1'-(2-methyl-1-propenylidene)bis-						
	CH ₃ CN	5.0 × 10 ⁶			PL/Ld-2	S = (C ₆ H ₅) ₂ CO.	91E697
3.27	Benzene, nitro-						
	CH ₂ Cl ₂	5.0 × 10 ³			PL/Ld-2	S = MPDME.	91E134
3.28	Benzene, pentamethoxy-						
	MeOH			293	CP/Ac,A'c-17	S = MB and RB; A' = 1,2,4,5-C ₆ H ₂ (CH ₃ O) ₄ ; meas. $k_r/k_r^{A'} = 0.19$.	72A020

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.49	Benzene, 1,2,3,4-tetramethoxy-						
	MeOH	$\leq 5 \times 10^6 (k_r)$		293	CP/Ac,A'c-17	S = MB and RB; A' = 1,2,4,5-C ₆ H ₂ (CH ₃ O) ₄ ; No measurable effect.	72A020
3.50	Benzene, 1,2,3,5-tetramethoxy-						
	MeOH			293	CP/Ac,A'c-17	S = MB and RB; A' = 1,2,4,5-C ₆ H ₂ (CH ₃ O) ₄ ; meas. $k_r/k_r^{A'} = 0.15$.	72A020
3.51	Benzene, 1,2,4,5-tetramethoxy-						
	MeOH			293	CP/Ac,A'c-17	S = MB and RB; A' = 1,2,4-C ₆ H ₃ (OCH ₃) ₃ ; meas. $k_r/k_r^{A'} = 12$.	72A020
3.52	Benzene, 1,2,3-trimethoxy-						
	MeOH	$\leq 5 \times 10^6 (k_r)$		293	CP/Ac,A'c-17	S = MB and RB; A' = 1,2,4,5-C ₆ H ₂ (CH ₃ O) ₄ ; No measurable effect.	72A020
3.53	Benzene, 1,2,4-trimethoxy-						
	MeOH	1.8×10^7	6.4×10^{-3}		CP/A'c-16	S = MB; A' = DPF; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	78A171
3.54	Benzene, 1,3,5-trimethoxy-						
	MeOH	$\leq 5 \times 10^6 (k_r)$		293	CP/Ac,A'c-17	S = MB and RB; A' = 1,2,4,5-C ₆ H ₂ (CH ₃ O) ₄ ; No measurable effect.	72A020
3.55	Benzinemethanol, α -methyl-						
	CCl ₄ / MeOH (98:2)	4.1×10^3		293	PL/A'd-8	S = MB; A' = DPBF.	83A371
3.56	Benzo[1,2,3- <i>k</i> :4,5,6- <i>k'</i>]dixanthene (BDX)						
	C ₆ H ₅ CH ₃	1.0×10^8	5.1×10^{-4}		CP/Ac-15	S = A; used $k_d = 5 \times 10^4$ s ⁻¹ .	84F197 82F068
	CS ₂	7.2×10^7		295	MP/Ld-2	S = TPP.	89A400
3.57	Benzoic acid, methyl ester						
	CCl ₄	3×10^3			PL/A'd-8	S = MB; A' = DPBF.	76F903
3.58	Benzophenone (BP)						
	C ₆ H ₆	$< 3 \times 10^4 (k_r)$			CP/A'c-17	S = A' = Rub; $k_A[A']/(k_A[A'] + k[BP] > 0.95$ when [BP] = 0.1 mol L ⁻¹ and [A'] = 5×10^{-4} mol L ⁻¹ .	81F364
	ClCF ₂ CCl ₂ F	4×10^3		298	FP/Ld-2	S = Per.	82A322
3.59	Biphenyl, 2-methoxy-						
	EtOH	2.3×10^6			PL/Ld-2	S = RB; 7.5×10^{-3} mol L ⁻¹ KOH.	93E536
3.60	Cyclopentadienone, tetraphenyl-						
	Dibutyl terephthalate				CP/Ac-17	S = N-[2-[(2-Bromo-4,6-dinitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)amino]-4-methoxyphenyl]acetamide; A' = BRH ₂ ; meas. $k_r/k_r^{A'} = 1.0$.	81F609
	C ₆ H ₅ CHOHCH ₃				CP/Ac-17	S = N-[2-[(2-Bromo-4,6-dinitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)amino]-4-methoxyphenyl]acetamide; A' = BRH ₂ ; meas. $k_r/k_r^{A'} = 0.6$.	81F609
	C ₆ H ₆	$1 \times 10^8 (k_r)$ $2.2 \times 10^8 (k_q)$	1.0×10^{-4}	298	CP/Ac-15,27	S = Cor; used $k_d = 3.7 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r = 1.6$.	82A179
	C ₆ H ₆			298	CP/Ac-17	S = A' = DPBF; meas. $k_r/k_r^{A'} = 3.8$.	82A179
	CH ₂ Cl ₂	$5 \times 10^7 (k_r)$ $6 \times 10^7 (k_q)$	1×10^{-4}		CR/Ac-31	used $k_d = 1.2 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from triphenyl phosphite ozonide; $f_r^A = 0.46$.	85M406
	CH ₃ CN	$6 \times 10^7 (k_r)$ $4 \times 10^8 (k_q)$		298	CP/Ac-15,27	S = Cor; used $k_d = 1.8 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r = 6.2$.	82A179

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.60 Cyclopentadienone, tetraphenyl- — Continued							
	CHCl ₃				CP/Ac-17	S = N-[2-[(2-Bromo-4,6-dinitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)amino]-4-methoxyphenyl]acetamide; A' = BRH ₂ ; meas. $k_r/k_r^A = <>0.06 \times k_r^A$ in dibutyl terephthalate.	81F609
3.61 Dibenz[a,h]anthracene							
	c-C ₆ H ₁₂	$9.5 \times 10^3 (k_r)$		298	CP/Ac,A'c-17	S = A' = An; used $k_r^A = 1.6 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^A = 0.59$.	747312
3.62 Dibenzo[fg,uv]naphtho[2,3-c]pentaphene-9,19-dione, 1,2,3,4,11,12,13,14-octahydro-							
	C ₆ H ₆	4.3×10^8	9.7×10^{-5}	298	CP/Ac-14	S = A; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	93F128
3.63 Dibenz[a,j]perylene							
	C ₆ H ₆	$>2 \times 10^8$			CP/Ac-14	S = A; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	79F473
3.64 Dibenzo[a,o]perylene, 7,16-diphenyl- (Mesodiphenylhelianthrene, MDH)							
	n-C ₅ H ₁₂	4×10^9	7×10^{-6}		CP/Pa-14	S = A; used $k_d = 2.9 \times 10^4$ s ⁻¹ .	87F480
	c-C ₆ H ₁₂	7.5×10^9	6.7×10^{-6}		CP/Ac-23	S = A; Q = Car; used $k_d = 5.0 \times 10^4$ s ⁻¹ .	83F406
	C ₆ H ₅ CH ₂ OH	2.6×10^9 $8 \times 10^8 (k_r)$	3×10^{-5}		CP/Pa-14	S = MB; used $k_d = 7.1 \times 10^4$ s ⁻¹ ; meas. $f_r^A = 0.35$.	87F479
	C ₆ H ₅ CH ₃	7.3×10^9	5.5×10^{-6}		CP/Ac-23	S = A; Q = Car; used $k_d = 4.0 \times 10^4$ s ⁻¹ .	83F406
	C ₆ H ₅ CH ₃ / MeOH (99:1)	5×10^9 $1.5 \times 10^9 (k_r)$	8.2×10^{-6}		CP/Pa-14	S = MB; used $k_d = 4.0 \times 10^4$ s ⁻¹ ; meas. $f_r^A = 0.29$.	87F479
	CH ₃ COCH ₃	5×10^9 $1.4 \times 10^9 (k_r)$	5.2×10^{-6}		CP/Pa-14	S = MB; used $k_d = 2.5 \times 10^4$ s ⁻¹ .	87F479
	CH ₃ COCH ₃	5.7×10^9	3.9×10^{-6}		CP/Ac-23	S = A; Q = Car; used $k_d = 2.2 \times 10^4$ s ⁻¹ .	83F406
	CHCl ₃	3.4×10^9	1.2×10^{-6}		CP/Ac-23	S = A; Q = Car; used $k_d = 4.1 \times 10^3$ s ⁻¹ .	83F406
	CS ₂	2.0×10^9		295	MP/Ld-2	S = TPP.	89A400
	CS ₂	1.1×10^9	6×10^{-7}		CP/Ac-23	S = A; Q = Car; used $k_d = 1.1 \times 10^3$ s ⁻¹ .	83F406
	EtOH	7.0×10^9		293	PL/Ad,Pb-5	S = An.	87F541
	MeOH/ C ₆ H ₅ CH ₃ (97:3)	4×10^9 $1.3 \times 10^9 (k_r)$	2.6×10^{-5}		CP/Pa-14	S = MB; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; meas. $f_r^A = 0.27$.	87F479
3.65 Diindeno[5,6-a:5',6'-j]perylene-8,17-dione, 1,2,3,10,11,12-hexahydro-							
	C ₆ H ₆	5.1×10^8	8.2×10^{-5}	298	CP/Ac-14	S = A; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	93F128
3.66 Fluorene, 9-(fluoren-9-ylidene)-							
	CH ₂ Cl ₂				CP/Ac,A'c-17	S = MB; A' = TME; meas. $k_r/k_r^A = 0.034$.	70F200
	dioxane/ MeOH (80:20)				CR/Ac,A'c-17	A' = TME; meas. $k_r/k_r^A = 0.034$; ${}^1\text{O}_2^*$ from H ₂ O ₂ /NaOCl.	70F200
3.67 Helianthrene (Dibenzo[a,o]perylene)							
	n-C ₅ H ₁₂	4×10^9	8×10^{-6}		CP/Pa-14	S = A; used $k_d = 2.9 \times 10^4$ s ⁻¹ .	87F480
	C ₆ H ₅ CH ₂ OH	$2 \times 10^9 (k_r)$	$3 \times 10^{-5} (\beta_r)$		CP/Pa-14	S = MB; used $k_d = 7.1 \times 10^4$ s ⁻¹ ; meas. $f_r^A = 0.99$.	87F479
	C ₆ H ₅ CH ₃ / MeOH (99:1)	$4.7 \times 10^9 (k_r)$	$8.6 \times 10^{-6} (\beta_r)$		CP/Pa-14	S = MB; used $k_d = 4.0 \times 10^4$ s ⁻¹ ; meas. $f_r^A = -1.0$.	87F479

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.67 Helianthrene (Dibenzo[<i>a,o</i>]perylene) — Continued							
	CH ₃ COCH ₃	4.6×10^9 (k_r)	5.5×10^{-6} (β_r)		CP/Pa-14	S = MB; used $k_d = 2.5 \times 10^4$ s ⁻¹ ; $f_r^A = -1.0$.	87F479
	MeOH/ C ₆ H ₅ CH ₃ (97:3)	5.1×10^9 (k_r)	2.2×10^{-5} (β_r)		CP/Pa-14	S = MB; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; meas. $f_r^A = -1.0$.	87F479
3.68 Heterocoerdianthrone (Dibenzoperylene-3,9-quinone, HCD)							
	1,2,4-C ₆ H ₃ Cl ₃	4×10^7	5×10^{-4}	298	CP/Ac-15	S = A; k estimated using $k_d(C_6H_5Br) = 2 \times 10^4$ s ⁻¹ .	767570
	C ₆ H ₅ CH ₃	6.8×10^7	5.9×10^{-4}		CP/Ac-15	S = A; used $k_d = 4.0 \times 10^4$ s ⁻¹ .	80F151
	C ₆ H ₆	3.6×10^8	1.2×10^{-4}	298	CP/Ac-14	S = A; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	93F128
	C ₆ H ₆	7×10^7	6×10^{-4}	298	CP/Ac-15	S = A; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	767570 777176
	CCl ₄	7×10^7	2×10^{-5}	298	CP/Ac-15	S = A; used $k_d = 1.4 \times 10^3$ s ⁻¹ .	767570
	CHCl ₃	7×10^7	3×10^{-4}	298	CP/Ac-15	S = A; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	767570
	CS ₂	6.6×10^7		295	MP/Ld-2	S = TPP.	89A400
	CS ₂	8×10^7	6×10^{-5}	298	CP/Ac-15	S = A; used $k_d = 5.0 \times 10^3$ s ⁻¹ .	767570 777176
3.69 Indene							
	MeOH		2.0	293	CP/Oc-15	S = MB; $E_a = 11$ kJ mol ⁻¹ .	68F288
	MeOH		1.5	293	CP/Oc-15	S = RB; $E_a = 13$ kJ mol ⁻¹ .	68F288
3.70 Indene, 1,2-diphenyl-							
	CH ₃ COCH ₃	8.2×10^5		248	PL/Ld-2	S = MPDME.	83F195
		8.4×10^5		224			
3.71 Indene, 1-methyl-2-phenyl-							
	CH ₃ COCH ₃	2×10^6		298	PL/Ld-2	S = MPDME.	83F195
		1.0×10^6		223			
	CH ₃ COCH ₃	2×10^5 (k_r)		298	CP/Ac-17	S = RB; $A' = 2M2P$; used $k_r^{A'} = 8.1 \times 10^5$ L mol ⁻¹ s ⁻¹ .	83F195
		1×10^6 (k_r)		195			
3.72 1,4-Methanonaphthalene, 1,2,3,4-tetrahydro-							
	CH ₂ Cl ₂	3.9×10^4			PR/A'd-8	S = An; $A' = DPBF$.	79A457
3.73 1,4-Methanonaphthalene, 1,2,3,4-tetrahydro-5,8-dimethoxy-							
	C ₆ H ₆ / MeOH (75:25)	6×10^6		298	CP/A'c-20	S = RB; $A' = c\text{-}C_6H_{10}$; used $k_d = 6.3 \times 10^4$ s ⁻¹ , $\beta_{A'} = 3.1$ mol L ⁻¹ .	78F586
	CH ₂ Cl ₂	1.3×10^7			PR/A'd-8	S = An; $A' = DPBF$.	79A457
3.74 Naphthalene							
	1-BuOH	5.2×10^8	1.0×10^{-4}	293	CP/Ac-15	S = MB; used $k_d = 5.2 \times 10^4$ s ⁻¹ .	78A266
3.75 Naphthalene, 1,2-dimethyl-							
	CH ₃ CN			288	CP/Oc-17	S = MB; $A' = 1,5\text{-}Np(CH_3)_2$; meas. $k_r/k_r^{A'} = 3.3$.	80F204
3.76 Naphthalene, 1,3-dimethyl-							
	CH ₃ CN			288	CP/Oc-17	S = MB; $A' = 1,5\text{-}Np(CH_3)_2$; meas. $k_r/k_r^{A'} = 1.0$.	80F204
3.77 Naphthalene, 1,4-dimethyl-							
	c-C ₆ H ₁₂	1.2×10^4 (k_r)		298	CP/Ac,A'c-17	S = A' = An; used $k_r^{A'} = 1.6 \times 10^5$ L mol ⁻¹ s ⁻¹ , meas. $k_r/k_r^{A'} = 0.075$.	747312
	CH ₃ CN			288	CP/Oc-17	S = MB; $A' = 1,5\text{-}Np(CH_3)_2$; meas. $k_r/k_r^{A'} = 10$.	80F204
	EtOH/ H ₂ O (86:14)	1.3×10^5 (k_r)		278	CP/A'c-14,27	S = RB; $A' = \alpha\text{-Terpinene}$; used $k_d = 8.8 \times 10^4$ s ⁻¹ .	86A216
	MeOH	4.2×10^4 (k_r)		278	CP/A'c-14,27	S = RB; $A' = \alpha\text{-Terpinene}$; used $k_d = 9.5 \times 10^4$ s ⁻¹ .	86A216

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.77	Naphthalene, 1,4-dimethyl-	— Continued					
	MeOH/H ₂ O (80:20)	$3.7 \times 10^5 (k_r)$		278	CP/A'c-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.3 \times 10^5$ s ⁻¹ .	86A216
	1-PrOH/H ₂ O (73:27)	$2.4 \times 10^5 (k_r)$		278	CP/A'c-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	86A216
3.78	Naphthalene, 1,6-dimethyl-						
	CH ₃ CN			288	CP/Oc-17	S = MB; A' = 1,5-Np(CH ₃) ₂ ; meas. $k_r/k_r^{A'} = 0.52$.	80F204
3.79	Naphthalene, 1,7-dimethyl-						
	CH ₃ CN			288	CP/Oc-17	S = MB; A' = 1,5-Np(CH ₃) ₂ ; meas. $k_r/k_r^{A'} = 0.54$.	80F204
3.80	Naphthalene, 1,8-dimethyl-						
	CH ₃ CN			288	CP/Oc-17	S = MB; A' = 1,5-Np(CH ₃) ₂ ; meas. $k_r/k_r^{A'} = 4.4$.	80F204
3.81	Naphthalene, 2,3-dimethyl-						
	CH ₃ CN			288	CP/Oc-17	S = MB; A' = 1,5-Np(CH ₃) ₂ ; meas. $k_r/k_r^{A'} = 0.22$.	80F204
3.82	Naphthalene, 2,6-dimethyl-						
	CH ₃ CN			288	CP/Oc-17	S = MB; A' = 1,5-Np(CH ₃) ₂ ; meas. $k_r/k_r^{A'} = 0.12$.	80F204
3.83	Naphthalene, 2,7-dimethyl-						
	CH ₃ CN			288	CP/Oc-17	S = MB; A' = 1,5-Np(CH ₃) ₂ ; meas. $k_r/k_r^{A'} = 0.12$.	80F204
3.84	Naphthalene, 1,1'-(1,2-dimethyl-1,2-diethenyl)bis-, (Z)-						
	CCl ₄			CP/Ac,A''c,Pa- 17	CP/Ac,A''c,Pa- 17	S = TPP; A' = (E)-2,3-Di(α -naphthyl)-2-butene; meas. $k_r/k_r^{A'} = 9.4$; A'' = TME.	79F051
3.85	Naphthalene, 2,2'-(1,2-dimethyl-1,2-diethenyl)bis-, (Z)-						
	CCl ₄			CP/Ac,A''c,Pa- 17	CP/Ac,A''c,Pa- 17	S = TPP; A' = (E)-2,3-Di(β -naphthyl)-2-butene; meas. $k_r/k_r^{A'} = 9.1$; A'' = TME.	79F051
3.86	Naphthalene, 1-methyl-						
	CH ₃ CN			288	CP/Oc-17	S = MB; A' = 1,5-Np(CH ₃) ₂ ; meas. $k_r/k_r^{A'} = 0.39$.	80F204
3.87	Naphthalene, 1-(2-methylpropenyl)-						
	C ₆ H ₆	1.9		277	CP/Oc-14	S = TPP.	81F582
	CCl ₄	0.14		277	CP/Oc-14	S = TPP.	81F582
	CH ₃ COCH ₂ CH ₃	1.0		277	CP/Oc-14	S = RB.	81F582
	CHCl ₃	0.23		277	CP/Oc-14	S = TPP.	81F582
	EtOH	3.3		277	CP/Oc-14	S = RB.	81F582
3.88	1,4-Naphthalenedipropionate ion (NDP)						
	EtOH/H ₂ O (86:14)	$6.0 \times 10^4 (k_r)$		278	CP/A'c-14,27	S = RB; A' = α -Terpinene; used $k_d = 8.8 \times 10^4$ s ⁻¹ ; counter ion Na ⁺ .	86A216
	H ₂ O pH = 9.5-11.5	0.16		293	CR/Ac-31	$^1\text{O}_2^*$ from MoO ₄ ²⁻ /H ₂ O ₂ .	89M015
	H ₂ O	$1.4 \times 10^6 (k_r)$		278	CP/A'c-17	S = RB; A' = Rubrene-2,3,8,9-tetracarboxylate ion; used $k_d = 2.4 \times 10^5$ s ⁻¹ , $\beta_{A'} = 1.5 \times 10^{-3}$ mol L ⁻¹ ; counter ion Na ⁺ .	86A216
	MeOH	$1.0 \times 10^4 (k_r)$		278	CP/A'c-14,27	S = RB; A' = α -Terpinene; used $k_d = 9.6 \times 10^4$ s ⁻¹ ; counter ion Na ⁺ .	86A216
	MeOH/H ₂ O (80:20)	$5.8 \times 10^4 (k_r)$		278	CP/A'c-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.3 \times 10^5$ s ⁻¹ ; counter ion Na ⁺ .	86A216
	1-PrOH/H ₂ O (73:27)	$1.3 \times 10^5 (k_r)$		278	CP/A'c-14,27	S = RB; A' = α -Terpinene; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; counter ion Na ⁺ .	86A216

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.89	1,3-Naphthalenedipropionic acid, 4-methyl-						
	CH ₃ CN	2×10^7		263	CP/Ac-15	S = MB; used $k_d = 1.8 \times 10^4$ s ⁻¹ .	93F361
3.90	1,3-Naphthalenedipropionic acid, 4-methyl-, dimethyl ester						
	CH ₃ CN	1×10^8		263	CP/Ac-15	S = MB; used $k_d = 1.8 \times 10^4$ s ⁻¹ .	93F361
3.91	1-Naphthalenepropionate ion, 4-methyl-						
	H ₂ O pH = 9.5-11.5	0.07		293	CR/Pa-31	${}^1\text{O}_2^*$ from MoO ₄ ²⁻ /H ₂ O ₂ .	89M015
3.92	1-Naphthalenepropionate ion, 4-methyl-, endoperoxide (MNPO ₂)						
	H ₂ O	9.0×10^8 (k_q)		308	CR/A'c-32	A' = BR ² ; used $k_d = 3 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from A.	83A317
3.93	1-Naphthalenepropionate ion, 4-methyl-, endoperoxide						
	CH ₂ Cl ₂	8.5×10^7 (k_q)		308	CR/A'c-32	A' = BRH ₂ ; used $k_d = 1 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from A.	83A317
3.94	Naphtho[1,8- <i>bc</i> :5,4- <i>b'</i> , <i>c'</i>]dipyran (1,6-Dioxapyrene)						
	CD ₃ OD	5.8×10^7			PL/Ld-2	S = HP.	93R310
3.95	Naphtho[1,2,3,4- <i>rstf</i>]pentaphene-5,8-dione, 3,10-dimethyl- (Dimethylhomocoerdianthrone, HOCD)						
	C ₆ H ₅ CH ₃		9×10^{-5}		CP/Ac-15	S = DBrA.	82F446 84F197
	CS ₂	3.4×10^8		295	MP/Ld-2	S = TPP.	89A400
3.96	Pentacene						
	C ₆ H ₆	4.2×10^9 (k_p)		298	CP/Ac,A'c-17	S = A; A' = DPBF; used $k_r^{A'} = 7.0 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 6.0$.	747312
3.97	Phenalen-1-one (PHO)						
	Perfluorodecalin	3.2×10^4		295	PL/Ld-2	S = A.	91E427
3.98	1,1'-Spirobi[indene]-5,5',6,6'-tetracarboxylic acid, 2,2',3,3'-tetrahydro-3,3',3'-tetramethyl-, tetramethyl ester						
	MeOH	9.0×10^7			CP/A'c-16	S = MB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	91F516
	MeOH	2.0×10^7 (k_p)			CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_r^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ .	91F516
3.99	Stilbene (Z)-						
	CCl ₄ / MeOH (96:4)	3×10^3			PL/A'd-8	S = MB; A' = DPBF.	777162
3.100	Stilbene, 4,4'-dimethoxy-						
	CH ₃ CN		1.4		CP/Oc-18	S = RB; A' = 2,5-DMF, $\beta_r^{A'}$ not given.	93A103
3.101	Stilbene, α,β -dimethoxy-						
	CH ₃ COCH ₃	9.7×10^6 (k_p)		281	?	S = ?; A' = 1,4-Dioxene; used $k_{A'} = 2.2 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 44$.	70F733
3.102	Stilbene, α,α' -dimethyl-, (E)-						
	CH ₃ CN	1.0×10^6			PL/Ld-2	S = (C ₆ H ₅) ₂ CO.	91E697
	CH ₃ CN		0.11	286	CP/Oc-14	S = RB.	86A099
3.103	Stilbene, α,α' -dimethyl-, (Z)-						
	CCl ₄				CP/Ac,A''c-Pa-17	S = TPP; A' = <i>trans</i> - α,α' -Dimethylstilbene; meas. $k_r/k_r^{A'} = 9.0$; A'' = TME; c.k. with TME.	79F051
	CH ₃ CN		0.014	286	CP/Oc-14	S = RB.	86A099
3.104	Stilbene, α -methyl-, (E)-						
	CCl ₄ / MeOH (96:4)	3×10^3			PL/A'd-8	S = MB; A' = DPBF.	777162

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.105	Stilbene, α -methyl-, (Z)- CCl ₄ / MeOH (96:4)	6×10^3			PL/A'd-8	S = MB; A' = DPBF.	777162
3.106	Stilbene-2,2'-disulfonate ion, 4,4'-diacetamido- H ₂ O	2×10^5			CP/A'c-23	S = MB, RB; A' = InH; used $k_{A'} = 7.7 \times 10^5$ L mol ⁻¹ s ⁻¹ ; counter ion Na ⁺ .	87F554
3.107	Stilbene-2,2'-disulfonate ion, 4,4'-diamino- H ₂ O	1.9×10^6			CP/A'c-23	S = MB, RB; A' = InH; used $k_{A'} = 7.7 \times 10^5$ L mol ⁻¹ s ⁻¹ ; counter ion Na ⁺ .	87F554
3.108	Styrene CCl ₄ / MeOH (96:4)	5×10^3			PL/A'd-8	S = MB; A' = DPBF.	777162
3.109	Styrene, 3-chloro- α , β , β -trimethyl- MeOH/ <i>tert</i> -BuOH (50:50)				CR/Pa,P'a-17	A' = 4, α , β , β -Tetramethylstyrene; meas. $k_t/k_t^{A'} = 0.44$; ${}^1\text{O}_2^*$ from H ₂ O ₂ /NaOCl.	71F577
	MeOH/ C ₅ H ₅ N (98:2)				CP/Pa,P'a-17	S = RB; A' = 4, α , β , β -Tetramethylstyrene; meas. $k_t/k_t^{A'} = 0.38$.	71F577
3.110	Styrene, 4-chloro- α , β , β -trimethyl- MeOH				CP/Pa,P'a-17	S = RB; A' = 2M2B; meas. $k_t/k_t^{A'} = 2.2$.	71F577
3.111	Styrene, 3-cyano- α , β , β -trimethyl- MeOH/ <i>tert</i> -BuOH (50:50)				CR/Pa,P'a-17	A' = 4, α , β , β -Tetramethylstyrene; meas. $k_t/k_t^{A'} = 0.25$; ${}^1\text{O}_2^*$ from H ₂ O ₂ /NaOCl.	71F577
	MeOH/ C ₅ H ₅ N (98:2)				CP/Pa,P'a-17	S = RB; A' = 4, α , β , β -Tetramethylstyrene; meas. $k_t/k_t^{A'} = 0.20$.	71F577
3.112	Styrene, 4-cyano- α , β , β -trimethyl- MeOH/ <i>tert</i> -BuOH (50:50)				CR/Pa,P'a-17	A' = 4, α , β , β -Tetramethylstyrene; meas. $k_t/k_t^{A'} = 0.15$; ${}^1\text{O}_2^*$ from H ₂ O ₂ /NaOCl.	71F577
	MeOH/ <i>tert</i> -BuOH (98:2)				CP/Pa,P'a-17	S = RB; A' = 4, α , β , β -Tetramethylstyrene; meas. $k_t/k_t^{A'} = 0.17$.	71F577
3.113	Styrene, α -cyclopropyl- β , β -trimethyl- CH ₃ COCH ₃				CP/Ac,A'c-17	S = Eos; A' = (Dicyclopropylmethyldene)cyclobutane; meas. $k_t/k_t^{A'} = 0.79$.	78F430
3.114	Styrene, β , β -dimethyl- C ₆ H ₆	0.33		277	CP/Oc-14	S = TPP.	81F582
	CCl ₄	0.05		277	CP/Oc-14	S = TPP.	81F582
	CH ₃ COCH ₂ CH ₃	0.25		277	CP/Oc-14	S = RB.	81F582
	CHCl ₃	0.08		277	CP/Oc-14	S = TPP.	81F582
	EtOH	0.40		277	CP/Oc-14	S = RB.	81F582
3.115	Styrene, 4-(dimethylamino)- α , β , β -trimethyl- MeOH/ <i>tert</i> -BuOH (50:50)				CR/Pa,P'a-17	A' = 4-CH ₃ OC ₆ H ₄ C(CH ₃)=C(CH ₃) ₂ ; meas. $k_t/k_t^{A'} = 2.9$; ${}^1\text{O}_2^*$ from H ₂ O ₂ /NaOCl.	71F577
	MeOH/ C ₅ H ₅ N (98:2)				CP/Pa,P'a-17	S = RB; A' = 4-CH ₃ OC ₆ H ₄ C(CH ₃)=C(CH ₃) ₂ ; meas. $k_t/k_t^{A'} = 2.0$.	71F577
3.116	Styrene, 4-methoxy- β -methyl- MeOH	~ 0.010		293	CP/Oc-15	S = RB; $E_a = 17$ kJ mol ⁻¹ .	68F288
3.117	Styrene, 3-methoxy- α , β , β -trimethyl- MeOH/ <i>tert</i> -BuOH (50:50)				CR/Pa,P'a-17	A' = 4, α , β , β -Tetramethylstyrene; meas. $k_t/k_t^{A'} = 0.72$; ${}^1\text{O}_2^*$ from H ₂ O ₂ /NaOCl.	71F577

TABLE V Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

No.	Substrate	λ ($\text{L mol}^{-1} \text{s}^{-1}$)	$\beta (k_d/k)$ (mol L^{-1})	T (K)	Method	Comment	Ref.
3.117	Styrene, 3-methoxy- α,β,β -trimethyl-	— Continued					
	MeOH/ $\text{C}_5\text{H}_5\text{N}$ (98:2)				CP/Pa,P'a-17	$S = RB; A' = 4,\alpha,\beta,\beta$ -Tetramethylstyrene; meas. $k_t/k_t^{A'} = 0.67$.	71F577
3.118	Styrene, 4-methoxy- α,β,β -trimethyl-				CR/Pa,P'a-17	$A' = 4,\alpha,\beta,\beta$ -Tetramethylstyrene; meas. $k_t/k_t^{A'} = 1.3$; $^1\text{O}_2^*$ from $\text{H}_2\text{O}_2/\text{NaOCl}$.	71F577
	MeOH/ <i>tert</i> -BuOH (50:50)				CP/Pa,P'a-17	$S = RB; A' = 4,\alpha,\beta,\beta$ -Tetramethylstyrene; meas. $k_t/k_t^{A'} = 1.4$.	71F577
3.119	Styrene, β -methyl-, (<i>E</i>)-				PL/A'd-8	$S = MB; A' = DPBF$.	777162
	CCl_4/MeOH (96:4) 2×10^3						
3.120	Styrene, β -methyl-, (<i>Z</i>)-				PL/A'd-8	$S = MB; A' = DPBF$.	777162
	CCl_4/MeOH (96:4) 5×10^3						
3.121	Styrene, 3, α,β,β -tetramethyl-				CR/Pa,P'a-17	$A' = p\text{-Chloro-}\alpha,\beta,\beta$ -trimethylstyrene; meas. $k_t/k_t^{A'} = 1.4$; $^1\text{O}_2^*$ from $\text{H}_2\text{O}_2/\text{NaOCl}$.	71F577
	MeOH/ <i>tert</i> -BuOH (50:50)				CP/Pa,P'a-17	$S = RB; A' = p\text{-Chloro-}\alpha,\beta,\beta$ -trimethylstyrene; meas. $k_t/k_t^{A'} = 1.5$.	71F577
3.122	Styrene, 4, α,β,β -tetramethyl-				CR/Pa,P'a-17	$A' = p\text{-Chloro-}\alpha,\beta,\beta$ -trimethylstyrene; meas. $k_t/k_t^{A'} = 1.9$; $^1\text{O}_2^*$ from $\text{H}_2\text{O}_2/\text{NaOCl}$.	71F577
	MeOH/ <i>tert</i> -BuOH (50:50)				CP/Pa,P'a-17	$S = RB; A' = p\text{-Chloro-}\alpha,\beta,\beta$ -trimethylstyrene; meas. $k_t/k_t^{A'} = 2.2$.	71F577
3.123	Styrene, α,β,β -trimethyl- (TMS)				CR/Pa,P'a-17	$A' = 4,\alpha,\beta,\beta$ -Tetramethylstyrene; meas. $k_t/k_t^{A'} = 0.83$; $^1\text{O}_2^*$ from $\text{H}_2\text{O}_2/\text{NaOCl}$.	71F577
	MeOH/ <i>tert</i> -BuOH (50:50)				CR/Pa,P'a-17	$A' = 4\text{-CH}_3\text{OC}_6\text{H}_4\text{C}(\text{CH}_3)=\text{C}(\text{CH}_3)_2$; meas. $k_t/k_t^{A'} = 0.57$; $^1\text{O}_2^*$ from $\text{H}_2\text{O}_2/\text{NaOCl}$.	71F577
	MeOH/ $\text{C}_5\text{H}_5\text{N}$ (98:2)				CP/Pa,P'a-17	$S = RB; A' = 4\text{-CH}_3\text{OC}_6\text{H}_4\text{C}(\text{CH}_3)=\text{C}(\text{CH}_3)_2$; meas. $k_t/k_t^{A'} = 0.51$.	71F577
3.124	Tetracene						
	C_6H_6	2.4×10^{-3}		298	CP/Ac-?	$S = A$.	767422
	C_6H_6	$1.2 \times 10^7 (k_t)$		298	CP/Ac,A'c-17	$S = A' = \text{Rub}$; used $k_t^{A'} = 4.2 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$; meas. $k_t/k_t^{A'} = 0.29$.	747312
	C_6H_6			298	CP/Pa,P'a-17	$S = A; A' = \text{DMA}$; meas. $k_t/k_t^{A'} = 0.18$.	706079
	C_6H_6	1.7×10^{-3}		298	CP/Ac-15	$S = A$.	706079
	C_6H_6			298	CP/Pa,P'a-17	$S = A; A' = \text{DMBA}$; meas. $k_t/k_t^{A'} = 0.46$.	706079
	CCl_4	$6.9 \times 10^6 (k_t)$			CP/Ac-14,27	$S = \text{PP}$; used $k_d = 36 \text{ s}^{-1}$.	82A421
	CCl_4	$5 \times 10^6 (k_t)$			CP/Ac-27,28	$S = \text{Ph a}$; used $k_d = 36 \text{ s}^{-1}; f_r^A = 1$.	79A010
	CCl_4	5×10^6			MP/LI-12	$S = \text{PP, Ph a or BChl a}$; used $k_d = 36 \text{ s}^{-1}$.	78E881
	CCl_4	7×10^6			MP/LI-12	$S = \text{TPP}$; used $k_d = 36 \text{ s}^{-1}$.	78E892
	CCl_4	5×10^6			MP/LI-12	$S = \text{Ret}$; used $k_d = 36 \text{ s}^{-1}$.	78F700
	CH_2Cl_2	1.8×10^7	4.0×10^{-4}		CP/Ac-14	$S = A$; used $k_d = 7.3 \times 10^3 \text{ s}^{-1}$.	75F654
3.125	Tetracene, 5,6,11,12-tetraphenyl- (Rubrene)						
	$\text{C}_5\text{H}_5\text{N}$				CP/Ac-17	$S = A; A' = \text{TME}$; meas. $k_t/k_t^{A'} = 2.1$.	66F041
	$\text{C}_5\text{H}_5\text{N}$				CP/Ac-17	$S = MB; A' = \text{TME}$; meas. $k_t/k_t^{A'} = 2.4$.	66F041
	C_6D_6	4.3×10^7			PL/Ad-5	$S = 3,6\text{-Bis(dibutylamino)phenothiazinium bromide}$.	81A287

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.125 Tetracene, 5,6,11,12-tetraphenyl- (Rubrene) — Continued							
	c-C ₆ H ₁₂	1.5×10^7	4.0×10^{-3}	297	CP/Ac-15	S = A; used $k_d = 5.9 \times 10^4$ s ⁻¹ .	74F207
	1,2,3-C ₆ H ₃ Cl ₃	5.0×10^7	4×10^{-4}	298	CP/Ac-15	S = A; used $k_d = 2.0 \times 10^4$ s ⁻¹ .	75E223
	C ₆ H ₅ Br	1.6×10^7	8×10^{-4}	297	CP/Ac-15	S = A; used $k_d = 1.3 \times 10^4$ s ⁻¹ .	74F207
	C ₆ H ₅ CH ₃	2.5×10^7		293	PL/Ad-5	S = A.	82E072
	C ₆ H ₅ CH ₃	5.5×10^7	9×10^{-4}	298	CP/Ac-15	S = A; used $k_d = 5.0 \times 10^4$ s ⁻¹ .	75E223
	C ₆ H ₅ CN	1.3×10^8			PL/Ad-5	S = A.	83F075
	C ₆ H ₆	3.4×10^7			CP/Ac-15	S = A; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	87E142
	C ₆ H ₆		1.0×10^{-3}	298	CP/Ac-?	S = A.	76F422
	C ₆ H ₆	4.7×10^7	9×10^{-4}	298	CP/Ac-15	S = A; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	75E223
	C ₆ H ₆	4.2×10^7	1.0×10^{-3}	298	CP/Ac-15	S = A; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	74F312
	C ₆ H ₆	4.3×10^7 (k_r)			CP/Ac,A'c-17	S = A; A' = 1,2-Dithiolane-3-pentanoic acid; used $k_r^{A'} = 1.0 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.43$; β derived assuming $f_r^{A'} = 2$.	74F641
	C ₆ H ₆		9×10^{-4}	298	CP/Ac-23	S = A; A' = 1,2-Dithiolane-3-pentanoic acid.	74F641
	C ₆ H ₆		3.0×10^{-4}	298	CP/Ac-15	S = A.	70E079
	C ₆ H ₆			298	CP/Pa,P'a-17	S = A; A' = DMA; meas. $k_r/k_r^{A'} = 1.0$.	70E079
	C ₆ H ₆			298	CP/Pa,P'a-17	S = A; A' = DMBA; meas. $k_r/k_r^{A'} = 2.8$.	70E079
	C ₆ H ₆		1.7×10^{-3}		CP/Ac-15	S = A.	55F004
	CCl ₄	2.3×10^7	6×10^{-5}	298	CP/Ac-15	S = A; used $k_d = 1.4 \times 10^3$ s ⁻¹ .	75E223
	CCl ₄	1.4×10^7	1×10^{-4}	297	CP/Ac-15	S = A; used $k_d = 1.4 \times 10^3$ s ⁻¹ .	74F207
	CD ₃ CN	8.0×10^7			PL/Ad-5	S = 2-ACN.	81A287
	CD ₃ COCD ₃	2.8×10^7			PL/Ad-5	S = 2-ACN.	81A287
	CDCl ₃	3.9×10^7			PL/Ad-5	S = 2-ACN.	81A287
	CH ₂ Cl ₂	5.4×10^7 (k_r)	2.2×10^{-4} (β_r)		CR/Ac-31	used $k_d = 1.2 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from triphenyl phosphite ozonide; $f_r^{A'} = 1$.	85M406
	CHCl ₃	5.3×10^7	3.1×10^{-4}		CP/Ac-15	S = A; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	77F486
	CHCl ₃	5.9×10^7	3×10^{-4}	298	CP/Ac-15	S = A; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	75E223
	CHCl ₃	4.3×10^7	4×10^{-4}	297	CP/Ac-15	S = A; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	74F207
	CS ₂	4.4×10^7		295	MP/Ld-2	S = TPP.	89A400
	CS ₂		1×10^{-6}	295	CP/Ac-14	S = A; [Rub] = $(8-16) \times 10^{-6}$ mol L ⁻¹ .	89A400
	CS ₂		9×10^{-6}	295	CP/Ac-14	S = A; [Rub] = $(6-15) \times 10^{-5}$ mol L ⁻¹ .	89A400
	CS ₂	5.6×10^7	9×10^{-5}	298	CP/Ac-15	S = A; used $k_d = 5.0 \times 10^3$ s ⁻¹ .	75E223
	CS ₂	2.5×10^7	2×10^{-4}	297	CP/Ac-15	S = A; used $k_d = 5.0 \times 10^3$ s ⁻¹ .	74F207
	EtOH	2.3×10^7	3.6×10^{-3}	297	CP/Ac-15	S = A; used $k_d = 8.3 \times 10^4$ s ⁻¹ .	74F207
	MeOH	3.1×10^7	4×10^{-3}	297	CP/Ac-15	S = A; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	74F207
3.126 Tetracene-2,3,8,9-tetracarboxylate, 5,6,11,12-tetraphenyl- (Rubrene-2,3,8,9-tetracarboxylate)							
	D ₂ O		9×10^{-5}		CL/Ac-15	S = A.	81F043
	H ₂ O		1.6×10^{-3}	298	CR/Pa-31	${}^1\text{O}_2^*$ from MoO ₄ ²⁻ /H ₂ O ₂ .	88M109
	pH = 10.5						
	H ₂ O		1.5×10^{-3}		CL/Ac-15	S = A.	81F043
3.127 Toluene							
	CCl ₄	3.9×10^3		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329
3.128 Toluene-d₈							
	CCl ₄	3.4×10^2		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329

TABLE 3. Rate constants for interaction of singlet oxygen with aromatic substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
3.129 Tribenzo[<i>a,f,j</i>]perylene, 9,18-diphenyl- (Mesodiphenylbenzheptanthrene)							
	C ₆ H ₅ CH ₂ OH	4 × 10 ⁹ 1.0 × 10 ⁸ (k_r)	2 × 10 ⁻⁵		CL/Ac-14	S = RB; used $k_d = 7.1 \times 10^4$ s ⁻¹ ; meas. $f_r^A = 0.028$.	87F479
	C ₆ H ₅ CH ₃ / MeOH (99:1)	7 × 10 ⁹ 1.7 × 10 ⁸ (k_r)	6.2 × 10 ⁻⁶		CL/Ac-14	S = RB; used $k_d = 4.0 \times 10^4$ s ⁻¹ ; meas. $f_r^A = 0.025$.	87F479
	MeOH/ C ₆ H ₅ CH ₃ (97:3)	6.2 × 10 ⁹ 1.6 × 10 ⁸ (k_r)	2 × 10 ⁻⁵		CL/Ac-14	S = RB; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; meas. $f_r^A = 0.026$.	87F479

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols.

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.1 Alanine, glycyl-L-tyrosyl-							
	EtOH/CH ₃ CN (80:20)	$3.0 \times 10^7 (k_t)$			CP/Ac,A'c-17	S = RB; A' = DMA; used $k_t^{A'} = 8.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	93F413
	EtOH/CH ₃ CN (80:20)	2.0×10^8			PL/Ld-2	S = RB; 0.01 mol L ⁻¹ KOH.	93F413
	H ₂ O pH = 11.5	2.0×10^9			CP/Ac-15	S = RB; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	93F413
	H ₂ O pH = 11.5	$1.0 \times 10^8 (k_t)$			CP/Oc-17	S = RB; A' = FFA; used $k_t^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93F413
4.2 Anthrone, 1,8-dihydroxy- (Anthralin)							
	C ₆ D ₆	2.8×10^4			PL/Ld-2	S = TPP.	90A300
	C ₆ H ₆	$\sim 4 \times 10^4$			PL/Ld-2	S = PPDME.	89E158
4.3 Anthrone, 1,8-dihydroxy-, conjugate base							
	CH ₃ CN	3.1×10^8	5.9×10^{-5}	295	CP/Ac-15	S = RB; used $k_d = 1.8 \times 10^4$ s ⁻¹ .	90A300
	CH ₃ CN	$2 \times 10^8 (k_t)$			CP/Ac-17	S = RB; A' = TME; used $k_t^{A'} = 3.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 6$.	90A300
	D ₂ O pD = 9.4-11.4	3×10^8			PL/Ld-2	S = RB.	90A300
4.4 1,2-Benzenediol (Catechol)							
	CD ₃ OD	1.7×10^6		293	PL/Ld-2	S = HP.	90F411
	CH ₃ CN	2.9×10^6			PL/Ld-2	S = RB.	91A341
	D ₂ O	5.4×10^7			PL/Ld-2	S = RB.	91A341
	dioxane	1.4×10^6			PL/Ld-2	S = ZnTPP.	91A341
4.5 1,2-Benzenediol, 4-(2-aminoethyl)- (Dopamine)							
	D ₂ O	2.5×10^7			PL/Ld-2	S = MB.	91R251
	MeOH	1.0×10^6	0.092		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9 \times 10^4$ s ⁻¹ .	86F226
4.6 1,2-Benzenediol, 4-(2-amino-1-hydroxyethyl)- (Noradrenaline)							
	MeOH	1.6×10^6	0.055		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9 \times 10^4$ s ⁻¹ .	86F226
4.7 1,2-Benzenediol, 3,5-bis(1,1-dimethylethyl)-							
	CD ₃ OD	2.5×10^7		293	PL/Ld-2	S = HP.	90F411
	CD ₃ COCD ₃	1.3×10^7			PL/Ld-2	S = MPDME.	87F676
4.8 1,2-Benzenediol, 4-(1,1-dimethylethyl)-							
	CH ₂ Cl ₂	4×10^6			PL/Tb-3	S = TPP.	83A050
4.9 1,2-Benzenediol, 4-(2-methylamino-1-hydroxyethyl)- (Adrenaline)							
	MeOH	2.3×10^6	0.040		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9 \times 10^4$ s ⁻¹ .	86F226
4.10 1,3-Benzenediol (Resorcinol)							
	CD ₃ OD	6.5×10^5		293	PL/Ld-2	S = HP.	90F411
	CH ₃ CN	3.9×10^6			PL/Ld-2	S = RB.	91A341
	CH ₃ CN	$2 \times 10^5 (k_t)$			CP/Ac,A'c-17	S = MPDME; A' = DMA; used $k_t^{A'} = 1.3 \times 10^8$ L mol ⁻¹ s ⁻¹ .	91A341
	D ₂ O	2×10^7			PL/Ld-2	S = RB.	91A341
	H ₂ O	$2 \times 10^7 (k_t)$		293	CP/Oc-17	S = Eos; A' = His; used $k_t^{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A341
	dioxane	5.5×10^5			PL/Ld-2	S = ZnTPP.	91A341
4.11 1,3-Benzenediol, 4-chloro-							
	CH ₃ CN	1.9×10^7			PL/Ld-2	S = RB.	91A341

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.11	1,3-Benzenediol, 4-chloro-	— Continued					
	CH ₃ CN	$5 \times 10^5 (k_r)$			CP/Ac,A'c-17	S = MPDME; A' = DMA; used $k_r^{A'} = 1.3 \times 10^8$ L mol ⁻¹ s ⁻¹ .	91A341
	D ₂ O	2.2×10^7			PL/Ld-2	S = RB.	91A341
	H ₂ O	$2.1 \times 10^7 (k_r)$		293	CP/Oc-17	S = Eos; A' = His; used $k_r^{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A341
	dioxane	6.5×10^5			PL/Ld-2	S = ZnTPP.	91A341
4.12	1,4-Benzenediol (Hydroquinone)						
	C ₅ H ₅ N	6.9×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 6.0 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	743112
	C ₆ H ₆ /MeOH (80:20)	7×10^7		298	CP/P'a-20	S = MB; A' = 2M2P; used $k_d = 1.0 \times 10^5$ s ⁻¹ , $\beta_{A'} = 0.04$ mol L ⁻¹ .	70F734
	CD ₃ OD	1.4×10^7		293	PL/Ld-2	S = HP.	90F411
	CH ₃ CN	$5 \times 10^5 (k_r)$			CP/Ac,A'c-17	S = MPDME; A' = DMA; used $k_r^{A'} = 1.3 \times 10^8$ L mol ⁻¹ s ⁻¹ .	91A341
	CH ₃ CN	2.8×10^7			PL/Ld-2	S = RB.	91A341
	D ₂ O	2.9×10^8			PL/Ld-2	S = RB.	91A341
	EtOH	1.4×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	93M161
	EtOH/H ₂ O (95:5)	$\sim 6 \times 10^8$			CR/LI-12	${}^1\text{O}_2^*$ from pyrogallol autoxidation by O ₂ /KOH; k_d not given.	78F605
	H ₂ O	$2.5 \times 10^7 (k_r)$		293	CP/Oc-17	S = Eos; A' = His; used $k_r^{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A341
	H ₂ O	$\sim 4 \times 10^7 (k_r)$		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; reported $k = (3.8 \pm 5.5) \times 10^7$ L mol ⁻¹ s ⁻¹ .	91F300
	dioxane	1.9×10^7			PL/Ld-2	S = ZnTPP.	91A341
4.13	1,4-Benzenediol, conjugate dibase (Hydroquinone dianion)						
	H ₂ O pH = 10.6	5.6×10^7		310	CR/LI-12	used $k_d = 3.2 \times 10^5$ s ⁻¹ ; soln. cont. 5×10^{-4} mol L ⁻¹ CoCl ₂ , ${}^1\text{O}_2^*$ from autoxidation of oxytetracycline.	92M228
4.14	1,4-Benzenediol, 2,5-bis(1,1-dimethylethyl)-						
	CD ₃ OD	1.9×10^8		293	PL/Ld-2	S = HP.	90F411
	MeOH	6.2×10^7	2.9×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	88F440
4.15	1,4-Benzenediol, 2,6-bis(1,1-dimethylethyl)-, 4-propanoate						
	MeOII	2.2×10^5	0.53		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	78A171
4.16	1,4-Benzenediol, 2,5-bis(2,2-dimethylpropyl)-						
	MeOH	6.9×10^7	2.6×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	88F440
4.17	1,4-Benzenediol, 2,5-bis-sec-dodecyl-						
	MeOH	6.4×10^7	2.8×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	88F440
4.18	1,4-Benzenediol, 2,5-bis-sec-hexadecyl-						
	MeOH	6.6×10^7	2.7×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	88F440
4.19	1,4-Benzenediol, 2,5-bis-sec-octyl-						
	MeOH	6.8×10^7	2.6×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	88F440
4.20	1,4-Benzenediol, chloro-						
	CH ₃ CN	1.6×10^7			PL/Ld-2	S = RB.	91A341
	CH ₃ CN	$3 \times 10^5 (k_r)$			CP/Ac,A'c-17	S = MPDME; A' = DMA; used $k_r^{A'} = 1.3 \times 10^8$ L mol ⁻¹ s ⁻¹ .	91A341

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.10 1,4-Benzenediol, chloro- — Continued							
	D ₂ O	6×10^7			PL/Ld-2	S = RB.	91A341
	H ₂ O	$1.7 \times 10^7 (k_t)$		293	CP/Oc-17	S = Eos; A' = His; used $k_t^{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A341
	dioxane	9.7×10^6			PL/Ld-2	S = ZnTPP.	91A341
4.11 1,4-Benzenediol, 2,3-dimethoxy-5-methyl-							
	EtOH	9.3×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	93M161
4.12 1,4-Benzenediol, 2,3-dimethyl-							
	EtOH	8.5×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	93M161
4.13 1,4-Benzenediol, 2,5-dimethyl-							
	EtOH	9.9×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	93M161
4.14 1,4-Benzenediol, 2,6-dimethyl-							
	EtOH	5.4×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	93M161
4.15 1,4-Benzenediol, 2-(3,7,11,15,19,23-hexamethyl-2,6,10,14,18,22-tetracosahexaenyl)-5,6-dimethoxy-3-methyl- (<i>all-E</i>) (Ubiquinol 10)							
	EtOH	1.6×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	93M161
4.16 1,4-Benzenediol, 2-(3-hydroxy-3,7,11,15-tetramethylhexadecyl)-3,6-dimethyl- (β-Tocopherol hydroquinone)							
	EtOH	1.2×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	93M161
4.17 1,4-Benzenediol, 2-(3-hydroxy-3,7,11,15-tetramethylhexadecyl)-5,6-dimethyl- (γ-Tocopherol hydroquinone)							
	EtOH	1.2×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	93M161
4.18 1,4-Benzenediol, 2-(3-hydroxy-3,7,11,15-tetramethylhexadecyl)-3,5,6-trimethyl- (α-Tocopherol hydroquinone)							
	EtOH	1.3×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	93M161
4.19 1,4-Benzenediol, 2-methyl-							
	EtOH	4.4×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	93M161
4.20 1,4-Benzenediol, phenyl- (2,5-Dihydroxybiphenyl)							
	H ₂ O pH = 8.0	$2.4 \times 10^7 (k_t)$			CP/Oc-17	S = RB; A' = Met; used $k_t^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	93E536
4.21 1,4-Benzenediol, trimethyl-							
	EtOH	1.2×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	93M161
4.22 Benzenemethanol, 3,5-bis(1,1-dimethylethyl)-4-hydroxy-							
	C ₆ H ₆	4.1×10^5			CP/P'a-19	S = ZnTPP; A' = 2M2P; used $k_{A'} = 7.5 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 0.55$.	78A171
	MeOH	1.5×10^6	0.079		CP/A'c-16	S = MB; A' = DPP; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	78A171
4.23 3-Benzenepropanoic acid, 3,5-bis(1,1-dimethylethyl)-4-hydroxy-, octadecyl ester							
	CCl ₄ / MeOH (98:2)	4.6×10^5			FP/A'd-5	S = MB; A' = DPBF.	78E238
	i-octane	$<1 \times 10^6$			CP/A'c-23	S = A' = Rub; No measurable effect.	732066

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.34	3-Benzenepropanoic acid, 4-hydroxy-						
	D ₂ O/ EtOH (75:25)	2.4 × 10 ⁷		295	PL/Ld-2	S = RB.	94A113
4.35	Benzoate ion, 2-hydroxy- (Salicylate ion)						
	H ₂ O (ves) pH = 7.5	1.4 × 10 ⁵ (k_t)		310	CP/Oc-17	S = RB; A' = His; used $k_t^{A'} = 2.8 \times 10^7$ L mol ⁻¹ s ⁻¹ ; DLPC liposomes.	93F231
	H ₂ O (ves) pH = 7.5	1.6 × 10 ⁷	0.016	310	CP/Oc-14	S = AlCl(tspc); used $k_d = 2.5 \times 10^5$ s ⁻¹ ; DLPC liposomes.	93F231
	H ₂ O (ves) pH = 7.5	2.0 × 10 ⁷	0.012	310	CP/Oc-14	S = MC 540; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; DLPC liposomes.	93F231
	H ₂ O (ves) pH = 7.5	2.5 × 10 ⁵ (k_t)		310	CP/Oc-17	S = MC 540; A' = His; used $k_t^{A'} = 2.8 \times 10^7$ L mol ⁻¹ s ⁻¹ ; DLPC liposomes.	93F231
	H ₂ O (ves) pH = 7.5	2.1 × 10 ⁵ (k_t)		310	CP/Oc-17	S = AlCl(tspc); A' = His; used $k_t^{A'} = 2.8 \times 10^7$ L mol ⁻¹ s ⁻¹ ; DLPC liposomes.	93F231
	H ₂ O (ves) pH = 7.5	1.1 × 10 ⁷	0.022	310	CP/Oc-14	S = RB; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; DLPC liposomes.	93F231
4.36	Benzofuran-5-ol, 2,3-dihydro-2,2,4,6,7-pentamethyl-						
	EtOH	3.6 × 10 ⁸		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	91F285
4.37	Benzofuran-5-ol, 2,3-dihydro-2,4,6,7-tetramethyl-						
	EtOH	2.5 × 10 ⁸		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	91F285
4.38	Benzoic acid, 3,5-bis(1,1-dimethylethyl)-4-hydroxy-, 2,4-bis(1,1-dimethylethyl)phenyl ester						
	C ₆ H ₅ Br	<1 × 10 ⁶		273	MD/A'c-33	A' = Rub; No measurable effect.	737333
	i-octane	<1 × 10 ⁶			CP/A'c-23	S = A' = Rub; No measurable effect.	732066
4.39	Benzoic acid, 3,5-bis(1,1-dimethylethyl)-4-hydroxy-, methyl ester						
	1-BuOH	9.2 × 10 ⁵		293	CP/A'c-23	S = MB; A' = Np; used $k_d = 5.2 \times 10^4$ s ⁻¹ , $k_{A'} = 5.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A266
4.40	Benzoic acid, 2-hydroxy-, methyl ester (Methyl salicylate)						
	H ₂ O	<2 × 10 ⁶ (k_t)		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_t^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; estd. using pK _a = 9.8.	87A180
4.41	Benzoic acid, 2-hydroxy-, methyl ester, conjugate base						
	H ₂ O pH = 10.0	1.6 × 10 ⁸ (k_t)		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_t^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; estd. using pK _a = 9.8.	87A180
4.42	Benzophenone, 4-dodecyloxy-2-hydroxy-						
	CCl ₄ / MeOH (98:2)	<2 × 10 ⁵			FP/A'd-5	S = MB; A' = DPBF.	78E238
	i-octane	<1 × 10 ⁶			CP/A'c-23	S = A' = Rub; No measurable effect.	732066
4.43	Benzophenone, 2-hydroxy-4-octyloxy-						
	C ₆ H ₅ Br	<1 × 10 ⁶		273	MD/A'c-33	A' = Rub; No measurable effect.	737333
	CCl ₄ / CHCl ₃ (90:10)	1.0 × 10 ⁴			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k/(k_{A'}[A'] + k_d) = 10.0$ L mol ⁻¹ at [A'] = 5 × 10 ⁻⁶ mol L ⁻¹ .	74F645
4.44	1-Benzopyran-2-carboxylate ion, 3,4-dihydro-6-hydroxy-2,5,7,8-tetramethyl- (Trolox C anion)						
	CD ₃ OD	1.2 × 10 ⁸		293	PL/Ld-2	S = HP.	90F411
	D ₂ O/ EtOH (50:50)	4.7 × 10 ⁸		310	CR/LI-12	used $k_d = 1 \times 10^5$ s ⁻¹ ; ¹ O ₂ * from NDPO ₂ .	90A184
	EtOH	8.1 × 10 ⁷		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	91F285

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.31 1-Benzopyran-6-ol, 5,7-diethyl-3,4-dihydro-2-methyl-2-(4,8,12-trimethyltridecyl)-							
	EtOH	9.5×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
4.32 1-Benzopyran-6-ol, 3,4-dihydro-2,8-dimethyl-2-(4,8,12-trimethyltridecyl)- (δ-Tocopherol)							
	n-C ₁₆ H ₃₄	5.0×10^6		298	MD/A'c-33	A' = Rub; used $k_d = 9 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88F415
	CHCl ₃ /EtOH (50:50)	1.6×10^8		310	CR/LI,Ac-12,31	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	90A184
	EtOH	5.3×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
	EtOH	1.0×10^8	8.3×10^{-4}	293	CP/P'a-22	S = MB; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; A' = Methyl linoleate; P' = Methyl linoleate hydroperoxide.	77F858
	EtOH	7×10^5 (k_r)		293	CP/P'a,Ac-17	S = MB; used $k_r^{A'} = 7.3 \times 10^4$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 11$; A' = Methyl linoleate; P' = Methyl linoleate hydroperoxide.	77F858
	MeOH	1×10^8	1×10^{-3}	298	CP/Ac-15	S = MB; used $k_d = 1 \times 10^5$ s ⁻¹ .	72A019
4.33 1-Benzopyran-6-ol, 3,4-dihydro-2-methyl-5,7-di(1-methylethyl)-2-(4,8,12-trimethyltridecyl)-							
	EtOH	1.1×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
4.34 1-Benzopyran-6-ol, 3,4-dihydro-1-methyl-1-(4,8,12-trimethyltridecyl)- (Tocol)							
	EtOH	2.8×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
4.35 1-Benzopyran-6-ol, 3,4-dihydro-2,2,5,7,8-pentamethyl-							
	c-C ₆ H ₁₂	2.4×10^7		PL/A'd-8	S = 2-ACN; A' = DPBF; $\Delta H^\ddagger = -8.8$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -130$ J K ⁻¹ mol ⁻¹ ; studied at 278-357 K.	84E492	
	C ₆ H ₅ CH ₃	1.5×10^8		PL/A'd-8	S = 2-ACN; A' = DPBF; $\Delta H^\ddagger = -15$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -138$ J K ⁻¹ mol ⁻¹ ; studied at 230-385 K.	84E492	
	CH ₃ CN	5.7×10^8		PL/A'd-8	S = 2-ACN; A' = DPBF; $\Delta H^\ddagger = -15$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -130$ J K ⁻¹ mol ⁻¹ ; studied at 230-357 K.	84E492	
	CH ₃ COCH ₃	2.6×10^8		PL/A'd-8	S = 2-ACN; A' = DPBF; $\Delta H^\ddagger = -8.4$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -109$ J K ⁻¹ mol ⁻¹ ; studied at 250-328 K.	84E492	
	D ₂ O (mic) pD = 7.4	4.1×10^8		PL/A'd-5	S = MB; A' = DPBF; 0.1 mol L ⁻¹ CTAB.	81N048	
	D ₂ O (mic) pD = 7.4	4.0×10^8		PL/A'd-5	S = MB; A' = DPBF; 0.1 mol L ⁻¹ SDS.	81N048	
	EtOH	2.0×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
	MeOH	4.0×10^7 (k_r)		CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_r^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ .	91F516	
	MeOH	1.6×10^8		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	91F516	
	MeOH	3.4×10^8		PL/A'd-8	S = 2-ACN; A' = DPBF; $\Delta H^\ddagger = -5.9$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -100$ J K ⁻¹ mol ⁻¹ ; studied at 250-328 K.	84E492	
4.36 1-Benzopyran-6-ol, 3,4-dihydro-2,5,7,8-tetramethyl-2-(4,8,12-trimethyltridecyl)- (α-Tocopherol)							
	n-C ₁₆ H ₃₄	4.2×10^7		298	MD/A'c-33	A' = Rub; used $k_d = 9 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88F415
	c-C ₆ H ₁₂	8.4×10^7		PL/A'd-8	S = 2-ACN; A' = DPBF.	84E492	
	c-C ₆ H ₁₂	1.1×10^6 (k_r)		298	CP/Ac,A'c-17	S = A' = DPA; used $k_r^{A'} = 1.2 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.9$.	743114
	c-C ₆ H ₁₂	9.0×10^7	6.8×10^{-4}	298	CP/A'c-23	S = A' = Rub; used $k_d = 5.9 \times 10^4$ s ⁻¹ , $\beta_{A'} = 1.4 \times 10^{-3}$ mol L ⁻¹ .	743114

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.50 1-Benzopyran-6-ol, 3,4-dihydro-2,5,7,8-tetramethyl-2-(4,8,12-trimethyltridecyl)- (α-Tocopherol) — Continued							
	C ₅ H ₅ N	2.5×10^8			CP/A'c-23	S = A' = Rub; used $k_d = 6.0 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	743112
	C ₅ H ₅ N	2.1×10^6 (k_r)			CP/Ac-14,28	S = PP; A' = Cholesterol; used $k_d = 6.0 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.9$ mol L ⁻¹ ; used $k_A = 2.5 \times 10^8$ L mol ⁻¹ s ⁻¹ .	743112
	C ₆ H ₅ CH ₃	2.2×10^8			PL/A'd-8	S = 2-ACN; A' = DPBF; $\Delta H^\ddagger = -14$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -138$ J K ⁻¹ mol ⁻¹ ; studied at 230-385 K.	84E492
	C ₆ H ₆	1×10^8 2.1×10^6 (k_r) 1.4×10^8 (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
	C ₆ H ₆	1.7×10^8	2.7×10^{-4}	298	CP/A'c-23	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $\beta_{A'} = 1.0 \times 10^{-3}$ mol L ⁻¹ .	743114
	CCl ₄	5.2×10^7			CL/LI-12	S = TPP; used $k_d = 1 \times 10^3$ s ⁻¹ .	85F667
	CCl ₄	1×10^8			MP/LI-12	S = PP, TPP, Ph a or BPh a; used $k_d = 36$ s ⁻¹ .	79A010 78E892
	CCl ₄	10^8			MP/LI-12	S = Ret; used $k_d = 36$ s ⁻¹ .	79F463 78F700
	CH ₂ Cl ₂	2.7×10^7		298	CP/P'a or Oc-19	S = Chl a; A' = Soybean oil; used $k_{A'} = 1.0 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 2.6 \times 10^2$; measured peroxide formation.	91U128
	CH ₃ CN	5.9×10^8			PL/A'd-8	S = 2-ACN; A' = DPBF; $\Delta H^\ddagger = -11$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -138$ J K ⁻¹ mol ⁻¹ ; studied at 230-357 K.	84E492
	CH ₃ COCH ₃	4.3×10^8			PL/A'd-8	S = 2-ACN; A' = DPBF; $\Delta H^\ddagger = -12$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -122$ J K ⁻¹ mol ⁻¹ ; studied at 250-328 K.	84E492
	CHCl ₃	1.1×10^8			CP/A'c-16	S = MB; A' = DPBF; used $k_d = 4.4 \times 10^3$ s ⁻¹ .	83A394
	CHCl ₃ / EtOH (50:50)	2.8×10^8 3.6×10^6 (k_r)		310	CR/LI,Ac-12,31	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	90A184
	ClCF ₂ CCl ₂ F	3×10^7		298	FP/Ld-2	S = Per.	82A322
	D ₂ O/ EtOH (50:50)	4.5×10^8		310	CR/LI-12	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	90A184
	D ₂ O (mic)	6.4×10^8			PL/A'd-5	S = 2-ACN; A' = DPBF; 0.1 mol L ⁻¹ SDS.	78E144
	EtOH		3.3×10^{-3}	298	CP/Ac-15	S = 8-MOP.	92F192
	EtOH	2.1×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
	EtOH	1×10^8			CP/A'c-19	S = RB; A' = Chl a; used $k_d = 1 \times 10^5$ s ⁻¹ .	78F404
	EtOH	2.6×10^8	3.2×10^{-4}	293	CP/P'a-22	S = MB; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; A' = Methyl linoleate; P' = Methyl linoleate hydroperoxide.	77F858
	EtOH	6.6×10^6 (k_r)		293	CP/P'a,Ac-17	S = MB; used $k_r^{A'} = 7.3 \times 10^4$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 91$; A' = Methyl linoleate; P' = Methyl linoleate hydroperoxide.	77F858
	MeOH	4.5×10^8		297	CR/P'a-16	$A' = \text{TEMPO}$; k_d not given; formn. of TEMPO monitored by esr; soln. cont. MeONa and CoCl ₂ , ${}^1\text{O}_2^*$ from autoxidation of adrenaline.	92D227
	MeOH	3.8×10^8	4.8×10^{-4}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	88F440
	MeOH	3.0×10^8			PL/A'd-8	S = 2-ACN; A' = DPBF; $\Delta H^\ddagger = -8$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -109$ J K ⁻¹ mol ⁻¹ ; studied at 250-328 K.	84E492
	MeOH	5×10^7 (k_r)			CP/Ac,A'c-17	S = MB; A' = DPF; used $k_r^{A'} = 1.6 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.29$.	743113
	MeOH	6.7×10^8	2.1×10^{-4}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	743113
	MeOH	1×10^9	1×10^{-4}	298	CP/Ac-15	S = MB; used $k_d = 1 \times 10^5$ s ⁻¹ .	72A019
	MeOH/ H ₂ O (95:5)	2.6×10^9		310	CR/LI-12	used $k_d = 1.8 \times 10^5$ s ⁻¹ ; soln. cont. 0.05 mol L ⁻¹ MeONa and 5 $\times 10^{-4}$ mol L ⁻¹ CoCl ₂ , ${}^1\text{O}_2^*$ from autoxidation of α,γ -tetracycline.	92M228

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.50 1-Benzopyran-6-ol, 3,4-dihydro-2,5,7,8-tetramethyl-2-(4,8,12-trimethyltridecyl)- (α-Tocopherol) — Continued						
<i>t</i> -octane	1×10^8			CP/A'c-23	$S = A' = \text{Rub}$; used $k_{A'} = 4 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$; k_d not given.	743112
4.51 1-Benzopyran-6-ol, 3,4-dihydro-2,5,7-trimethyl-2-(4,8,12-trimethyltridecyl)-						
EtOH	9.7×10^7		308	CR/A'c-16	$A' = \text{DPBF}$; used $k_d = 8.3 \times 10^4 \text{ s}^{-1}$; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
4.52 1-Benzopyran-6-ol, 3,4-dihydro-2,5,8-trimethyl-2-(4,8,12-trimethyltridecyl)- (β-Tocopherol)						
<i>n</i> -C ₁₆ H ₃₄	2.3×10^7		298	MD/A'c-33	$A' = \text{Rub}$; used $k_d = 9 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 7.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	88F415
CHCl ₃ / EtOH (50:50)	2.7×10^8 $2.3 \times 10^5 (k_r)$		310	CR/LI,Ac-12,31	used $k_d = 1 \times 10^5 \text{ s}^{-1}$; ${}^1\text{O}_2^*$ from NDPO ₂ .	90A184
EtOH	1.5×10^8		308	CR/A'c-16	$A' = \text{DPBF}$; used $k_d = 8.3 \times 10^4 \text{ s}^{-1}$; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
MeOH	3.6×10^8	2.8×10^{-4}	298	CP/Ac-15	$S = \text{MB}$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$.	72A019
4.53 1-Benzopyran-6-ol, 3,4-dihydro-2,7,8-trimethyl-2-(4,8,12-trimethyltridecyl)- (γ-Tocopherol)						
<i>n</i> -C ₁₆ H ₃₄	1.1×10^7		298	MD/A'c-33	$A' = \text{Rub}$; used $k_d = 9 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 7.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	88F415
CHCl ₃ / EtOH (50:50)	2.3×10^8 $2.8 \times 10^6 (k_r)$		310	CR/LI,Ac-12,31	used $k_d = 1 \times 10^5 \text{ s}^{-1}$; ${}^1\text{O}_2^*$ from NDPO ₂ .	90A184
EtOH	1.4×10^8		308	CR/A'c-16	$A' = \text{DPBF}$; used $k_d = 8.3 \times 10^4 \text{ s}^{-1}$; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
EtOH	$2.6 \times 10^6 (k_r)$		293	CP/P'a,Ac-17	$S = \text{MB}$; used $k_r^{A'} = 7.3 \times 10^4 \text{ L mol}^{-1} \text{ s}^{-1}$; meas. $k_r/k_r^{A'} = 36$; $A' = \text{Methyl linoleate}$; P' = Methyl linoleate hydroperoxide.	77F858
EtOH	1.8×10^8	4.5×10^{-4}	293	CP/P'a-22	$S = \text{MB}$; used $k_d = 8.3 \times 10^4 \text{ s}^{-1}$; $A' = \text{Methyl linoleate}$; P' = Methyl linoleate hydroperoxide.	77F858
MeOH	1.9×10^8	5.4×10^{-4}	298	CP/Ac-15	$S = \text{MB}$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$.	72A019
4.54 1-Benzopyran-6-ol, 7-(1,1-dimethylethyl)-3,4-dihydro-2,2-dimethyl-						
MeOH	$4.7 \times 10^7 (k_r)$			CP/Ac,A'c-17	$S = \text{MB}$; $A' = \text{DPBF}$, used $k_r^{A'} = 8 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	91F516
MeOH	1.9×10^8			CP/A'c-16	$S = \text{MB}$; $A' = \text{DPBF}$; used $k_d = 1.8 \times 10^5 \text{ s}^{-1}$.	91F516
4.55 1-Benzopyran-6-ol, 7-(1,1-dimethylethyl)-3,4-dihydro-2-methyl-2-(4,8,12-trimethyltridecyl)-						
EtOH	1.2×10^8		308	CR/A'c-16	$A' = \text{DPBF}$; used $k_d = 8.3 \times 10^4 \text{ s}^{-1}$; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
4.56 1-Benzopyran-6-ol, 8-(1,1-dimethylethyl)-3,4-dihydro-2-methyl-2-(4,8,12-trimethyltridecyl)-						
EtOH	1.6×10^8		308	CR/A'c-16	$A' = \text{DPBF}$; used $k_d = 8.3 \times 10^4 \text{ s}^{-1}$; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
4.57 1-Benzopyran-4-one, 2-[3,4-bis(2-hydroxyethoxy)phenyl]-3-[6-O-(6-deoxy-α-mannopyranosyl-β-D-glycopyranosyl]oxy]-5-hydroxy-7-(2-hydroxyethoxy)- (Troxerutin)						
CD ₃ OD	7.0×10^4			CP/LI-12	$S = \text{RB}$; used $k_d = 3.7 \times 10^3 \text{ s}^{-1}$.	93A326
MeOD	$1.0 \times 10^4 (k_r)$			CP/Ac-14	$S = \text{RB}$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$; MDH as actinometer.	93A326
4.58 1-Benzopyran-4-one, 3-[[6-O-(6-deoxy-α-mannopyranosyl)-β-glucopyranosyl]oxy]-2-(3,4-dihydroxyphenyl)-5,7-dihydroxy- (Rutin)						
CD ₃ OD	1.6×10^6			CP/LI-12	$S = \text{RB}$; used $k_d = 4.0 \times 10^3 \text{ s}^{-1}$.	93A326
MeOD	$1.1 \times 10^5 (k_r)$			CP/Ac-14	$S = \text{RB}$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$; MDH as actinometer.	93A326
4.59 1-Benzopyran-4-one, 2-(2,3-dihydroxyphenyl)-2,3-dihydro-5,7-dihydroxy-, (Eriodictyol)						
CD ₃ OD	1.4×10^6			CP/LI-12	$S = \text{RB}$; used $k_d = 3.7 \times 10^3 \text{ s}^{-1}$.	93A326

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.60 1-Benzopyran-4-one, 2-(3,4-dihydroxyphenyl)-5,7-dihydroxy- (Luteolin)							
	CD ₃ OD	1.3×10^6			CP/LI-12	S = RB; used $k_d = 3.7 \times 10^3$ s ⁻¹ .	93A326
	MeOD	1.8×10^4 (k_r)			CP/Ac-14	S = RB; used $k_d = 1 \times 10^5$ s ⁻¹ ; MDH as actinometer.	93A326
4.61 1-Benzopyran-4-one, 2-(3,4-dihydroxyphenyl)-3,5,7-trihydroxy- (Quercetin)							
	CH ₃ CN	9.3×10^3 (k_r)		295	CP/Ac,A'c-17	S = A' = An; used $k_r^{A'} = 2.8 \times 10^5$ L mol ⁻¹ s ⁻¹ .	87A371
	CH ₃ COCH ₃	3.1×10^5		295	PL/Ld-2	S = An.	87A371
	CD ₃ OD	2.4×10^6			CP/LI-12	S = RB; used $k_d = 4.0 \times 10^3$ s ⁻¹ .	93A326
	MeOD	8.9×10^5 (k_r)			CP/Ac-14	S = RB; used $k_d = 1 \times 10^5$ s ⁻¹ ; MDH as actinometer.	93A326
4.62 1-Benzopyran-4-one, 5,6,7,8-tetramethoxy-2-(4-methoxyphenyl)- (Tangeretin)							
	CD ₃ OD	2.4×10^5			CP/LI-12	S = RB; used $k_d = 3.7 \times 10^3$ s ⁻¹ .	93A326
	MeOD	1.0×10^4 (k_r)			CP/Ac-14	S = RB; used $k_d = 1 \times 10^5$ s ⁻¹ ; MDH as actinometer.	93A326
4.63 1-Benzopyran-4-one, 2,3-dihydro-3,5,7-trihydroxy-2-(3,4-dihydroxyphenyl)- (Taxifolin)							
	CD ₃ OD	1.1×10^6			CP/LI-12	S = RB; used $k_d = 3.7 \times 10^3$ s ⁻¹ .	93A326
4.64 1-Benzopyran-4-one, 3,5,7-trihydroxy-2-(4-hydroxyphenyl)- (Kaempferol)							
	CD ₃ OD	7.1×10^5			CP/LI-12	S = RB; used $k_d = 3.7 \times 10^3$ s ⁻¹ .	93A326
	MeOD	4.8×10^5 (k_r)			CP/Ac-14	S = RB; used $k_d = 1 \times 10^5$ s ⁻¹ ; MDH as actinometer.	93A326
4.65 1-Benzopyran-4-one, 3,5,7-trihydroxy-2-phenyl- (Galangin)							
	CD ₃ OD	1.2×10^6			CP/LI-12	S = RB; used $k_d = 3.7 \times 10^3$ s ⁻¹ .	93A326
	MeOD	7.4×10^5 (k_r)			CP/Ac-14	S = RB; used $k_d = 1 \times 10^5$ s ⁻¹ ; MDH as actinometer.	93A326
4.66 Benzopyran-2-propanoic acid, 3,4-dihydro-6-hydroxy-2,5,7,8-tetramethyl-							
	EtOH	1.8×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
4.67 1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro- (2<i>R</i>-trans)- (Catechin)							
	CD ₃ OD	5.8×10^6			CP/LI-12	S = RB; used $k_d = 3.7 \times 10^3$ s ⁻¹ .	93A326
4.68 Cinnamic acid, 3,4-dihydroxy-							
	CD ₃ OD	5.4×10^5		293	PL/Ld-2	S = HP.	88R236 90F411
4.69 Cinnamic acid, 2-hydroxy-							
	CD ₃ OD	2.1×10^4		293	PL/Ld-2	S = HP.	90F411
4.70 Cinnamic acid, 4-hydroxy-3-methoxy-							
	CD ₃ OD	3.5×10^5		293	PL/Ld-2	S = HP.	88R236 90F411
4.71 5-S-Cysteinyl-dopa							
	D ₂ O	2.9×10^7			PL/Ld-2	S = MB.	91R251
4.72 Flavanone, 4',5,7-trihydroxy- (Naringenin)							
	CD ₃ OD	5.0×10^4			CP/LI-12	S = RB; used $k_d = 3.7 \times 10^3$ s ⁻¹ .	93A326
4.73 Flavone, 5,7-dihydroxy- (Chrysins)							
	CD ₃ OD	2.4×10^5			CP/LI-12	S = RB; used $k_d = 3.7 \times 10^3$ s ⁻¹ .	93A326
	MeOD	6×10^3 (k_r)			CP/Ac-14	S = RB; used $k_d = 1 \times 10^5$ s ⁻¹ ; MDH as actinometer.	93A326

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

#	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.74 Flavone, 3-hydroxy-							
	C ₅ H ₅ N	7.5×10^7			PL/Ld-2	S = RB, TPP or H ₂ TPPS ⁴ .	93U041 89A322
	C ₆ D ₆	1.6×10^4			PL/Ld-2	S = RB, TPP or H ₂ TPPS ⁴ ; 95%-d.	93U041 89A322
	C ₆ H ₆	1.9×10^4			PL/Ld-2	S = RB, TPP or H ₂ TPPS ⁴ .	93U041 89A322
	CCl ₄	1.7×10^4			PL/Ld-2	S = RB, TPP or H ₂ TPPS ⁴ .	93U041 89A322
	CD ₃ CN	1.1×10^5			PL/Ld-2	S = RB, TPP or H ₂ TPPS ⁴ .	93U041 89A322
	CD ₃ OD	9.0×10^4			PL/Ld-2	S = RB or H ₂ TPPS ⁴ .	93U041 89A322
	CH ₃ CN	1.1×10^5			PL/Ld-2	S = RB or H ₂ TPPS ⁴ .	93U041 89A322
	D ₂ O pD = 3	2.5×10^5			PL/Ld-2	S = RB or H ₂ TPPS ⁴ .	93U041 89A322
	D ₂ O pD = 12	1.9×10^8			PL/Ld-2	S = RB or H ₂ TPPS ⁴ .	93U041 89A322
	H ₂ O pH = 12	2.3×10^8			PL/Ld-2	S = RB or H ₂ TPPS ⁴ .	93U041 89A322
	MeOH	2.3×10^5			PL/Ld-2	S = RB or H ₂ TPPS ⁴ .	93U041 89A322
4.75 Flavone, 3,3',4',7-tetrahydroxy- (Fisetin)							
	CD ₃ OD	3.1×10^6			CP/LI-12	S = RB; used $k_d = 4.0 \times 10^3$ s ⁻¹ .	93A326
	MeOD	1.1×10^6 (k_t)			CP/Ac-14	S = RB; used $k_d = 1 \times 10^5$ s ⁻¹ ; MDH as actinometer.	93A326
4.76 Glycine, L-tyrosyl-							
	EtOH/ CH ₃ CN (80:20)	4.3×10^6			PL/Ld-2	S = ZnTPP.	91A252
	EtOH/ CH ₃ CN (80:20)	2.1×10^6 (k_t)			CP/Ac,A'c- 14,28	S = ZnTPP; A' = DMA; used $k_d = 5.3 \times 10^4$ s ⁻¹ ; used $k_A = 4.3 \times 10^6$ L mol ⁻¹ s ⁻¹ .	91A252
	EtOH/ CH ₃ CN (80:20)	1.2×10^8			PL/Ld-2	S = ZnTPP; 0.01 mol L ⁻¹ KOH.	91A252
	EtOH/ CH ₃ CN (80:20)	1×10^7 (k_t)			CP/Ac,A'c- 14,28	S = ZnTPP; A' = DMA; used $k_d = 5.3 \times 10^4$ s ⁻¹ ; used $k_A = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	91A252
	EtOH/ CH ₃ CN (80:20)	4.2×10^6			PL/Ld-2	S = ZnTPP; 0.01 mol L ⁻¹ HCl.	91A252
	EtOH/ CH ₃ CN (80:20)	$<5 \times 10^4$ (k_t)			CP/Ac,A'c- 14,28	S = ZnTPP; A' = DMA; used $k_d = 5.3 \times 10^4$ s ⁻¹ ; used $k_A = 4.2 \times 10^6$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ HCl.	91A252
	H ₂ O pH = 10	3×10^7 (k_t)			CP/Oc-17	S = RB; A' = Met; used $k_t^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A252
	H ₂ O pH = 11.5	4.3×10^7 (k_t)			CP/Oc-17	S = Eos; A' = Met; used $k_t^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A252
4.77 1,6-Heptadiene-3,5-dione, 1,7-bis(4-hydroxy-3-methoxyphenyl)- (Curcumin)							
	C ₆ D ₆	2.5×10^5		295	CP/LI-12	S = TPP; used $k_d = 1.5 \times 10^3$ s ⁻¹ .	94E153
	CH ₃ CN	$<7 \times 10^6$			CP/LI-12	S = RB; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	94F069
4.78 Melanin (from DOPA)							
	D ₂ O pD = 7.5-8.1	1×10^8			PL/Ld-2	S = ?; rate per monomer unit.	87R227

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.78	Melanin (from DOPA) — Continued						
	D ₂ O pD = 7.6 ?	$\leq 2 \times 10^7$	$>6.7 \times 10^{-3}$	296	CP/Oc-15	S = RB; used $k_d = 1.9 \times 10^4$ s ⁻¹ ; based on monomer unit of MW 150.	84F330
4.79	5,12-Naphthacenedione, 8-acetyl-10[(3-amino-2,3,6-trideoxy-hexopyranosyloxy)-tetrahydro-6,8,11-trihydroxy-1-methoxy- (Daunomycin)						
	D ₂ O pH = 7.4	$\sim 10^8$			PL/Ld-2	S = AlCl(tspc); cor. for aggregation of substrate.	89B024
	H ₂ O pH = 7.8	$2 \times 10^7 (k_r)$ $1 \times 10^9 (k_q)$		308	CR/Ac-31	used $k_d = 2.4 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ ; 15% dimer.	89M096
	H ₂ O (mic) pH = 7.8	1.7×10^9		308	CP/Ac-16	S = ?; A' = DPBF; used $k_d = 2.4 \times 10^5$ s ⁻¹ ; 15% dimer; 0.1 mol L ⁻¹ CTAB.	89M096
4.80	5,12-Naphthacenedione, 10-[(3-amino-2,3,6-trideoxy- α -L-lyxo-hexopyranosyl)oxy]-7,8,9,10-tetrahydro-6,8,11-trihydroxy-8-(hydroxyacetyl)-1-methoxy- (Adriamycin)						
	D ₂ O pH = 7.4	$\sim 10^8$			PL/Ld-2	S = AlCl(tspc); cor. for aggregation of substrate.	89B024
	H ₂ O pH = 7.8	$2 \times 10^7 (k_r)$ $9 \times 10^8 (k_q)$		308	CR/Ac-31	used $k_d = 2.4 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ ; 15% dimer.	89M096
	H ₂ O (mic) pH = 7.8	2.1×10^9		308	CP/Ac-16	S = ?; A' = DPBF; used $k_d = 2.4 \times 10^5$ s ⁻¹ ; 15% dimer; 0.1 mol L ⁻¹ CTAB.	89M096
4.81	12-Naphthaceneone, 8-acetyl-10[(3-amino-2,3,6-trideoxyhexopyranosyloxy)-tetrahydro-6,7,11-trihydroxy-5-imino-1-methoxy-						
	D ₂ O pH = 7.4	$\sim 10^7$			PL/Ld-2	S = AlCl(tspc).	89B024
	H ₂ O pH = 7.8	$2 \times 10^7 (k_r)$ $3 \times 10^9 (k_q)$		308	CR/Ac-31	used $k_d = 2.4 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ ; 15% dimer.	89M096
	H ₂ O (mic) pH = 7.8	3.5×10^9		308	CP/Ac-16	S = ?; A' = DPBF; used $k_d = 2.4 \times 10^5$ s ⁻¹ ; 15% dimer; 0.1 mol L ⁻¹ CTAB.	89M096
4.82	Naphthalene, 1,5-dihydroxy-						
	CD ₃ OD	6.1×10^6 $1.6 \times 10^6 (k_r)$			CP/LI,Pa-12,28	S = RB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; P = 5-Hydroxy-1,4-naphthoquinone.	90F065
4.83	Naphthalene, 1,6-dihydroxy-						
	CD ₃ OD	3.2×10^6 $1.2 \times 10^6 (k_r)$			CP/LI,Pa-12,28	S = RB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; P = 6-Hydroxy-1,4-naphthoquinone.	90F065
4.84	Naphthalene, 1,7-dihydroxy-						
	CD ₃ OD	3.6×10^6 $6.8 \times 10^5 (k_r)$			CP/LI,Pa-12,28	S = RB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; P = 6-Hydroxy-1,4-naphthoquinone.	90F065
4.85	Naphthalene, 1,8-dihydroxy-						
	CD ₃ OD	5.3×10^7 $3.0 \times 10^7 (k_r)$			CP/LI,Pa-12,28	S = RB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; P = 5-Hydroxy-1,4-naphthoquinone.	90F065
4.86	Naphthalene, 2,6-dihydroxy-						
	CD ₃ OD	1.9×10^6			CP/LI-12	S = RB; used $k_d = 4.4 \times 10^3$ s ⁻¹ .	90F065
4.87	Naphthalene, 2,7-dihydroxy-						
	CD ₃ OD	2.1×10^6			CP/LI-12	S = RB; used $k_d = 4.4 \times 10^3$ s ⁻¹ .	90F065
4.88	1-Naphthol						
	1-BuOH	7.6×10^6		293	PL/A'd-8	S = PP; A' = DPBF.	88F462
	1-BuOH	3.2×10^7		293	CP/A'c-23	S = MB; A' = Np; used $k_d = 5.2 \times 10^4$ s ⁻¹ , $k_{A'} = 5.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A266
	C ₆ H ₆	$\leq 5 \times 10^5$		293	PL/A'd-8	S = PP; A' = DPBF.	88F462 85A124

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.90	1-Naphthol						
	1-BuOH	3.2×10^7		293	PL/A'd-8	S = PP; A' = DPBF.	88F462
	1-BuOH	7.6×10^6		293	CP/A'c-23	S = MB; A' = Np; used $k_d = 5.2 \times 10^4$ s ⁻¹ , $k_{A'} = 5.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A266
	C ₆ H ₅ Cl	3.9×10^4		295	PL/Ld-2	S = Zn(pc).	87A371
	C ₆ H ₆	$\leq 1 \times 10^5$		293	PL/A'd-8	S = PP; A' = DPBF.	88F462 85A124
	C ₆ H ₆	1.8×10^4		295	PL/Ld-2	S = Zn(pc).	87A371
	CH ₃ CN	9.3×10^4		295	PL/Ld-2	S = Zn(pc).	87A371
	CH ₃ COCH ₃	4.0×10^4		295	PL/Ld-2	S = Zn(pc).	87A371
	CHCl ₃	7.1×10^4		295	PL/Ld-2	S = Zn(pc).	87A371
	2-PrOH	3.5×10^4		295	PL/Ld-2	S = Zn(pc).	87A371
4.91	1,4-Naphthoquinone, 5-hydroxy-						
	CD ₃ OD	1.8×10^7			CP/LI-12	S = RB; used $k_d = 4.4 \times 10^3$ s ⁻¹ .	90F065
4.92	1,4-Naphthoquinone, 6-hydroxy-						
	CD ₃ OD	5.9×10^6			CP/LI-12	S = RB; used $k_d = 4.4 \times 10^3$ s ⁻¹ .	90F065
4.92	Phenol						
	C ₆ H ₆ / MeOH (60:40)	$<10^4$			CP/A'c-23	S = A' = Rub; k_d and $k_{A'}$ not given.	88F650
	CD ₃ OD	2.5×10^4		293	PL/Ld-2	S = HP.	88R236 90F411
	D ₂ O	1.3×10^6			PL/Ld-2	S = HP or H ₂ TPPS ⁴⁻ ; calcd. from pH dependence (7-14) and pK _a .	90F411
	H ₂ O pH = 8.0	1×10^6 (k_r)			CP/Oc-17	S = RB; A' = Met; used $k_r^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	93E536
	pH = 9.0	1.6×10^6 (k_r)					
	pH = 10.0	5.0×10^7 (k_r)					
	H ₂ O	$\sim 3 \times 10^6$ (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a ; reported $k = (2.6 \pm 4.0) \times 10^6$ L mol ⁻¹ s ⁻¹ .	91F300
	H ₂ O	$<10^6$ (k_r)		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence (8-11.5) and pK _a = 9.9.	87A180
	H ₂ O pH = 8	2.0×10^7	0.026		CP/Ac-15	S = RB; used $k_d = 5 \times 10^5$ s ⁻¹ .	82A464
	pH = 9	6.1×10^7	8.2×10^{-3}				
	pH = 10.3	2.7×10^8	1.9×10^{-3}				
4.93	Phenol, 4-acetyl-						
	H ₂ O	1.5×10^6 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
4.94	Phenol, 4-acetyl-2,6-bis(1,1-dimethylethyl)-						
	1-BuOH	1.0×10^6		293	CP/A'c-23	S = MB; A' = Np; used $k_d = 5.2 \times 10^4$ s ⁻¹ , $k_{A'} = 5.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A266
4.95	Phenol, 4-amino-						
	D ₂ O pH = 7	1×10^8			PL/Ld-2	S = AlCl(tspc).	89R092
4.96	Phenol, 2-(benzotriazol-2-yl)-4,6-bis(1,1-dimethylpropyl)-						
	<i>i</i> -octane	$<1 \times 10^6$			CP/A'c-23	S = A' = Rub; No measurable effect.	732066

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k ($\text{L mol}^{-1} \text{s}^{-1}$)	$\beta (k_d/k)$ (mol L^{-1})	T (K)	Method	Comment	Ref.
4.97	Phenol, 2-(benzotriazol-2-yl)-6-chloro-4-(1,1-dimethylethyl)-						
	$\text{CCl}_4/\text{CHCl}_3$ (90:10)	1.0×10^4			CP/A'c-23	$S = A' = \text{Rub}$; used $k_d = 1.4 \times 10^3 \text{ s}^{-1}$, $k_{A'} = 7 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$; Measured $k/(k_A[A'] + k_d) = 10.0 \text{ L mol}^{-1} \text{ s}^{-1}$ at $[A'] = 5 \times 10^{-6} \text{ mol L}^{-1}$.	74F645
4.98	Phenol, 2-benzyl-						
	$\text{C}_6\text{H}_5/\text{MeOH}$ (80:20)	1.4×10^5			PL/Ld-2	$S = \text{PrPor}$.	93E536
	C_6H_6	4.3×10^4			PL/Ld-2	$S = \text{PrPor}$.	93E536
	EtOH	2.0×10^5			PL/Ld-2	$S = \text{RB}$.	93E536
4.99	Phenol, 2,6-bis(1,1-dimethylethyl)-						
	1-BuOH	1.7×10^6		293	CP/A'c-23	$S = \text{MB}$; $A' = \text{Np}$; used $k_d = 5.2 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.2 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	78A266
	C_6H_6	$2.0 \times 10^4 (k_r)$			CP/Ac,A'c-17	$S = \text{HA}$; $A' = \text{BHT}$; used $k_r^{A'} = 1.2 \times 10^5 \text{ L mol}^{-1} \text{ s}^{-1}$.	89F350
	C_6H_6	3.2×10^4			CP/P'a-20	$S = \text{ZnTPP}$; $A' = 2\text{M2P}$; used $k_d = 4.2 \times 10^4 \text{ s}^{-1}$, $\beta_{A'} = 0.053 \text{ mol L}^{-1}$; Measured $(k[A] + k_d)/k_{A'} = 6.17 \times 10^{-2} \text{ mol L}^{-1}$ at $[A] = 0.22 \text{ mol L}^{-1}$.	78A171
	CD_3OD	2.6×10^5		293	PL/Ld-2	$S = \text{HP}$.	88R236 90F411
	CH_2Cl_2	7.5×10^5			CP/A'c-33	$S = A' = \text{Rub}$; used $k_{A'} = 7.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$; k_d not given.	84F199
	MeOH	$3.0 \times 10^4 (k_r)$			CP/Ac,A'c-17	$S = \text{HA}$; $A' = \text{BHT}$; used $k_r^{A'} = 2.2 \times 10^5 \text{ L mol}^{-1} \text{ s}^{-1}$.	89F350
	MeOH	1.0×10^6	0.12		CP/A'c-16	$S = \text{MB}$; $A' = \text{DPF}$; used $k_d = 1.2 \times 10^5 \text{ s}^{-1}$.	78A171
4.100	Phenol, 2,4-bis(1,1-dimethylethyl)-5-methoxy-						
	MeOH				CR/Pa,P'a-17	$A' = \text{TME}$; meas. $k_r/k_r^{A'} = 6.7 \times 10^{-3}$; ${}^1\text{O}_2^*$ from $\text{H}_2\text{O}_2/\text{NaOCl}$.	70F454
	MeOH				CP/Pa,P'a-17	$S = \text{RB}$; $A' = \text{TME}$; meas. $k_r/k_r^{A'} = 7.0 \times 10^{-3}$.	70F454
4.101	Phenol, 3,5-bis(1,1-dimethylethyl)-4-methoxy-						
	1-BuOH	2.6×10^7		293	CP/Ac-15	$S = \text{MB}$; used $k_d = 5.2 \times 10^4 \text{ s}^{-1}$.	78A266
	1-BuOH	2.7×10^7		293	CP/A'c-23	$S = \text{MB}$; $A' = \text{Np}$; used $k_d = 5.2 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.2 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	78A266
4.102	Phenol, 2,6-bis(1,1-dimethylethyl)-4-methyl- (BHT)						
	1-BuOH	7.6×10^6		293	CP/A'c-23	$S = \text{MB}$; $A' = \text{Np}$; used $k_d = 5.2 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.2 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	78A266
	C_6H_6	1.3×10^6 $4 \times 10^5 (k_r)$ $7.7 \times 10^5 (k_q)$		298	CP/Ac,A'c-23	$S = A' = \text{Rub}$; used $k_d = 4.2 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 4.2 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	90F360
	C_6H_6	6.6×10^5			CP/P'a-20	$S = \text{ZnTPP}$; $A' = 2\text{M2P}$; used $k_{A'} = 7.5 \times 10^5 \text{ L mol}^{-1} \text{ s}^{-1}$; meas. $k_A/k_{A'} = 0.88$.	78A171
	C_6H_6	2.5×10^6	0.016		CP/A'c-16	$S = \text{ZnTPP}$; $A' = \text{DPF}$; used $k_d = 4.2 \times 10^4 \text{ s}^{-1}$.	78A171
	C_6H_6	$1.2 \times 10^5 (k_r)$			CP/Pa,P'a-17	$S = \text{ZnTPP}$; $A' = 2\text{M2P}$; used $k_r^{A'} = 7.5 \times 10^5 \text{ L mol}^{-1} \text{ s}^{-1}$; meas. $k_r/k_{A'} = 0.16$.	78A171
	C_6H_6	8.2×10^5			CP/Pa-20	$S = \text{ZnTPP}$; $A' = \text{Car}$; used $k_{A'} = 1.2 \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$; Measured $(k_A/k_{A'}) = 1.46$.	78A171
	C_6H_6	6.1×10^5	0.06		CP/Pa-15	$S = \text{ZnTPP}$; used $k_d = 4.2 \times 10^4 \text{ s}^{-1}$.	78A171
	CCl_4	3.7×10^7			CL/LI-12	$S = \text{TPP}$; used $k_d = 1 \times 10^3 \text{ s}^{-1}$.	85F667
	CCl_4	$9.8 \times 10^6 (k_r)$			CP/Ac-17	$S = A' = \text{An}$; used $k_r^{A'} = 3.5 \times 10^5 \text{ L mol}^{-1} \text{ s}^{-1}$.	80F718
	CHCl_3	6.5×10^6			PL/Ld-2	$S = \text{MPDEE}$ or PdMPDEE .	81A327

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.102 Phenol, 2,6-bis(1,1-dimethylethyl)-4-methyl- (BHT) — Continued							
	EtOH	3.4×10^6		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
	MeOH	5.0×10^6	0.036		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	88F440
	MeOH	2.2×10^5 (k_r)			CP/Pa,P'a-17	S = MB; A' = 2M2P; used $k_r^{A'} = 6.3 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_d^{A'} = 0.36$.	78A171
	MeOH	5.6×10^6	0.021		CP/A'c-16	S = MB; A' = DPF; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	78A171
	MeOH	4.2×10^6	0.03		CP/Pa-15	S = MB; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	78A171
4.103 Phenol, 2,6-bis(1,1-dimethylethyl)-4-phenyl-							
	1-BuOH	3.9×10^6		293	CP/A'c-23	S = MB; A' = Np; used $k_d = 5.2 \times 10^4$ s ⁻¹ , $k_{A'} = 5.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A266
	MeOH	7.9×10^6	0.015		CP/A'c-16	S = MB; A' = DPF; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	78A171
4.104 Phenol, 2,6-bis(1,1-dimethylethyl)-4-(phenylmethyl)-							
	MeOH	3.8×10^6	0.031		CP/A'c-16	S = MB; A' = DPF; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	78A171
4.105 Phenol, 4-bromo-2,6-bis(1,1-dimethylethyl)-							
	1-BuOH	1.5×10^6		293	CP/A'c-23	S = MB; A' = Np; used $k_d = 5.2 \times 10^4$ s ⁻¹ , $k_{A'} = 5.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A266
	MeOH	8.4×10^5	0.14		CP/A'c-16	S = MB; A' = DPF; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	78A171
4.106 Phenol, 2-chloro-							
	H ₂ O	$\sim 9 \times 10^6$ (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a ; reported $k = (9.2 \pm 9.4) \times 10^6$ L mol ⁻¹ s ⁻¹ .	91F300
4.107 Phenol, 3-chloro-							
	H ₂ O	5.4×10^6 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
4.108 Phenol, 4-chloro-							
	H ₂ O	$\sim 6 \times 10^6$ (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
4.109 Phenol, 2-(5-chlorobenzotriazol-2-yl)-4,6-bis(1,1-dimethylethyl)-							
	C ₆ H ₅ Br	2.6×10^6		273	MD/A'c-33	A' = Rub; used $k_d = 1.3 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k/(k_d/[A'] + k_{A'}) = 2.0 \times 10^{-2}$ at $[A'] = 1.5 \times 10^{-4}$ mol L ⁻¹ .	73J333
4.110 Phenol, 4-chloro-2,6-di(1,1-dimethylethyl)-							
	1-BuOH	3.2×10^6		293	CP/A'c-23	S = MB; A' = Np; used $k_d = 5.2 \times 10^4$ s ⁻¹ , $k_{A'} = 5.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A266
4.111 Phenol, 3-chloro-5-methoxy-							
	CH ₃ CN	1.2×10^7			PL/Ld-2	S = RB.	91A341
	H ₂ O	$<1 \times 10^3$ (k_r)		293	CP/Oc-17	S = Eos; A' = His; used $k_r^{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A341
	dioxane	5.3×10^5			PL/Ld-2	S = ZnTPP.	91A341
4.112 Phenol, 4-cyano-							
	H ₂ O	$\sim 2 \times 10^5$ (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a ; reported $k = (2.4 \pm 4.4) \times 10^5$ L mol ⁻¹ s ⁻¹ .	91F300
4.113 Phenol, 2,4-dichloro-							
	C ₆ H ₆ / MeOH (60:40)	$<10^4$			CP/A'c-23	S = A' = Rub.	88F650

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.113	Phenol, 2,4-dichloro-	—Continued					
	H ₂ O	$\sim 5 \times 10^6$ (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a ; reported $k = (5.1 \pm 4.7) \times 10^6$ L mol ⁻¹ s ⁻¹ .	91F300
	H ₂ O	$< 5 \times 10^5$ (k_r)		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. over range of pH = 5.5 to 9.6; pK _a = 7.8.	87A180
4.114	Phenol, 2,6-dichloro-						
	C ₆ H ₆ / MeOH (60:40)	$< 10^4$			CP/A'c-23	S = A' = Rub.	88F650
4.115	Phenol, 2,6-dimethoxy-						
	H ₂ O	3.6×10^7 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
4.116	Phenol, 2,4-dimethyl-						
	C ₆ H ₆	3.4×10^4 (k_r)			CP/Ac,A'c-17	S = HA; A' = BHT; used $k_r^{A'} = 1.2 \times 10^5$ L mol ⁻¹ s ⁻¹ .	89F350
	MeOH	5.0×10^4 (k_r)			CP/Ac,A'c-17	S = HA; A' = BHT; used $k_r^{A'} = 2.2 \times 10^5$ L mol ⁻¹ s ⁻¹ .	89F350
4.117	Phenol, 2,6-dimethyl-						
	C ₆ H ₆	8×10^3 (k_r)			CP/Ac,A'c-17	S = HA; A' = BHT; used $k_r^{A'} = 1.2 \times 10^5$ L mol ⁻¹ s ⁻¹ .	89F350
	MeOH	1.0×10^4 (k_r)			CP/Ac,A'c-17	S = HA; A' = BHT; used $k_r^{A'} = 2.2 \times 10^5$ L mol ⁻¹ s ⁻¹ .	89F350
4.118	Phenol, 3,4-dimethyl-						
	C ₆ H ₆	2.2×10^4 (k_r)			CP/Ac,A'c-17	S = HA; A' = BHT; used $k_r^{A'} = 1.2 \times 10^5$ L mol ⁻¹ s ⁻¹ .	89F350
	MeOH	3.9×10^4 (k_r)			CP/Ac,A'c-17	S = HA; A' = BHT; used $k_r^{A'} = 2.2 \times 10^5$ L mol ⁻¹ s ⁻¹ .	89F350
4.119	Phenol, 4-(1,1-dimethylethoxy)-2,6-bis(1,1-dimethylethyl)-						
	1-BuOH	2.4×10^7		293	CP/A'c-23	S = MB; A' = Np; used $k_d = 5.2 \times 10^4$ s ⁻¹ , $k_{A'} = 5.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A266
4.120	Phenol, 2-(1,1-dimethylethyl)-						
	CD ₃ OD	2.6×10^5		293	PL/Ld-2	S = HP.	88R236 90F411
4.121	Phenol, 4-(1,1-dimethylethyl)-						
	H ₂ O	$\sim 10^7$ (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a ; reported $k = (1.2 \pm 1.3) \times 10^7$ L mol ⁻¹ s ⁻¹ .	91F300
4.122	Phenol, 3-(1,1-dimethylethyl)-4-methoxy-						
	CD ₃ OD	2.2×10^7		293	PL/Ld-2	S = HP.	88R236 90F411
4.123	Phenol, 2,4-dinitro-						
	C ₆ H ₆	$< 1.4 \times 10^6$ (k_r)			CP/A'c-23	S = A' = Rub; used $k_d = \geq 3.3 \times 10^4$ s ⁻¹ , $k_{A'} = 3.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87E142
4.124	Phenol, 2,5-dinitro-						
	C ₆ H ₆	$< 1.6 \times 10^6$			CP/A'c-23	S = A' = Rub; used $k_d = \geq 3.3 \times 10^4$ s ⁻¹ , $k_{A'} = 3.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87E142

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.125	Phenol, 2,6-dinitro-						
	C ₆ H ₆	<2.0 × 10 ⁶			CP/A'c-23	S = A' = Rub; used $k_d = 2.3 \times 10^4$ s ⁻¹ , $k_{A'} = 3.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87E142
4.126	Phenol, 4-ethoxy-2,3,5,6-tetramethyl-						
	MeOH	7.3 × 10 ⁷	1.6 × 10 ⁻³		CP/A'c-16	S = MB; A' = DPF; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	78A171
4.127	Phenol, 4-[1-hydroxy-2,2-dimethyl)propyl]-2-methoxy-						
	CH ₃ CH(OCH ₃) ₂	8.5 × 10 ⁵	0.037		MP/Pa-15	S = 4-HOC ₆ H ₃ (3-OCH ₃)COCH ₃ ; used $k_d = 3.1 \times 10^4$ s ⁻¹ .	84F194
	CH ₃ CH(OCH ₃) ₂	1.1 × 10 ⁸			CP/Ac-32	S = 4-CH ₃ OC ₆ H ₄ COC ₂ H ₅ ; Q = Car; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $k_Q = 1.3 \times 10^{10}$ L mol ⁻¹ s ⁻¹ .	84F194
	CH ₃ CH(OCH ₃) ₂	1.2 × 10 ⁶	0.025		MP/Ac-15	S = 4-HOC ₆ H ₃ (3-OCH ₃)COCH ₃ ; used $k_d = 3.1 \times 10^4$ s ⁻¹ .	84F194
	CH ₃ CH(OCH ₃) ₂	3.4 × 10 ⁵	0.16		MP/Ac-16	S = 4-HOC ₆ H ₃ (3-OCH ₃)COCH ₃ ; A' = DABCO; used $k_d = 3.1 \times 10^4$ s ⁻¹ ; unexplained discrepancy in these data.	84F194
4.128	Phenol, 2-(2-hydroxyphenyl)- (2,2'-Dihydroxybiphenyl)						
	D ₂ O pD = 8.5	2.7 × 10 ⁷			PL/Ld-2	S = RB.	93E536
	EtOH	7.5 × 10 ⁶			PL/Ld-2	S = RB.	93E536
4.129	Phenol, 2-methoxy-						
	H ₂ O	~6 × 10 ⁶ (k_t)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_t^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a ; reported $k = (0.6 \pm 1.8) \times 10^7$ L mol ⁻¹ s ⁻¹ .	91F300
4.130	Phenol, 3-methoxy-						
	H ₂ O	~1 × 10 ⁷ (k_t)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_t^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a ; reported $k = (1.3 \pm 1.0) \times 10^7$ L mol ⁻¹ s ⁻¹ .	91F300
4.131	Phenol, 4-methoxy-						
	CD ₃ OD	5.8 × 10 ⁶		293	PL/Ld-2	S = HP.	88R236 90F411
	H ₂ O	~2 × 10 ⁷ (k_t)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_t^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a ; reported $k = (2.2 \pm 2.2) \times 10^7$ L mol ⁻¹ s ⁻¹ .	91F300
4.132	Phenol, 4-methoxy-2,6-dimethyl-						
	EtOH	2.5 × 10 ⁷		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	91F285
4.133	Phenol, 4-methoxy-2,3,5,6-tetramethyl-						
	EtOH	1.4 × 10 ⁷		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	91F285
4.134	Phenol, 4-methoxy-2,3,6-trimethyl-						
	EtOH	6.8 × 10 ⁷		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	91F285
4.135	Phenol, 4-methyl-						
	D ₂ O/ EtOH (75:25)	2.4 × 10 ⁷		295	PL/Ld-2	S = RB.	94A113
	H ₂ O	1 × 10 ⁷ (k_t)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_t^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
	H ₂ O	<5 × 10 ⁶ (k_t)		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_t^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. over the range pH = 8.3 to 11.5; pK _a = 10.2.	87A180

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.136	Phenol, 2,2'-methylenebis(6- <i>tert</i> -butyl-4-methyl-						
	MeOH	2.6×10^7	7.0×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	88F440
4.137	Phenol, 2-nitro-						
	H ₂ O	-1×10^6 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a ; reported $k = (1.3 \pm 1.1) \times 10^6$ L mol ⁻¹ s ⁻¹ .	91F300
4.138	Phenol, 3-nitro-						
	H ₂ O	-3×10^6 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a ; reported $k = (2.7 \pm 1.8) \times 10^6$ L mol ⁻¹ s ⁻¹ .	91F300
4.139	Phenol, 4-nitro-						
	H ₂ O	-3×10^5 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a ; reported $k = (2.6 \pm 2.3) \times 10^5$ L mol ⁻¹ s ⁻¹ .	91F300
4.140	Phenol, pentachloro-						
	H ₂ O	-2×10^5 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a ; reported $k = (0.2 \pm 5.5) \times 10^6$ L mol ⁻¹ s ⁻¹ .	91F300
4.141	Phenol, 2-phenyl- (2-Hydroxybiphenyl)						
	C ₆ H ₆ / MeOH (60:40)	7.3×10^4			PL/Ld-2	S = PrPor.	93E536
	C ₆ H ₆	2.3×10^4			PL/Ld-2	S = PrPor.	93E536
4.142	Phenol, 3-(phenylamino)-						
	CH ₃ CN/ H ₂ O (96:4)	3.0×10^7	1.1×10^{-3}		CP/Pa-14	S = MB; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	80F427
4.143	Phenol, 2,3,5,6-tetrafluoro-						
	CD ₃ OD	3.0×10^4		293	PL/Ld-2	S = HP.	88R236 90F411
4.144	Phenol, 4-(1,1,3,3-tetramethylbutyl)-, salicylate						
	<i>i</i> -octane	$<1 \times 10^6$			CP/A'c-23	S = A' = Rub; No measurable effect.	732066
4.145	Phenol, 2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)-]						
	<i>i</i> -octane	$<1.0 \times 10^6$			CP/A'c-23	S = A' = Rub; No measurable effect.	732066
4.146	Phenol, 2,4,6-trichloro-						
	C ₆ H ₆ / MeOH (60:40)	$<10^4$			CP/A'c-16	S = A' = Rub.	88F650
	H ₂ O	2×10^7 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
	H ₂ O	$<5 \times 10^5$ (k_r)		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. over the range pH = 4.2 to 9.0; pK _a = 6.1.	87A180
4.147	Phenol, 2,4,6-tri(1,1-dimethylethyl)-						
	1-BuOH	3.7×10^6		293	CP/A'c-23	S = MB; A' = Np; used $k_d = 5.2 \times 10^4$ s ⁻¹ , $k_{A'} = 5.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A266
	c-C ₆ H ₁₂	5.8×10^5			PL/A'd-8	S = 2-ACN; A' = DPBF; $\Delta H^\ddagger = -0.8$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -147$ J K ⁻¹ mol ⁻¹ ; studied at 278–357 K.	84E492
	C ₆ H ₅ CH ₃	1.4×10^6			PL/A'd-8	S = 2-ACN; A' = DPBF.	84E492
	C ₆ H ₆	2.6×10^5			CP/P'a-19	S = ZnTPP; A' = 2M2P; used $k_{A'} = 7.5 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 0.34$.	78A171

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.147 Phenol, 2,4,6-tri(1,1-dimethylethyl). — Continued							
	C ₆ H ₆	1.4×10^6	0.030		CP/A'c-16	S = ZnTPP; A' = DPF; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	78A171
	CH ₃ CN	2.5×10^6			PL/A'd-8	S = 2-ACN; A' = DPBF.	84E492
	CH ₃ COCH ₃	2.4×10^6			PL/A'd-8	S = 2-ACN; A' = DPBF.	84E492
	MeOH	2.8×10^6			PL/A'd-8	S = 2-ACN; A' = DPBF; $\Delta H^\ddagger = -3.4$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -135$ J K ⁻¹ mol ⁻¹ ; studied at 250-328 K.	84E492
	MeOH	3.4×10^6	0.034		CP/A'c-16	S = MB; A' = DPF; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	78A171
	MeOH	$6.3 \times 10^4 (k_r)$			CP/Pa,P'a-17	S = RB; A' = BHT; used $k_r^{A'} = 2.2 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.29$.	78A171
4.148 Phenol, 2,4,6-trimethyl-							
	EtOH	3.8×10^6		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	91F285
4.149 Phenol, 2,4,6-triphenyl-							
	C ₆ H ₆	2.2×10^7	1.9×10^{-3}		CP/A'c-16	S = ZnTPP; A' = DPF; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	78A171
	CH ₃ CN	1.5×10^7	2.3×10^{-3}		CP/A'c-16	S = MB; A' = DPF; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	78A171
	MeOH	2.5×10^8	4.6×10^{-4}		CP/A'c-16	S = MB; A' = DPF; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	78A171
4.150 Phenoxide ion							
	C ₆ H ₆ /MeOH (60:40)	7.0×10^7			CP/A'c-23	S = A' = Rub; k_d and $k_{A'}$ not given; 6×10^{-3} mol L ⁻¹ KOH.	88F650
	D ₂ O	2.8×10^8			PL/Ld-2	S = HP or H ₂ TPPS ⁴⁻ ; calcd. from pH dependence (7-14) and pK _a .	90F411
	H ₂ O	$1.6 \times 10^8 (k_r)$		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
	H ₂ O	$1.8 \times 10^8 (k_r)$		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. over the range pH = 8.0 to 11.5.	87A180
4.151 Phenoxide ion, 4-acetyl-							
	H ₂ O	$2.4 \times 10^7 (k_r)$		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
4.152 Phenoxide ion, 2-benzyl-							
	EtOH	1.6×10^8			PL/Ld-2	S = RB; 7.5×10^{-3} mol L ⁻¹ KOH.	93E536
	H ₂ O pH = 11	$4.4 \times 10^7 (k_r)$			CP/Oc-17	S = RB; A' = Met; used $k_r^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	93E536
4.153 Phenoxide ion, 4-bromo-							
	C ₆ H ₅ COCH ₃ /MeOH (50:50)	6×10^7		295	CP/A'c-23	S = A' = Rub; used $k_d = 3 \times 10^4$ s ⁻¹ ; $k_{A'}$ not given; 0.01 mol L ⁻¹ KOH.	93F328
	C ₆ H ₆ /MeOH (80:20)	$1.5 \times 10^7 (k_r)$		295	CP/Ac,A'c-17	S = Rub; A' = DMA; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	93F328
	C ₆ H ₆ /MeOH (80:20)	1.5×10^7		295	CP/A'c-23	S = A' = Rub; used $k_d = 3.8 \times 10^4$ s ⁻¹ ; $k_{A'}$ not given; 0.01 mol L ⁻¹ KOH.	93F328
	CH ₃ CN/MeOH	6×10^7			CP/A'c-16	S = A' = Rub; used $k_d = 3 \times 10^5$ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	94F095
	H ₂ O pH = 11	1.1×10^9			CP/Ac-14	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	94F095
	H ₂ O pH = 11	$1.5 \times 10^8 (k_r)$			CP/Ac,A'c-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	94F095
	H ₂ O pH = 11	$1.5 \times 10^8 (k_r)$		295	CP/Oc-17	S = Eos; A' = Met; used $k_r^{A'} = 2.0 \times 10^7$ L mol ⁻¹ s ⁻¹ .	93F328
	H ₂ O pH = 11	1.1×10^9		295	CP/Oc-15	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	93F328

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.153 Phenoxide ion, 4-bromo- — Continued							
	MeOH/ C ₆ H ₆ (60:40)	1.6×10^7 (k_r)			CP/Ac,A'c-17	S = RB; A' = DMA; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	94F095
	MeOH/ C ₆ H ₆ (60:40)	1.5×10^7			CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	94F095
4.154 Phenoxide ion, 4-bromo-2,6-dimethyl-							
	CH ₃ CN/MeOH	2.1×10^8			CP/A'c-16	S = A' = Rub; used $k_d = 3 \times 10^5$ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	94F095
	H ₂ O pH = 11	2.6×10^9			CP/Ac-14	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	94F095
	H ₂ O pH = 11	3.9×10^8 (k_r)			CP/Ac,A'c-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	94F095
	MeOH/ C ₆ H ₆ (60:40)	8×10^7			CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	94F095
	MeOH/ C ₆ H ₆ (60:40)	5×10^7 (k_r)			CP/Ac,A'c-17	S = RB; A' = DMA; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	94F095
4.155 Phenoxide ion, 2-chloro-							
	H ₂ O	1.9×10^8 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
	H ₂ O pH = 11.5	1.7×10^7 (k_r)			CP/Oc-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F518
	H ₂ O pH = 11.5	2.3×10^8			CP/Ac,Oc-14	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	90F518
	MeOH	5.0×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^5$ s ⁻¹ , $k_{A'} = 5 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	90F518
4.156 Phenoxide ion, 3-chloro-							
	H ₂ O	1.6×10^8 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
	H ₂ O pH = 11.5	1.5×10^7 (k_r)			CP/Oc-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F518
	H ₂ O pH = 11.5	1.1×10^9			CP/Ac,Oc-14	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	90F518
	MeOH	1.5×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^5$ s ⁻¹ , $k_{A'} = 5 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	90F518
4.157 Phenoxide ion, 4-chloro-							
	C ₆ H ₅ COCH ₃ / MeOH (50:50)	6×10^7		295	CP/A'c-23	S = A' = Rub; used $k_d = 3 \times 10^4$ s ⁻¹ , $k_{A'} =$ not given; 0.01 mol L ⁻¹ KOH.	93F328
	C ₆ H ₆ / MeOH (80:20)	3×10^7		295	CP/A'c-23	S = A' = Rub; used $k_d = 3.8 \times 10^4$ s ⁻¹ , $k_{A'} =$ not given; 0.01 mol L ⁻¹ KOH.	93F328
	C ₆ H ₆ / MeOH (80:20)	3×10^7 (k_r)		295	CP/Ac,A'c-17	S = Rub; A' = DMA; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	93F328
	H ₂ O pH = 11	8.0×10^8		295	CP/Oc-15	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	93F328
	H ₂ O pH = 11	1.0×10^8 (k_r)		295	CP/Oc-17	S = Eos; A' = Met; used $k_r^{A'} = 2.0 \times 10^7$ L mol ⁻¹ s ⁻¹ .	93F328
	H ₂ O	1.9×10^8 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
	H ₂ O pH = 11.5	3.1×10^7 (k_r)			CP/Oc-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F518
	H ₂ O pH = 11.5	4×10^8			CP/Ac,Oc-14	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	90F518

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.157 Phenoxide ion, 4-chloro- — Continued							
	MeOH	3.3×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^5$ s ⁻¹ , $k_{A'} = 5 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	90F518
4.158 Phenoxide ion, 4-cyano-							
	H ₂ O	6.2×10^6 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
4.159 Phenoxide ion, 2,4-dichloro-							
	C ₆ H ₆ / MeOH (60:40)	5.7×10^7			CP/A'c-23	S = A' = Rub; k_d and $k_{A'}$ not given; 6×10^{-3} mol L ⁻¹ KOH.	88F650
	C ₆ H ₆ / MeOH (60:40)	4.5×10^6 (k_r)			CP/Ac,A'c-17	S = Eos; A' = DMA; used $k_r^{A'} = 4.8 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_r = 11.5$; 6×10^{-3} mol L ⁻¹ KOH.	88F650
	H ₂ O	1.7×10^8 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
	H ₂ O	1.2×10^8 (k_r)		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. over the range pH = 5.5 to 9.6.	87A180
	H ₂ O pH = 10	3.1×10^9			CP/Ac-14	S = RB; used $k_d = 5 \times 10^5$ s ⁻¹ .	87F537
	H ₂ O pH = 10	2.1×10^9			CP/Ac,Oc-14	S = Eos; used $k_d = 5 \times 10^5$ s ⁻¹ .	87F537
	H ₂ O pH = 10	1.1×10^8 (k_r)			CP/Oc-17	S = Eos; A' = Met; used $k_r^{A'} = 5 \times 10^6$ L mol ⁻¹ s ⁻¹ ; used $k_A = 2.1 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87F537
4.160 Phenoxide ion, 2,6-dichloro-							
	C ₆ H ₆ / MeOH (60:40)	3.7×10^6 (k_r)			CP/Ac,A'c-17	S = Eos; A' = DMA; used $k_r^{A'} = 4.8 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_r = 14.5$; 6×10^{-3} mol L ⁻¹ KOH.	88F650
	C ₆ H ₆ / MeOH (60:40)	5.4×10^7			CP/A'c-23	S = A' = Rub; k_d and $k_{A'}$ not given; 6×10^{-3} mol L ⁻¹ KOH.	88F650
	H ₂ O pH = 10	1×10^9			CP/Ac-14	S = Eos; used $k_d = 5 \times 10^5$ s ⁻¹ .	87F537
	H ₂ O pH = 10	2×10^9			CP/Ac-14	S = RB; used $k_d = 5 \times 10^5$ s ⁻¹ .	87F537
	H ₂ O pH = 10	1.5×10^9			CP/Oc-14	S = Eos; used $k_d = 5 \times 10^5$ s ⁻¹ .	87F537
	H ₂ O pH = 10	5.7×10^7 (k_r)			CP/Oc-17	S = Eos; A' = Met; used $k_r^{A'} = 5 \times 10^6$ L mol ⁻¹ s ⁻¹ ; used $k_A = 1.5 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87F537
4.161 Phenoxide ion, 2,6-dimethoxy-							
	H ₂ O	8.6×10^8 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
4.162 Phenoxide ion, 2,6-dimethyl-							
	CH ₃ CN/MeOH	2×10^8			CP/A'c-16	S = A' = Rub; used $k_d = 3 \times 10^5$ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	94F095
	H ₂ O pH = 11	2.8×10^9			CP/Ac-14	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	94F095
	H ₂ O pH = 11	2.0×10^8 (k_r)			CP/Ac,A'c-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	94F095
	MeOH/ C ₆ H ₆ (60:40)	3×10^7			CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	94F095
	MeOH/ C ₆ H ₆ (60:40)	1.2×10^7 (k_r)			CP/Ac,A'c-17	S = RB; A' = DMA; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	94F095

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.163	Phenoxyde ion, 4-(1,1-dimethylethyl)-						
	H ₂ O	3.6×10^8 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
4.164	Phenoxyde ion, 2,4-dinitro-						
	H ₂ O	4×10^5 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; pK _a of 2,4-dinitrophenol = 4.07.	91F300
4.165	Phenoxyde ion, 4-fluoro-						
	C ₆ H ₆ / MeOH (80:20)	2.0×10^8		295	CP/A'c-23	S = A' = Rub; used $k_d = 3.8 \times 10^4$ s ⁻¹ ; $k_{A'}$ not given; 0.01 mol L ⁻¹ KOH.	93F328
	C ₆ H ₆ / MeOH (80:20)	5.7×10^7 (k_r)		295	CP/Ac,A'c-17	S = Rub; A' = DMA; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	93F328
	H ₂ O pH = 11	2.5×10^9		295	CP/Oc-15	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	93F328
	H ₂ O pH = 11	3.5×10^8 (k_r)		295	CP/Oc-17	S = Eos; A' = Met; used $k_r^{A'} = 2.0 \times 10^7$ L mol ⁻¹ s ⁻¹ .	93F328
4.166	Phenoxyde ion, 2-(2-hydroxyphenyl)- (2,2'-Dihydroxybiphenyl, conjugate base)						
	D ₂ O pD = 11.8	1.2×10^8			PL/Ld-2	S = RB.	93E536
	EtOH	7×10^6 (k_r)			CP/Oc,A'c-17	S = RB; A' = DMA; used $k_r^{A'} = 4.4 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 7.5×10^{-3} mol L ⁻¹ KOH.	93E536
	EtOH	1.1×10^8			PL/Ld-2	S = RB; 7.5×10^{-3} mol L ⁻¹ KOH.	93E536
	EtOH	9.4×10^7			PL/Ld-2	S = RB; 7.5×10^{-3} mol L ⁻¹ KOH.	93E536
	H ₂ O pH = 11	2.2×10^7 (k_r)			CP/Oc-17	S = RB; A' = Met; used $k_r^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	93E536
4.167	Phenoxyde ion, 5-hydroxy-2-phenyl- (2,5-Dihydroxybiphenyl, conjugate base)						
	EtOH	9.9×10^7			PL/Ld-2	S = RB; 7.5×10^{-3} mol L ⁻¹ KOH.	93E536
4.168	Phenoxyde ion, 4-iodo-						
	C ₆ H ₆ / MeOH (80:20)	1.4×10^7		295	CP/A'c-23	S = A' = Rub; used $k_d = 3.8 \times 10^4$ s ⁻¹ ; $k_{A'}$ not given; 0.01 mol L ⁻¹ KOH.	93F328
	C ₆ H ₆ / MeOH (80:20)	1.0×10^7 (k_r)		295	CP/Ac,A'c-17	S = Rub; A' = DMA; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	93F328
	H ₂ O pH = 11	8×10^7 (k_r)		295	CP/Oc-17	S = Eos; A' = Met; used $k_r^{A'} = 2.0 \times 10^7$ L mol ⁻¹ s ⁻¹ .	93F328
	H ₂ O pH = 11	1.0×10^9		295	CP/Oc-15	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	93F328
4.169	Phenoxyde ion, 2-methoxy-						
	H ₂ O	4.4×10^8 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
4.170	Phenoxyde ion, 3-methoxy-						
	H ₂ O	2.9×10^8 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
4.171	Phenoxyde ion, 4-methoxy-						
	H ₂ O	6.7×10^8 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
4.172	Phenoxyde ion, 4-methyl-						
	H ₂ O	3.5×10^8 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
	H ₂ O	3.7×10^8 (k_r)		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. over the range pH = 8.3 to 11.5.	87A180

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.	
4.173	Phenoxyde ion, 2-methyl-4,6-dinitro-							
	H ₂ O	$1.3 \times 10^5 (k_r)$		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; pK _a of 2-methyl-4,6-dinitrophenol = 4.7.	91F300	
4.174	Phenoxyde ion, 4-methyl-2,6-dinitro-							
	H ₂ O	$1.4 \times 10^7 (k_r)$		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; pK _a of 4-methyl-2,6-dinitrophenol = 4.23.	91F300	
4.175	Phenoxyde ion, 4-methyl-2-nitro-							
	CH ₃ CN/H ₂ O (90:10)	$6 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = RB; A' = 9-Anthracenemethanol; $k_r^{A'}$ not given; 0.01 mol L ⁻¹ KOH.	92N330	
	CH ₃ CN/H ₂ O (90:10)	3.4×10^8			PL/Ld-2	S = RB; 0.01 mol L ⁻¹ KOH.	92N330	
	C ₆ H ₆ /MeOH (60:40)	$2.9 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = RB; A' = DMA; $k_r^{A'}$ not given; 0.01 mol L ⁻¹ KOH.	92N330	
	C ₆ H ₆ /MeOH (60:40)	4.5×10^7			PL/Ld-2	S = RB; 0.01 mol L ⁻¹ KOH.	92N330	
	H ₂ O pH = 10	$2.8 \times 10^7 (k_r)$			CP/Oc-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F518 94F095	
	H ₂ O pH = 10	4.3×10^9			CP/Ac,Oc-14	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	90F518 94F095	
	MeOH/C ₆ H ₆ (60:40)	$2.9 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = RB; A' = DMA; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	94F095	
	MeOH/C ₆ H ₆ (60:40)	4.5×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 5 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH; cor. for quenching of singlet rubrene by A.	90F518 94F095	
4.176	Phenoxyde ion, 2-nitro-							
	CH ₃ CN/H ₂ O (90:10)	3.3×10^7			PL/Ld-2	S = RB; 0.01 mol L ⁻¹ KOH.	92N330	
	CH ₃ CN/H ₂ O (90:10)	$5.8 \times 10^6 (k_r)$			CP/Ac,A'c-17	S = RB; A' = 9-Anthracenemethanol; $k_r^{A'}$ not given; 0.01 mol L ⁻¹ KOH.	92N330	
	C ₆ H ₆ /MeOH (60:40)	$3.2 \times 10^6 (k_r)$			CP/Ac,A'c-17	S = RB; A' = DMA; $k_r^{A'}$ not given; 0.01 mol L ⁻¹ KOH.	92N330	
	C ₆ H ₆ /MeOH (60:40)	3.1×10^6			PL/Ld-2	S = RB; 0.01 mol L ⁻¹ KOH.	92N330	
	H ₂ O	$3.4 \times 10^7 (k_r)$		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300	
	H ₂ O pH = 10	1.5×10^9			CP/Ac,Oc-14	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	90F518 94F095	
	H ₂ O pH = 10	$1.6 \times 10^7 (k_r)$			CP/Oc-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F518 94F095	
	MeOH/C ₆ H ₆ (60:40)	$3 \times 10^6 (k_r)$			CP/Ac,A'c-17	S = RB; A' = DMA; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	94F095	
	MeOH/C ₆ H ₆ (60:40)	3.0×10^6			CP/A'c-23	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 5 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH; cor. for quenching of singlet rubrene by A.	90F518	
4.177	Phenoxyde ion, 3-nitro-							
	H ₂ O	$6.1 \times 10^7 (k_r)$			300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
	H ₂ O pH = 10	$1.0 \times 10^7 (k_r)$			CP/Oc-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F518	
	H ₂ O pH = 10	2.0×10^9			CP/Ac,Oc-14	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	90F518	

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.177	Phenoxide ion, 3-nitro-	— Continued					
	MeOH/ C ₆ H ₆ (60:40)	9.0×10^6			CP/A'c-23	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 5 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH; cor. for quenching of singlet rubrene by A.	90F518
4.178	Phenoxide ion, 4-nitro-						
	H ₂ O	5.1×10^6 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
	H ₂ O pH = 10	6×10^8			CP/Ac,Oc-14	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	90F518
	H ₂ O pH = 10	3×10^6 (k_r)			CP/Oc-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F518
	H ₂ O pH = 8.8	3×10^6 (k_r)		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; pK _a of 2-nitrophenol = 7.2.	87A180
	MeOH/ C ₆ H ₆ (60:40)	<10 ³			CP/A'c-23	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 5 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH; cor. for quenching of singlet rubrene by A.	90F518
4.179	Phenoxide ion, pentachloro-						
	H ₂ O	9.4×10^7 (k_r)		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
4.180	Phenoxide ion, 2-phenyl- (2-Hydroxybiphenyl, conjugate base)						
	D ₂ O pD = 9.0	2×10^7			PL/Ld-2	S = RB.	93E536
	pD = 10.0	3.8×10^7					
	pD = 11.8	2.3×10^8					
	EtOH	2.2×10^8			PL/Ld-2	S = RB; 7.5×10^{-3} mol L ⁻¹ KOH.	93E536
	EtOH	4.3×10^7 (k_r)			CP/Oc,A'c-17	S = RB; A' = DMA; used $k_r^{A'} = 4.4 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 7.5×10^{-3} mol L ⁻¹ KOH.	93E536
	H ₂ O pH = 8.0	3.2×10^5 (k_r)			CP/Oc-17	S = RB; A' = Met; used $k_r^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	93E536
	pH = 9.0	3.2×10^6 (k_r)					
	pH = 10.0	5.9×10^6 (k_r)					
	pH = 11	4.4×10^7 (k_r)					
4.181	Phenoxide ion, 4-phenyl- (4-Hydroxybiphenyl, conjugate base)						
	C ₆ H ₆ / MeOH (80:20)	1×10^7 (k_r)		295	CP/Ac,A'c-17	S = Rub; A' = DMA; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	93F328
	C ₆ H ₆ / MeOH (80:20)	4×10^7		295	CP/A'c-23	S = A' = Rub; used $k_d = 3.8 \times 10^4$ s ⁻¹ ; k _{A'} not given; 0.01 mol L ⁻¹ KOH.	93F328
	D ₂ O pD = 11.8	7.8×10^7			PL/Ld-2	S = RB.	93E536
	D ₂ O pD = 11.8	8×10^7		295	CP/Oc-15	S = Eos; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	93F328
	EtOH	1.3×10^8			PL/Ld-2	S = RB; 7.5×10^{-3} mol L ⁻¹ KOH.	93E536
	H ₂ O pH = 11	3.8×10^7 (k_r)			CP/Oc-17	S = RB; A' = Met; used $k_r^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	93E536
	H ₂ O pH = 11	4×10^7 (k_r)		295	CP/Oc-17	S = Eos; A' = Met; used $k_r^{A'} = 2.0 \times 10^7$ L mol ⁻¹ s ⁻¹ .	93F328
4.182	Phenoxide ion, 2,4,6-trichloro-						
	C ₆ H ₆ / MeOH (60:40)	4.0×10^6 (k_r)			CP/Ac,A'c-17	S = Eos; A' = DMA; used $k_r^{A'} = 4.8 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_d/k_r = 4.5$; 6×10^{-3} mol L ⁻¹ KOH.	88F650
	C ₆ H ₆ / MeOH (60:40)	2.2×10^7			CP/A'c-23	S = A' = Rub; k _d and k _{A'} not given; 6×10^{-3} mol L ⁻¹ KOH.	88F650

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
Phenoxyde ion, 2,4,6-trichloro- — Continued							
	H ₂ O	$1.7 \times 10^8 (k_r)$		300	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. from pH dependence and pK _a .	91F300
	H ₂ O	$1.2 \times 10^8 (k_r)$		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; calcd. over the range pH = 4.2 to 9.0.	87A180
	H ₂ O pH = 10	6×10^8			CP/Ac-14	S = Eos; used $k_d = 5 \times 10^5$ s ⁻¹ .	87F537
	H ₂ O pH = 10	4×10^8			CP/Ac-14	S = RB; used $k_d = 5 \times 10^5$ s ⁻¹ .	87F537
	H ₂ O pH = 10	7.0×10^8			CP/Oc-14	S = Eos; used $k_d = 5 \times 10^5$ s ⁻¹ .	87F537
	H ₂ O pH = 10	$5.9 \times 10^7 (k_r)$			CP/Oc-17	S = Eos; A' = Met; used $k_r^{A'} = 5 \times 10^6$ L mol ⁻¹ s ⁻¹ ; used $k_A = 7.0 \times 10^8$ L mol ⁻¹ s ⁻¹ .	87F537
		$6.4 \times 10^8 (k_r)$					
L-Phenylalanine, 3,4-dihydroxy- (DOPA)							
	D ₂ O	2.2×10^7			PL/Ld-2	S = MB.	91R251
	D ₂ O pD = 7.6	3×10^7	6×10^{-4}	296	CP/Oc-15	S = HP; used $k_d = 1.8 \times 10^4$ s ⁻¹ .	84F075
	H ₂ O	9×10^8	5.6×10^{-4}		CP/Ac-15	S = MB; used $k_d = 5 \times 10^5$ s ⁻¹ .	79F314
	MeOH/ H ₂ O (58:42)		1.2×10^{-3}	293	CP/Ac-15	S = NAZ.	79F315
	MeOH/ H ₂ O (58:42)		1.3×10^{-3}		CP/Ac-15	S = 5,8-DMOP.	81R052
	MeOH/ H ₂ O (58:42)		1.1×10^{-3}		CP/Ac-15	S = 5-MOP.	81R052
	MeOH/ H ₂ O (58:42)		1.3×10^{-3}		CP/Ac-15	S = Pso.	81R052
	MeOH/ H ₂ O (58:42)		1.3×10^{-3}		CP/Ac-15	S = TMPs.	81R052
	MeOH/ H ₂ O (58:42)		1×10^{-3}		CP/Ac-15	S = 8-MOP.	79F314
	MeOH/ H ₂ O (58:42)		1.2×10^{-3}		CP/Ac-15	S = MB.	79F314
2,2'-Spirobi[1-benzopyran]-6,6'-diol, 7,7'-bis(1,1-dimethylethyl)-3,3',4,4'-tetrahydro-4,4',4'-tetramethyl-							
	MeOH	9.2×10^7			CP/A'c-16	S = MB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	91F516
	MeOH	$2.2 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_r^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ .	91F516
	MeOH	9.8×10^7	1.8×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	88F440
2,2'-Spirobi[1-benzopyran]-6,6'-diol, 3,3',4,4'-tetrahydro-4,4',4',7,7',7'-hexamethyl-							
	MeOH	1.2×10^8			CP/A'c-16	S = MB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	91F516
	MeOH	$3.1 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_r^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ .	91F516
	MeOH	9.2×10^7	2.0×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	88F440
Tyramine							
	D ₂ O/ EtOH (75:25)	5×10^6		295	PL/Ld-2	S = RB.	94A113
	H ₂ O pH = 10	2.8×10^8		298	CP/Ac-?	S = MB; A' = N ₃ ⁻ ; k and k_r by computer fit of rate parameters to experimental ϕ_{ox} versus [A].	777489
Tyrosine							
	D ₂ O pD = 8.4	$8 \times 10^6 (k_r)$		293	CL/Ad-35	used $k_d = 2.9 \times 10^4$ s ⁻¹ ; high pressure O ₂ (0.195 mol L ⁻¹); statistical error of 30% and systematic error of the same order.	79A112

TABLE 4. Rate constants for interaction of singlet oxygen with phenols and naphthols. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
4.187 Tyrosine — Continued							
	D ₂ O/ EtOH (75:25)	$\sim 2 \times 10^6$		295	PL/Ld-2	S = RB.	94A113
	EtOH/ CH ₃ CN (80:20)	6.2×10^6			PL/Ld-2	S = ZnTPP.	91A252
	EtOH/ CH ₃ CN (80:20)	3.1×10^6 (k_r)			CP/Ac,A'c-14,28	S = ZnTPP; A' = DMA; used $k_d = 5.3 \times 10^4$ s ⁻¹ ; used $k_A = 6.2 \times 10^6$ L mol ⁻¹ s ⁻¹ .	91A252
	EtOH/ CH ₃ CN (80:20)	5.5×10^6			PL/Ld-2	S = ZnTPP; 0.01 mol L ⁻¹ HCl.	91A252
	EtOH/ CH ₃ CN (80:20)	$<5 \times 10^4$ (k_r)			CP/Ac,A'c-14,28	S = ZnTPP; A' = DMA; used $k_d = 5.3 \times 10^4$ s ⁻¹ , $k_A = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; used $k_A = 5.5 \times 10^6$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ HCl.	91A252
	EtOH/ CH ₃ CN (80:20)	1.8×10^8			PL/Ld-2	S = ZnTPP; 0.01 mol L ⁻¹ KOH.	91A252
	EtOH/ CH ₃ CN (80:20)	3×10^7 (k_r)			CP/Ac,A'c-14,28	S = ZnTPP; A' = DMA; used $k_d = 5.3 \times 10^4$ s ⁻¹ ; used $k_A = 1.8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	91A252
	H ₂ O pH = 10	3.8×10^7 (k_r)			CP/Oc-17	S = RB; A' = Met; used $k_r^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A252
	H ₂ O pH = 11.5	5.2×10^7 (k_r)			CP/Oc-17	S = Eos; A' = Met; used $k_r^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A252
	H ₂ O pH = 7.0		0.017	283	CP/Oc-15	S = MB; The mechanism of oxidation is not clear.	65F029
4.188 L-Tyrosine, N-acetyl-, ethyl ester	D ₂ O/ EtOH (75:25)	8×10^6		295	PL/Ld-2	S = RB.	94A113
4.189 Tyrosine, glycyl-							
	D ₂ O/ EtOH (75:25)	1.1×10^7		295	PL/Ld-2	S = RB.	94A113
	EtOH/ CH ₃ CN (80:20)	3.5×10^6			PL/Ld-2	S = ZnTPP.	91A252
	EtOH/ CH ₃ CN (80:20)	3×10^5 (k_r)			CP/Ac,A'c-14,28	S = ZnTPP; A' = DMA; used $k_d = 5.3 \times 10^4$ s ⁻¹ ; used $k_A = 3.5 \times 10^6$ L mol ⁻¹ s ⁻¹ .	91A252
	EtOH/ CH ₃ CN (80:20)	9.4×10^7			PL/Ld-2	S = ZnTPP; 0.01 mol L ⁻¹ KOH.	91A252
	EtOH/ CH ₃ CN (80:20)	2.7×10^6 (k_r)			CP/Ac,A'c-14,28	S = ZnTPP; A' = DMA; used $k_d = 5.3 \times 10^4$ s ⁻¹ ; used $k_A = 9.4 \times 10^7$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ KOH.	91A252
	EtOH/ CH ₃ CN (80:20)	5.5×10^6			PL/Ld-2	S = ZnTPP; 0.01 mol L ⁻¹ HCl.	91A252
	EtOH/ CH ₃ CN (80:20)	$<5 \times 10^4$ (k_r)			CP/Ac,A'c-14,28	S = ZnTPP; A' = DMA; used $k_d = 5.3 \times 10^4$ s ⁻¹ ; used $k_A = 5.5 \times 10^6$ L mol ⁻¹ s ⁻¹ ; 0.01 mol L ⁻¹ HCl.	91A252
	H ₂ O pH = 10	1.4×10^7 (k_r)			CP/Oc-17	S = RB; A' = Met; used $k_r^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A252
	H ₂ O pH = 11.5	1.9×10^7 (k_r)			CP/Oc-17	S = Eos; A' = Met; used $k_r^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A252
4.190 Tyrosine, methyl ester	D ₂ O/ EtOH (75:25)	5×10^6		295	PL/Ld-2	S = RB.	94A113

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds.

	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
1.1 Angelicin							
	CCl ₄	6.5×10^3			MP/LI-12	S = TPP; used $k_d = 39$ s ⁻¹ .	83E813
1.2 Benzo[1,2- <i>b</i> :4,3- <i>b'</i>]dipyrany, 1,2,3,8,9,10-hexahydro-3,3,5,6,8,8-hexamethyl-							
	EtOH	2.8×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	91F285
1.3 Benzo[1,2- <i>b</i> :4,5- <i>b'</i>]dipyrany, 2,3,4,7,8,9-hexahydro-2,2,5,7,7,10-hexamethyl-							
	EtOH	3.4×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	91F285
1.4 1-Benzopyran, 3,4-dihydro-6-methoxy-2,5,7,8-tetramethyl-2-(4,8,12-trimethyltridecyl)- (α -Tocopherol methyl ether)							
	CHCl ₃ / EtOH (50:50)	9.4×10^5 (k_r)		310	CR/Ac-31	used $k_d = 1 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from NDPO ₂ .	90A184
1.5 1-Benzopyran, 6-ethoxy-3,4-dihydro-2,5,7,8-tetramethyl-2-(4,8,12-trimethyltridecyl)- (α -Tocopherol ethyl ether)							
	CHCl ₃ / EtOH (50:50)	1.6×10^5 (k_r)		310	CR/Ac-31	used $k_d = 1 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from NDPO ₂ .	90A184
1.6 Coumarin, 7-(diethylamino)-							
	C ₆ H ₆	9×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 3.7 \times 10^4$ s ⁻¹ ; $k_{A'}$ not given.	87F569
1.7 Coumarin, 7-(diethylamino)-4-methyl-							
	C ₆ H ₆	9×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 3.7 \times 10^4$ s ⁻¹ ; $k_{A'}$ not given.	87F569
1.8 Cyclopenta[<i>b</i>][1]benzopyran, 2-phenyl-							
	C ₆ H ₆	7.8×10^5	0.054		CP/Ac-15	S = A; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	78F438
1.9 2-Furaldehyde							
	CH ₂ Cl ₂	5.1×10^4			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	EtOH			293	CP/Oc-17	S = RB; A' = FFA; meas. $k_r/k_r^{A'} = 0.22$.	86F510
	MeOH	8.4×10^4 (k_r)	1.7 (β_r)	286	CP/Oc-14	S = RB or TPP; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	85F145
	MeOH		0.080		CP/A'c-16	S = MB; A' = DPBF.	72F510
	MeOH		0.60	293	CP/Oc-15	S = RB; $E_a = 21$ kJ mol ⁻¹ .	68F288
1.10 2-Furaldehyde, 5-(2-hydroxyethoxy)methyl-							
	EtOH			293	CP/Oc-17	S = RB; A' = FFA; meas. $k_r/k_r^{A'} = 0.43$.	86F510
1.11 2-Furaldehyde, 5-(hydroxymethyl)-							
	EtOH			293	CP/Oc-17	S = RB; A' = FFA; meas. $k_r/k_r^{A'} = 0.38$.	86F510
1.12 2-Furaldehyde, 5-(methoxymethyl)-							
	EtOH			293	CP/Oc-17	S = RB; A' = FFA; meas. $k_r/k_r^{A'} = 0.35$.	86F510
1.13 2-Furaldehyde, 5-methyl-							
	CH ₂ Cl ₂	5.3×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	EtOH			293	CP/Oc-17	S = RB; A' = FFA; meas. $k_r/k_r^{A'} = 0.59$.	86F510
	MeOH	1.7×10^6 (k_r)	0.084 (β_r)	286	CP/Oc-14	S = RB or TPP; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	85F145
1.14 Furan							
	C ₆ H ₅ CH ₃	1.1×10^7			PR/A'd-5	S = Np; A' = DPBF; $\Delta H^\ddagger = 0$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -118$ J K ⁻¹ mol ⁻¹ ; studied at 223-353 K.	79A106
	CH ₂ Cl ₂	1.4×10^7			CP/A'c-16	S = MPDME; A' = DPBF; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F477
	CH ₂ Cl ₂	1.2×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
5.14 Furan — Continued							
	EtOH			293	CP/Oc-17	S = RB; A' = FFA; meas. $k_r/k_r^{A'} = 1$.	86F510
	MCH			298	PL/A'd-8	S = An; A' = DPBF; $\Delta V\ddagger = -17 \text{ cm}^3 \text{ mol}^{-1}$; studied at 0.1–400 MPa.	92A014
	MeOH	8.9×10^6 (k_r)	0.016 (β_r)	286	CP/Oc-14	S = RB or TPP; used $k_d = 1.4 \times 10^6 \text{ s}^{-1}$.	85F145
	MeOH		2.6×10^{-3}		CP/A'c-16	S = MB; A' = DPBF.	72F510
	MeOH		4.5×10^{-3}	293	CP/Oc-15	S = RB; $E_a = 0.84 \text{ kJ mol}^{-1}$.	68F288
5.15 Furan, 2-acetyl-							
	MeOH		0.022		CP/A'c-16	S = MB; A' = DPBF.	72F510
5.16 Furan, 2,5-bis(4-bromophenyl)-							
	CH ₂ Cl ₂	2.5×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 3.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	84F477
5.17 Furan, 2,5-bis(4-chlorophenyl)-							
	C ₆ H ₅ CH ₃	5.8×10^7			PR/A'd-5	S = Np; A' = DPBF; $\Delta H\ddagger = 0 \text{ kJ mol}^{-1}$, $\Delta S\ddagger = -105 \text{ J K}^{-1} \text{ mol}^{-1}$; studied at 223–353 K.	79A106
	CH ₂ Cl ₂	2.9×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 3.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	84F477
5.18 Furan, 2,5-bis(hydroxymethyl)- (BMHF)							
	EtOH			293	CP/Oc-17	S = RB; A' = FFA; meas. $k_r/k_r^{A'} = 1.2$.	86F510
	H ₂ O pH = 7		4.0×10^{-3}		CP/Ac-15	S = RB.	92N305
5.19 Furan, 2,5-bis(4-methoxyphenyl)-							
	CH ₂ Cl ₂	9.9×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 3.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	84F477
5.20 Furan, 2,5-bis(4-methylphenyl)-							
	CH ₂ Cl ₂	7.0×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 3.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	84F477
5.21 Furan, 2-bromo-							
	CH ₂ Cl ₂	6.7×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 3.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	84F477
5.22 Furan, 2-(4-bromophenyl)-							
	CH ₂ Cl ₂	1.0×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 3.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	84F477
5.23 Furan, 3-(4-bromophenyl)-							
	MeOH		4.4×10^{-3}		CP/A'c-16	S = MB; A' = DPBF.	72F510
5.24 Furan, 2-<i>tert</i>-butyl-							
	CH ₂ Cl ₂	6.1×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 3.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	84F477
	CH ₂ Cl ₂	6.4×10^7			CP/A'c-16	S = MPDME; A' = DPBF; used $k_d = 1 \times 10^4 \text{ s}^{-1}$.	84F477
5.25 Furan, 2-(4-chlorophenyl)-							
	1-BuOH		3.4×10^{-3} (β_r)	298	CP/Ac,A'c-17	S = RB; A' = DPF; used $\beta_r^{A'} = 9.5 \times 10^{-4} \text{ mol L}^{-1}$; meas. $\beta_r^{A'}/\beta_r^{A'}(\text{MeOH}) = 3.58$.	717398
	<i>tert</i> -BuOH		0.035 (β_r)	298	CP/Ac,A'c-17	S = RB; A' = DPF; used $\beta_r^{A'} = 9.5 \times 10^{-4} \text{ mol L}^{-1}$; meas. $\beta_r^{A'}/\beta_r^{A'}(\text{MeOH}) = 36.8$.	717398
	CH ₂ Cl ₂	1.0×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 3.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	84F477
	MeOH		2.9×10^{-3}		CP/A'c-16	S = MB; A' = DPBF.	72F510

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
§.35 Furan, 2-(4-chlorophenyl)- — Continued							
	MeOH	2.7×10^{-3} (β_r)		298	CP/Ac,A'c-17	$S = RB; A' = DPF$; used $\beta_r^{A'} = 9.5 \times 10^{-4}$ mol L ⁻¹ .	71F398
§.36 Furan, 2-(4-cyanophenyl)-							
	CH ₂ Cl ₂	4×10^6			CP/A'c-33	$S = A' = Rub$; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
§.37 Furan, 2,5-dibromo-							
	CH ₂ Cl ₂	2.0×10^6			CP/A'c-33	$S = A' = Rub$; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
§.38 Furan, 2,5-di-(1,1-dimethylethyl)-							
	C ₆ H ₅ CH ₃	2.3×10^8			PR/A'd-5	$S = Np; A' = DPBF; \Delta H^\ddagger = 0$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -94.1$ J K ⁻¹ mol ⁻¹ ; studied at 223-353 K.	79A106
	CH ₂ Cl ₂	1.5×10^8			CP/A'c-16	$S = MPDME; A' = DPBF$; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F477
	CH ₂ Cl ₂	9.4×10^7			CP/A'c-33	$S = A' = Rub$; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
§.39 Furan, 2,3-dihydro-							
	C ₆ H ₆	2.1×10^6 (k_r)	0.020 (β_r)	286	CP/Oc-29	$S = RB$ or TPP; $A' = 2,5\text{-DMF}$; used $k_d = 4.2 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 32% [2+2] cycloaddition.	91F163
	CCl ₄	1.6×10^6 (k_r)	9.2×10^{-4} (β_r)	286	CP/Oc-29	$S = TPP; A' = 2,5\text{-DMF}$; used $k_d = 1.4 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 24% [2+2] cycloaddition.	91F163
	CH ₂ Cl ₂	3.2×10^6 (k_r)	3.0×10^{-3} (β_r)	286	CP/Oc-29	$S = TPP; A' = 2,5\text{-DMF}$; used $k_d = 9.3 \times 10^5$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 74% [2+2] cycloaddition.	91F163
	CH ₃ CN	4.7×10^6 (k_r)	7.1×10^{-3} (β_r)	286	CP/Oc-29	$S = RB; A' = 2,5\text{-DMF}$; used $k_d = 3.3 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 70% [2+2] cycloaddition.	91F163
	CH ₃ COCH ₃	2.8×10^6 (k_r)	0.014 (β_r)	286	CP/Oc-29	$S = RB; A' = 2,5\text{-DMF}$; used $k_d = 3.8 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 71% [2+2] cycloaddition.	91F163
	CHCl ₃	1.0×10^6 (k_r)	3.9×10^{-3} (β_r)	286	CP/Oc-29	$S = TPP; A' = 2,5\text{-DMF}$; used $k_d = 4.0 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 73% [2+2] cycloaddition.	91F163
	MeOH	2.5×10^6 (k_r)	0.058 (β_r)	286	CP/Oc-29	$S = RB; A' = 2,5\text{-DMF}$; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$.	91F163
§.40 Furan, 2,3-dihydro-4,5-dimethyl-							
	MeOH	1.1×10^8 (k_r)		286	CP/Oc-29	$S = RB; A' = 2,5\text{-DMF}$; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 80% [2+2] cycloaddition.	91F163
§.41 Furan, 2,3-dihydro-5-methyl-							
	C ₆ H ₆	1.5×10^7 (k_r)	2.9×10^{-3} (β_r)	286	CP/Oc-29	$S = RB$ or TPP; $A' = 2,5\text{-DMF}$; used $k_d = 4.2 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 52% [2+2] cycloaddition.	91F163
	CH ₃ CN	3.4×10^7 (k_r)	9.7×10^{-4} (β_r)	286	CP/Oc-29	$S = RB; A' = 2,5\text{-DMF}$; used $k_d = 3.3 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 85% [2+2] cycloaddition.	91F163
	CH ₃ COCH ₃	2.8×10^7 (k_r)	1.4×10^{-3} (β_r)	286	CP/Oc-29	$S = RB; A' = 2,5\text{-DMF}$; used $k_d = 3.8 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 78% [2+2] cycloaddition.	91F163
	MeOH	1.4×10^7 (k_r)	0.010 (β_r)	286	CP/Oc-29	$S = RB; A' = 2,5\text{-DMF}$; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$.	91F163
§.42 Furan, 2-(dimethoxymethyl)-5-(methoxymethyl)-							
	EtOH			293	CP/Oc-17	$S = RB; A' = FFA$; meas. $k_r/k_r^{A'} = 1.0$.	86F510
§.43 Furan, 2,4-dimethyl-							
	MeOH	2.0×10^{-3}		293	CP/Oc-15	$S = RB; E_a = 6.3$ kJ mol ⁻¹ .	68F288

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
5.34 Furan, 2,5-dimethyl- (2,5-DMF)							
	C ₆ H ₅ CH ₃	6.3×10^8			PR/A'd-5	S = Np; A' = DPBF; $\Delta H^\ddagger = 0$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -84.5$ J K ⁻¹ mol ⁻¹ ; studied at 223-353 K.	79A106
	CH ₂ Cl ₂	2.6×10^8			CP/A'c-16	S = MPDME; A' = DPBF; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F477
	CH ₂ Cl ₂	1.3×10^8			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	CH ₃ COCH ₃	2.1×10^8	1.8×10^{-4}		CP/Oc-?	S = MB; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	75F652
	EtOH			293	CP/Oc-17	S = RB; A' = FFA; meas. $k_r/k_r^{A'} = 1.3$.	86F510
	EtOH	2.6×10^8			CP/Oc-20	S = BXP; Q = DABCO; used $k_d = 8.3 \times 10^4$ s ⁻¹ , $k_Q = 1.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; soln. contg. TEMPO.	83F323
	EtOH	3.8×10^8	2.2×10^{-4}	298	CP/Ac-15	S = MB; used $k_d = 8.3 \times 10^4$ s ⁻¹ .	78F061
	EtOH		1.5×10^{-4}		CP/Ac-15	S = MB.	78F201
	EtOH		1.6×10^{-4}		CP/Ac-15	S = Ret.	78F201
	EtOH		3.3×10^{-3}	273	CP/Oc-20	S = MB; A' = 1-Cyclohexylamino-4-phenylaminobenzene.	72F518
	H ₂ O pH = 7.0	$8.2 \times 10^8 (k_r)$		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	87A180
	H ₂ O	1.1×10^9			PL/P'a-16	S = Ery; A' = TMPD; used $k_d = 3.3 \times 10^5$ s ⁻¹ ; P' = [TMPD] ⁺ .	82A080
	H ₂ O		$3.3 \times 10^{-4} (\beta_r)$		CP/Ac-14	S = N-[5-Hydroxy-8-[[2-(methylsulfonyl)-4-nitrophenyl]jazo]-1-naphthalenyl-1,3-benzenedisulfonamide, conjugate base.	79F412
	H ₂ O		5.5×10^{-4}	298	CR/Oc-15	¹ O ₂ * from H ₂ O ₂ /NaOCl.	78A227
	H ₂ O	1.6×10^9	3.2×10^{-4}	298	CP/Ac-15	S = MB; used $k_d = 5.0 \times 10^5$ s ⁻¹ .	78F061
	H ₂ O	1.6×10^9	3.1×10^{-4}	298	CP/Ac-15	S = MB; used $k_d = 5.0 \times 10^5$ s ⁻¹ .	747044
	MeOH	$1.8 \times 10^8 (k_r)$		286	CP/Oc-14	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	91F163
	MeOH	3.8×10^8			PL/Tb-3	S = Ery.	83A050
	MeOH	9×10^7			PL/Tb-3	S = CH ₃ COCOCH ₃ .	80E606
	MeOH	3.9×10^8	2.8×10^{-4}	298	CP/Ac-15	S = MB; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	78F061
	MeOH	1.4×10^8	1.0×10^{-3}		CP/Oc-?	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	75F652
	MeOH	4×10^8			PL/A'd-5	S = MB; A' = DPBF.	727260
	MeOH		2.8×10^{-4}		CP/A'c-16	S = MB; A' = DPBF.	72F510
	MeOH		$\sim 2 \times 10^{-4}$	293	CP/Oc-15	S = RB; $E_a = 4.2$ kJ mol ⁻¹ .	68F288
	MeOH/ <i>tert</i> -BuOH (50:50)		1.6×10^{-4}		CP/A'c-16	S = RB; A' = DPBF.	717398
	MeOH/ <i>tert</i> -BuOH (50:50)				CP/Ac,A'c-17	S = RB; A' = TME; meas. $k_r/k_r^{A'} = 2.4$.	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)				CR/Ac,A'c-17	A' = TME; meas. $k_r/k_r^{A'} = 1.5$; ¹ O ₂ * from H ₂ O ₂ /Ca(OCl) ₂ .	68F292
	MeOH/ <i>tert</i> -BuOH (50:50)				CR/Ac,A'c-17	A' = TME; meas. $k_r/k_r^{A'} = 5.2$; ¹ O ₂ * from H ₂ O ₂ /NaOCl.	68F292
5.35 Furan, 2-[(1,1-dimethylethoxy)methyl]-							
	EtOH			293	CP/Oc-17	S = RB; A' = FFA; meas. $k_r/k_r^{A'} = 1.0$.	86F510
5.36 Furan, 2-[2-(1,3-dioxolanyl)-							
	EtOH			293	CP/Oc-17	S = RB; A' = FFA; meas. $k_r/k_r^{A'} = 0.94$.	86F510
5.37 Furan, 2,3-diphenyl-							
	C ₆ H ₅ CH ₃	6.2×10^7			PR/A'd-5	S = Np; A' = DPBF; $\Delta H^\ddagger = 0$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -104$ J K ⁻¹ mol ⁻¹ ; studied at 223-353 K.	79A106

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
Furan, 2,5-diphenyl-							
	1-BuOH	5.0×10^7	1.0×10^{-3}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 5.2 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
	1-BuOH		$9.8 \times 10^{-4} (\beta_r)$	298	CP/Ac-17	S = RB; meas. $\beta_r/\beta_r(\text{MeOH}) = 1.03$, used $\beta_r(\text{MeOH}) = 9.5 \times 10^{-4}$ mol L ⁻¹ .	717398
	tert-BuOH	3.6×10^7	8.3×10^{-4}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 3.0 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
	tert-BuOH		$8.0 \times 10^{-4} (\beta_r)$	298	CP/Ac-17	S = RB; meas. $\beta_r/\beta_r(\text{MeOH}) = 0.842$, used $\beta_r(\text{MeOH}) = 9.5 \times 10^{-4}$ mol L ⁻¹ .	717398
	C ₅ H ₅ N	6.1×10^7	9.9×10^{-4}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 6.0 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
	c-C ₆ H ₁₁ OH	5.3×10^7	1.2×10^{-3}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 6.2 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
	C ₆ H ₅ CH ₃	1.0×10^8			PR/A'd-5	S = Np; A' = DPBF; $\Delta H^\ddagger = 0$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -100$ J K ⁻¹ mol ⁻¹ ; studied at 223-353 K.	79A106
	C ₆ H ₅ CO ₂ CH ₃	4.1×10^7	6.1×10^{-4}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 2.5 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
	CH ₂ Cl ₂	7.0×10^7			CP/A'c-16	S = MPDME; A' = DPBF; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F477
	CH ₂ Cl ₂	5.2×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	CH ₂ Cl ₂	5.1×10^7	1.4×10^{-4}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 7.1 \times 10^3$ s ⁻¹ .	74F646
	CH ₃ CN	3.7×10^8	8.3×10^{-5}	293	CP/Pa-18	S = MB; A' = DPBF; used $k_d = 3 \times 10^4$ s ⁻¹ , $\beta_{A'} = 1.3 \times 10^{-5}$ mol L ⁻¹ ; P = 2,5-Diphenylfuran endoperoxide.	87F440
	CH ₃ CN	1.4×10^8	1.3×10^{-4}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 1.8 \times 10^4$ s ⁻¹ .	74F646
	CH ₃ CO ₂ C ₂ H ₅	3.0×10^7	6.9×10^{-4}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 2.1 \times 10^4$ s ⁻¹ .	74F646
	CH ₃ COCH ₃	1.4×10^8	1.7×10^{-4}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 2.4 \times 10^4$ s ⁻¹ .	74F646
	CHCl ₃	2.2×10^7	2.0×10^{-4}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 4.4 \times 10^3$ s ⁻¹ .	74F646
	D ₂ O		$7.7 \times 10^{-5} (\beta_r)$	296	CP/Ac-17	S = MB; meas. $\beta_r/\beta_r(\text{H}_2\text{O}) = 0.115$, used $\beta_r(\text{H}_2\text{O}) = 4.6 \times 10^{-4}$ mol L ⁻¹ .	74R214
	D ₂ O pD = 7.1		$1.2 \times 10^{-4} (\beta_r)$	296	CP/Ac-17	S = MB; meas. $\beta_r/\beta_r(\text{H}_2\text{O}) = 0.182$, used $\beta_r(\text{H}_2\text{O}) = 4.6 \times 10^{-4}$ mol L ⁻¹ .	74R214
	D ₂ O pD = 7.1		$1.4 \times 10^{-4} (\beta_r)$	296	CP/Ac-17	S = MB; meas. $\beta_r/\beta_r(\text{H}_2\text{O}) = 0.206$, used $\beta_r(\text{H}_2\text{O}) = 4.6 \times 10^{-4}$ mol L ⁻¹ . Solvent contains 960 μM H ₂ O ₂ .	74R214
	D ₂ O pD = 7.1		$1.1 \times 10^{-4} (\beta_r)$	296	CP/Ac-17	S = MB; meas. $\beta_r/\beta_r(\text{H}_2\text{O}) = 0.165$, used $\beta_r(\text{H}_2\text{O}) = 4.6 \times 10^{-4}$ mol L ⁻¹ . 1 μl/ml of catalase present in reaction medium.	74R214
	H ₂ O		$4.6 \times 10^{-4} (\beta_r)$		CP/Ac-17	S = RB; meas. $\beta_r/\beta_r(\text{MeOH}) = 0.481$, used $\beta_r(\text{MeOH}) = 9.5 \times 10^{-4}$ mol L ⁻¹ .	717398
	HOCH ₂ CH ₂ OH		$3.5 \times 10^{-4} (\beta_r)$		CP/Ac-17	S = RB; meas. $\beta_r/\beta_r(\text{MeOH}) = 0.366$, used $\beta_r(\text{MeOH}) = 9.5 \times 10^{-4}$ mol L ⁻¹ .	717398
	MeOH	1.0×10^8	8.8×10^{-4}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	74F646
	MeOH	4.6×10^7			PL/A'd-8	S = RB; A' = DPBF.	737014
	MeOH	$\sim 5 \times 10^5 (k_d)$			CP/Ac-14	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; used $\phi_{isc}(\text{RB}) = 0.76$.	73F660
	MeOH		9.5×10^{-4}	298	CP/A'c-16	S = RB; A' = DPBF.	717398
	MeOH/H ₂ O (12:88)		$4.2 \times 10^{-4} (\beta_r)$		CP/Ac-17	S = RB; meas. $\beta_r/\beta_r(\text{MeOH}) = 0.439$, used $\beta_r(\text{MeOH}) = 9.5 \times 10^{-4}$ mol L ⁻¹ ; solvent mixt. in mole %.	717398
	MeOH/H ₂ O (40:60)		$4.2 \times 10^{-4} (\beta_r)$		CP/Ac-17	S = RB; meas. $\beta_r/\beta_r(\text{MeOH}) = 0.446$, used $\beta_r(\text{MeOH}) = 9.5 \times 10^{-4}$ mol L ⁻¹ ; solvent mixt. in mole %.	717398

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
5.38 Furan, 2,5-diphenyl- — Continued							
	MeOH/ H ₂ O (66:34)	5.9×10^{-4} (β_r)			CP/Ac-17	S = RB; meas. $\beta_r/\beta_r(\text{MeOH}) = 0.621$, used $\beta_r(\text{MeOH}) = 9.5 \times 10^{-4}$ mol L ⁻¹ ; solvent mixt. in mole %.	717398
	MeOH/ H ₂ O (81:19)	7.7×10^{-4} (β_r)			CP/Ac-17	S = RB; meas. $\beta_r/\beta_r(\text{MeOH}) = 0.806$, used $\beta_r(\text{MeOH}) = 9.5 \times 10^{-4}$ mol L ⁻¹ ; solvent mixture in mole %.	717398
	MeOH/ HOCH ₂ CH ₂ OH (75:25)	5.5×10^{-4} (β_r)			CP/Ac-17	S = RB; meas. $\beta_r/\beta_r(\text{MeOH}) = 0.578$, used $\beta_r(\text{MeOH}) = 9.5 \times 10^{-4}$ mol L ⁻¹ .	717398
	MeOH/ HOCH ₂ CH ₂ OH (50:50)	3.4×10^{-4} (β_r)			CP/Ac-17	S = RB; meas. $\beta_r/\beta_r(\text{MeOH}) = 0.355$, used $\beta_r(\text{MeOH}) = 9.5 \times 10^{-4}$ mol L ⁻¹ .	717398
	MeOH/ HOCH ₂ CH ₂ OH (25:75)	2.6×10^{-4} (β_r)			CP/Ac-17	S = RB; meas. $\beta_r/\beta_r(\text{MeOH}) = 0.278$, used $\beta_r(\text{MeOH}) = 9.5 \times 10^{-4}$ mol L ⁻¹ .	717398
	2-PrOH	4.8×10^7	1.0×10^{-3}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 5.0 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
	THF	7.5×10^7	5.9×10^{-4}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 4.3 \times 10^4$ s ⁻¹ .	74F646
	dioxane	2.7×10^7	1.1×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 2.9 \times 10^4$ s ⁻¹ .	74F646
5.39 Furan, 3,4-diphenyl-							
	MeOH		1.7×10^{-3}		CP/A'c-16	S = RB or MB; A' = DPBF.	72F510
5.40 Furan, 2-[(4-ethoxycarbonyl)phenyl]							
	CH ₂ Cl ₂	7×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
5.41 Furan, 3-(4-fluorophenyl)-							
	MeOH		3.4×10^{-3}		CP/A'c-16	S = MB; A' = DPBF.	72F510
5.42 Furan, 2-methoxy-							
	CH ₂ Cl ₂	1.6×10^8			CP/A'c-16	S = MPDME; A' = DPBF; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F477
	CH ₂ Cl ₂	1.2×10^8			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	MeOH	1.6×10^8 (β_r)	9.2×10^{-4} (β_r)	286	CP/Oc-14	S = RB or TPP; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	85F145
	MeOH		4.0×10^{-4}		CP/A'c-16	S = MB; A' = DPBF.	72F510
5.43 Furan, 2-(methoxymethyl)-							
	MeOH		3.4×10^{-3}	293	CP/Oc-15	S = RB; $E_a = 5.0$ kJ mol ⁻¹ .	68F288
5.44 Furan, 2-(4-methoxyphenyl)-							
	1-BuOH		1.9×10^{-3} (β_r)		CP/Ac,A'c-17	S = RB or MB; A' = DPF; used $\beta_r^{A'} = 9.5 \times 10^{-4}$ mol L ⁻¹ ; meas. $\beta_r^{A'}/\beta_r^{A'}(\text{MeOH}) = 2.0$.	72F510
	CH ₂ Cl ₂	3.7×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	H ₂ O		5.9×10^{-4} (β_r)		CP/Ac,A'c-17	S = MB or RB; A' = DPF; used $\beta_r^{A'} = 9.5 \times 10^{-4}$ mol L ⁻¹ ; meas. $\beta_r^{A'}/\beta_r^{A'}(\text{MeOH}) = 0.618$.	72F510
	HOCH ₂ CH ₂ OH		3.4×10^{-4} (β_r)		CP/Ac,A'c-17	S = MB or RB; A' = DPF; used $\beta_r^{A'} = 9.5 \times 10^{-4}$ mol L ⁻¹ ; meas. $\beta_r^{A'}/\beta_r^{A'}(\text{MeOH}) = 0.36$.	72F510
	MeOH				CP/Ac,A'c-17	S = MB or RB; A' = DPF; meas. $k_r/k_r^{A'} = 0.92$.	72F510
5.45 Furan, 3-(4-methoxyphenyl)-							
	MeOH		1.9×10^{-3}		CP/A'c-16	S = MB; A' = DPBF.	72F510
5.46 Furan, 2-methyl-							
	1-BuOH		1.2×10^{-3}	296	CP/A'c-16	S = MB; A' = DPBF.	717398

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
§.46 Furan, 2-methyl- — Continued							
	tert-BuOH		6.9×10^{-4}	296	CP/A'c-16	S = MB; A' = DPBF.	717398
	C ₆ H ₅ Cl/ 2-PrOH (90:10)	1.3×10^8		273	CR/LI-12	used $k_d = 4.1 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from hydrotrioxide, e.g. (CH ₃) ₂ C(OH)OOOH.	90M125
	CH ₂ Cl ₂	6.2×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	CH ₂ Cl ₂	8.4×10^7			CP/A'c-16	S = MPDME; A' = DPBF; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F477
	EtOH			293	CP/Oc-17	S = RB; A' = FFA; meas. $k_r/k_t^{A'} = 1.2$.	86F510
	H ₂ O pH = 7.0	6.0×10^7			CP/Oc-19	S = HP; Q = N ₃ ⁻ ; used $k_Q = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A278
	H ₂ O (mic) pH = 7.0	1.0×10^8			CP/Oc-19	S = Chl a; Q = N ₃ ⁻ ; used $k_Q = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; 1, 2, or 5% Triton X-100.	78A278
	H ₂ O (mic) pH = 7.0	7.0×10^7			CP/Oc-19	S = HP; Q = N ₃ ⁻ ; used $k_Q = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; 1% Triton X-100.	78A278
	MeOH	9.9×10^7 (k_r)	1.4×10^{-3} (β_r)	286	CP/Oc-14	S = RB or TPP; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	85F145
	MeOH		1.1×10^{-3}	296	CP/A'c-16	S = MB; A' = DPBF.	72F510 717398
	MeOH		3.8×10^{-3}	293	CP/Oc-15	S = RB; $E_a = 1.7$ kJ mol ⁻¹ .	68F288
§.47 Furan, 2-(4-methylphenyl)-							
	CH ₂ Cl ₂	2.4×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	MeOH				CP/Ac,A'c-17	S = MB or RB; A' = DPF; meas. $k_r/k_t^{A'} = 0.68$.	72F510
§.48 Furan, 3-(4-methylphenyl)-							
	MeOH		2.4×10^{-3}		CP/A'c-16	S = RB or MB; A' = DPBF.	72F510
§.49 Furan, 2-(4-nitrophenyl)-							
	CH ₂ Cl ₂	2.4×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
§.50 Furan, 2-phenyl-							
	1-BuOH		2.0×10^{-3} (β_r)	298	CP/Ac,A'c-17	S = RB; A' = DPF; used $\beta_r^{A'} = 9.5 \times 10^{-4}$ mol L ⁻¹ ; meas. $\beta_r^{A'}/\beta_r^{A'}(\text{MeOH}) = 2.1$.	717398
	tert-BuOH		3.4×10^{-3} (β_r)	298	CP/Ac,A'c-17	S = RB; A' = DPF; used $\beta_r^{A'} = 9.5 \times 10^{-4}$ mol L ⁻¹ ; meas. $\beta_r^{A'}/\beta_r^{A'}(\text{MeOH}) = 3.58$.	717398
	CH ₂ Cl ₂	1.8×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
	MeOH		1.7×10^{-3}	298	CP/Ac,A'c-17	S = MB or RB; A' = DPF; used $\beta_{A'} = 9.5 \times 10^{-4}$ mol L ⁻¹ .	72F510 717398
§.51 Furan, 3-phenyl-							
	MeOH		2.8×10^{-3}		CP/A'c-16	S = MB; A' = DPBF.	72F510
§.52 Furan, tetraphenyl-							
	C ₆ H ₅ CH ₃	2.8×10^8			PR/A'd-5	S = Np; A' = DPBF; $\Delta H^\ddagger = 0$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -91.2$ J K ⁻¹ mol ⁻¹ ; studied at 223-353 K.	79A106
	C ₆ H ₅ Cl/ 2-PrOH (90:10)	3.6×10^8		273	CR/LI-12	used $k_d = 4.1 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from hydrotrioxide, e.g. (CH ₃) ₂ C(OH)OOOH.	90M125
	MeOH		-0.030	293	CP/Oc-15	S = RB; $E_a = 4.2$ kJ mol ⁻¹ .	68F288
§.53 Furan, 2-vinyl-							
	MeOH		1.8×10^{-3}	293	CP/Oc-15	S = RB; $E_a = 3.8$ kJ mol ⁻¹ .	68F288
§.54 2-Furancarboxylic acid							
	MeOH		0.012		CP/A'c-16	S = MB; A' = DPBF.	72F510

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
5.55	2-Furancarboxylic acid, 5-bromo-, ethyl ester						
	CH ₂ Cl ₂	1.5×10^4			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
5.56	2-Furancarboxylic acid, ethyl ester						
	CH ₂ Cl ₂	1.3×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
5.57	2-Furancarboxylic acid, methyl ester						
	CH ₂ Cl ₂	8.5×10^4			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
5.58	3-Furancarboxylic acid, 4,5-dihydro-2-methyl-, methyl ester						
	C ₆ H ₆	2.0×10^5 (k_r)	0.21 (β_r)	286	CP/Oc-29	S = RB or TPP; A' = 2,5-DMF; used $k_d = 4.2 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 4% [2+2] cycloaddition.	91F163
	CCl ₄	2.0×10^5 (k_r)	7.2×10^{-3} (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; used $k_d = 1.4 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 3% [2+2] cycloaddition.	91F163
	CH ₂ Cl ₂	2.4×10^5 (k_r)	0.041 (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; used $k_d = 9.5 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 32% [2+2] cycloaddition.	91F163
	CH ₃ CN	8.8×10^5 (k_r)	0.038 (β_r)	286	CP/Oc-29	S = RB; A' = 2,5-DMF; used $k_d = 3.3 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 44% [2+2] cycloaddition.	91F163
	CH ₃ COCH ₃	6.1×10^5 (k_r)	0.063 (β_r)	286	CP/Oc-29	S = RB; A' = 2,5-DMF; used $k_d = 3.8 \times 10^4$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 39% [2+2] cycloaddition.	91F163
	CHCl ₃	7.0×10^4 (k_r)	0.057 (β_r)	286	CP/Oc-29	S = TPP; A' = 2,5-DMF; used $k_d = 4.0 \times 10^3$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 36% [2+2] cycloaddition.	91F163
	MeOH	7.3×10^5 (k_r)	0.19 (β_r)	286	CP/Oc-29	S = RB; A' = 2,5-DMF; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$; 29% [2+2] cycloaddition.	91F163
5.59	2,5-Furandicarboxaldehyde						
	CH ₂ Cl ₂	2.4×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
5.60	2,5-Furandicarboxylic acid, diethyl ester						
	CH ₂ Cl ₂	4.5×10^4			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
5.61	2,5-Furandicarboxylic acid, dimethyl ester						
	CH ₂ Cl ₂	2.0×10^4			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
5.62	3,4-Furandicarboxylic acid, diethyl ester						
	MeOH	~0.20		293	CP/Oc-15	S = RB; $E_a = 12.6$ kJ mol ⁻¹ .	68F288
5.63	2-Furanitrile						
	CH ₂ Cl ₂	1.1×10^4			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
5.64	2-Furanitrile, 5-methyl-						
	CH ₂ Cl ₂	1.5×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	84F477
5.65	2-Furanmethanamine, N-methyl-						
	MeOH	0.013		293	CP/Oc-15	S = RB; $E_a = 4.2$ kJ mol ⁻¹ .	68F288
5.66	2-Furanmethanediol, diacetate						
	C ₆ H ₅ Cl/ 2-PrOH (90:10)	2.3×10^6		273	CR/LI-12	used $k_d = 4.1 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from hydrotrioxide, e.g. $(\text{CH}_3)_2\text{C}(\text{OH})\text{OOH}$.	90M125

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
5.67	2-Furanmethanol, α,α -diphenyl-						
	MeOH	9.9×10^6	9.2×10^{-3}	298	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
5.68	2-Furanmethanol, α -(diphenylmethyl)-						
	MeOH	1.1×10^7	8.0×10^{-3}	298	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
5.69	2-Furanmethanol, α -methyl-						
	MeOH	1.0×10^8	8.7×10^{-4}	298	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
5.70	2-Furanmethanol, α -phenyl-						
	MeOH	5.2×10^6	0.018	298	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
5.71	2-Furanmethanol, α -(2-phenylethyl)-						
	MeOH	2.5×10^7	3.6×10^{-3}	298	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
5.72	2-Furanmethanol, α -(phenylmethyl)-						
	MeOH	1.1×10^7	8.3×10^{-3}	298	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
5.73	2-Furanmethanol, α -(3-phenylpropyl)-						
	MeOH	8.9×10^8	1.0×10^{-4}	298	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
5.74	3-Furanmethanol, 5-(phenylmethyl)-						
	MeOH	3.0×10^7		298	CP/Oc-15	S = RB; $\Delta S^\ddagger = -102.7$ J K ⁻¹ mol ⁻¹ ; $E_a = -0$ kJ mol ⁻¹ ; studied at 293-308 K; k_d not given.	92F465
	MeOH/ H ₂ O (50:50)	1.6×10^8		298	CP/Oc-15	S = RB; $\Delta S^\ddagger = -87.2$ J K ⁻¹ mol ⁻¹ ; $E_a = -0$ kJ mol ⁻¹ ; studied at 293-308 K; k_d not given.	92F465
5.75	3-Furanmethanol, 5-(phenylmethyl)-, acetate						
	MeOH	6.4×10^7		298	CP/Oc-15	S = RB; $\Delta S^\ddagger = -96.5$ J K ⁻¹ mol ⁻¹ ; $E_a = -0$ kJ mol ⁻¹ ; studied at 293-308 K; k_d not given.	92F465
	MeOH/ H ₂ O (50:50)	8.2×10^8		298	CP/Oc-15	S = RB; $\Delta S^\ddagger = -74.8$ J K ⁻¹ mol ⁻¹ ; $E_a = -0$ kJ mol ⁻¹ ; studied at 293-308 K; k_d not given.	92F465
5.76	3-Furanmethanol, 5-(phenylmethyl)-, ester with 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylic acid (Bioresmethrin)						
	MeOH	5.3×10^7		298	CP/Oc-15	S = RB; $\Delta S^\ddagger = -97.9$ J K ⁻¹ mol ⁻¹ ; $E_a = -0$ kJ mol ⁻¹ ; studied at 293-308 K; k_d not given.	92F465
5.77	[2,2](2,5)Furanophane						
	MeOH	1.6×10^8	5.5×10^{-4}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
				298			
5.78	Furfuryl alcohol (FFA)						
	CD ₃ OD	3.0×10^7 (k_r) $<10^4$ (k_q)		295	CP/Oc-15,27	S = RB; used $k_d = 4.0 \times 10^3$ s ⁻¹ .	91F332
	H ₂ O	1.4×10^8 (k_r)		298	CP/Oc-14	S = H ₂ TPPS ⁴⁺ ; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; $E_a = 22.7$ kJ mol ⁻¹ ; log(A) = 12.1; studied at 288-318 K; phosphate buffer contg. 1.6% NaCl.	91R053
	H ₂ O pH = 10.0, 11.5	1.2×10^8 (k_r)		292	MP/Ac-17	S = RB; rel. to $k(FFA) = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ in H ₂ O at pH = 7.0.	87A180
	H ₂ O		3.0×10^{-3}	295	CP/Oc-14	S = HPD; 0.9% NaCl.	85F332
	H ₂ O	1.2×10^8	2.3×10^{-3}	295	CP/Oc-15	S = RB; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	84F381
	H ₂ O pH = 3.0, 8.0		1.8×10^{-3}	310	CP/Oc-14	S = PF.	617008
	MeOH		4.7×10^{-3}		CP/A'c-16	S = MB; A' = DPBF.	72F510
	MeOH		3.3×10^{-3}	293	CP/Oc-15	S = RB; $E_a = 2.9$ kJ mol ⁻¹ .	68F288
5.79	Furfurylamine						
	MeOH	0.030			CP/A'c-16	S = MB; A' = DPBF.	72F510

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
5.79	Furfurylamine — Continued						
	MeOH	9.0×10^{-3}		293	CP/Oc-15	S = RB; $E_a = 2.9$ kJ mol ⁻¹ .	68F288
5.80	Isobenzofuran						
	MeOH	9.2×10^7	9.8×10^{-4}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F517
5.81	Isobenzofuran, 1,3-bis[4-(11-carboxynonyl)phenyl]-4,7-dihydro-5,6-dimethyl-, dianion						
	D ₂ O/ H ₂ O (95:5) pH = 10	3×10^8 (k_r)			CP/Ac,A'c-17	S = HP; A' = His; used $k_r^{A'} = 5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82F379
	H ₂ O pH = 10	7.1×10^7 (k_r)			CP/Ac,A'c-17	S = HP; A' = His; used $k_r^{A'} = 5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82F379
5.82	Isobenzofuran, 5,6-dimethyl-1,3-diphenyl-						
	MeOH	5.2×10^8 (k_r)			CP/Ac-14	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; used ϕ_{isc} (RB) = 0.76.	73F660
5.83	Isobenzofuran, 1,3-diphenyl- (DPBF)						
	C ₆ H ₅ COCH ₃	9.0×10^8		293	PL/Ad-8	S = MB.	83A006 83A371
	1-BuOH	8.3×10^8 (k_r)			CP/Ac,A'c-17	S = MB; A' = DPF; used $k_r^{A'} = 5.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 17$; Solvent contained 1% MeOH.	74F646
	1-BuOH	8×10^8	7×10^{-5}		CP/Ac-15	S = MB; used $k_d = 5.2 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
	1-BuOH	8×10^8			PL/Ad-8	S = MB.	73F014
	1-BuOH		7.1×10^{-5} (β_r)	298	CP/Ac,A'c-17	S = RB; A' = DPF; used $\beta_r^{A'} = 9.5 \times 10^{-4}$ mol L ⁻¹ ; meas. $\beta_r/\beta_r^{A'}(\text{MeOH}) = 0.075$.	71F398
	tert-BuOH	6.2×10^8 (k_r)			CP/Ac,A'c-17	S = MB; A' = DPF; used $k_r^{A'} = 3.6 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 17$; Solvent contained 1% MeOH.	74F646
	tert-BuOH	2.3×10^8	1×10^{-4}		CP/Ac-15	S = MB; used $k_d = 3.0 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
	tert-BuOH	5.7×10^8			PL/Ad-8	S = MB; Solvent contained 1% MeOH.	73F014
	tert-BuOH		4.9×10^{-5} (β_r)	298	CP/Ac,A'c-17	S = RB; A' = DPF; used $\beta_r^{A'} = 9.5 \times 10^{-4}$ mol L ⁻¹ ; meas. $\beta_r/\beta_r^{A'}(\text{MeOH}) = 0.052$.	71F398
	Epibromohydrin	7.8×10^8			PL/Ad-8	S = MB.	74F646
	Propylene carbonate	1×10^9		293	PL/Ad-5	S = MB.	87E690
	C ₅ H ₅ N	6.5×10^8 (k_r)	9.1×10^{-5} (β_r)	298	CP/Ac-14	S = A; used $k_d = 5.9 \times 10^4$ s ⁻¹ .	84F240
	C ₅ H ₅ N	4.8×10^8 (k_r)			CP/Ac,A'c-17	S = RB; A' = DPF; used $k_r^{A'} = 6.1 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 7.9$; Solvent contained 1% MeOH.	74F646
	C ₅ H ₅ N	5×10^8			PL/Ad-8	S = RB; Solvent contained 1% MeOH.	74F646
	C ₅ H ₅ N	2.1×10^9			PL/Ad-8	S = MB.	73F014
	C ₅ H ₅ N				CP/Ac,A'c-17	S = A' = Rub; meas. $k_r/k_r^{A'} = 22$.	66F041
	c-C ₆ H ₁₁ OH	6.5×10^8 (k_r)			CP/Ac,A'c-17	S = MB; A' = DPF; used $k_r^{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 12$; Solvent contained 1% MeOH.	74F646
	c-C ₆ H ₁₁ OH	1×10^8			PL/Ad-8	S = MB; Solvent contained 1% MeOH.	74F646
	c-C ₆ H ₁₁ OH	2×10^8	3×10^{-4}		CP/Ac-15	S = MB; used $k_d = 6.3 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
	c-C ₆ H ₁₂	3.4×10^8			PL/Ad-5	S = 2-ACN; used $k_d = 5.0 \times 10^4$ s ⁻¹ .	84F005
	c-C ₆ H ₁₂	5.0×10^8 (k_r)	1.2×10^{-4} (β_r)	298	CP/Ac-14	S = A; used $k_d = 5.9 \times 10^4$ s ⁻¹ .	84F240
	c-C ₆ H ₁₂	5.4×10^8 (k_r)	1.1×10^{-4} (β_r)	298	CP/Ac-14	S = A; used $k_d = 5.9 \times 10^4$ s ⁻¹ .	84P629

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
Isobenzofuran, 1,3-diphenyl- (DPBF) — Continued							
	c-C ₆ H ₁₂	4.5×10^8			PL/Ad-8	S = An.	82A349
	c-C ₆ H ₁₂	5.3×10^8		303	CP/Ac-20	S = Py; used $k_d = 5.9 \times 10^5$ s ⁻¹ .	79N020
	c-C ₆ H ₁₂ (mic)	5.3×10^8			PL/Tb-3	S = Ery; reverse micelles contg. 0.04 mol L ⁻¹ DAP and 0.1 mol L ⁻¹ water.	83A050
	c-C ₆ H ₁₂ (mic)	5.3×10^8			CP/Ac-14	S = Fl ²⁻ ; used $k_d = 5.9 \times 10^4$ s ⁻¹ ; DAP reverse micelles.	80N021
	n-C ₆ H ₁₄	2.6×10^8		298	PL/Ad-8	S = An; $\Delta V^\ddagger = -14$ cm ³ mol ⁻¹ ; studied at 0.1-400 MPa.	92A014 90A018
	n-C ₆ H ₁₄	4.4×10^8 (k_r)	1.4×10^{-4} (β_r)	298	CP/Ac-14	S = A; used $k_d = 6.3 \times 10^4$ s ⁻¹ .	84F240
	n-C ₆ H ₁₄	1.2×10^9			CP/Ac-14	S = An; used $k_d = 3 \times 10^4$ s ⁻¹ .	80E761
	n-C ₆ H ₁₄		1.5×10^{-5}		CP/Pa-15	S = I ₂ .	72F521
	C ₆ H ₅ Br	6×10^8	2×10^{-5}		CP/Ac-15	S = MB; used $k_d = 1.3 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
	C ₆ H ₅ Br	5×10^8			PL/Ad-8	S = MB; Solvent contained 1% MeOH.	74F646
	C ₆ H ₅ Br/ MeOH (80:20)	7.0×10^8			PL/Ad-8	S = MB.	73T014
	C ₆ H ₅ Br/ MeOH (80:20)	5×10^8	9×10^{-5}		CP/Ac-15	S = MB; used $k_d = 4.3 \times 10^4$ s ⁻¹ .	74F646
	C ₆ H ₅ CH ₂ CH ₃	6.7×10^8		293	PL/Ad-8	S = MB.	83A006 83A371
	C ₆ H ₅ CH ₃	8.9×10^8			PL/Ld-2	S = PPDME; $\Delta H^\ddagger = -5$ (4) kJ mol ⁻¹ ; $\Delta S^\ddagger = -109$ (-70) J K ⁻¹ mol ⁻¹ ; $E_a = -3$ (7) kJ mol ⁻¹ ; studied at 183-363 K; pre-exciplex-equilibrium limit (and diffusion limit).	88A427
	C ₆ H ₅ CH ₃				CP/Ac-15	S = Tetr.	84F197
	C ₆ H ₅ CH ₃	4.3×10^8 (k_r)	1.2×10^{-4} (β_r)	298	CP/Ac-14	S = A; used $k_d = 5.0 \times 10^4$ s ⁻¹ .	84F240
	C ₆ H ₅ CH ₃	6.7×10^8			PL/Ad-8	S = An.	82A349
	C ₆ H ₅ CH ₃	8.1×10^8			PR/Ad-5	S = Np; $\Delta H^\ddagger = 0$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -82.4$ J K ⁻¹ mol ⁻¹ ; studied at 223-353 K.	79A106
	C ₆ H ₅ CHOHCH ₃	1.0×10^9		293	PL/Ad-8	S = MB.	83A006 83A371
	C ₆ H ₅ CO ₂ CH ₃	2.8×10^8	8.9×10^{-5}		CP/Ac-15	S = MB; used $k_d = 2.5 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
	C ₆ H ₅ CO ₂ CH ₃	4×10^8			PL/Ad-8	S = MB; Solvent contained 1% MeOH.	74F646
	C ₆ H ₅ Cl	7.0×10^8		293	PL/Ad-8	S = MB.	83A006 83A371
	C ₆ H ₅ Cl/ 2-PrOH (90:10)	7×10^8		273	CR/LI-12	used $k_d = 4.1 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from hydrotrioxide, e.g. (CH ₃) ₂ C(OH)OOOH.	90M125
	C ₆ D ₆	2.8×10^7			PL/Ad-5	S = 2-ACN.	81A287
	C ₆ H ₆	8.0×10^8			PL/Ad-5	S = 2-ACN; used $k_d = 4.0 \times 10^4$ s ⁻¹ .	84F005
	C ₆ H ₆		6.0×10^{-5}		CP/Ac-15	S = HCD.	84F197
	C ₆ H ₆	7.0×10^8 (k_r)	6.0×10^{-5} (β_r)	298	CP/Ac-14	S = A; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	84F240
	C ₆ H ₆	6.7×10^8			PL/Ad-5	S = 2-ACN.	81A287
	C ₆ H ₆		4.0×10^{-5}		CP/Ac-15	S = Rub.	78E036
	C ₆ H ₆	9.4×10^8			PR/Ad-5	S = Np, An, BP, Benzil.	78E263
	C ₆ H ₆	7.0×10^8 (k_r)	6.0×10^{-5}	298	CP/Ac,A'c-17	S = A' = Rub; used $k_r^{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 17$.	747312
	C ₆ H ₆	5×10^8	8×10^{-5}		CP/Ac-15	S = MB; used $k_d = 3.8 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
	C ₆ H ₆	4×10^8			PL/Ad-8	S = MB; Solvent contained 1% MeOH.	74F646

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
5.83 Isobenzofuran, 1,3-diphenyl- (DPBF) — Continued							
	C ₆ H ₆	6.0 × 10 ⁻⁵		298	CP/Ac-15	S = A.	74F649
	C ₆ H ₆	6.0 × 10 ⁻⁵ (β_r)		298	CP/Ac,A'c-17	S = A' = Rub; meas. $k_r/k_r^{A'} = 17$; used $\beta_r^{A'} = 1.0 \times 10^{-3}$ mol L ⁻¹ .	74F649
	C ₆ H ₆	2 × 10 ⁹		298	PL/Ad-8	S = An.	737438
	C ₆ H ₆ / MeOH (80:20)	2 × 10 ⁹	2 × 10 ⁻⁵		CP/Ac-15	S = MB; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	74F646
	C ₆ H ₆ / MeOH (80:20)	1.0 × 10 ⁹ (k_r)			CP/Ac,A'c-17	S = MB; A' = DPF; $k_r^{A'}$ not given.	74F646
	C ₆ H ₆ / MeOH (80:20)	9 × 10 ⁸			PL/Ad-8	S = MB.	737014
	CCl ₃ CH ₂ OH	3 × 10 ⁸			PL/Ad-8	S = MB; Solvent contained 1% MeOH.	74F646
	CCl ₄	8 × 10 ⁸			MP/LI-12	S = PP, Ph a or BPh a; used $k_d = 36$ s ⁻¹ .	79A010
	CCl ₄	8.0 × 10 ⁸			MP/LI-12	S = Ret; used $k_d = 36$ s ⁻¹ .	79F463
	CCl ₄	1.1 × 10 ⁸	1.2 × 10 ⁻⁵	298	CP/Ac-15	S = MB; used $k_d = 1.4 \times 10^3$ s ⁻¹ .	78F061
	CCl ₄	2.6 × 10 ⁸			PL/Ad-8	S = MB; Solvent contained 1% MeOH.	74F646
	CCl ₄ / MeOH (99:1)	7.3 × 10 ⁸		298	FP/Ad-5	S = MB.	78E238
	CCl ₄ / MeOH (98:2)	2.9 × 10 ⁸		293	PL/Ad-8	S = MB.	83A006 83A371
	CCl ₄ / MeOH (98:2)	4.8 × 10 ⁸		298	FP/Ad-5	S = MB.	78E238
	CCl ₄ / MeOH (97:3)	5.4 × 10 ⁸		298	FP/Ad-5	S = MB.	78E238
	CCl ₄ / MeOH (96:4)	5.2 × 10 ⁸		298	FP/Ad-5	S = MB.	78E238
	CCl ₄ / MeOH (95:5)	6.0 × 10 ⁸		298	FP/Ad-5	S = MB.	78E238
	CCl ₄ / MeOH (94:6)	2.5 × 10 ⁸			PL/Ad-8	S = MB.	74F646
	CCl ₄ / MeOH (90:10)	3 × 10 ⁸			PL/Ad-8	S = MB.	74F646
	CCl ₄ / MeOH (86:14)	2.6 × 10 ⁸			PL/Ad-8	S = MB.	74F646
	CCl ₄ / MeOH (83:17)	2.6 × 10 ⁸			PL/Ad-8	S = MB.	74F646
	CF ₃ CH ₂ OH	1.2 × 10 ⁸ (k_r)			CP/Ac,A'c-17	S = MB; A' = DPF; $k_r^{A'}$ not given. Solvent contained 1% MeOH.	74F646
	CF ₃ CH ₂ OH	2 × 10 ⁸	1 × 10 ⁻⁴		CP/Ac-15	S = MB; used $k_d = 2.3 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
	CF ₃ CH ₂ OH	6 × 10 ⁸			PL/Ad-8	S = MB; Solvent contained 1% MeOH.	74F646
	CH ₂ Cl ₂	1.6 × 10 ⁹	1.0 × 10 ⁻⁵	298	CP/Ac-15	S = MB; used $k_d = 1.6 \times 10^4$ s ⁻¹ .	78F061
	CH ₂ Cl ₂	1.1 × 10 ⁹		298	PL/Ad-5	S = MB.	78F061
	CH ₂ Cl ₂	8 × 10 ⁸			PL/Ad-8	S = MB.	74F646
	CH ₂ Cl ₂	6.3 × 10 ⁸ (k_r)			CP/Ac,A'c-17	S = MB; A' = DPF; used $k_r^{A'} = 8.6 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 7.3$.	74F646
	CH ₂ FC ₂ Cl ₂ OH	5 × 10 ⁸			PL/Ad-8	S = MB; Solvent contained 1% MeOH.	74F646
	CH ₃ CHCl ₂	6.0 × 10 ⁸		298	PL/Ad-5	S = MB.	78F061
	CH ₃ CHCl ₂	1.0 × 10 ⁹	1.5 × 10 ⁻⁵	298	CP/Ac-15	S = MB; used $k_d = 1.5 \times 10^4$ s ⁻¹ .	78F061
	CH ₃ CN	1.4 × 10 ⁹		298	PL/Ad-8	S = An; $\Delta V \ddagger = -15$ cm ³ mol ⁻¹ ; studied at 0.1–400 MPa.	92A014 90A018
	CH ₃ CN		1.3 × 10 ⁻⁵	293	CP/Ac-14	S = MB.	87F440
	CH ₃ CN	1.1 × 10 ⁹			PL/Ad-8	S = An.	82A349
	CH ₃ CN	1.3 × 10 ⁹			PL/Ad-5	S = 2-ACN.	81A287
	CH ₃ CN	1 × 10 ⁹	2 × 10 ⁻⁵		CP/Ac-15	S = MB; used $k_d = 1.8 \times 10^4$ s ⁻¹ .	74F646

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
1.1 Isobenzofuran, 1,3-diphenyl- (DPBF) — Continued							
	CH ₃ CN	6.6×10^8			PL/Ad-8	S = MB.	74F646
	CH ₃ CN	$1.1 \times 10^9 (k_r)$			CP/Ac,A'c-17	S = MB; A' = DPF; used $k_r^{A'} = 2.0 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 5.6$.	74F646
	CH ₃ CO ₂ C ₂ H ₅	8×10^8			PL/Ad-8	S = MB.	74F646
	CH ₃ CO ₂ C ₂ H ₅	$3.5 \times 10^8 (k_r)$			CP/Ac,A'c-17	S = MB; A' = DPF; used $k_r^{A'} = 3.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 12$.	74F646
	CH ₃ COCH ₃	$4.1 \times 10^8 (k_r)$	$9.4 \times 10^{-5} (\beta_r)$	298	CP/Ac-14	S = A; used $k_d = 3.9 \times 10^4$ s ⁻¹ .	84F240
	CH ₃ COCH ₃	6.4×10^8			PL/Ad-8	S = An.	82A349
	CH ₃ COCH ₃	1.0×10^9			PL/Ad-5	S = 2-ACN.	81A287
	CH ₃ COCH ₃	5×10^8	5×10^{-5}		CP/Ac-15	S = MB; used $k_d = 2.4 \times 10^4$ s ⁻¹ .	74F646
	CH ₃ COCH ₃	6×10^8			PL/Ad-8	S = MB.	74F646
	CH ₃ COCH ₃	$1.7 \times 10^9 (k_r)$			CP/Ac,A'c-17	S = MB; A' = DPF; used $k_r^{A'} = 1.8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 9.4$.	74F646
	CHCl ₃	4.9×10^8			PL/Ad-5	S = 2-ACN.	81A287
	CHCl ₃	9.0×10^8	1.0×10^{-5}	298	CP/Ac-15	S = MB; used $k_d = 9.0 \times 10^3$ s ⁻¹ .	78F061
	CHCl ₃	7.0×10^8		298	PL/Ad-5	S = MB.	78F061
	CHCl ₃	$1.4 \times 10^8 (k_r)$			CP/Ac,A'c-17	S = MB; A' = DPF; used $k_r^{A'} = 4.9 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 2.9$.	74F646
	CHCl ₃	3.6×10^8			PL/Ad-8	S = MB.	74F646
	CHCl ₃ /MeOH (90:10)	5×10^8	8×10^{-5}		CP/Ac-15	S = RB; used $k_d = 3.9 \times 10^4$ s ⁻¹ .	753071
	CHCl ₃ /MeOH (90:10)	$5 \times 10^8 (k_r)$			CP/Ac-14	S = RB; k_r derived using $\phi_{isc}(RB) = 0.66$ and $k_A = 5 \times 10^8$ L mol ⁻¹ s ⁻¹ .	753071
	CS ₂		7.5×10^{-6}		CP/Ac-15	S = Rub.	84F197
	CS ₂ /MeOH (98:2)	3.0×10^8			FP/Ad-5	S = MB.	737334
	Cl ₂ CHCH ₂ OH	5.7×10^8			PL/Ad-8	S = MB.	74F646
	Cl ₂ CHCHCl ₂	4.4×10^8		298	PL/Ad-5	S = MB.	78F061
	Cl ₂ CHCHCl ₂	5.9×10^8	1.4×10^{-5}	298	CP/Ac-15	S = MB; used $k_d = 8.3 \times 10^3$ s ⁻¹ .	78F061
	ClCF ₂ CCl ₂ F	9×10^8		293	PL/Ad-10	high pressure O ₂ .	79A113
	ClCF ₂ CCl ₂ F	$5 \times 10^8 (k_r)$		293	CL/Ad-36	high pressure O ₂ ; used $k_{O_2} = 2.5 \times 10^3$ L mol ⁻¹ s ⁻¹ .	79A113
	D ₂ O (mic)	1.0×10^9			PL/Ad-5	S = HYP; BRIJ 35 micelles.	88N343
	D ₂ O (mic)	6.2×10^8		295	PL/Ad-5	S = 2-ACN; 4.0×10^{-2} mol L ⁻¹ DDAB.	82N027
	D ₂ O (mic) pD = 7.4	1.0×10^9			PL/Ad-5	S = MB; 0.1 mol L ⁻¹ SDS.	81N048
	D ₂ O (mic) pD = 7.4	6.7×10^8			PL/Ad-5	S = MB; 0.1 mol L ⁻¹ CTAB.	81N048
	D ₂ O (mic)	1.3×10^9		293	PL/Ad-8	S = MB; 0.1 mol L ⁻¹ SDS.	80N018
	D ₂ O (mic)	6.7×10^8			PL/Ad-5	S = MB; 0.1 mol L ⁻¹ CTAB.	79N041
	D ₂ O (mic)	6.3×10^8			PL/Ad-5	S = 2-ACN; 0.1 mol L ⁻¹ Igepal CO-630.	79N041
	D ₂ O (mic)	6.6×10^8			PL/Ad-5	S = MB; 0.1 mol L ⁻¹ Igepal CO-660.	79N041
	D ₂ O (mic)	9.8×10^8			PL/Ad-5	S = 2-ACN; 0.05 mol L ⁻¹ sodium laurate.	79N041
	D ₂ O (mic)	6.6×10^8			PL/Ad-5	S = 2-ACN; 0.1 mol L ⁻¹ Brij 35.	79N041
	D ₂ O (mic)	1.0×10^9			PL/Ad-5	S = MB; 0.1 mol L ⁻¹ SDS.	79N041
	D ₂ O (mic)	1.1×10^9			PL/Ad-5	S = 2-ACN; 0.1 mol L ⁻¹ SDS.	79N041
	D ₂ O (mic)	6.5×10^8			PL/Ad-5	S = 2-ACN; 0.1 mol L ⁻¹ Igepal CO-660.	79N041
	D ₂ O (mic)	6.2×10^8			PL/Ad-5	S = 2-ACN; 0.1 mol L ⁻¹ CTAB.	79N041
	D ₂ O (mic)	8.4×10^8			PL/Ad-5	S = 2-ACN; 0.1 mol L ⁻¹ SDS.	78E144

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
5.83 Isobenzofuran, 1,3-diphenyl- (DPBF) — Continued							
	D ₂ O (ves) pD = 7	2.3×10^{10}	1.1×10^{-6}	293	CP/Ac-15	S = HYP; used $k_d = 2.5 \times 10^4$ s ⁻¹ ; 5×10^{-4} mol L ⁻¹ DPPC.	92N201
	DMF	1.1×10^9			CP/Ac-15	S = HPDME; used $k_d = 1.0 \times 10^5$ s ⁻¹ .	92F505
	DMF		2.0×10^{-5} (β_r)		CP/Ac-14	S = RB.	79F412
	DMSO	$\geq 2.8 \times 10^8$	1.2×10^{-4}		CP/Ac-15	S = CuTCPc; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	89F260
	DMSO/H ₂ O (75:25)	6.9×10^8			CP/Ac-15	S = MB; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	86R210
	EtOH	8.8×10^8			PL/Ad-5	S = HP.	91R206
	EtOH	1×10^9		291	CP/A'c-15	S = Zn(pc)(py) ₂ ; used $k_d = 7.1 \times 10^4$ s ⁻¹ .	88A284
	EtOH	1.1×10^9		291	PL/Ld-2	S = Zn(pc)(py) ₂ .	88A284
	EtOH	4.3×10^8 (k_r)	1.2×10^{-4} (β_r)	298	CP/Ac-14	S = A; used $k_d = 5.3 \times 10^4$ s ⁻¹ .	84F240
	EtOH	1.3×10^9	6.5×10^{-5}	298	CP/Ac-15	S = Eos; used $k_d = 8.3 \times 10^4$ s ⁻¹ .	78F061
	EtOH	1.0×10^9			PL/Ad-8	S = MB.	74F646
	EtOH		2×10^{-4}		CP/Ac-15	S = MB.	74F646
	H ₂ O (mic)	6.2×10^9		303	CP/Ac-20	S = HA; Q = N ₃ ⁻ ; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; 2.3×10^{-4} mol L ⁻¹ TDPB.	92N179
	H ₂ O (mic)	2.3×10^9		303	CP/Ac-20	S = HA; Q = N ₃ ⁻ ; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; 2.2×10^{-4} mol L ⁻¹ Triton X-100.	92N179
	H ₂ O (mic)	5.1×10^9		303	CP/Ac-20	S = HA; Q = N ₃ ⁻ ; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; 2.9×10^{-4} mol L ⁻¹ SDS.	92N179
	H ₂ O (mic)	2.2×10^{10}		313	CP/Ac-23	S = Py; Q = N ₃ ⁻ ; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; S and A' solubilized in SDS micelles.	78A174
	H ₂ O (mic)	2.8×10^{10}		313	CP/Ac-23	S = Py; Q = N ₃ ⁻ ; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; S and A' solubilized in DTAC micelles.	78A174
	H ₂ O (mic)	4.2×10^9	1.2×10^{-4}	298	CP/Ac-15	S = Eos; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; A solubilized in SDS micelles.	78F061
	H ₂ O (mic)	4.7×10^9	1.1×10^{-4}	298	CP/Ac-15	S = MB; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; S and A solubilized in SDS micelles.	78F061
	H ₂ O (mic)	5.2×10^9	9.7×10^{-5}		CP/Ac-15	S = MB; used $k_d = 5 \times 10^5$ s ⁻¹ ; A solubilized in SDS micelles.	767247
	H ₂ O (ves) pH = 7	2.5×10^{10}	1×10^{-5}	293	CP/Ac-15	S = HYP; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; 5×10^{-4} mol L ⁻¹ DPPC.	92N201
	H ₂ O (ves)	$>3 \times 10^6$		291	PL/Ld-2	S = Zn(pc)(py) ₂ ; unilamellar DPPC vesicles.	88A284
	H ₂ O (ves)	1×10^7		291	CP/A'c-15	S = Zn(pc)(py) ₂ ; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; unilamellar DPPC vesicles.	88A284
	HCONH ₂ / D ₂ O (90:10)	1.1×10^9			PL/Ad-6	S = HP.	83E667
	HMN	3.9×10^8		298	PL/Ad-8	S = An; $\Delta V^\ddagger = -6$ cm ³ mol ⁻¹ ; studied at 0.1-400 MPa.	92A014
	HOCH ₂ CH ₂ OH	1.5×10^9	3.2×10^{-4}		PL/Ad-8,16	S = MB or RB; k derived using β and $k_D = 5.1 \times 10^{-5}$ s ⁻¹ at [DPBF] = 1.7×10^{-5} mol L ⁻¹ .	737014
	MCH	3.7×10^8		298	PL/Ad-8	S = An; $\Delta V^\ddagger = -19$ cm ³ mol ⁻¹ ; studied at 0.1-400 MPa.	92A014 90A018
	MeOH	7.2×10^8		298	PL/Ad-8	S = An; $\Delta V^\ddagger = -20$ cm ³ mol ⁻¹ ; studied at 0.1-400 MPa.	92A014 90A018
	MeOH		8.0×10^{-5}		CP/Ac-14	S = MB.	90F157
	MeOH		7.8×10^{-5}		CP/Ac-14	S = RB.	90F157
	MeOH	9×10^8		293	PL/Ad-5	S = MB.	87E690
	MeOH		7×10^{-5}		CP/Ac-14	S = A; $f_r^A = \sim 1.0$.	87F479
	MeOH	8.1×10^8			PL/Ad-5	S = 2-ACN.	84F005

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
Isobenzofuran, 1,3-diphenyl-(DPBF) — Continued						
MeOH	$6.0 \times 10^8 (k_r)$	$2.4 \times 10^{-4} (\beta_r)$	298	CP/Ac-14	S = A; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	84F240
MeOH	8.3×10^8			PL/Tb-3	S = Ery.	83A050
MeOH	8.1×10^8			PL/Ad-8	S = An.	82A349
MeOH	7.8×10^8			CP/Ac-14	S = F_1^{2-} ; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	80N021
MeOH		$8.0 \times 10^{-5} (\beta_r)$		CP/Ac-14	S = N-[5-Hydroxy-8-[[2-(methylsulfonyl)-4-nitrophenyl]azol]-1-naphthalenyl-1,3-benzenedisulfonamide, conjugate base.	79F412
MeOH	7.8×10^8			CP/Ac-23	S = Py; Q = N ₃ ⁻ ; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	78A174
MeOH	1.6×10^9	7.1×10^{-5}	298	CP/Ac-15	S = Ery; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	78F061
MeOH	1.6×10^9		298	PL/Ad-5	S = MB.	78F061
MeOH	1.6×10^9	6.8×10^{-5}	298	CP/Ac-15	S = MB; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	78F061
MeOH	1.0×10^9	1.1×10^{-4}	293	CP/Ac-14	S = Acridine Orange; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; Complicated kinetic treatment.	767105
MeOH		6.4×10^{-5}		CP/Ac-15	S = MB.	767247
MeOH	1.3×10^9	7×10^{-5}		CP/Ac-15	S = MB; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	74F646
MeOH	$1.4 \times 10^9 (k_r)$			CP/Ac,A'c-17	S = MB; A' = DPF; used $k_r^{A'} = 1.1 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 12$.	74F646
MeOH	1×10^9			PL/Ad-8	S = RB.	737014
MeOH	1.3×10^9			PL/Ad-8	S = MB.	737014
MeOH	$6.1 \times 10^8 (k_r)$	$2.3 \times 10^{-4} (\beta_r)$		CP/Ac-14	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; used $\phi_{isc}(RB) = 0.76$.	73F660
MeOH	8×10^8			PL/Ad-5	S = MB.	727260
MeOH		$7.3 \times 10^{-5} (\beta_r)$	298	CP/Ac,A'c-17	S = RB; A' = DPF; used $\beta_r^{A'} = 9.5 \times 10^{-4}$ mol L ⁻¹ .	717398
MeOH	7.0×10^8			PL/Ad-5	S = MB.	719325
MeOH/ D ₂ O (90:10)	1.6×10^9			PL/Ad-6	S = HP.	83E667
MeOH/ H ₂ O (50:50)	5.1×10^9	5.5×10^{-5}		PL/Ad-8,16	S = RB or MB; k derived from β and $k_D = (3.7 \pm 1.2) \times 10^{-5}$ s ⁻¹ at [DPBF] = 1.7×10^{-5} mol L ⁻¹ .	737014
MeOH/ HOCH ₂ CH ₂ OH (50:50)	1.9×10^9	6.3×10^{-5}		PL/Ad-8,16	S = MB or RB; k derived from β and $k_{D_1} = 1.6 \times 10^{-5}$ s ⁻¹ at [DPBF] = 1.9×10^{-5} mol L ⁻¹ .	737014
CH ₃ OCH ₂ CH ₂ OH	1.2×10^9		293	PL/Ad-5	S = MB.	87E690
2-PrOH	$6.1 \times 10^8 (k_r)$			CP/Ac,A'c-17	S = MB; A' = DPF; used $k_r^{A'} = 4.8 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 13$; Solvent contained 1% MeOH.	74F646
2-PrOH	5×10^8	1×10^{-4}		CP/Ac-15	S = MB; used $k_d = 5.0 \times 10^4$ s ⁻¹ ; Solvent contained 1% MeOH.	74F646
2-PrOH	1.5×10^8			PL/Ad-8	S = MB; Solvent contained 1% MeOH.	74F646
THF	$5.2 \times 10^8 (k_r)$	$8.4 \times 10^{-5} (\beta_r)$	298	CP/Ac-14	S = A; used $k_d = 4.4 \times 10^4$ s ⁻¹ .	84F240
THF	$8.3 \times 10^8 (k_r)$			CP/Ac,A'c-17	S = RB; A' = DPF; used $k_r^{A'} = 7.5 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 11$.	74F646
THF	5×10^8			PL/Ad-8	S = RB.	74F646
dioxane	$3.1 \times 10^8 (k_r)$			CP/Ac,A'c-17	S = RB; A' = DPF; used $k_r^{A'} = 2.7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 11$.	74F646
dioxane	8.4×10^8			PL/Ad-8	S = RB.	74F646
dioxane	1.2×10^9			PL/Ad-8	S = MB.	737014
i-octane	$4.4 \times 10^8 (k_r)$	$1.2 \times 10^{-4} (\beta_r)$	298	CP/Ac-14	S = A; used $k_d = 5.0 \times 10^4$ s ⁻¹ .	84F240

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
5.84	Isobenzofuran, hexaphenyl-						
	MeOH	6.9×10^8 (k_r)			CP/Ac-14	$S = RB$; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; used $\phi_{isc}(RB) = 0.76$.	73F660
5.85	Isobenzofuran, 1,3,4,7-tetraphenyl-						
	MeOH	6.6×10^8 (k_r)			CP/Ac-14	$S = RB$; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; used $\phi_{isc}(RB) = 0.76$.	73F660
5.86	Psoralen						
	CCl ₄	5.6×10^3			MP/LI-12	$S = TPP$; used $k_d = 39$ s ⁻¹ .	83E813
5.87	Psoralen, 8-methoxy-						
	CCl ₄	9.4×10^3			MP/LI-12	$S = TPP$; used $k_d = 39$ s ⁻¹ .	83E813
	H ₂ O	2×10^9			CP/Ac-14	$S = A$; used $k_d = 2.6 \times 10^5$ s ⁻¹ .	83F188
5.88	Pyran, 5-acetyl-3,4-dihydro-6-methyl-						
	CCl ₄	1.2×10^5		283	CP/A'c-33	$S = A' = Rub$; used $k_d = 1.1 \times 10^3$ s ⁻¹ ; $k_{A'}$ not given.	90F329
	CH ₃ CN	6.8×10^3		283	CP/A'c-33	$S = MB$; $A' = DPBF$; used $k_d = 1.9 \times 10^4$ s ⁻¹ ; $k_{A'}$ not given.	90F329
	CHCl ₃	1.5×10^5		298	CP/A'c-33	$S = A' = Rub$; used $k_d = 7.1 \times 10^3$ s ⁻¹ ; $\Delta H^\ddagger = -22.6$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -221.6$ J K ⁻¹ mol ⁻¹ ; studied at 283–323 K; $k_{A'}$ not given.	90F329
5.89	Pyran, 5-benzoyl-3,4-dihydro-6-phenyl-						
	CH ₃ CN	1.9×10^3		283	CP/A'c-33	$S = MB$; $A' = DPBF$; used $k_d = 7.1 \times 10^3$ s ⁻¹ .	90F329
5.90	Pyran, 3,4-dihydro-						
	CCl ₄	7.9×10^4		283	CP/A'c-33	$S = A' = Rub$; used $k_d = 1.1 \times 10^3$ s ⁻¹ ; $k_{A'}$ not given.	90F329
	CH ₃ CN	1.8×10^5		283	CP/A'c-33	$S = MB$; $A' = DPBF$; used $k_d = 7.1 \times 10^3$ s ⁻¹ ; $k_{A'}$ not given.	90F329
	CH ₃ COCH ₃	7.2×10^4 (k_r)		281	?	$S = RB$; $A' = 1,4$ -Dioxene; used $k_{A'} = 2.2 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 0.33$.	70F733
	CHCl ₃	7.1×10^4		298	CP/A'c-33	$S = A' = Rub$; used $k_d = 7.1 \times 10^3$ s ⁻¹ ; $k_{A'}$ not given.	90F329
		6.1×10^4		323			
5.91	Pyran, 3,4-dihydro-5,6-dimethyl-						
	MeOH	7.2×10^7 (k_r)		286	CP/Oc-29	$S = RB$; $A' = 2,5$ -DMF; used $k_d = 1.4 \times 10^5$ s ⁻¹ , $k_{A'} = 1.8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_r \leq 0.1$, 24% [2+2] cycloaddition.	91F163
5.92	Pyran-4- <i>d</i> , 2,3-dihydro-4-methyl-						
	C ₆ H ₆		?			$S = TPP$; $A' = 2,3$ -Dihydro-4-methylpyran; meas. $k_r^{A'}/k_r = 1.087$.	777645
	CH ₃ CN		?			$S = MB$; $A' = 2,3$ -Dihydro-4-methylpyran; meas. $k_r^{A'}/k_r = 1.787$.	777645
5.93	Pyran-4- <i>t</i> , 2,3-dihydro-4-methyl-						
	C ₆ H ₆			CP/Ac,A'c-17		$S = TPP$; $A' = 2,3$ -Dihydro-4-methylpyran; meas. $k_r^{A'}/k_r = 1.335$.	777645
	CH ₃ CN			Ac,A'c-17		$S = MB$; $A' = 2,3$ -Dihydro-4-methylpyran; meas. $k_r^{A'}/k_r = 1.211$.	777645
5.94	Pyran-5- <i>t</i> , 2,3-dihydro-4,4-dimethyl-						
	C ₆ H ₆			CP/Ac,A'c-17		$S = TPP$; $A' = 2,3$ -Dihydro-4,4-dimethylpyran; meas. $k_r^{A'}/k_r = 0.897$.	777645
	CH ₃ CN			CP/Ac,A'c-17		$S = MB$; $A' = 2,3$ -Dihydro-4,4-dimethylpyran; meas. $k_r^{A'}/k_r = 0.897$.	777645

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
§.95 Pyran-5- <i>t</i> , 2,3-dihydro-4-methyl-							
	C ₆ H ₆				CP/Ac,A'c-17	S = TPP; A' = 2,3-Dihydro-4-methylpyran; meas. $k_t^{A'}/k_t = 0.908$.	777645
	CH ₃ CN				CP/Ac,A'c-17	S = MB; A' = 2,3-Dihydro-4-methylpyran; meas. $k_t^{A'}/k_t = 0.866$.	777645
§.96 Pyran-6- <i>t</i> , 2,3-dihydro-4,4-dimethyl-							
	C ₆ H ₆				CP/Ac,A'c-17	S = TPP; A' = 2,3-Dihydro-4,4-dimethylpyran; meas. $k_t^{A'}/k_t = 0.994$.	777645
	CH ₃ CN				CP/Ac,A'c-17	S = MB; A' = 2,3-Dihydro-4,4-dimethylpyran; meas. $k_t^{A'}/k_t = 1.0$.	777645
§.97 Pyran-6- <i>t</i> , 2,3-dihydro-4-methyl-							
	C ₆ H ₆				CP/Ac,A'c-17	S = TPP; A' = 2,3-Dihydro-4-methylpyran; meas. $k_t^{A'}/k_t = 0.98$.	777645
	CH ₃ CN				CP/Ac,A'c-17	S = MB; A' = 2,3-Dihydro-4-methylpyran; meas. $k_t^{A'}/k_t = 1.07$.	777645
§.98 Pyran-5-carboxylic acid, 3,4-dihydro-6-methyl-, ethyl ester							
	C ₆ D ₆	4.3×10^4		298	CP/Oc-17	S = TPP; A' = DPBF; rel. to $k_A = 1 \times 10^5$ L mol ⁻¹ s ⁻¹ measured in CH ₃ CN.	90F329
	C ₆ H ₆	2.4×10^4		298	CP/Oc-17	S = TPP; A' = DPBF; rel. to $k_A = 1 \times 10^5$ L mol ⁻¹ s ⁻¹ measured in CH ₃ CN.	90F329
	CCl ₄	3.9×10^4		298	CP/A'c-33	S = A' = Rub; used $k_d = 1.1 \times 10^3$ s ⁻¹ ; $\Delta H^\ddagger = -22.6$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -191.5$ J K ⁻¹ mol ⁻¹ ; studied at 283-323 K; k_A' not given.	90F329
	CCl ₄	4.3×10^4		298	CP/Oc-17	S = TPP; A' = DPBF; rel. to $k_A = 1 \times 10^5$ L mol ⁻¹ s ⁻¹ measured in CH ₃ CN.	90F329
	CD ₃ CN	1.1×10^5		298	CP/Oc-17	S = MB; A' = DPBF; rel. to $k_A = 1 \times 10^5$ L mol ⁻¹ s ⁻¹ measured in CH ₃ CN.	90F329
	CH ₂ Cl ₂	5.4×10^4		298	CP/Oc-17	S = TPP; A' = DPBF; rel. to $k_A = 1 \times 10^5$ L mol ⁻¹ s ⁻¹ measured in CH ₃ CN.	90F329
	CH ₃ CN	1.0×10^5		298	CP/A'c-33	S = MB; A' = DPBF; used $k_d = 1.9 \times 10^4$ s ⁻¹ ; $\Delta H^\ddagger = -22.8$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -222.5$ J K ⁻¹ mol ⁻¹ ; studied at 283-323 K; k_A' not given.	90F329
	CHCl ₃	4.0×10^4		298	CP/A'c-33	S = A' = Rub; used $k_d = 7.1 \times 10^3$ s ⁻¹ ; $\Delta H^\ddagger = -19.1$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -220.9$ J K ⁻¹ mol ⁻¹ ; studied at 283-323 K; k_A' not given.	90F329 91F497
§.99 Pyran-5-carboxylic acid, 3,4-dihydro-6-(1-methylethyl)-, ethyl ester							
	CCl ₄	5.5×10^3		298	CP/A'c-33	S = A' = Rub; k_d and k_A' not given.	91F497
	CH ₃ CN	1.1×10^5		298	CP/A'c-33	S = MB; A' = DPBF; k_d and k_A' not given.	91F497
	CHCl ₃	4.3×10^4		298	CP/A'c-33	S = A' = Rub; $\Delta H^\ddagger = -17.4$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -215$ J K ⁻¹ mol ⁻¹ ; studied at 273-323 K; k_d and k_A' not given.	91F497
§.100 Pyran-5-carboxylic acid, 3,4-dihydro-6-phenyl-, ethyl ester							
	CCl ₄	2.8×10^4		283	CP/A'c-33	S = A' = Rub; used $k_d = 1.1 \times 10^3$ s ⁻¹ ; k_A' not given.	90F329
	CH ₃ CN	1.1×10^5		283	CP/A'c-33	S = MB; A' = DPBF; used $k_d = 1.9 \times 10^4$ s ⁻¹ ; k_A' not given.	90F329
	CHCl ₃	1.2×10^4		298	CP/A'c-33	S = A' = Rub; used $k_d = 7.1 \times 10^3$ s ⁻¹ ; k_A' not given.	90F329 91F497
§.101 Pyran-5-carboxylic acid, 6-ethyl-3,4-dihydro-, ethyl ester							
	CH ₃ CN	4.1×10^4		298	CP/A'c-33	S = MB; A' = DPBF; k_d and k_A' not given.	91F497

TABLE 5. Rate constants for interaction of singlet oxygen with furans, pyrans and other heterocyclic oxygen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
5.101 Pyran-5-carboxylic acid, 6-ethyl-3,4-dihydro-, ethyl ester — Continued							
	CHCl ₃	2.7×10^4		298	CP/A'c-33	S = A' = Rub; $\Delta H^\ddagger = -12.9$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -204.7$ J K ⁻¹ mol ⁻¹ ; studied at 273-323 K; k_d and $k_{A'}$ not given.	91F497
5.102 Sydnone, 3,4-diphenyl-							
	C ₆ H ₆	5.3×10^6		296	PL/A'd-5	S = NMTA; A' = DPBF.	86A203
	MeOH	2.8×10^6		296	PL/A'd-5	S = NMTA; A' = DPBF.	86A203
5.103 Sydnone, 4-methyl-3-(4-methylphenyl)-							
	c-C ₆ H ₁₂	$\sim 1.0 \times 10^7$		296	PL/A'd-5	S = CQ; A' = DPBF.	86A203
	C ₆ H ₆	4.0×10^7		296	PL/A'd-5	S = NMTA; A' = DPBF.	86A203
	CCl ₄	1.8×10^7		296	PL/A'd-5	S = CQ; A' = DPBF.	86A203
	CH ₃ CN	4.9×10^7		296	PL/A'd-5	S = CQ; A' = DPBF.	86A203
	CH ₃ COCH ₃	4.2×10^7		296	PL/A'd-5	S = NMTA; A' = DPBF.	86A203
	MeOH	1.7×10^7		296	PL/A'd-5	S = NMTA; A' = DPBF.	86A203
5.104 Sydnone, 3-(4-methylphenyl)-							
	C ₆ H ₆	$\leq 5 \times 10^5$		296	PL/A'd-5	S = CQ; A' = DPBF.	86A203
	MeOH	$\leq 5 \times 10^5$		296	PL/A'd-5	S = NMTA; A' = DPBF.	86A203
5.105 Sydnone, 3-methyl-4-phenyl-							
	C ₆ H ₆	2.7×10^6		296	PL/A'd-5	S = NMTA; A' = DPBF.	86A203
	MeOH	2.2×10^6		296	PL/A'd-5	S = NMTA; A' = DPBF.	86A203
5.106 Sydnone, 3-(4-methylphenyl)-4-phenyl-							
	C ₆ H ₆	5.4×10^6		296	PL/A'd-5	S = NMTA; A' = DPBF.	86A203
	MeOH	3.5×10^6		296	PL/A'd-5	S = NMTA; A' = DPBF.	86A203
5.107 Sydnone, 3-phenyl-							
	C ₆ H ₆	$\leq 5 \times 10^5$		296	PL/A'd-5	S = CQ; A' = DPBF.	86A203
	MeOH	$\leq 5 \times 10^5$		296	PL/A'd-5	S = NMTA; A' = DPBF.	86A203

TABLE 6. Rate constants for interaction of singlet oxygen with pyrroles, oxazoles, pyridines and other heterocyclic nitrogen compounds.

Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
1,8-Acridinedione, 3,4,6,7,9,10-hexahydro-3,3,6,6-tetramethyl-						
CHCl ₃	2.0×10^6			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327
Carbazole, 2,3,4,9-tetrahydro-9-methyl-						
MeOH			293	CP/Ac-17	S = RB; A' = 9-Acetyl-2,3,4,9-tetrahydrocarbazole; meas. $k_t/k_t^{A'} = 59$.	79F434
Carbazole, 2,3,4,9-tetrahydro-9-methyl-6-nitro-						
MeOH			293	CP/Ac-17	S = RB; A' = 9-Acetyl-2,3,4,9-tetrahydrocarbazole; meas. $k_t/k_t^{A'} = 4.8$.	79F434
Cyclopenta[b]quinoline, 4-methyl-1,2-diphenyl-						
C ₆ H ₆	3.0×10^6	0.014		CP/Ac-15	S = A; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	78F438
Histamine						
H ₂ O pH = 7.1	2.8×10^7		298	CP/Oc-19	S = Phenosafranine; Q = N ₃ ⁻ ; used $k_Q = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A360
H ₂ O	2.0×10^8	2.5×10^{-3}	298	CP/Ac-15	S = MB; used $k_d = 5.0 \times 10^5$ s ⁻¹ .	78F061
Imidazole						
D ₂ O pD = 7	2×10^7			PL/Ld-2	S = Carboxyanthracene.	92E225
D ₂ O/EtOH (75:25)	1.4×10^7 (k_t)		295	CP/Ac-14,28	S = RB; used $k_d = 4.8 \times 10^4$ s ⁻¹ ; DMA as actinometer, used $k_A = 2.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	94A113
D ₂ O/EtOH (75:25)	2.4×10^7		295	PL/Ld-2	S = RB.	94A113
H ₂ O	3.4×10^7 (k_t)		298	CP/Oc-14	S = H ₂ TPPS ⁴⁻ ; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; $E_a = 30.2$ kJ mol ⁻¹ ; log(A) = 12.8; studied at 288-318 K; phosphate buffer contg. 1.6% NaCl.	91R053
H ₂ O pH = 7.0	3.4×10^7			CP/Oc-19	S = HP; Q = N ₃ ⁻ ; used $k_Q = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A278
H ₂ O pH = 7.1	4×10^7		298	CP/Oc-19	S = Phenosafranine; Q = N ₃ ⁻ ; used $k_Q = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A360
H ₂ O (mic) pH = 7.0	2.0×10^7			CP/Oc-19	S = Chl a; Q = N ₃ ⁻ ; used $k_Q = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; 1% Triton X-100; $k = 2.9 \times 10^7$ and 3.9×10^7 L mol ⁻¹ s ⁻¹ in 2% and 5% Triton X-100, resp.	78A278
H ₂ O (mic) pH = 7.0	3.6×10^7			CP/Oc-19	S = HP; Q = N ₃ ⁻ ; used $k_Q = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; 1% Triton X-100.	78A278
Imidazole, 4-methyl-						
D ₂ O/EtOH (75:25)	1.3×10^8		295	PL/Ld-2	S = RB.	94A113
D ₂ O/EtOH (75:25)	1.1×10^8 (k_t)		295	CP/Ac-14,28	S = RB; used $k_d = 4.8 \times 10^4$ s ⁻¹ ; DMA as actinometer, used $k_A = 1.3 \times 10^8$ L mol ⁻¹ s ⁻¹ .	94A113
Indole						
C ₆ H ₅ CH ₃	7.7×10^5			PR/A'd-5	S = Np; A' = DPBF; $\Delta H^\ddagger = 0$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -141$ J K ⁻¹ mol ⁻¹ ; studied at 223-353 K.	79A106
CH ₃ CN	$<10^4$ (k_t)			CP/Ac,A'c-17	S = ZnTPP; A' = An; used $k_t^{A'} = 1.2 \times 10^6$ L mol ⁻¹ s ⁻¹ .	90F347
CH ₃ CN	2.5×10^6			PL/Ld-2	S = ZnTPP.	90F347
D ₂ O/EtOH (75:25)	1.9×10^7		295	PL/Ld-2	S = RB.	94A113
D ₂ O/EtOH (75:25)	9×10^6 (k_t)		295	CP/Ac-14,28	S = RB; used $k_d = 4.8 \times 10^4$ s ⁻¹ ; DMA as actinometer, used $k_A = 1.9 \times 10^7$ L mol ⁻¹ s ⁻¹ .	94A113
H ₂ O pH = 7.2	7×10^7			CP/Ac-14	S = RB; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	88A385
Indole, 5,6-dimethoxy-						
CHCl ₃	4.3×10^7	2.3×10^{-4}		CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F670

TABLE 6. Rate constants for interaction of singlet oxygen with pyrroles, oxazoles, pyridines and other heterocyclic nitrogen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
6.10	Indole, 1,3-dimethyl-						
	MeOH			293	CP/Ac-17	S = RB; A' = 9-Acetyl-2,3,4,9-tetrahydrocarbazole; meas. $k_r/k_r^{A'} = 13$.	79F434
6.11	Indole, 2,3-dimethyl-						
	C ₆ H ₅ CH ₃	1.0×10^8			PR/A'd-5	S = Np; A' = DPBF; $\Delta H^\ddagger = 0$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -99.2$ J K ⁻¹ mol ⁻¹ ; studied at 223-353 K.	79A106
	D ₂ O (ves)	4.8×10^7		295	PL/A'd-5	S = 2-ACN; A' = DPBF; 4.0×10^{-2} mol L ⁻¹ DDAB.	82N027
	D ₂ O (mic) pD = 7.4	7.0×10^8			PL/A'd-5	S = MB; A' = DPBF; 0.1 mol L ⁻¹ SDS.	81N048
	D ₂ O (mic) pD = 7.4	4.4×10^8			PL/A'd-5	S = MB; A' = DPBF; 0.1 mol L ⁻¹ CTAB.	81N048
6.12	Indole, 5-hydroxy-						
	CHCl ₃	$<2 \times 10^5$	>0.05		CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F670
6.13	Indole, 5-methoxy-						
	CHCl ₃	$<2 \times 10^5$	>0.05		CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F670
6.14	Indole, 5-methoxy-1,2,3-trimethyl-						
	MeOH			293	CP/Ac-17	S = RB; A' = 9-Acetyl-2,3,4,9-tetrahydrocarbazole; meas. $k_r/k_r^{A'} = 98$.	79F434
6.15	Indole, 1-methyl-						
	CH ₃ CN	3.7×10^6			PL/Ld-2	S = ZnTPP.	90F347
	CH ₃ CN	$<10^4$ (k_r)			CP/Ac,A'c-17	S = ZnTPP; A' = An; used $k_r^{A'} = 1.2 \times 10^6$ L mol ⁻¹ s ⁻¹ .	90F347
6.16	Indole, 3-methyl-						
	C ₆ H ₅ CH ₃	9.2×10^6			PR/A'd-5	S = Np; A' = DPBF; $\Delta H^\ddagger = 0$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -120$ J K ⁻¹ mol ⁻¹ ; studied at 223-353 K.	79A106
	CH ₂ Cl ₂	6×10^6			PL/Tb-3	S = TPP.	83A050
	CH ₃ CN	6.8×10^7			PL/Ld-2	S = ZnTPP.	90F347
	CH ₃ CN	2.3×10^6 (k_r)			CP/Ac,A'c-17	S = ZnTPP; A' = An; used $k_r^{A'} = 1.2 \times 10^6$ L mol ⁻¹ s ⁻¹ .	90F347
	D ₂ O/ EtOH (75:25)	1.6×10^8		295	PL/Ld-2	S = RB.	94A113
	D ₂ O/ EtOH (75:25)	1.2×10^8 (k_r)		295	CP/Ac-14,28	S = RB; used $k_d = 4.8 \times 10^4$ s ⁻¹ ; DMA as actinometer, used $k_A = 1.6 \times 10^8$ L mol ⁻¹ s ⁻¹ .	94A113
6.17	Indole, 3-methyl-2-phenyl-						
	C ₆ H ₅ CH ₃	2.1×10^7			PR/A'd-5	S = Np; A' = DPBF; $\Delta H^\ddagger = 0$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -113$ J K ⁻¹ mol ⁻¹ ; studied at 223-353 K.	79A106
6.18	Indole, 1,2,3-trimethyl-						
	MeOH			293	CP/Ac-17	S = RB; A' = 9-Acetyl-2,3,4,9-tetrahydrocarbazole; meas. $k_r/k_r^{A'} = 70$.	79F434
6.19	Indole, 1,2,3-trimethyl-5-nitro-						
	MeOH			293	CP/Ac-17	S = RB; A' = 9-Acetyl-2,3,4,9-tetrahydrocarbazole; meas. $k_r/k_r^{A'} = 5.0$.	79F434
6.20	Indole-3-acetate ion						
	H ₂ O pH = 7	9.2×10^7 (k_r)			CP/Oc,A'c-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91F359
	H ₂ O pH = 7.0	1.0×10^9	1.0×10^{-4}		CP/Ac-18	S = FMN; Q = N ₃ ⁻ ; used $k_d = 1.0 \times 10^5$ s ⁻¹ .	86A406

TABLE 6. Rate constants for interaction of singlet oxygen with pyrroles, oxazoles, pyridines and other heterocyclic nitrogen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
6.21 Indole-3-acetic acid							
	CH ₃ CN	$4.6 \times 10^5 (k_r)$			CP/Ac-29	S = ZnTPP; A' = An; used $k_{A'} = 1.2 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_r = 40 \pm 6$; Used $k_A = 1.9 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F347
	CH ₃ CN	1.9×10^7			PL/Ld-2	S = ZnTPP.	90F347
	D ₂ O/EtOH (75:25)	7.8×10^7		295	PL/Ld-2	S = RB.	94A113
	H ₂ O pH = 4				CP/A'c-29	S = RB; A' = Met; used $k_{A'} = 2.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_r = 12 \pm 5$.	90F347
	H ₂ O pH = 4.5	7.7×10^8	1.3×10^{-4}		CP/Ac-18	S = FMN; Q = N ₃ ⁻ ; used $k_d = 1.0 \times 10^5$ s ⁻¹ .	86A406
	H ₂ O pH = 3.0	5.9×10^8	1.7×10^{-4}		CP/Ac-18	S = FMN; Q = N ₃ ⁻ ; used $k_d = 1.0 \times 10^5$ s ⁻¹ .	86A406
6.22 Indole-3-butyrate ion							
	H ₂ O pH = 7	$7.2 \times 10^7 (k_r)$			CP/Oc-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91F359
6.23 3-Indolebutyric acid							
	CH ₃ CN	$1.3 \times 10^6 (k_r)$			CP/A'c-29	S = ZnTPP; A' = An; used $k_{A'} = 1.2 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_r = 32 \pm 3$; Used $k_A = 4.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F347
	CH ₃ CN	4.5×10^7			PL/Ld-2	S = ZnTPP.	90F347
	H ₂ O pH = 4				CP/A'c-29	S = RB; A' = Met; used $k_{A'} = 2.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_r = 11 \pm 5$.	90F347
6.24 Indole-3-carbinol							
	CH ₃ CN	$6.1 \times 10^5 (k_r)$			CP/Ac,A'c-17	S = ZnTPP; A' = An; used $k_r^{A'} = 1.2 \times 10^6$ L mol ⁻¹ s ⁻¹ .	90F347
	CH ₃ CN	1.7×10^7			PL/Ld-2	S = ZnTPP.	90F347
6.25 Indole-2-carboxylic acid							
	CH ₃ CN	$<10^4 (k_r)$			CP/Ac,A'c-17	S = ZnTPP; A' = An; used $k_r^{A'} = 1.2 \times 10^6$ L mol ⁻¹ s ⁻¹ .	90F347
	CH ₃ CN	3.6×10^5			PL/Ld-2	S = ZnTPP.	90F347
6.26 Indole-3-carboxylate ion							
	H ₂ O pH = 7	$1.4 \times 10^7 (k_r)$			CP/Oc,A'c-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91F359
6.27 Indole-3-carboxylic acid							
	CH ₃ CN	$1.3 \times 10^5 (k_r)$			CP/Ac,A'c-17	S = ZnTPP; A' = An; used $k_r^{A'} = 1.2 \times 10^6$ L mol ⁻¹ s ⁻¹ .	90F347
	CH ₃ CN	5.2×10^5			PL/Ld-2	S = ZnTPP.	90F347
6.28 Indole-3-propionamide							
	D ₂ O/EtOH (75:25)	7.7×10^7		295	PL/Ld-2	S = RB.	94A113
6.29 Indole-3-propionate ion							
	H ₂ O pH = 7	$7.3 \times 10^7 (k_r)$			CP/Oc-17	S = RB; A' = His; used $k_r^{A'} = 8.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91F359
6.30 Indole-3-propionic acid							
	CH ₃ CN	$6.1 \times 10^5 (k_r)$			CP/A'c-29	S = ZnTPP; A' = An; used $k_{A'} = 1.2 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_r = 54 \pm 8$; Used $k_A = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F347
	CH ₃ CN	3.3×10^7			PL/Ld-2	S = ZnTPP.	90F347
	D ₂ O/EtOH (75:25)	$6.6 \times 10^7 (k_r)$		295	CP/Ac-14,28	S = RB; used $k_d = 4.8 \times 10^4$ s ⁻¹ ; DMA as actinometer, used $k_A = 9.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	94A113

TABLE 6. Rate constants for interaction of singlet oxygen with pyrroles, oxazoles, pyridines and other heterocyclic nitrogen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
6.30	Indole-3-propionic acid — Continued						
	D ₂ O/ EtOH (75:25)	9.8×10^7		295	PL/Ld-2	S = RB.	94A113
	H ₂ O pH = 4				CP/A'c-29	S = RB; A' = Met; used $k_{A'} = 2.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_r = 15 \pm 4$.	90F347
6.31	Oxazole, 2,5-diphenyl-						
	H ₂ O/ D ₂ O (50:50 (mic))	1.6×10^8	2.2×10^{-3}	298	CP/Ac-15	S = MB; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; A solubilized in DTAC micelles; solvent mixt. in mole %.	78F061
	H ₂ O/ D ₂ O/ MeOH (20:50:30)	1.6×10^8	1.6×10^{-3}	298	CP/Ac-15	S = MB; used $k_d = 2.6 \times 10^5$ s ⁻¹ ; solvent mixt. in mole %.	78F061
6.32	Oxazole, 2,5-diphenyl-4-methyl-						
	C ₆ H ₆	3.9×10^7	2.0×10^{-3}		CP/Ac-14	S = ?; used $k_d = 8.0 \times 10^4$ s ⁻¹ .	81F622
	CH ₃ (CH ₂) ₄ OH		3.0×10^{-3}		CP/Ac-14	S = ?.	81F622
	DMSO		1.5×10^{-3}		CP/Ac-14	S = ?.	81F622
	EtOH	3.3×10^7	5.5×10^{-3}		CP/Ac-14	S = ?; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	81F622
	MeOH	3.5×10^7	5.8×10^{-3}		CP/Ac-14	S = ?; used $k_d = 2.0 \times 10^5$ s ⁻¹ .	81F622
	dioxane	1.6×10^7	1.8×10^{-3}		CP/Ac-14	S = ?; used $k_d = 2.9 \times 10^4$ s ⁻¹ .	81F622
6.33	Oxazole, 4-methyl-2-(3-chlorophenyl)-5-phenyl-						
	MeOH	1.7×10^7	0.012		CP/Ac-14	S = ?; used $k_d = 2.0 \times 10^5$ s ⁻¹ .	81F622
6.34	Oxazole, 4-methyl-2-(4-chlorophenyl)-5-phenyl-						
	MeOH	1.9×10^7	0.010		CP/Ac-14	S = ?; used $k_d = 2.0 \times 10^5$ s ⁻¹ .	81F622
6.35	Oxazole, 4-methyl-2-(4-methoxyphenyl)-5-phenyl-						
	MeOH	5.3×10^7	3.8×10^{-3}		CP/Ac-14	S = ?; used $k_d = 2.0 \times 10^5$ s ⁻¹ .	81F622
6.36	Oxazole, 4-methyl-2-(4-methylphenyl)-5-phenyl-						
	MeOH	3.7×10^7	5.3×10^{-3}		CP/Ac-14	S = ?; used $k_d = 2.0 \times 10^5$ s ⁻¹ .	81F622
6.37	Oxazole, 4-methyl-2-(4-nitrophenyl)-5-phenyl-						
	MeOH	1.2×10^7	0.017		CP/Ac-14	S = ?, used $k_d = 2.0 \times 10^5$ s ⁻¹ .	81F622
6.38	Phenoxyazine						
	C ₆ H ₅ Br/ MeOH (67:33)	1.0×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 4.9 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777240
6.39	Pyrazolo[1',2':2,3][1,2,3]triazolo[4,5-a]phenazin-4-iium, 1,3-dimethyl-						
	c-C ₆ H ₁₂	4.5×10^7 (k_r)			CP/Ac-14	S = A; used $k_d = 5.9 \times 10^4$ s ⁻¹ .	80F548
	CHCl ₃	3.7×10^7 (k_r)			CP/Ac-14	S = A; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	80F548
	CHCl ₃ / MeOH (80:20)	6.3×10^7 (k_r)			CP/Ac-14	S = RB; used $k_d = 5.4 \times 10^4$ s ⁻¹ ; Used ϕ_A for RB = 0.76.	80F548
	CHCl ₃ / MeOH (80:20)	4.2×10^7 (k_r)			CP/Ac-17	S = RB; A' = DPBF; used $k_r^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ .	80F548
	CHCl ₃ / MeOH (80:20)	6.4×10^7 (k_r)			CP/Ac-14	S = A; used $k_d = 5.8 \times 10^4$ s ⁻¹ .	80F548
6.40	Pyridine						
	CCl ₄	2×10^3			PL/A'd-8	S = MB; A' = DPBF.	76F903
6.41	Pyridine, 3,5-diacetyl-1,4-dihydro-						
	CHCl ₃	4.7×10^6			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327
6.42	Pyridine, 3,5-diacetyl-1,4-dihydro-2,6-dimethyl-						
	CHCl ₃	5.1×10^6			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327

TABLE 6. Rate constants for interaction of singlet oxygen with pyrroles, oxazoles, pyridines and other heterocyclic nitrogen compounds. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
6.41	Pyridine, 3,5-dibenzoyl-1,4-dihydro-2,6-dimethyl-						
	CHCl ₃	3.8×10^6			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327
6.44	Pyridine-3,5-dicarboxamide, 1,4-dihydro-2,6-dimethyl-N,N-diphenyl-						
	CHCl ₃	7.5×10^7			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327
6.45	Pyridine-3,5-dicarboxamide, 1,4-dihydro-2,4,6-trimethyl-N,N-diphenyl-						
	CHCl ₃	6.5×10^6			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327
6.46	Pyridine-3,5-dicarboxylic acid, 1,4-dihydro-2,6-dimethyl-, dicyclohexyl ester						
	CHCl ₃	6.0×10^6			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327
6.47	Pyridine-3,5-dicarboxylic acid, 1,4-dihydro-2,6-dimethyl-, diethyl ester						
	CHCl ₃	3.4×10^6			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327
6.48	Pyridine-3,5-dicarboxylic acid, 1,4-dihydro-2,6-dimethyl-, dihexyl ester						
	CHCl ₃	5.0×10^6			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327
6.49	Pyridine-3,5-dicarboxylic acid, 1,4-dihydro-2,6-dimethyl-, di(1-methylethyl) ester						
	CHCl ₃	6.6×10^6			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327
6.50	Pyridine-3,5-dicarboxylic acid, 1,4-dihydro-2,6-diphenyl-, diethyl ester						
	CHCl ₃	<10 ⁵			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327
6.51	Pyridine-3,5-dicarboxylic acid, 1,4-dihydro-2,4,6-trimethyl-, diethyl ester						
	CHCl ₃	<10 ⁵			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327
6.52	Pyridine-3,5-dicarboxylic acid, 2,6-dimethyl-, diethyl ester						
	CHCl ₃	<10 ⁵			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327
6.53	Pyrnidinium, 2,3-dihydro-1-methyl-4-phenyl-						
	D ₂ O	1.7×10^6			PL/Ld-2	S = MB.	91R251
6.54	Pyridinium, 1-methyl-						
	D ₂ O	6.5×10^5			PL/Ld-2	S = MB.	91R251
6.55	Pyridinium, 1-methyl-4-phenyl-						
	D ₂ O	8.9×10^5			PL/Ld-2	S = MB.	91R251
6.56	Pyrido[4,3- <i>b</i>]indole, 2,3,4,4a,5,9b-hexahydro-2,8-dimethyl-, <i>cis</i> -(-), (Stobadine)						
	D ₂ O pD = 7.4	1.3×10^8		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	92A252
6.57	Pyrido[4,3- <i>b</i>]indole, <i>N</i> -acetyl-2,3,4,4a,5,9b-hexahydro-2,8-dimethyl-						
	D ₂ O pD = 7.4	1×10^6		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	92A252
6.58	Pyrido[4,3- <i>b</i>]indole, 1,2,3,4-tetrahydro-2,8-dimethyl- (Dehydrostobadine)						
	D ₂ O pD = 7.4	1.7×10^8		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	92A252
6.59	Pyrrole						
	C ₅ H ₅ N	1.4×10^9		295	PL/Ld-2	S = An.	93E684
	C ₅ H ₅ N	2×10^8			CP/A'c-18	S = TPBC; A' = Tetr; used $k_d = 6.3 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^6$ L mol ⁻¹ s ⁻¹ ; cor. for $k_q(S) = 10^8$ L mol ⁻¹ s ⁻¹ .	86F670
	1,2-C ₆ H ₄ (CH ₃) ₂	1.8×10^8		295	PL/Ld-2	S = An.	93E684
	C ₆ H ₅ CH ₃	1.7×10^8		295	PL/Ld-2	S = An.	93E684
	C ₆ H ₅ CN	2.1×10^8		295	PL/Ld-2	S = An.	93E684

TABLE 6. Rate constants for interaction of singlet oxygen with pyrroles, oxazoles, pyridines and other heterocyclic nitrogen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
6.59 Pyrrole — Continued							
	C ₆ H ₆	1.5 × 10 ⁸		295	PL/Ld-2	S = An.	93E684
	C ₆ H ₆	<1 × 10 ⁶			CP/A'c-18	S = TPBC; A' = Tetr; used $k_d = 3.3 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^6$ L mol ⁻¹ s ⁻¹ ; cor. for $k_q(S) = 10^8$ L mol ⁻¹ s ⁻¹ .	86F670
	CH ₃ COCH ₃	5.4 × 10 ⁸		295	PL/Ld-2	S = An.	93E684
	CH ₃ COCH ₃	6 × 10 ⁷			CP/A'c-18	S = TPBC; A' = Tetr; used $k_d = 2.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^6$ L mol ⁻¹ s ⁻¹ ; cor. for $k_q(S) = 10^8$ L mol ⁻¹ s ⁻¹ .	86F670
6.60 Pyrrole, 2,5-dimethyl-							
	MeOH	0.16		293	CP/Oc-15	S = RB; $E_a = 7.1$ kJ mol ⁻¹ .	68F288
6.61 Pyrrole, 1-(1,1-dimethylethyl)-							
	MeOH	1.2 × 10 ⁸	1.2 × 10 ⁻⁴		CP/Oc-?	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	75F652
	CH ₃ COCH ₃	3.9 × 10 ⁷	9.7 × 10 ⁻⁴		CP/Oc-?	S = MB; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	75F652
6.62 Pyrrole, 2-(1,1-dimethylethyl)-							
	CH ₃ COCH ₃	4.2 × 10 ⁷	9.0 × 10 ⁻⁴		CP/Oc-?	S = MB; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	75F652
	MeOH	1.5 × 10 ⁸	9.3 × 10 ⁻⁴		CP/Oc-?	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	75F652
6.63 Pyrrole, 3-(1,1-dimethylethyl)-							
	CH ₃ COCH ₃	2.9 × 10 ⁷	1.3 × 10 ⁻³		CP/Oc-?	S = MB; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	75F652
	MeOH	1.8 × 10 ⁸	7.8 × 10 ⁻⁴		CP/Oc-?	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	75F652
6.64 Pyrrole, 1-methyl-							
	C ₆ H ₆	3.7 × 10 ⁸		295	PL/Ld-2	S = An.	93E684
	CCl ₄	1 × 10 ⁹	5.5 × 10 ⁻⁸		CP/LI-12	S = PdTTPP; used $k_d = 55$ s ⁻¹ .	87A202
	CH ₃ COCH ₃	2.9 × 10 ⁸		295	PL/Ld-2	S = An.	93E684
6.65 Pyrrole-2,4-dicarboxylic acid, 5,5'-methylenebis[3-methyl-, tetraethyl ester							
	CCl ₄	2 × 10 ⁶	2.8 × 10 ⁻⁵		CP/LI-12	S = PdTTPP; used $k_d = 55$ s ⁻¹ .	87A202
6.66 Pyrrole-3,5-dicarboxylic acid, 2,4-dimethyl-, diethyl ester							
	CCl ₄	3 × 10 ⁶	1.8 × 10 ⁻⁵		CP/LI-12	S = PdTTPP; used $k_d = 55$ s ⁻¹ .	87A202
6.67 Pyrrolo[2,3,4-<i>k,l</i>]acridine-10-one, 1-(4-bromophenyl)-2,3,5,7,8,9-hexahydro-4,4,8,8-tetramethyl-							
	CHCl ₃	2.6 × 10 ⁸			PL/Ld-2	S = MPDEE.	89F368
6.68 Pyrrolo[2,3,4-<i>k,l</i>]acridine-10-one, 2,3,5,7,8,9-hexahydro-1-(4-methoxyphenyl)-4,4,8,8-tetramethyl-							
	CHCl ₃	2.5 × 10 ⁸			PL/Ld-2	S = MPDEE.	89F368
6.69 Pyrrolo[2,3,4-<i>k,l</i>]acridine-10-one, 2,3,5,7,8,9-hexahydro-4,4,8,8-tetramethyl-1-phenyl-							
	CHCl ₃	4.2 × 10 ⁸			PL/Ld-2	S = MPDEE.	89F368
6.70 Pyrrolo[3',2':3,4]cyclopenta[1,2-<i>b</i>]pyridine, 2-(4-ethoxycarbonylphenyl)-1,8-dihydro- (EPCP)							
	CH ₂ Cl ₂	3.6 × 10 ⁸ (k_r)			CP/Ac-14	S = Rubi; used $k_d = 1 \times 10^4$ s ⁻¹ ; used $\phi_\Delta(\text{Rubi}) = 0.31$.	92F270
	CH ₂ Cl ₂	4.0 × 10 ⁸			PL/Ld-2	S = MB.	92F270
6.71 Quinoline							
	EtOH	<1 × 10 ⁹		273	CP/Oc-23	S = MB; A' = 2,5-DMF; No measurable effect.	72F518
6.72 Quinoline, 1,2-dihydro-2,2,4-trimethyl-, homopolymer (Permanax 45)							
	EtOH	9.6 × 10 ⁸		273	CP/Oc-23	S = MB; A' = 2,5-DMF; used $k_d = 7.9 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^8$ L mol ⁻¹ s ⁻¹ ; k calcd. in [81Z251].	72F518

TABLE 6. Rate constants for interaction of singlet oxygen with pyrroles, oxazoles, pyridines and other heterocyclic nitrogen compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
6.73 Quinoline, 8-hydroxy-							
	CHCl ₃	1.1×10^8	9.5×10^{-5}		CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F670
6.74 Quinoline, 6-methoxy-							
	CHCl ₃	<2 × 10 ⁵	>0.05		CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F670

TABLE 7. Rate constants for the interaction of singlet oxygen with porphyrins, phthalocyanines and polypyrroles.

No.	Solvent	k ($\text{L mol}^{-1} \text{s}^{-1}$)	$\beta (k_d/k)$ (mol L^{-1})	T (K)	Method	Comment	Ref.
7.1 Bacteriochlorophyll <i>a</i>							
	$(\text{C}_2\text{H}_5)_2\text{O}$	2.2×10^8			PL/Ld-2	$S = A$.	90E324
	$(\text{C}_2\text{H}_5)_2\text{O}$				CL/Ac,A'c-17	$S = A' = \text{Rub}$; meas. $k_r/k_r^{A'} = 0.035$.	84F193
	$\text{C}_5\text{H}_5\text{N}$	5.9×10^8			PL/Ld-2	$S = A$.	90E324
	$\text{C}_5\text{H}_5\text{N}$				CL/Ac,A'c-17	$S = A' = \text{Rub}$; meas. $k_r/k_r^{A'} = 0.1$.	84F193
	C_6H_6				CL/Ac,A'c-17	$S = A' = \text{Rub}$; meas. $k_r/k_r^{A'} = 0.82$.	84F193
	CCl_4	9×10^8 $3 \times 10^7 (k_r)$			MP/LI-12,27	$S = \text{TPP}$; $A' = \text{Tetr}$; used $k_d = 39 \text{ s}^{-1}$.	85F517
	CCl_4	$2 \times 10^7 (k_r)$			CP/Ac,A'c-14,27	$S = \text{TPP}$, TPBC or MP; $A' = \text{Tetr}$; k_r derived using $k_A = 1 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$, $f_r^{A'} = 1$.	82A421
	CCl_4	10^9			MP/LI-12	$S = \text{PP}$ or Ph a; used $k_d = 36 \text{ s}^{-1}$; also see [78E881, 79A010].	77E617
	CH_2Cl_2				CL/Ac,A'c-17	$S = A' = \text{Rub}$; meas. $k_r/k_r^{A'} = 1.1$.	84F193
	CH_3CN				CL/Ac,A'c-17	$S = A' = \text{Rub}$; meas. $k_r/k_r^{A'} = 1.0$.	84F193
	CH_3COCH_3				CL/Ac,A'c-17	$S = A' = \text{Rub}$; meas. $k_r/k_r^{A'} = 0.41$.	84F193
	THF				CL/Ac,A'c-17	$S = A' = \text{Rub}$; meas. $k_r/k_r^{A'} = 0.015$.	84F193
	dioxane				CL/Ac,A'c-17	$S = A' = \text{Rub}$; meas. $k_r/k_r^{A'} = 0.044$.	84F193
7.2 Bacteriochlorophyll <i>b</i>							
	$(\text{C}_2\text{H}_5)_2\text{O}$	5.8×10^8			PL/Ld-2	$S = A$.	90E324
	$\text{C}_5\text{H}_5\text{N}$	1×10^9			PL/Ld-2	$S = A$.	90E324
	CCl_4	2×10^9 $2 \times 10^8 (k_r)$			MP/LI-12,27	$S = \text{TPP}$; $A' = \text{Tetr}$; used $k_d = 39 \text{ s}^{-1}$.	85F517
7.3 Biline-1,19-dione, 3,8,12,17-tetraethyl-22,24-dihydro-2,7,13,18-tetramethyl- (Etiobiliverdin-IVγ)							
	CHCl_3	1.8×10^9 $4.3 \times 10^6 (k_r)$			CP/Ac-15	$S = \text{RBCE}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$; k_r derived using $\phi_{isc}(S) = 0.36$.	78F701
	MeOH	5.5×10^9 $3.5 \times 10^6 (k_r)$			CP/Ac-15	$S = \text{RB}$; used $k_d = 1.4 \times 10^5 \text{ s}^{-1}$; k_r derived using $\phi_{isc}(S) = 0.76$.	78F701
7.4 Biline-1,19-dione, 3,8,12,17-tetraethyl-10,22,23,24-tetrahydro-2,7,13,18-tetramethyl- (Etiobilirubin-IVγ)							
	CHCl_3	3.0×10^9 $2.3 \times 10^9 (k_r)$			CP/Ac-15	$S = \text{RBCE}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$; k_r derived using $\phi_{isc}(S) = 0.36$.	78F701
	MeOH/ CHCl_3 (90:10)	2.2×10^9 $8.5 \times 10^8 (k_r)$			CP/Ac-15	$S = \text{RB}$; used $k_d = 1.4 \times 10^5 \text{ s}^{-1}$; k_r derived using $\phi_{isc}(S) = 0.76$.	78F701 79F104
7.5 Biline-8,12-dipropanoic acid, 2,17-diethenyl-1,10,19,22,23,24-hexahydro-3,7,13,18-tetramethyl-1,19-dioxo- (Bilirubin, BR)							
	$\text{C}_6\text{H}_5\text{CHOHCH}_3$				CP/Ac-17	$S = N\text{-}[2\text{-}[(2\text{-Bromo-4,6-dinitrophenyl)azo}\text{-}5\text{-}[(2\text{-cyanoethyl)(2-hydroxyethyl)amino]\text{-}4\text{-methoxyphenyl]acetamide}}$; $k_A = 0.5$ rel. to k_A in dibutyl terephthalate.	81F609
	<i>o</i> -Terphenyl/ $(\text{C}_6\text{H}_5)_2\text{O}$ (50:50)				CP/Ac-17	$S = N\text{-}[2\text{-}[(2\text{-Bromo-4,6-dinitrophenyl)azo}\text{-}5\text{-}[(2\text{-cyanoethyl)(2-hydroxyethyl)amino]\text{-}4\text{-methoxyphenyl]acetamide}}$; $k_A = 0.47$ rel. to k_A in dibutyl terephthalate.	81F609
	$(\text{C}_6\text{H}_5)_2/\text{(C}_6\text{H}_5)_2\text{O}$ (25:75)				CP/Ac-17	$S = N\text{-}[2\text{-}[(2\text{-Bromo-4,6-dinitrophenyl)azo}\text{-}5\text{-}[(2\text{-cyanoethyl)(2-hydroxyethyl)amino]\text{-}4\text{-methoxyphenyl]acetamide}}$; $k_A = 0.3$ rel. to k_A in dibutyl terephthalate.	81F609
	C_6H_6	$2 \times 10^8 (k_r)$			CP/A'c-17	$S = A' = \text{Rub}$; used $k_r^{A'} = 4.2 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$; meas. $k_r/k_r^{A'} = 4$.	763011
	CCl_4	$2 \times 10^8 (k_r)$			CP/A'c-17	$S = A' = \text{Rub}$; used $k_r^{A'} = 4.2 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$; meas. $k_r/k_r^{A'} = 4$.	763011
	CCl_4	$2 \times 10^8 (k_r)$ $2 \times 10^9 (k_q)$			CP/A'c-24	$S = A' = \text{Rub}$; used $k_r^{A'} = 4.2 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$; k_r and k_q derived using $(k_r/k_r^{A'}) = 54$, $(k_r/k_r^{A'}) = 4$.	763011

TABLE 7. Rate constants for the interaction of singlet oxygen with porphyrins, phthalocyanines and polypyrroles. — Continued

Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
Bilane-8,12-dipropanoic acid, 2,17-diethenyl-1,10,19,22,23,24-hexahydro-3,7,13,18-tetramethyl-1,19-dioxo- (Bilirubin, BR) — Continued						
CHCl ₃	2.8×10^9 $3.8 \times 10^8 (k_r)$			CP/Ac-15	S = RBCE; k_r derived using $\phi_{isc}(S) = 0.36$.	79F104
CHCl ₃	$2.1 \times 10^8 (k_r)$			CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_r^{A'} = 7 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.30$.	753071
CHCl ₃	1.3×10^9	8×10^{-6}	296	CP/Ac-15	S = MB; used $k_d = 1.0 \times 10^4$ s ⁻¹ .	753071
CHCl ₃	1.5×10^9	6.7×10^{-6}		CP/Ac'-16	S = MB; A' = DPBF; used $k_d = 1.0 \times 10^4$ s ⁻¹ .	753071
CHCl ₃	$4.3 \times 10^8 (k_r)$			CP/Ac,P'a-17	S = MB; A' = TME; used $k_r^{A'} = 4.8 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 9.0$.	753071
CHCl ₃ /MeOH (90:10)	1.3×10^9	3.1×10^{-5}	296	CP/Ac-15	S = RB; used $k_d = 3.9 \times 10^4$ s ⁻¹ .	753071
CHCl ₃ /MeOH (90:10)	$4.3 \times 10^8 (k_r)$			CP/Ac-14	S = RB; k_r derived using $k_A = 2.5 \times 10^9$ L mol ⁻¹ s ⁻¹ and $\phi_{isc}(S) = 0.66$.	753071
CH ₂ Cl ₂	$5.0 \times 10^8 (k_r)$ $5.0 \times 10^9 (k_d)$		308	CR/Ac-19	A' = MNPO ₂ ; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 8.5 \times 10^7$ L mol ⁻¹ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	83A317
ClCF ₂ CCl ₂ F	$1 \times 10^8 (k_r)$			CL/Ad-36	high pressure O ₂ ; k_r derived using $k_{O_2} = 2.7 \times 10^3$ L mol ⁻¹ s ⁻¹ ; Previous value by same workers [747103] low due to overdepletion of A in solution exposed directly to laser beam.	777129
D ₂ O pD = 8.4	1.3×10^9		293	PL/Ad-5	S = RB.	79A111
D ₂ O pD = 8.4	$<4 \times 10^8 (k_r)$		293	CP/Ac,A'c-17	S = RB; A' = DPBF; used $k_r^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.4$.	79A111
D ₂ O (mic) pD = 8.4			293	CP/Ac,A'c-17	S = RB; A' = DPBF; meas. $k_r/k_r^{A'} = 0.44$; 0.1 mol L ⁻¹ SDS.	80N018
D ₂ O (mic) pD = 8.4			293	CL/Ad-37	A' = DPBF; meas. $k_A/k_{A'} = 0.13$; high pressure oxygen (0.195 mol L ⁻¹); 0.1 mol L ⁻¹ SDS.	80N018
D ₂ O/EtOH	3.2×10^9		310	CR/LI-12	used $k_d = 1 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from NDPO ₂ .	89R188
H ₂ O	$1 \times 10^9 (k_r)$ $4.7 \times 10^9 (k_d)$		308	CR/Ac-19	A' = MNPO ₂ ; used $k_d = 3 \times 10^5$ s ⁻¹ , $k_{A'} = 9.0 \times 10^8$ L mol ⁻¹ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ ; in the presence of excess albumin $k_q = 1.9 \times 10^9$ and $k_r = 8.4 \times 10^8$ L mol ⁻¹ s ⁻¹ .	83A317
MeOH	2.1×10^9 $2.8 \times 10^8 (k_r)$			CP/Ac-15	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_{isc}(S) = 0.76$.	79F104
7.6 Bilane-8,12-dipropanoic acid, 2,17-diethenyl-1,10,19,22,23,24-hexahydro-3,7,13,18-tetramethyl-1,19-dioxo-, dimethyl ester (Bilirubin IX dimethyl ester)						
CHCl ₃	1.8×10^9 $6.7 \times 10^8 (k_r)$			CP/Ac-15	S = RBCE; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; k_r derived using $\phi_{isc}(S) = 0.36$.	78F701
MeOH	2.3×10^9 $5.5 \times 10^8 (k_r)$			CP/Ac-15	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_{isc}(S) = 0.76$.	78F701
7.7 Bilane-8,12-dipropanoic acid, 2,17-diethenyl-1,19,22,24-tetrahydro-3,7,13,18-tetramethyl-1,19-dioxo- (Biliverdin)						
CH ₂ Cl ₂	4.6×10^9		308	CR/Ac-19	A' = BRH ₂ ; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 5.0 \times 10^8$ L mol ⁻¹ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	83A317
CHCl ₃	1.7×10^9 $1.9 \times 10^6 (k_r)$			CP/Ac-15	S = RBCE; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; k_r derived using $\phi_{isc}(S) = 0.36$.	78F701
CHCl ₃	$\leq 3 \times 10^6 (k_r)$			CP/Ac,A'c-17	S = A' = Rub; used $k_r^{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = \leq 0.07$.	763011
CHCl ₃	3.3×10^9	5×10^{-6}		CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^4$ s ⁻¹ .	763011
D ₂ O pD = 11.2	9.1×10^8		310	CR/A'c-32	A' = DPBF; used $k_d = 1.5 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from Dopamine/H ₂ O ₂ .	89M038
D ₂ O pD = 11.2	1.1×10^9		310	CR/A'c-32	A' = DPBF; used $k_d = 1.5 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from DOPA/H ₂ O ₂ .	89M038

TABLE 7. Rate constants for the interaction of singlet oxygen with porphyrins, phthalocyanines and polypyroles. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
7.7 Biline-8,12-dipropionic acid, 2,17-diethenyl-1,19,22,24-tetrahydro-3,7,13,18-tetramethyl-1,19-dioxo- (Biliverdin) — Continued							
	D ₂ O pD = 8.4	5.1 × 10 ¹⁰			CL/A'd-16	A' = BR ²⁻ ; used $k_d = 5.0 \times 10^4$ s ⁻¹ ; high pressure O ₂ .	777129
	D ₂ O pD = 8.4	3.0 × 10 ⁸ (k_t)			CL/Ad-35	used $k_d = 5.0 \times 10^4$ s ⁻¹ ; high pressure O ₂ ; k_t derived using $k_A = 5.1 \times 10^{10}$ L mol ⁻¹ s ⁻¹ .	777129
	D ₂ O pD = 13.4	5 × 10 ⁸ (k_t)			CL/Ad-35	used $k_d = 5.0 \times 10^4$ s ⁻¹ ; high pressure O ₂ ; k_t derived using $k_A = 1.5 \times 10^{10}$ L mol ⁻¹ s ⁻¹ .	777129
	D ₂ O pD = 13.4	1.5 × 10 ¹⁰			CL/A'd-16	A' = BR ²⁻ ; used $k_d = 5.0 \times 10^4$ s ⁻¹ ; high pressure O ₂ .	777129
	D ₂ O pD = 11.8	4.0 × 10 ⁸ (k_t)			CL/Ad-35	used $k_d = 5.0 \times 10^4$ s ⁻¹ ; high pressure O ₂ ; k_t derived using $k_A = 6 \times 10^{10}$ L mol ⁻¹ s ⁻¹ .	777129
	D ₂ O pD = 11.8	6 × 10 ¹⁰			CL/A'd-16	A' = BR ²⁻ ; used $k_d = 5.0 \times 10^4$ s ⁻¹ ; high pressure O ₂ .	777129
	D ₂ O/EtOH	2.3 × 10 ⁹		310	CR/LI-12	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	89R188
	H ₂ O pH = 10.5	5 × 10 ⁹			CR/LI-12	used $k_d = 5 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NaOCl/H ₂ O ₂ .	87M341
	MeOH	8.4 × 10 ⁹ 2.4 × 10 ⁶ (k_t)			CP/Ac-15	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; k_t derived using $\phi_{isc}(S) = 0.76$.	78F701
	MeOH/H ₂ O (95:5)	4 × 10 ⁹		310	CR/LI-12	used $k_d = 1.8 \times 10^5$ s ⁻¹ ; soln. cont. 0.05 mol L ⁻¹ MeONa and 5 × 10 ⁻⁴ mol L ⁻¹ CoCl ₂ ; ${}^1\text{O}_2^*$ from autoxidation of oxytetracycline.	92M228
7.8 Biline-8,12-dipropionic acid, 2,17-diethenyl-1,19,22,24-tetrahydro-3,7,13,18-tetramethyl-1,19-dioxo-, dimethyl ester (Biliverdin dimethyl ester)							
	CHCl ₃	1.4 × 10 ⁹ 1.2 × 10 ⁶ (k_t)			CP/Ac-15	S = RBCE; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; k_t derived using $\phi_{isc}(S) = 0.36$.	78F701
	ClCF ₂ CCl ₂ F	6 × 10 ⁵ (k_t) 8 × 10 ⁸ (k_q)			CL/A'd-35	A' = BRH ₂ ; high pressure O ₂ ; k derived using $k_O_2 = 2.7 \times 10^3$ L mol ⁻¹ s ⁻¹ .	777129
	ClCF ₂ CCl ₂ F	9 × 10 ⁸			CL/A'd-16	A' = BRH ₂ ; high pressure O ₂ ; k derived using $k_O_2 = 2.7 \times 10^3$ L mol ⁻¹ s ⁻¹ .	777129
	MeOH	3.3 × 10 ⁹ 1.6 × 10 ⁶ (k_t)			CP/Ac-15	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; k_t derived using $\phi_{isc}(S) = 0.76$.	78F701
7.9 Biline-8,12-dipropionic acid, 2,17-diethyl-1,10,19,22,23,24-hexahydro-3,7,13,18-tetramethyl-1,19-dioxo- (Mesobilirubin)							
	CHCl ₃	2.8 × 10 ⁹ 5.9 × 10 ⁸ (k_t)			CP/Ac-15	S = RBCE; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; k_t derived using $\phi_{isc}(S) = 0.36$.	78F701
	MeOH	2.5 × 10 ⁹ 7.9 × 10 ⁸ (k_t)			CP/Ac-15	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; Soln. contg. 2% NH ₄ OH; k_t derived using $\phi_{isc}(S) = 0.76$.	78F701
7.10 Bilirubin ditaurate							
	D ₂ O/EtOH	1.2 × 10 ⁹		310	CR/LI-12	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	89R188
7.11 Chlorophyll <i>a</i>							
	(C ₂ H ₅) ₂ O	1 × 10 ⁸			PL/Ld-2	S = A.	90E324
	C ₅ D ₅ N	9 × 10 ⁷			PL/Ld-2	S = A.	90E324
	C ₅ H ₅ N	5 × 10 ⁷	0.02	293	CP/Oc-15	S = A; used $k_d = 1 \times 10^5$ s ⁻¹ .	75F669
	C ₆ H ₆	2.0 × 10 ⁸			PL/Ld-2	S = TPP.	93R231
	C ₆ H ₆	7.3 × 10 ⁸			CP/Oc-18	S = A; A' = 2M2P; used $k_d = 4 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.049$ mol L ⁻¹ .	88R136
	C ₆ H ₆ /EtOH (67:33)	1.2 × 10 ⁷			CP/Ac-19	S = RB; A' = Car; uscd $k_d = 1 \times 10^5$ s ⁻¹ .	78F404
	CCl ₄	2 × 10 ⁶ (k_t)			CP/Ac,A'c-14,27	S = TPP, TPBC or MP; A' = Tetr; k_t derived using $k_A = 7 \times 10^8$ L mol ⁻¹ s ⁻¹ , $f_t^{A'} = 1$.	82A421
	CCl ₄	4 × 10 ⁶ (k_t)			CP/Ac,A'c-14,27	S = A; A' = Tetr; k_t derived using $k_A = 7 \times 10^8$ L mol ⁻¹ s ⁻¹ , $f_t^{A'} = 1$, $\phi_\Delta = 0.55$.	79A010

TABLE 7. Rate constants for the interaction of singlet oxygen with porphyrins, phthalocyanines and polypyroles. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
7.11 Chlorophyll <i>a</i> — Continued							
	CCl ₄	7×10^8			MP/LI-12	S = PP, TPP, Ph a or BChl a; used $k_d = 36$ s ⁻¹ ; also see [78E881, 78E892, 79A010].	77E617
	EtOH	2.0×10^7	5.0×10^{-3}		CP/Ac-15	S = RB; used $k_d = 1 \times 10^5$ s ⁻¹ .	78F404
7.12 Chlorophyll <i>b</i>							
	C ₆ H ₆	4.2×10^8			CP/Oc-18	S = A; A' = 2M2P; used $k_d = 4 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.049$ mol L ⁻¹ .	88R136
	CCl ₄	6×10^5 (k_r)			CP/Ac,A'c-14,27	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 3 \times 10^8$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421
7.13 Cyanohemoglobin							
	H ₂ O pH = 7.0	1.2×10^9			CP/Oc-19	S = MB; Q = N ₃ ⁻ ; used $k_Q = 5 \times 10^8$ L mol ⁻¹ s ⁻¹ ; $k = 1.1 \times 10^9$ L mol ⁻¹ s ⁻¹ for subunit α and 2.0×10^9 L mol ⁻¹ s ⁻¹ for subunit β .	83A395
7.14 Hematoporphyrin derivative							
	MeOH	5.1×10^8			CP/Oc-16	S = A; A' = FFA; used $k_d = 9.6 \times 10^4$ s ⁻¹ .	88R142
7.15 Hematoporphyrin-human serum albumin complex							
	D ₂ O pH = 7.4	7×10^8			PL/Ld-2	S = A.	87A064
7.16 Methyl acetal of oxidized octaethylpurpurin ethyl ester							
	MeOH	1.2×10^8			PL/Ld-2	S = A.	90E491
7.17 Naphthalocyanine, bis(tribenzylsiloxy)silicon							
	CH ₂ Cl ₂	1.5×10^8			PL/Ld-2	S = A.	93E520
7.18 Naphthalocyanine, bis(trihexyloxsiloxy)silicon							
	C ₆ H ₆	1.1×10^{10}			PL/Ld-2	S = A.	88E657
	CHCl ₃	3.9×10^8			PL/Ld-2	S = TPP.	90E731 92R076
7.19 Naphthalocyanine, bis(trihexylsiloxy)tin							
	CHCl ₃	5.3×10^9			PL/Ld-2	S = TPP.	90E731 92R076
7.20 Naphthalocyanine, bis(triisobutylsiloxy)silicon							
	CH ₂ Cl ₂	1.5×10^8			PL/Ld-2	S = A.	93E520
7.21 Naphthalocyanine, 1,6,10,15,19,24,28,33-octabutoxy-, palladium(II)							
	C ₆ H ₆	8.8×10^9		295	PL/Ld-2	S = PPDME; energy transfer.	93A380
7.22 Naphthalocyanine, 2,11,20,29-tetrakis(1,1-dimethylethyl)-							
	CHCl ₃	1.0×10^{10}			PL/Ld-2	S = TPP.	90E731 92R076
7.23 Naphthalocyanine, 1,10,19,28-tetraphenyl-, hydroxyaluminum(III)							
	CHCl ₃	2.0×10^9			PL/Ld-2	S = TPP.	90E731 92R076
7.24 Naphthalocyanine, trihexylsiloxyaluminum							
	CHCl ₃	3.3×10^9			PL/Ld-2	S = TPP.	90E731 92R076
7.25 Naphthalocyanine, trihexylsiloxygallium							
	CHCl ₃	1.1×10^{10}			PL/Ld-2	S = TPP.	90E731 92R076

TABLE 7. Rate constants for the interaction of singlet oxygen with porphyrins, phthalocyanines and polypyrroles. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
7.26	Pheophytin <i>a</i>						
	C ₆ H ₆	7.4 × 10 ⁷			CP/Oc-18	S = A; A' = 2M2P; used $k_d = 4 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.049$ mol L ⁻¹ .	88R136
	CCl ₄	1.4 × 10 ⁴ (k_r)		14,27	CP/Ac,A'c-	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421
	CCl ₄	1 × 10 ⁴ (k_r)		14,27	CP/Ac,A'c-	S = A; A' = Tetr; k_r derived using $k_A = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	79A010
	CCl ₄	2 × 10 ⁷			MP/LI-12	S = PP or BChl a; used $k_d = 36$ s ⁻¹ ; also see [78E881, 79A010].	77E617
7.27	Pheophytin <i>b</i>						
	C ₆ H ₆	3.0 × 10 ⁷			CP/Oc-18	S = A; A' = 2M2P; used $k_d = 4 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.049$ mol L ⁻¹ .	88R136
	CCl ₄	10 ³ (k_r)		14,27	CP/Ac,A'c-	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 5 \times 10^6$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421
7.28	20-Phorbinecarboxylic acid, 3,4-didehydro-3,4,8,9,13,14,18,19-octaethyl-18,19-dihydro-, ethyl ester (NT2H2)						
	MeOH	1.5 × 10 ⁸			PL/Ld-2	S = A.	90E491
7.29	20-Phorbinecarboxylic acid, 3,4-didehydro-3,4,8,9,13,14,18,19-octaethyl-18,19-dihydro-, ethyl ester, dichlorotin(IV)						
	MeOH	1 × 10 ⁷			PL/Ld-2	S = A.	90E491
7.30	20-Phorbinecarboxylic acid, 3,4,20,21-tetrahydro-3,4,8,9,13,14,18,19-octaethyl-18,19-dihydro-, ethyl ester (NT2)						
	C ₆ H ₆	~6 × 10 ⁷			PL/Ld-2	S = A.	88R200
	MeOH	5 × 10 ⁷			PL/Ld-2	S = A.	90E491
7.31	20-Phorbinecarboxylic acid, 3,4,20,21-tetrahydro-4,9,14,19-tetraethyl-18,19-dihydro-3,8,13,18-tetramethyl-, ethyl ester, zinc(II) (ZnET2)						
	C ₆ H ₆	5.4 × 10 ⁸			PL/Ld-2	S = A.	88R200
7.32	3-Phorbinepropanoic acid, 9-acetyl-14-ethyl-13,14-dihydro-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, 3,7,11,15-tetramethyl-2-hexadecenyl ester (Bacteriopheophytin <i>a</i>)						
	(C ₂ H ₅) ₂ O	3 × 10 ⁷			PL/Ld-2	S = A.	90E324
	CCl ₄	3 × 10 ⁴ (k_r)		14,27	CP/Ac,A'c-	S = TPP; A' = Tetr; k_r derived using $k_A = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	85F517
	CCl ₄	1.2 × 10 ⁷			MP/LI-12	S = TPP; used $k_d = 39$ s ⁻¹ .	85F517
	CCl ₄	10 ⁷			MP/LI-12	S = PP, Ph a or BChl a; used $k_d = 36$ s ⁻¹ ; also see [78E881, 79A010].	77E617
7.33	3-Phorbinepropanoic acid, 9-acetyl-14-ethylidene-13,14-dihydro-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, 3,7,11,15-tetramethyl-2-hexadecenyl ester (Bacteriopheophytin <i>b</i>)						
	(C ₂ H ₅) ₂ O	1.9 × 10 ⁸			PL/Ld-2	S = A.	90E324
	C ₅ H ₅ N	2.8 × 10 ⁸			PL/Ld-2	S = A.	90E324
	CCl ₄	3 × 10 ⁶ (k_r)		14,27	CP/Ac,A'c-	S = TPP; A' = Tetr; k_r derived using $k_A = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	85F517
	CCl ₄	2 × 10 ⁸			MP/LI-12,27	S = TPP; used $k_d = 39$ s ⁻¹ .	85F517
7.34	3-Phorbinepropanoic acid, 3,4-didehydro-4,9-diethenyl-14-ethyl-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, magnesium(II), 3,7,11,15-tetramethyl-2-hexadecenyl ester (4-Vinylprotochlorophyll)						
	CCl ₄	≤3 × 10 ⁵ (k_r)		14,27	CP/Ac,A'c-	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = \le 1 \times 10^8$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421
7.35	3-Phorbinepropanoic acid, 3,4-didehydro-4,9-diethenyl-14-ethyl-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, 3,7,11,15-tetramethyl-2-hexadecenyl ester (4-Vinylprotopheophytin)						
	CCl ₄	-2 × 10 ⁵ (k_r)				S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421

TABLE 7. Rate constants for the interaction of singlet oxygen with porphyrins, phthalocyanines and polypyroles. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
7.36 3-Phorbinepropanoic acid, 3,4-didehydro-9-ethenyl-14-ethyl-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, 2-amino-2-methoxycarbonyl)ethyl ester, magnesium(II),							
	C ₆ H ₆	2.4×10^8			PL/Ld-2	S = TPP.	93R231
	D ₂ O	1.0×10^8			PL/Ld-2	S = Ru(bpy) ₃ ²⁺ .	93R231
7.37 3-Phorbinepropanoic acid, 3,4-didehydro-9-ethenyl-14-ethyl-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, magnesium(II) (Protoporphobilinogen)							
	C ₆ H ₆	2.0×10^8			PL/Ld-2	S = TPP.	93R231
	D ₂ O	1.6×10^8			PL/Ld-2	S = Ru(bpy) ₃ ²⁺ .	93R231
7.38 3-Phorbinepropanoic acid, 3,4-didehydro-9-ethenyl-14-ethyl-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, 3,7,11,15-tetramethyl-2-hexadecenyl ester (Protopheophytin)							
	CCl ₄	$\sim 10^8 (k_r)$		CP/Ac,A'c-14,27	S = A; A' = Tetr; k _r derived using $k_A = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	79A010	
	CCl ₄	2×10^8		MP/LI-12	S = PP, Ph a or BChl a; used $k_d = 36$ s ⁻¹ ; also see [78E881, 79A010].	77E617	
7.39 3-Phorbinepropanoic acid, 3,4-didehydro-9-ethenyl-14-ethyl-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, magnesium(II), 3,7,11,15-tetramethyl-2-hexadecenyl ester (Protopchlorophyll)							
	CCl ₄	$\leq 10^6 (k_r)$		CP/Ac,A'c-14,27	S = A; A' = Tetr; k _r derived using $k_A = \leq 1 \times 10^8$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	79A010	
	CCl ₄	$\leq 10^8$		MP/LI-12	S = PP, Ph a or BChl a; used $k_d = 36$ s ⁻¹ ; also see [78E881, 79A010].	77E617	
7.40 Phthalocyanine, bis(tribenzylsiloxy)silicon							
	CH ₂ Cl ₂	1.4×10^7		PL/Ld-2	S = A.	93E520	
7.41 Phthalocyanine, bis(tripropylsiloxy)silicon							
	CH ₂ Cl ₂	1.6×10^7		PL/Ld-2	S = A.	93E520	
7.42 Phthalocyanine, 1,4,8,11,15,18,22,25-octabutoxy-							
	CHCl ₃	5.9×10^9		PL/Ld-2	S = TPP.	90E731 92R076	
7.43 Phthalocyanine, sulfo-, chloroaluminum(III) [AlCl(spc)]							
	D ₂ O pH = 7	$< 10^8$		PL/Ld-2	S = A; No measurable effect; soln. contg. 2×10^{-3} mol L ⁻¹ phosphate buffer and 1% NaCl wt/wt.	90A022	
7.44 Phthalocyanine, sulfo-, zinc(II)							
	D ₂ O pH = 7	$< 10^8$		PL/Ld-2	S = A; No measurable effect; soln. contg. 2×10^{-3} mol L ⁻¹ phosphate buffer and 1% NaCl wt/wt.	90A022	
7.45 Phthalocyanine, tetracarboxy-, copper(II)							
	DMSO	$\geq 1.4 \times 10^9$		CP/Ac-16	S = A; A' = DPRF; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	89F260	
7.46 Phthalocyanine, 2,9,16,23-tetra(1,1-dimethylethyl)-							
	C ₆ F ₆	2×10^8		PL/Ld-2	S = Fullerene-C ₇₀ .	93E301	
	CHCl ₃	2.9×10^8		PL/Ld-2	S = TPP.	90E731 92R076	
7.47 Porphine, (acetato)-5,10,15,20-tetraphenyl-, iron(III)							
	CCl ₄	$< 10^4 (k_r)$		CP/Ac,A'c-14,27	S = TPP, TPBC or MP; A' = Tetr; k _r derived using $k_A = 1.5 \times 10^9$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421	
	CCl ₄	1.5×10^9		MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	78E892 79A010 80E548	

TABLE 7. Rate constants for the interaction of singlet oxygen with porphyrins, phthalocyanines and polypyrroles. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
7.48	Porphine, (acetato)-5,10,15,20-tetraphenyl-, manganese(III)						
	CCl ₄	2.5×10^8			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	80E548
7.49	Porphine, (chloro)-5,10,15,20-tetraphenyl-, iron(III)						
	CCl ₄	1×10^9			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	80E548
7.50	Porphine, 2,3-dihydro-5,10,15,20-tetraphenyl- (Tetraphenylchlorin)						
	CCl ₄	$1.2 \times 10^2 (k_t)$		14,27	CP/Ac,A'c-	S = TPP, TPBC or MP; A' = Tetr; k_t derived using $k_A = 2 \times 10^6$ L mol ⁻¹ s ⁻¹ , $f_t^{A'} = 1$.	82A421
	CCl ₄	2×10^6			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	79F824
7.51	Porphine, 2,3-dihydro-5,10,15,20-tetraphenyl-, cadmium(II)						
	CCl ₄	7×10^9			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	79F824
7.52	Porphine, 2,3-dihydro-5,10,15,20-tetraphenyl-, copper(II)						
	CCl ₄	2×10^7			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	79F824
7.53	Porphine, 2,3-dihydro-5,10,15,20-tetraphenyl-, zinc(II)						
	C ₅ H ₅ N	6×10^8 $<1 \times 10^5 (k_t)$		16,17	CP/A'c-16,17	S = TPBC; A' = Tetr; used $k_d = 6.3 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^6$ L mol ⁻¹ s ⁻¹ .	86F610
	C ₆ H ₆	6×10^9 $2.5 \times 10^7 (k_t)$			CP/A'c-16,17	S = TPBC; A' = Tetr; used $k_d = 3.3 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^6$ L mol ⁻¹ s ⁻¹ .	86F610
	CCl ₄	4×10^9 $2 \times 10^8 (k_t)$			CP/A'c-16,17	S = TPBC; A' = Tetr; used $k_d = 36$ s ⁻¹ , $k_{A'} = 7 \times 10^6$ L mol ⁻¹ s ⁻¹ .	86F610
	CCl ₄	$2 \times 10^8 (k_t)$		14,27	CP/Ac,A'c-14,27	S = TPP, TPBC or MP; A' = Tetr; k_t derived using $k_A = 4.0 \times 10^9$ L mol ⁻¹ s ⁻¹ , $f_t^{A'} = 1$.	82A421
	CCl ₄	$2 \times 10^8 (k_t)$		14,27	CP/Ac,A'c-14,27	S = PP, Ph a or BChl a; A' = Tetr; used $k_d = 36$ s ⁻¹ , $k_{A'} = 5 \times 10^6$ L mol ⁻¹ s ⁻¹ ; k_t derived using $k_A = 4.0 \times 10^9$ L mol ⁻¹ s ⁻¹ , $f_t^{A'} = 1$.	79A010
	CCl ₄	4×10^9			CP/LI-12	S = PP, TPP, Ph a or BChl a; used $k_d = 36$ s ⁻¹ .	78E892 79A010
	CH ₃ COCH ₃	1×10^9 $8 \times 10^5 (k_t)$			CP/A'c-16,17	S = TPBC; A' = Tetr; used $k_d = 2.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^6$ L mol ⁻¹ s ⁻¹ .	86F610
7.54	Porphine, 7,8,17,18-tetrahydro-5,10,15,20-tetraphenyl-, (E) (Tetraphenylbacteriochlorin- <i>trans</i> , TPBC)						
	CCl ₄	1×10^8			MP/LI-12	S = PP, TPP, Ph a or BChl a; used $k_d = 36$ s ⁻¹ ; also see [78E892, 79A010].	78E881
	CCl ₄	$1.5 \times 10^5 (k_t)$		14,27	CP/Ac,A'c-14,27	S = TPP, TPBC or MP; A' = Tetr; k_t derived using $k_A = 1 \times 10^8$ L mol ⁻¹ s ⁻¹ , $f_t^{A'} = 1$.	82A421
	CCl ₄	$2 \times 10^5 (k_t)$		14,27	CP/Ac,A'c-14,27	S = PP, Ph a or BChl a; A' = Tetr; k_t derived using $k_A = 1 \times 10^8$ L mol ⁻¹ s ⁻¹ , $f_t^{A'} = 1$.	78E892 79A010
7.55	Porphine, 5,10,15,20-tetrakis(4-iodophenyl)-, copper(II)						
	CCl ₄	2×10^7			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	79F824
7.56	Porphine, 5,10,15,20-tetrakis(4-methoxyphenyl)-, copper(II)						
	CCl ₄	2×10^7			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	79F824
7.57	Porphine, 5,10,15,20-tetrakis(4-sulfonatophenyl)- (H ₂ TPPS ⁴⁻)						
	D ₂ O pH = 7	$<10^8$			PL/Ld-2	S = A; No measurable effect; soln. contg. 2×10^{-3} mol L ⁻¹ phosphate buffer and 1% NaCl wt/wt.	90A022
7.58	Porphine, 5,10,15,20-tetrakis(4-sulfonatophenyl)-, dichlorotin(IV)						
	D ₂ O pH = 7	$<10^8$			PL/Ld-2	S = A; No measurable effect; soln. contg. 2×10^{-3} mol L ⁻¹ phosphate buffer and 1% NaCl wt/wt.	90A022

TABLE 7. Rate constants for the interaction of singlet oxygen with porphyrins, phthalocyanines and polypyroles. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
7.59 Porphine, 5,10,15,20-tetraphenyl-, (TPP)							
	C ₆ D ₆	6×10^7			PL/Ld-2	S = MPDME.	83E235
	C ₆ H ₆	4.4×10^7			CP/Oc-18	S = A; A' = 2M2P; used $k_d = 4 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.049$ mol L ⁻¹ .	88R136
	CCl ₄	9×10^6		286	CP/Oc-29	S = A; A' = 2,3-Dihydrofuran; used $k_{A'} = 1.5 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 6.0$.	91F163
	CCl ₄	2.9×10^7		286	CP/Oc-29	S = A; A' = TME; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $\beta_{A'} = 1.0 \times 10^{-4}$ mol L ⁻¹ .	84F065
	CCl ₄	2.2×10^7		286	CP/Oc-29	S = A; A' = DMHD; meas. $k_A/k_{A'} = 220$, used $k_{A'} = 1.0 \times 10^5$ L mol ⁻¹ s ⁻¹ .	84F335
	CCl ₄	30 (k_r)			CP/Ac,A'c-14,27	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 9 \times 10^5$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421
	CCl ₄	10^6			MP/LI-12	S = PP, TPP, Ph a or BChl a; used $k_d = 36$ s ⁻¹ ; also see [78E892, 79A010].	78E881
	CCl ₄	3×10^2 (k_r)			CP/Ac,A'c-14,27	S = A; A' = Tetr; used $k_d = 36$ s ⁻¹ ; k_r derived using $k_A = 1 \times 10^6$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	78E892 79A010
	CH ₂ Cl ₂	1.4×10^7		286	CP/Oc-29	S = A; A' = 2,3-Dihydrofuran; used $k_{A'} = 3.2 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 4.5$.	91F163
	CH ₂ Cl ₂	1.7×10^7		286	CP/Oc-29	S = A; A' = TME; used $k_d = 9.5 \times 10^3$ s ⁻¹ , $\beta_{A'} = 4.0 \times 10^{-4}$ mol L ⁻¹ .	84F065
	CH ₂ Cl ₂	2.4×10^7		286	CP/Oc-29	S = A; A' = DMHD; meas. $k_A/k_{A'} = 8$; meas. $k_A/k_{A'} = 28$, used $k_{A'} = 8.6 \times 10^5$ L mol ⁻¹ s ⁻¹ .	84F335
	CHCl ₃	2.1×10^7		286	CP/Oc-29	S = A; A' = 2,3-Dihydrofuran; used $k_A = 1.0 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 20$.	91F163
	CHCl ₃	3.0×10^7		286	CP/Oc-29	S = A; A' = TME; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $\beta_{A'} = 3.6 \times 10^{-4}$ mol L ⁻¹ .	84F065
	CHCl ₃	1.9×10^7		286	CP/Oc-29	S = A; A' = DMHD; meas. $k_A/k_{A'} = 40$, used $k_{A'} = 4.5 \times 10^5$ L mol ⁻¹ s ⁻¹ .	84F335
7.60 Porphine, 5,10,15,20-tetraphenyl-, cadmium(II)							
	CCl ₄	3×10^6 (k_r)			CP/Ac,A'c-14,27	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 6.0 \times 10^8$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421
	CCl ₄	6×10^8			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	79F824
7.61 Porphine, 5,10,15,20-tetraphenyl-, chloroaluminum(III)							
	CCl ₄	1×10^7			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	80E548
7.62 Porphine, 5,10,15,20-tetraphenyl-, cobalt(II)							
	CCl ₄	< 10^4 (k_r)			CP/Ac,A'c-14,27	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 2.0 \times 10^9$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421
	CCl ₄	2×10^9			MP/LI-12	S = PP, Ph a or BChl a; used $k_d = 36$ s ⁻¹ .	79A010
	CCl ₄	2×10^9			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	78E892
	CH ₃ COCH ₃ / H ₂ O (95:5)	3×10^8			PL/Ld-2	S = PdMP.	82A412
7.63 Porphine, 5,10,15,20-tetraphenyl-, copper(II)							
	CCl ₄	<40 (k_r)			CP/Ac,A'c-14,27	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 4 \times 10^6$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421
	CCl ₄	5×10^6			MP/LI-12	S = PP, TPP, Ph a or BChl a; used $k_d = 36$ s ⁻¹ .	78E892 79A010
7.64 Porphine, 5,10,15,20-tetraphenyl-, dichlororotin(IV)							
	CCl ₄	2×10^6			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	80E548

TABLE 7. Rate constants for the interaction of singlet oxygen with porphyrins, phthalocyanines and polypyrroles. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
7.65	Porphine, 5,10,15,20-tetraphenyl-, magnesium(II)						
	C ₆ H ₆	5.0 × 10 ⁸			CP/Oc-18	S = A; A' = 2M2P; used $k_d = 4 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.049$ mol L ⁻¹ .	88R136
	C ₆ H ₆	1.2 × 10 ⁹			MP/Ac-33	S = A; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ ; A' = α -Tocopherol.	80F020
	CCl ₄	6 × 10 ⁶ (k_r)		14,27	CP/Ac,A'c-	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 8.0 \times 10^8$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421
	CCl ₄	8 × 10 ⁸			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	79F824
	CH ₂ Cl ₂	1 × 10 ⁸			MP/Ac-14	S = A; used $k_d = 9.5 \times 10^3$ s ⁻¹ .	80F020
	CH ₂ Cl ₂	2.3 × 10 ⁹			MP/Ac-33	S = A; Q = Car; used $k_d = 9.5 \times 10^3$ s ⁻¹ , $k_Q = 8.5 \times 10^9$ L mol ⁻¹ s ⁻¹ .	80F020
	CH ₂ Cl ₂	4 × 10 ⁷ (k_r)			MP/Ac-17	S = A; A' = TME; used $k_r^{A'} = 4.8 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.8$.	80F020
7.66	Porphine, 5,10,15,20-tetraphenyl-, manganese(III)						
	CCl ₄	2.5 × 10 ⁹			PL/Ld-2	S = PdMP.	82A412
7.67	Porphine, 5,10,15,20-tetraphenyl-, nickel(II)						
	CCl ₄	<10 ⁴ (k_r)		14,27	CP/Ac,A'c-	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 1.5 \times 10^9$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421
	CCl ₄	2 × 10 ⁹			MP/LI-12	S = PP, TPP, Ph a or BChl a; used $k_d = 36$ s ⁻¹ .	78E892 79A010
7.68	Porphine, 5,10,15,20-tetraphenyl-, zinc(II)						
	C ₆ H ₆	1.5 × 10 ⁸			CP/Oc-18	S = A; A' = 2M2P; used $k_d = 4 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.049$ mol L ⁻¹ .	88R136
	CCl ₄	8 × 10 ³ (k_r)		14,27	CP/Ac,A'c-	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421
	CCl ₄	4 × 10 ⁷			MP/LI-12	S = PP, TPP, Ph a or BChl a; used $k_d = 36$ s ⁻¹ ; see also [78E892, 79A010].	78E881
	CCl ₄	1 × 10 ⁴ (k_r)		14,27	CP/Ac,A'c-	S = PP, Ph a or BChl a; A' = Tetr; used $k_d = 36$ s ⁻¹ ; k_r derived using $k_A = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	79A010
7.69	Porphine, 5,10,15,20-tetraphenyl[μ-oxobis-, iron(II)]						
	CCl ₄	4 × 10 ⁷			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	80E548
7.70	Porphine, 5,10,15,20-tetraphenyl(thiocyanato-S)-, iron(III)						
	CCl ₄	1 × 10 ⁹			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	80E548
7.71	Porphine, zinc(II)						
	CCl ₄	3 × 10 ⁷			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	79F824
7.72	Porphine-2,18-dipropanoic acid, 7,12-bis(1-hydroxyethyl)-3,8,13,17-tetramethyl- (Hematoporphyrin)						
	MeOH	7.7 × 10 ⁸			CP/Oc-16	S = A; A' = FFA; used $k_d = 9.6 \times 10^4$ s ⁻¹ .	88R142
	MeOH/ H ₂ O (90:10)	8.0 × 10 ⁸			CP/Oc-16	S = A; A' = FFA; used $k_d = 1.7 \times 10^5$ s ⁻¹ .	88R142
7.73	Porphine-2,18-dipropanoic acid, 7,12-diethenyl-3,8,13,17-tetramethyl- (Protoporphyrin, PP)						
	DMAA	1.1 × 10 ⁷			PL/Ld-2	S = A.	92E274
7.74	Porphine-2,18-dipropanoic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, acetatoferate(III), dimethyl ester [FeMPDME(OAc)]						
	C ₅ H ₅ N	1 × 10 ⁹			CP/LI-12	S = TPP; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	80E548
	CCl ₄	1 × 10 ⁹			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	80E548
7.75	Porphine-2,18-dipropanoic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, cadmium(II) (CdPP)						
	CCl ₄	4 × 10 ⁸			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	79F824

TABLE 7. Rate constants for the interaction of singlet oxygen with porphyrins, phthalocyanines and polypyroles. — Continued

#	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
7.78 Porphine-2,18-dipropionic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, cobalt(II) (CoPP)							
	CCl ₄	3.0×10^9			PL/Ld-2	S = PdMP.	82A412
	CCl ₄	3×10^9			MP/LI-12	S = TPP; used $k_d = 36 \text{ s}^{-1}$.	78E892
7.79 Porphine-2,18-dipropionic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, copper(II) (CuPP)							
	CCl ₄	5×10^6			MP/LI-12	S = TPP; used $k_d = 36 \text{ s}^{-1}$.	79F824
7.80 Porphine-2,18-dipropionic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, dimethyl ester (PPDME)							
	C ₆ H ₅ Cl	1.0×10^9			PL/Ld-2	S = TPP.	88A507
	C ₆ H ₆	9.1×10^7			CP/Oc-18	S = A; A' = 2M2P; used $k_d = 4 \times 10^4 \text{ s}^{-1}$, $\beta_{A'} = 0.049 \text{ mol L}^{-1}$.	88R136
	CCl ₄	$5 \times 10^3 (k_r)$		14,27	CP/Ac,A'-c-	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 5 \times 10^5 \text{ L mol}^{-1} \text{ s}^{-1}$, $f_r^{A'} = 1$.	82A421
	CCl ₄	5×10^5			MP/LI-12	S = TPP; used $k_d = 36 \text{ s}^{-1}$.	79F824
	CH ₂ Cl ₂	3.0×10^6			PL/Ld-2	S = A.	92E274
	CH ₂ Cl ₂	8.6×10^5			MP/Pa-25	S = A; A' = Cholesterol; used $k_d = 1.4 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 6.7 \times 10^4 \text{ L mol}^{-1} \text{ s}^{-1}$; P = Dimethyl 7-(formylimethylene)-7,8-dihydro-8-hydroxy-3,8,13,17-tetramethyl-12-vinyl-2,18-porphinedipropionate; and Dimethyl 7-ethenyl-13-hydroxy-3,8,13,17-tetramethyl-12-(oxoethylidene)porphine-2,18-dipropionate.	82F018
	CH ₂ Cl ₂	8.3×10^5			MP/Pa-25	S = A; A' = DABCO; used $k_d = 1.4 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 3.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$, see above.	82F018
	CH ₃ CN	$1.5 \times 10^7 (k_r)$			CP/Ac,A'-c-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.1 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$.	88A507
	CH ₃ CN	4.7×10^8			PL/Ld-2	S = TPP.	88A507
	CH ₃ CO ₂ C ₂ H ₅	6.7×10^6			PL/Ld-2	S = A.	92E274
	CH ₃ COCH ₃	6.3×10^6			PL/Ld-2	S = A.	92E274
	CHCl ₃	3.9×10^6			PL/Ld-2	S = A.	92E274
	DMAA	1.2×10^7			PL/Ld-2	S = A.	92E274
	DMF	1.2×10^7			PL/Ld-2	S = A.	92E274
	2-ProH	3.9×10^8			PL/Ld-2	S = TPP.	88A507
7.81 Porphine-2,18-dipropionic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, dimethyl ester, nickel(II) (NiPPDME)							
	CCl ₄	2×10^9			MP/LI-12	S = TPP; used $k_d = 36 \text{ s}^{-1}$.	78E892
7.82 Porphine-2,18-dipropionic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, nickel(II) (NiPP)							
	CCl ₄	2.0×10^9			PL/Ld-2	S = PdMP.	82A412
	CCl ₄	10^9			MP/LI-12	S = TPP; used $k_d = 36 \text{ s}^{-1}$.	79F824
7.83 Porphine-2,18-dipropionic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, zinc(II) (ZnPP)							
	CCl ₄	$2 \times 10^3 (k_r)$		14,27	CP/Ac,A'-c-	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 2 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$, $f_r^{A'} = 1$.	82A421
	CCl ₄	2×10^7			MP/LI-12	S = TPP; used $k_d = 36 \text{ s}^{-1}$.	79F824
7.84 Porphine-2,18-dipropionic acid, 7,12-diethyl-3,8,13,17-tetramethyl-, cobalt(II) (CoMP)							
	CCl ₄	3.0×10^9			PL/Ld-2	S = PdMP.	82A412
	CH ₃ COCH ₃ / H ₂ O (95:5)	1.5×10^9			PL/Ld-2	S = PdMP.	82A412
	CHCl ₃	3.0×10^9			PL/Ld-2	S = MPDEE or PdMPDEE.	81E472
	CHCl ₃	2×10^9			PL/Ld-2	S = PdMP.	82A412

TABLE 7. Rate constants for the interaction of singlet oxygen with porphyrins, phthalocyanines and polypyrroles. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
7.83 Porphine-2,18-dipropanoic acid, 7,12-diethyl-3,8,13,17-tetramethyl-, di[4-(diphenylmethylaminocarbonyl-2-nitrophenylmethyl] ester (MP-Nbb-NO₂)							
	CH ₂ Cl ₂	2.0×10^7			PL/Ld-2	S = Zn(pc)(py) ₂ .	91E134
	CH ₂ Cl ₂	1.9×10^7			PL/Ld-2	S = A.	91E134
7.84 Porphine-2,18-dipropanoic acid, 7,12-diethyl-3,8,13,17-tetramethyl-, diethyl ester, manganese(III) (MnMPDEE)							
	CH ₃ COCH ₃ / H ₂ O (95:5)	1.5×10^9			PL/Ld-2	S = PdMP.	82A412
7.85 Porphine-2,18-dipropanoic acid, 7,12-diethyl-3,8,13,17-tetramethyl-, dimethyl ester (MPDME)							
	CCl ₄	4×10^2 (k_r)			CP/Ac,A'c-14,27	S = TPP, TPBC or MP; A' = Tetr; k_r derived using $k_A = 2 \times 10^6$ L mol ⁻¹ s ⁻¹ , $f_r^{A'} = 1$.	82A421
	CCl ₄	3×10^6			MP/LI-12	S = PP, Ph a or BChl a; used $k_d = 36$ s ⁻¹ .	79A010
	CCl ₄	2×10^6			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	78E892
	CD ₃ CN	2.2×10^7			PL/Ld-2	S = A.	83E235
	CD ₃ COCD ₃	1×10^7			PL/Ld-2	S = A.	83E235
	CH ₂ Cl ₂	1.4×10^7			PL/Ld-2	S = Zn(pc)(py) ₂ .	91E134
	CH ₃ CN	2.5×10^7			PL/Ld-2	S = A.	83E235
	CH ₃ COCH ₃	4×10^7			PL/Ld-2	S = A.	83E235
	DCON(CD ₃) ₂	1.0×10^7			PL/Ld-2	S = A.	92E274
	DMAA- <i>d</i> ₉	1.4×10^7			PL/Ld-2	S = A.	92E274
7.86 Porphine-2,18-dipropanoic acid, 7,12-diethyl-3,8,13,17-tetramethyl-, dimethyl ester, cobalt(II) (CoMPDME)							
	CCl ₄	3.0×10^9		295	PL/Ld-2	S = PdMPDME or MPDME; decay at 1588 nm; decay at 1272 nm gave 3.5×10^9 .	80E558
7.87 Porphine-2,18-dipropanoic acid, 7,12-diethyl-3,8,13,17-tetramethyl-, nickel(II) (NiMP)							
	CCl ₄	2.0×10^9			PL/Ld-2	S = PdMP.	82A412
	CCl ₄	2×10^9			MP/LI-12	S = TPP; used $k_d = 36$ s ⁻¹ .	78E892
	CH ₃ COCH ₃ / H ₂ O (95:5)	1.0×10^9			PL/Ld-2	S = PdMP.	82A412
7.88 Porphine-2,18-dipropanoic acid, 7-[2-(dimethylamino)-2-oxoethyl]-8-ethyl-7,8-dihydro-3,7,12,17-tetramethyl, dimethyl ester, (Z) (CHLII)							
	CHCl ₃	2×10^7			CL/LI-12	S = A; used $k_d = 4.8 \times 10^3$ s ⁻¹ .	90R006
7.89 Porphine-2,18-dipropanoic acid, 7-[2-(dimethylamino)-2-oxoethyl]-8-ethylenedide-7,8-dihydro-3,7,12,17-tetramethyl, dimethyl ester (CHLI)							
	CDCl ₃	2.7×10^8			CL/LI-12	S = A; used $k_d = 1.1 \times 10^2$ s ⁻¹ .	90R006
	CHCl ₃	3.2×10^8			CL/LI-12	S = A; used $k_d = 4.8 \times 10^3$ s ⁻¹ .	90R006
7.90 Porphine-2,18-dipropanoic acid, 8-ethenyl-13-ethyl-3,7,12,17-tetramethyl- (EVD)							
	DMAA	2.3×10^7			PL/Ld-2	S = A.	92E274
	DMAA- <i>d</i> ₉	2.2×10^7			PL/Ld-2	S = A.	92E274
7.91 Porphine-2,18-dipropanoic acid, 8-ethenyl-13-ethyl-3,7,12,17-tetramethyl-, dimethyl ester (EVDDME)							
	DMAA	2.2×10^7			PL/Ld-2	S = A.	92E274
7.92 Porphine-2,18-dipropanoic acid, 7-ethenyl-12-(1-hydroxyethyl)-3,8,13,17-tetramethyl- (HVD)							
	CH ₃ CO ₂ C ₂ H ₅	9.7×10^6			PL/Ld-2	S = A.	92E274
	CH ₃ COCH ₃	9.2×10^6			PL/Ld-2	S = A.	92E274
	DMAA	1.4×10^7			PL/Ld-2	S = A.	92E274
	DMAA- <i>d</i> ₉	1.4×10^7			PL/Ld-2	S = A.	92E274
	DMF	1.3×10^7			PL/Ld-2	S = A.	92E274
7.93 Porphine-2,18-dipropanoic acid, 7-ethenyl-3,8,13,17-tetramethyl- (VD)							
	DMAA	1.7×10^7			PL/Ld-2	S = A.	92E274

TABLE 7. Rate constants for the interaction of singlet oxygen with porphyrins, phthalocyanines and polypyrroles. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
7.94 Porphine-2,18-dipropanoic acid, 3,7,12,17-tetramethyl-, dimethyl ester (DPDME)							
	CD ₃ COCD ₃	6.9×10^6			PL/Ld-2	S = A.	92E274
	CDCl ₃	1.7×10^7			PL/Ld-2	S = A.	92E274
	DMAA- <i>d</i> ₉	1.1×10^7			PL/Ld-2	S = A.	92E274
7.95 Porphine-2,18-dipropanoic acid, 3,7,12,17-tetramethyl-, nickel(II) (NiDP)							
	CCl ₄	2.0×10^9			PL/Ld-2	S = MP.	82A412
7.96 Porphine-2,18-dipropanoic acid, 3,7,12,17-tetramethyl-, nickel(II), bis(piperidine)							
	CCl ₄ /c-C ₅ H ₁₀ NH (80:20)	2.0×10^8			PL/Ld-2	S = MP.	82A412
7.97 5-Porphinopropenoic acid, octaethyl-, ethyl ester (NTW)							
	MeOH	4.6×10^8			PL/Ld-2	S = A.	90E491
7.98 [22]Porphyrin-(2,2.2.2), octaethyl-, (<i>di-trans</i>)							
	C ₆ H ₆	8×10^9			PL/Ld-2	S = A; reversible energy transfer, $k_{\text{reverse}} = 1.9 \times 10^9$ L mol ⁻¹ s ⁻¹ .	92E555
7.99 [26] Porphyrin							
	CHCl ₃	4.2×10^8			CL/LI-12	S = A; used $k_d = 4.8 \times 10^3$ s ⁻¹ .	90E530
7.100 [26]Porphyrin-(2,4.2.4), 2,3,10,11,16,17,24,25-octaethyl-5,6,7,8,19,20,21,22-octadehydro-							
	C ₆ H ₆	1.2×10^{10}			PL/Ld-2	S = ZnTPP.	92E555
7.101 Pyrrol-2-one, 3,4-diethyl-1,5-dihydro-5-[(5-methylpyrrol-2-yl)methylene]-							
	CHCl ₃	1.5×10^9 $8.0 \times 10^8 (k_r)$			CP/Ac-15	S = RBCE; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; k_r derived using $\phi_{\text{isc}}(S) = 0.36$.	79F104
	MeOH	8×10^8 $2.1 \times 10^8 (k_r)$			CP/Ac-15	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_{\text{isc}}(S) = 0.76$.	79F104
7.102 Pyrrol-2-one, 3,4-diethyl-5-[(4-ethyl-3,5-dimethylpyrrol-2-yl)methylene]-1,5-dihydro-							
	CHCl ₃	4.2×10^9 $3.2 \times 10^9 (k_r)$			CP/Ac-15	S = RBCE; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; k_r derived using $\phi_{\text{isc}}(S) = 0.36$.	78F701 79F104
	ClCF ₂ CCl ₂ F	$7.5 \times 10^8 (k_r)$	293	CL/Ad-36		high pressure O ₂ ; used $k_{O_2} = 2.5 \times 10^3$ L mol ⁻¹ s ⁻¹ .	79A113
	ClCF ₂ CCl ₂ F	6.6×10^8	293	PL/Ad-5		high pressure O ₂ .	79A113
	MeOH	2.7×10^9 $1.4 \times 10^9 (k_r)$			CP/Ac-15	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_{\text{isc}}(S) = 0.76$.	78F701 79F104
7.103 Pyrrol-2-one, 5-[(3,5-dimethylpyrrol-2-yl)methylene]-4-ethyl-1,5-dihydro-3-methyl-							
	CHCl ₃	2.4×10^9 $1.6 \times 10^9 (k_r)$			CP/Ac-15	S = RBCE; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; k_r derived using $\phi_{\text{isc}}(S) = 0.36$.	79F104
	MeOH	1.1×10^9 $6 \times 10^8 (k_r)$			CP/Ac-15	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_{\text{isc}}(S) = 0.76$.	79F104
7.104 Pyrrol-2-one, 5-[(4,5-dimethylpyrrol-2-yl)methylene]-4-ethyl-1,5-dihydro-3-methyl-							
	CHCl ₃	2.5×10^9 $1.7 \times 10^9 (k_r)$			CP/Ac-15	S = RBCE; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; k_r derived using $\phi_{\text{isc}}(S) = 0.36$.	79F104
	MeOH	2.3×10^9 $9 \times 10^8 (k_r)$			CP/Ac-15	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_{\text{isc}}(S) = 0.76$.	79F104
7.105 Pyrrol-2-one, 3-ethenyl-5-[(4-ethyl-3,5-dimethylpyrrol-2-yl)methylene]-1,5-dihydro-4-methyl-							
	CHCl ₃	2.2×10^9 $1.9 \times 10^9 (k_r)$			CP/Ac-15	S = RBCE; k_r derived using $\phi_{\text{isc}}(S) = 0.36$.	79F104
	MeOH	3.1×10^9 $1.2 \times 10^9 (k_r)$			CP/Ac-15	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	79F104

TABLE 7. Rate constants for the interaction of singlet oxygen with porphyrins, phthalocyanines and polypyrroles. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
7.106 Pyrrol-2-one, 4-ethenyl-5-[(4-ethyl-3,5-dimethylpyrrol-2-yl)methylene]-1,5-dihydro-3-methyl-							
	ClCF ₂ CCl ₂ F	6.7×10^8 (k_t)		293	CL/Ad-36	high pressure O ₂ ; used $k_{O_2} = 2.5 \times 10^3$ L mol ⁻¹ s ⁻¹ .	79A113
	ClCF ₂ CCl ₂ F	1×10^9		293	PL/Ad-5	high pressure O ₂ .	79A113
7.107 Pyrrol-2-one, 4-ethyl-1,5-dihydro-3-methyl-5-[(5-methylpyrrol-2-yl)methylene]-							
	ClCF ₂ CCl ₂ F	3.4×10^8 (k_t)		293	CL/Ad-36	high pressure O ₂ ; used $k_{O_2} = 2.5 \times 10^3$ L mol ⁻¹ s ⁻¹ .	79A113
7.108 Pyrrol-2-one, 5-[(4-ethyl-3,5-dimethylpyrrol-2-yl)methylene]-1,5-dihydro-							
	CHCl ₃	4.4×10^9			CP/Ac-15	S = RBCE; uscd $k_d = 1.7 \times 10^4$ s ⁻¹ ; k_t derived using $\phi_{isc}(S) = 0.36$.	79F104
		3.0×10^9 (k_t)					
	ClCF ₂ CCl ₂ F	7×10^8		293	PL/Ad-5	high pressure O ₂ .	79A113
	ClCF ₂ CCl ₂ F	7×10^8 (k_t)		293	CL/Ad-36	high pressure O ₂ ; used $k_{O_2} = 2.5 \times 10^3$ L mol ⁻¹ s ⁻¹ .	79A113
	MeOH	2×10^8			CP/Ac-15	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; k_t derived using $\phi_{isc}(S) = 0.76$.	79F104
		1×10^8 (k_t)					

TABLE 8. Rate constants for the interaction of singlet oxygen with aliphatic and alicyclic amines.

No.	Solvent	k ($\text{L mol}^{-1} \text{s}^{-1}$)	$\beta (k_d/k)$ (mol L^{-1})	T (K)	Method	Comment	Ref.
8.1	8-Azabicyclo[3.2.1]octan-3-ol, 8-methyl-, α -(hydroxymethyl)benzeneacetate (Atropine)						
	CHCl ₃	4.3×10^7	2.3×10^{-4}		CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^4 \text{ s}^{-1}$.	84F670
8.2	Benzeneethanol, β -phenyl- α -pyrrolidinyl-						
	C ₆ H ₆	1.4×10^3			CP/Pa-16	S = RB; A' = DPBF; used $k_d = 4 \times 10^4 \text{ s}^{-1}$; P = Benzaldehyde.	89F238
	CD ₃ OD	6.5×10^6			PL/Ld-2	Unpublished data, E. Oliveros, M.T. Maurette and A. Braun.	89F238
8.3	[1]Benzopyrano[6,7,8-ij]quinolizin-11-one, 2,3,6,7-tetrahydro-						
	C ₆ H ₆	8×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 3.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} \text{ not given}$.	87F569
8.4	[1]Benzopyrano[6,7,8-ij]quinolizin-11-one, 2,3,6,7-tetrahydro-9-methyl-				CP/A'c-33	S = A' = Rub; used $k_d = 3.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} \text{ not given}$.	87F569
8.5	Benzinemethanamine (Benzylamine)						
	CH ₃ CN	1.7×10^6			PL/Ld-2	S = RB.	89E324
	EtOH	$<1 \times 10^9$		273	CP/Oc-23	S = MB; A' = 2,5-DMF; No measurable effect.	72F518
	MeOH	2.6×10^5	0.34		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4 \text{ s}^{-1}$.	736061
	2-PrOH	9.6×10^4			PL/Ld-2	S = RB.	89E324
8.6	Benzinemethanamine, <i>N,N</i> -di(phenylmethyl)- (Tribenzylamine)						
	MeOH/C ₆ H ₆ (67:33)		$1.4 \times 10^{-5} (\beta_r)$	298	CR/A'c-17	A' = Rub; used $\beta_r^{A'} = 1.4 \times 10^{-3} \text{ mol L}^{-1}$, ${}^1\text{O}_2^*$ from $(\text{PhO})_3\text{PO}_3$.	80M378
8.7	1,1'-Bipiperidine						
	C ₆ H ₆	1.7×10^7			PL/Ld-2	S = Ac.	90E297
	CF ₃ CH ₂ OH	2.4×10^5			PL/Ld-2	S = RB.	90E297
	CH ₃ CN	2.8×10^7			PL/Ld-2	S = RB.	90E297
	2-PrOH	3.2×10^6			PL/Ld-2	S = RB.	90E297
8.8	1,1'-Bipyrrolidine						
	C ₆ H ₆	7.2×10^8			PL/Ld-2	S = Ac.	90E297
	CF ₃ CH ₂ OH	2.2×10^6			PL/Ld-2	S = RB.	90E297
	CH ₃ CN	3.5×10^8			PL/Ld-2	S = RB.	90E297
	2-PrOH	1.8×10^8			PL/Ld-2	S = RB.	90E297
8.9	Brucine						
	CHCl ₃	3.9×10^8	2.6×10^{-5}		CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^4 \text{ s}^{-1}$.	84F670
8.10	1,4-Butanediamine, <i>N</i> -(3-aminopropyl)- (Spermidine)						
	C ₅ H ₅ N	$1.0 \times 10^7 (k_r)$			CP/A'c-17	S = A' = Rub; rel. to $k = 2.5 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$ for DABCO.	92R313
8.11	1,4-Butanediamine, <i>N,N</i> -bis(3-aminopropyl)- (Spermine)						
	C ₅ H ₅ N	$1.2 \times 10^7 (k_r)$			CP/A'c-17	S = A' = Rub; rel. to $k = 2.5 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$ for DABCO.	92R313
8.12	Butylamine						
	CHCl ₃	2.4×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	777486
	EtOH	$<1 \times 10^9$		73	CP/Oc-23	S = MB; A' = 2,5-DMF; No measurable effect.	72F518
8.13	Butylamine, <i>N,N</i> -dibutyl-						
	c-C ₆ H ₁₂	3×10^7			PL/Ld-2	S = 2-ACN.	90N078

TABLE 8. Rate constants for the interaction of singlet oxygen with aliphatic and alicyclic amines. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
8.13 Butylamine, <i>N,N</i>-dibutyl. — Continued							
	CHCl ₃	5.8×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486
	EtOH	$<1 \times 10^9$		273	CP/Oc-23	S = MB; A' = 2,5-DMF; No measurable effect.	72F518
8.14 Butylamine, <i>N,N</i>-dimethyl-							
	C ₆ H ₆	4.4×10^8			PL/Ld-2	S = Ac.	89E324
	CF ₃ CH ₂ OH	1.9×10^5			PL/Ld-2	S = RB.	89E324
	CH ₃ CN	2.7×10^8			PL/Ld-2	S = RB.	89E324
	CH ₃ COCH ₃	3.5×10^8			PL/Ld-2	S = RB.	89E324
	2-ProH	4.5×10^7			PL/Ld-2	S = RB.	89E324
8.15 <i>tert</i>-Butylamine							
	MeOH	6.6×10^5	0.14		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	736061
8.16 Chimasorb 944							
	C ₆ H ₆ / MeOH (80:20)	2.2×10^6			PL/A'd-8	S = MB; A' = DPBF; k calcd. for repeating unit.	84A167
	CHCl ₃	3.2×10^6		293	CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91P158
8.17 Cyclohexylamine							
	c-C ₆ H ₁₂ (mic)	4.6×10^7		313	CP/A'c-16	S = Fl ²⁻ ; A' = DPBF; used $k_d = 5.9 \times 10^4$ s ⁻¹ ; DAP reverse micelles.	80N021
	MeOH	2.9×10^7		313	CP/A'c-16	S = Fl ²⁻ ; A' = DPBF; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	80N021
	MeOH	8.0×10^4	1.1		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	736061
8.18 2,3-Diazabicyclo[2.2.1]heptane, 2,3-dimethyl-							
	CH ₃ CN	5.5×10^8			PL/Ld-2	S = RB.	90E297
8.19 1,4-Diazabicyclo[2.2.2]octane (DABCO)							
	1-BuOH	4.5×10^6	0.012		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 5.2 \times 10^4$ s ⁻¹ .	72F514
	C ₂ H ₅ CO ₂ CH ₃	2.5×10^8		293	PL/Ld-2	S = Pz and 2-ACN.	92E220
	C ₅ H ₅ N	3×10^8			CP/A'c-23	S = A' = Rub; used $k_d = 6.0 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	743112
	C ₅ H ₅ N		3.2×10^{-4}		CP/Oc-20	S = RB; A' = (C ₂ H ₅) ₃ N.	72F512
	C ₅ H ₅ N		2.3×10^{-4}		CP/Oc-21	S = RB; A' = 2M2P.	72F512
	c-C ₆ H ₁₂	2.1×10^7			PL/Ld-2	S = 2-ACN.	90N078
	c-C ₆ H ₁₂ (mic)	5.0×10^8		313	CP/A'c-16	S = Fl ²⁻ ; A' = DPBF; used $k_d = 5.9 \times 10^4$ s ⁻¹ ; DAP reverse micelles.	80N021
	1,2-C ₆ H ₄ Cl ₂				MD/A'c-23	A' = Rub; meas. $\beta_A/\beta_{A'} = 6.2$.	68F285
	C ₆ H ₅ Br	2.6×10^7		273	MD/A'c-33	A' = Rub; used $k_d = 1.3 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k/[k_d/[A'] + k_{A'}]$ = 0.2 at [A'] = 1.5×10^{-4} mol L ⁻¹ .	737333
	C ₆ H ₅ Br				MD/A'c-23	A' = DPBF; meas. $\beta_A/\beta_{A'} = 0.68$.	68F285
	C ₆ H ₅ CH ₃	2.4×10^8		293	PL/Ld-2	S = Pz and 2-ACN.	92E220
	C ₆ H ₅ CH ₃	2.2×10^8		298	PL/Ld-2	S = TPP; $\Delta V^\ddagger = -17$ cm ³ mol ⁻¹ ; studied at 0.1-120 MPa.	91A271
	C ₆ H ₅ CH ₃	2.1×10^8			PL/Ld-2	S = 2-ACN; $\Delta H^\ddagger = -7$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -126$ J K ⁻¹ mol ⁻¹ ; $E_a = -4$ kJ mol ⁻¹ ; studied at 183-363 K; pre-exciplex-equilibrium limit activation parameters.	88A427 84E066
	C ₆ H ₅ CH ₃	6.7×10^8			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063

TABLE 8. Rate constants for the interaction of singlet oxygen with aliphatic and alicyclic amines. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
1,4-Diazabicyclo[2.2.2]octane (DABCO) — Continued							
	C ₆ H ₅ CH ₃				CP/A'c-23	S = A' = Rub; meas. $\beta_A/\beta_{A'} = 0.23$.	68F285
	C ₆ H ₅ CN	3.7×10^8		298	PL/Ld-2	S = MB; $\Delta V_f = -10 \text{ cm}^3 \text{ mol}^{-1}$; studied at 0.1-120 MPa.	91A271
	C ₆ H ₆	2.9×10^8			PL/Ld-2	S = 2-ACN.	93N051
	C ₆ H ₆	2.9×10^8			PL/Ld-2	S = TPP and ZnTPP.	92F251
	C ₆ H ₆	2.6×10^8		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 4.2 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 4.2 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	90F360
	C ₆ H ₆	2.7×10^8 (k_q)					
	C ₆ H ₆	2.6×10^8			PL/Ld-2	S = 2-ACN.	84E066
	C ₆ H ₆	1.4×10^7	2.9×10^{-3}		CP/Ac-16	S = A' = 1,2-Diphenyl-4-methylcyclopenta[b]quinoline; used $k_d = 4.2 \times 10^4 \text{ s}^{-1}$.	78F438
	C ₆ H ₆					S = ZnTPP; A' = 2M2P.	727028
	C ₆ H ₆ /MeOH (80:20)	1.6×10^7		298	CP/P'a-20	S = MB; A' = 2M2P; used $k_d = 1.0 \times 10^4 \text{ s}^{-1}$, $\beta_{A'} = 0.040 \text{ mol L}^{-1}$.	70F734
	CCl ₄	4.7×10^7		295	PL/Ld-2	S = PdMPDME or MPDME; decay at 1588 nm; decay at 1272 nm gave $5 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	80E558
	CCl ₄	7×10^6			MP/LI-12	S = PP, Ph a or BPh a; used $k_d = 36 \text{ s}^{-1}$.	79A010
	CCl ₄	6.6×10^6			MP/LI-12	S = Ret; used $k_d = 36 \text{ s}^{-1}$.	79F463
						78F700	
	CF ₃ CH ₂ OH	4.4×10^4			PL/Ld-2	S = RB.	89E324
	CH ₂ Cl ₂	8.3×10^7		298	PL/Ld-2	S = MB; $\Delta V_f = -15 \text{ cm}^3 \text{ mol}^{-1}$; studied at 0.1-120 MPa.	91A271
	CH ₂ Cl ₂ /MeOH/C ₅ H ₅ N (90:5:5)	3.3×10^7		298	CR/A'c-32	A' = Rub; used $k_d = 8 \times 10^3 \text{ s}^{-1}$, $k_{A'} = 7.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$; ${}^1\text{O}_2^*$ from (PhO) ₃ PO ₃ .	747341
	CH ₂ Cl ₂ /MeOH/C ₅ H ₅ N (94:3:3)	3.3×10^7		298	CR/A'c-32	A' = Rub; used $k_d = 7.3 \times 10^3 \text{ s}^{-1}$, $k_{A'} = 7 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$; ${}^1\text{O}_2^*$ from (PhO) ₃ PO ₃ .	727319
	CH ₃ CN	4.0×10^8		298	PL/Ld-2	S = MB; $\Delta V_f = -11 \text{ cm}^3 \text{ mol}^{-1}$; studied at 0.1-120 MPa.	91A271
	CH ₃ CN	5.5×10^8		298	CP/P'a-17	S = DCA; A' = 2,3-Diphenyl-1,4-dioxene; used $k_d = 1.7 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	91A311
	CH ₃ CN	4.9×10^8			PL/Ld-2	S = RB.	89E324
	CH ₃ CN	4.0×10^8			PL/Ld-2	S = 2-ACN.	84E066
	CH ₃ COCH ₃	1.8×10^8		298	PL/Ld-2	S = MB; $\Delta V_f = -15 \text{ cm}^3 \text{ mol}^{-1}$; studied at 0.1-120 MPa.	91A271
	CH ₃ COCH ₃	3.8×10^8			PL/Ld-2	S = 2-ACN.	84E066
	CH ₃ COCH ₃	4.5×10^8			PL/Ld-2	S = MPDME.	83E235
	CHCl ₃	3.5×10^7		298	PL/Ld-2	S = MB; $\Delta V_f = -19 \text{ cm}^3 \text{ mol}^{-1}$; studied at 0.1-120 MPa.	91A271
	CHCl ₃	1.8×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	88P313
	CHCl ₃	5.8×10^7	1.7×10^{-4}		CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^4 \text{ s}^{-1}$.	84F670
	CHCl ₃	2.0×10^7			PL/Ld-2	S = MPDEE or PdMPDEE.	81A327
	CHCl ₃	4.7×10^7			PL/Ld-2	S = MPDEE or PdMPDEE.	81E472
	CHCl ₃	5.2×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	777486
	CS ₂		1.7×10^{-4}		CP/P'a-20	S = ZnTPP; A' = 2M2P.	727028
	ClCF ₂ CCl ₂ F	2.1×10^7		298	FP/Ld-2	S = Per.	82A322
	D ₂ O (mic) pD = 7.4	1.4×10^6		294	PL/A'd-5	S = 2-ACN; A' = DPBF; 0.1 mol L ⁻¹ SDS or CTAB.	81N048
	EtOH	5.0×10^6			PL/Ld-2	S = 2-ACN.	84E066

TABLE 8. Rate constants for the interaction of singlet oxygen with aliphatic and alicyclic amines. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
8.19 1,4-Diazabicyclo[2.2.2]octane (DABCO) — Continued							
	EtOH	6.4×10^6			CP/P'a-19	S = RB; A' = 2,2,6,6-Tetramethyl-4-piperidinol; used $k_{A'} = 8 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 8$; P' = 2,2,6,6-Tetramethyl-4-hydroxypiperidine-1-oxyl detected by esr.	80D076
	EtOH	3.1×10^7			CP/A'c-19	S = RB; A' = Chl a; used $k_d = 1 \times 10^5$ s ⁻¹ .	78F404
	EtOH	5.3×10^6			CP/Ac-23	S = RB; A' = Hexamethylenedithiocarbamate; used $k_d = 1 \times 10^4$ s ⁻¹ , $k_{A'} = 1.5 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_{A'}/(k_d + k[A]) = 58$ at [A] = 4.48×10^{-4} mol L ⁻¹ .	72T116
	H ₂ O pH = 8	2.8×10^8			CP/P'a-19	S = RB; A' = 2,2,6,6-Tetramethyl-4-piperidinol; meas. $k_A/k_{A'} = 7$; P' = 2,2,6,6-Tetramethyl-4-hydroxypiperidine-1-oxyl detected by esr.	80D076
	i-octane/ MeOH/ C ₅ H ₅ N (90:5:5)	3.9×10^7		298	CR/A'c-32	A' = Rub; used $k_d = 4.7 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from (PhO) ₃ PO ₃ .	74T341
	i-octane/ MeOH/ C ₅ H ₅ N (94:3:3)	3.5×10^7		298	CR/A'c-32	A' = Rub; used $k_d = 5.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from (PhO) ₃ PO ₃ .	72T319
	MeOH	5.4×10^8		313	CP/A'c-16	S = Fl ²⁻ ; A' = DPBF; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	80N021
	MeOH		6.5×10^{-3}		CP/P'a-20	S = ZnTPP; A' = 2M2P.	72T028
	MeOH	7.3×10^6	0.012		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F514
	2-PrOH	3.6×10^6			PL/Ld-2	S = RB.	89E324
	i-octane	3.0×10^7		298	CP/A'c-23	S = A' = Rub; used $k_d = 4.7 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	74T341
8.20 Dibenzo[<i>de,g</i>]quinoline-2,9-diol, 5,6,6a,7-tetrahydro-1,10-dimethoxy-6-methyl- (Boldine)							
	CHCl ₃	2.4×10^7	4.2×10^{-4}		CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F670
8.21 Dibenzo[<i>de,g</i>]quinoline-2,9-diol, 5,6,6a,7-tetrahydro-1,2,9,10-tetramethoxy-6-methyl- (Glaucine)							
	CHCl ₃	3.1×10^7	3.2×10^{-4}		CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F670
8.22 1,9:3,5-Dimethanocyclopenta[<i>d</i>]cyclopenta[3,4][1,2]diazeto[1,2-<i>a</i>]pyridazine, decahydro-							
	CH ₃ CN	2.8×10^8			PL/Ld-2	S = RB.	90E297
8.23 Ethanol, 2-(diethylamino)-							
	CHCl ₃	3.0×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	77T486
8.24 Ethylamine, <i>N,N</i>-diethyl-							
	C ₅ H ₅ N	2.5×10^7 (k_t)			CP/A'c-17	S = A' = Rub; rel. to $k = 2.5 \times 10^7$ L mol ⁻¹ s ⁻¹ for DABCO.	92R313
	C ₅ H ₅ N		2.7×10^{-3} (β_t)		CP/Oc-14,27	S = RB; A' = 2M2P.	72F512
	C ₅ H ₅ N		2.9×10^{-4} (β_q)		CP/Oc-14,27	S = RB; A' = 2M2P.	72F512
	C ₅ H ₅ N		4.2×10^{-4}		CP/Oc-20	S = RB; A' = 2M2P.	72F512
	c-C ₆ H ₁₂ (mic)	6.5×10^8		313	CP/A'c-16	S = Fl ²⁻ ; A' = DPBF; used $k_d = 5.9 \times 10^4$ s ⁻¹ ; DAP reverse micelles.	80N021
	n-C ₆ H ₁₄	4.8×10^7			CP/A'c-16	S = A' = Rub; k_d not given.	87F639
	C ₆ H ₆	1.8×10^8			CP/A'c-16	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	87F639
	CF ₃ CH ₂ OH	1.7×10^4			PL/Ld-2	S = RB.	89E324
	CH ₃ CN	3.3×10^8			PL/Ld-2	S = RB.	89E324
	CH ₃ CN	2.0×10^8			CP/A'c-16	S = RB; A' = DMA; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	87F639
	CH ₃ COCH ₃	3.1×10^8			CP/A'c-16	S = RB; A' = DMA; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	87F639
	CHCl ₃	7.4×10^7			CP/A'c-16	S = RB; A' = DMA; k_d not given.	87F639
	CHCl ₃	6.5×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	77T486

TABLE 8. Rate constants for the interaction of singlet oxygen with aliphatic and alicyclic amines. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_q/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
8.24 Ethylamine, <i>N,N</i>-diethyl- — Continued							
	EtOH	2.7×10^6 (k_r)	0.31 (β_r)		CP/Oc-19	$S = RB$; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; used (k_q/k_r) = 9.3.	75F655
	HCONH ₂	7.9×10^6			CP/A'c-16	$S = RB$; $A' = DMA$; k_d not given.	87F639
	MeOH	1.3×10^7			CP/A'c-16	$S = RB$; $A' = DMA$; used $k_d = 2.0 \times 10^4$ s ⁻¹ .	87F639
	MeOH	4.0×10^8		313	CP/A'c-16	$S = Fl^2$; $A' = DPBF$; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	80N021
	MeOH	9.3×10^6	9.7×10^{-3}		CP/A'c-16	$S = RB$; $A' = DPBF$; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	736061
	MeOH	1.2×10^7	7.6×10^{-3}		CP/A'c-16	$S = RB$; $A' = DPBF$; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F514
	MeOH/ C ₆ H ₆ (67:33)		4.9×10^{-4} (β_r)	298	CR/A'c-17	$A' = Rub$; used $\beta_r A' = 1.4 \times 10^{-3}$ mol L ⁻¹ ; $^1O_2^*$ from (PhO) ₃ PO ₃ .	80M378
	2-PrOH	3.6×10^7			PL/Ld-2	$S = RB$.	89E324
8.25 Ethylamine, <i>N,N</i>-diethyl-2-methoxy-							
	CHCl ₃	3.8×10^7			CP/A'c-33	$S = A' = Rub$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486
8.26 Ethylamine, 1,1-dimethyl-<i>N,N</i>-bis(2-hydroxyethyl)-							
	CHCl ₃	1.1×10^6			CP/A'c-33	$S = A' = Rub$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486
8.27 Ethylamine, <i>N</i>-ethyl-							
	C ₅ H ₅ N	5×10^6 (k_r)			CP/A'c-17	$S = A' = Rub$; rel. to $k = 2.5 \times 10^7$ L mol ⁻¹ s ⁻¹ for DABCO.	92R313
	c-C ₆ H ₁₂ (mic)	3.1×10^8		313	CP/A'c-16	$S = Fl^2$; $A' = DPBF$; used $k_d = 5.9 \times 10^4$ s ⁻¹ ; DAP reverse micelles.	80N021
	n-C ₆ H ₁₄	4.8×10^6			CP/A'c-16	$S = A' = Rub$; k_d not given.	87F639
	C ₆ H ₆	4.6×10^7			CP/A'c-16	$S = A' = Rub$; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	87F639
	C ₆ H ₆ / MeOH (75:25)	2.4×10^6		298	CP/A'c-20	$S = RB$; $A' = c-C_6H_{10}$; used $k_d = 6.3 \times 10^4$ s ⁻¹ , $\beta_{A'} = 3.1$ mol L ⁻¹ .	78F586
	CH ₃ CN	1.5×10^8			PL/Ld-2	$S = RB$.	89E324
	CH ₃ CN	8.6×10^7			CP/A'c-16	$S = RB$; $A' = DMA$; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	87F639
	CH ₃ COCH ₃	1.0×10^8			CP/A'c-16	$S = RB$; $A' = DMA$; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	87F639
	CHCl ₃	1.2×10^7			CP/A'c-16	$S = RB$; $A' = DMA$; k_d not given.	87F639
	CHCl ₃	1.5×10^7			CP/A'c-33	$S = A' = Rub$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486
	HCONH ₂	2.3×10^6			CP/A'c-16	$S = RB$; $A' = DMA$; k_d not given.	87F639
	MeOH	2.2×10^6			CP/A'c-16	$S = RB$; $A' = DMA$; used $k_d = 2.0 \times 10^4$ s ⁻¹ .	87F639
	MeOH	1.7×10^8		313	CP/A'c-16	$S = Fl^2$; $A' = DPBF$; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	80N021
	MeOH	1.9×10^6	0.048		CP/A'c-16	$S = RB$; $A' = DPBF$; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	736061
	MeOH/ C ₆ H ₆ (67:33)		4.0×10^{-4} (β_r)	298	CR/A'c-17	$A' = Rub$; used $\beta_r A' = 1.4 \times 10^{-3}$ mol L ⁻¹ ; $^1O_2^*$ from (PhO) ₃ PO ₃ .	80M378
	2-PrOH	3.8×10^6			PL/Ld-2	$S = RB$.	89E324
8.28 Ethylamine, 1-methyl-							
	MeOH	4.6×10^4	1.9		CP/A'c-16	$S = RB$; $A' = DPBF$; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	736061
8.29 Ethylamine, 1-methyl-<i>N,N</i>-(1-methylethyl)-							
	c-C ₆ H ₁₂	8.8×10^5			PL/Ld-2	$S = 2-ACN$.	90N078
	CHCl ₃	1.8×10^6			CP/A'c-33	$S = A' = Rub$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486
8.30 Ethylenediamine, tetraethyl-							
	CHCl ₃	3×10^8			CR/A'c-33	$A' = Rub$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; $^1O_2^*$ from DMNO ₂ .	81E003

TABLE 8. Rate constants for the interaction of singlet oxygen with aliphatic and alicyclic amines. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
8.31	1,7-Heptanediamine, <i>N,N</i> -diethyl-						
	CHCl ₃	6.1×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486
8.32	Hexamethylenetetramine						
	MeOH	1.9×10^5	0.46		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	736061
8.33	Hydrazine, 1,2-diethyl-1,2-dimethyl-						
	CH ₃ CN	6.8×10^7			PL/Ld-2	S = RB.	90E297
8.34	Hydrazine, 1,2-dimethyl-1,2-dibutyl-						
	C ₆ H ₆	3.9×10^7			PL/Ld-2	S = Ac.	90E297
	CH ₃ CN	3.9×10^7			PL/Ld-2	S = RB.	90E297
8.35	Hydrazine, 1,1-dimethyl-2,2-di(2,2-dimethylpropyl)-						
	CH ₃ CN	1.3×10^6			PL/Ld-2	S = RB.	90E297
8.36	Hydrazine, 1,2-dimethyl-1,2-di(2,2-dimethylpropyl)-						
	CH ₃ CN	5.3×10^6			PL/Ld-2	S = RB.	90E297
8.37	Hydrazine, 1,1-dimethyl-2,2-di(2-methylpropyl)-						
	CH ₃ CN	2.2×10^7			PL/Ld-2	S = RB.	90E297
8.38	Hydrazine, 1,2-dimethyl-1,2-di(2-methylpropyl)-						
	C ₆ H ₆	5.9×10^6			PL/Ld-2	S = Ac.	90E297
	CH ₃ CN	1.3×10^7			PL/Ld-2	S = RB.	90E297
8.39	Hydrazine, 1,2-dimethyl-1,2-dipentyl-						
	C ₆ H ₆	9.3×10^6			PL/Ld-2	S = Ac.	90E297
	CH ₃ CN	4.7×10^7			PL/Ld-2	S = RB.	90E297
8.40	Hydrazine, 1,2-dimethyl-1,2-dipropyl-						
	CH ₃ CN	4.4×10^7			PL/Ld-2	S = RB.	90E297
8.41	Hydrazine, tetramethyl-						
	C ₆ H ₆	1.8×10^8			PL/Ld-2	S = Ac.	90E297
	CF ₃ CH ₂ OH	1.2×10^6			PL/Ld-2	S = RB.	90E297
	CH ₃ CN	1.9×10^8			PL/Ld-2	S = RB.	90E297
	2-PrOH	2.1×10^7			PL/Ld-2	S = RB.	90E297
8.42	Hydrazine, tetra(methyl- <i>d</i> ₃)-						
	CH ₃ CN	1.9×10^8			PL/Ld-2	S = RB.	90E297
8.43	Hydrazine, tetra(2-methylpropyl)-						
	CH ₃ CN	3.0×10^5			PL/Ld-2	S = RB.	90E297
8.44	Hydrazine, 1,1,2-trimethyl-2-(2,2-dimethylpropyl)-						
	CH ₃ CN	1.1×10^8			PL/Ld-2	S = RB.	90E297
8.45	Hydrazine, 1,1,2-trimethyl-2-(2-methylpropyl)-						
	C ₆ H ₆	6.8×10^7			PL/Ld-2	S = Ac.	90E297
	CH ₃ CN	9.6×10^7			PL/Ld-2	S = RB.	90E297
8.46	Methylamine, <i>N,N</i> -bis(2-hydroxyethyl)-						
	CHCl ₃	2.1×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486

TABLE 8. Rate constants for the interaction of singlet oxygen with aliphatic and alicyclic amines. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	$\cdot T$ (K)	Method	Comment	Ref.
8.47	Methylamine, <i>N,N</i> -dimethyl-						
	MeOH	1.3×10^7	6.7×10^{-3}		CP/A'c-16	$S = MB; A' = DPBF$; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
8.48	Phenethylamine						
	c-C ₆ H ₁₂ (mic)	4.0×10^7		313	CP/A'c-16	$S = Fl^2$; $A' = DPBF$; used $k_d = 5.9 \times 10^4$ s ⁻¹ ; DAP reverse micelles.	80N021
	MeOH	5.3×10^7		313	CP/A'c-16	$S = Fl^2$; $A' = DPBF$; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	80N021
	MeOH		1.5		CP/A'c-16	$S = RB; A' = DPBF$.	736061
8.49	4-Phenylbutylamine						
	MeOH		0.98		CP/A'c-16	$S = RB; A' = DPBF$.	736061
8.50	3-Phenylpropylamine						
	MeOH		1.3		CP/A'c-16	$S = RB; A' = DPBF$.	736061
8.51	Piperazine						
	c-C ₆ H ₁₂	1.7×10^6			PL/Ld-2	$S = 2-ACN$.	90N078
	MeOH	1.2×10^6	0.073		CP/A'c-16	$S = RB; A' = DPBF$; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	736061
8.52	1-Piperidinamine, <i>N,N</i> -dimethyl-						
	CH ₃ CN	1.6×10^8			PL/Ld-2	$S = RB$.	90E297
8.53	Piperidine						
	2-BuOH	1.0×10^6			PL/Ld-2	$S = RB$.	89E324
	C ₆ H ₆ / MeOH (80:20)	3.5×10^6			PL/A'd-8	$S = MB; A' = DPBF$.	84A167
	CF ₃ CH ₂ OH	6.5×10^3			PL/Ld-2	$S = RB$.	89E324
	CH ₃ CN	5.6×10^7			PL/Ld-2	$S = RB$.	89E324
	CH ₃ COCH ₃	7.8×10^7			PL/Ld-2	$S = RB$.	89E324
	CHCl ₃	3.6×10^6			PL/Ld-2	$S = TPP$.	89E324
	CHCl ₃	5.8×10^6			CP/A'c-33	$S = A' = Rub$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486
	MeOH	9.2×10^5	0.098		CP/A'c-16	$S = RB; A' = DPBF$; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	736061
	2-PrOH	9.1×10^5			PL/Ld-2	$S = RB$.	89E324
8.54	Piperidine, 4-amino-2,2,6,6-tetramethyl-						
	C ₆ H ₆ / MeOH (80:20)	1.8×10^6			PL/A'd-8	$S = MB; A' = DPBF$.	84A167
8.55	Piperidine, 1-butoxy-2,2,6,6-tetramethyl-						
	CH ₂ Cl ₂	3×10^5			CP/A'c-18	$S = A' = Rub$; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88F059
8.56	Piperidine, 1-cyclohexyl-						
	MeOH	4.5×10^7	2.0×10^{-3}		CP/A'c-16	$S = MB; A' = DPBF$; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
8.57	Piperidine, 1,2-dimethyl-						
	C ₆ H ₆ / MeOH (80:20)	2.8×10^7			PL/A'd-8	$S = MB; A' = DPBF$.	84A167
8.58	Piperidine, 2,6-dimethyl-						
	C ₆ H ₆ / MeOH (80:20)	1.2×10^6			PL/A'd-8	$S = MB; A' = DPBF$.	84A167
	CHCl ₃	3.0×10^6			CP/A'c-33	$S = A' = Rub$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486

TABLE 8. Rate constants for the interaction of singlet oxygen with aliphatic and alicyclic amines. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
8.59	Piperidine, 2-methyl-						
	C ₆ H ₆ / MeOH (80:20)	1.5×10^6			PL/A'd-8	S = MB; A' = DPBF.	84A167
8.60	Piperidine, N-methyl-						
	C ₆ H ₆ / MeOH (80:20)	2.9×10^7			PL/A'd-8	S = MB; A' = DPBF.	84A167
	CHCl ₃	5.3×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486
8.61	Piperidine, 1,2,2,6,6-pentamethyl-						
	C ₆ H ₆ / MeOH (80:20)	6.0×10^7			PL/A'd-8	S = MB; A' = DPBF.	84A167
8.62	Piperidine, 1-(1-pyrrolidinyl)-						
	CH ₃ CN	2.6×10^8			PL/Ld-2	S = RB.	90E297
8.63	Piperidine, 2,2,6,6-tetramethyl-						
	C ₆ H ₆ / MeOH (80:20)	4.0×10^6			PL/A'd-8	S = MB; A' = DPBF.	84A167
	C ₆ H ₆	5.3×10^5	0.061		CP/Pa-15	S = HA; used $k_d = 3.2 \times 10^4$ s ⁻¹ ; P = 2,2,6,6-Tetramethylpiperidine-N-oxyl; P monitored by esr.	90R162
	CH ₂ Cl ₂	3.9×10^5			PL/Ld-2	S = TPP.	93E090
	CH ₂ Cl ₂	3.8×10^5			CP/A'c-33	S = A' = Rub; used $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; k_d not given.	84F199
	EtOH	1.6×10^5	0.51		CP/Pa-15	S = BXP; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; P = TEMPO monitored by esr.	83F323
	CH ₂ Cl ₂	4×10^5			CP/A'c-18	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88F059
8.64	Piperidine, 2,2,6,6-tetramethyl-4,4'-[1,6-hexanediaminyl]bis-						
	C ₆ H ₆ / MeOH (80:20)	8.0×10^6			PL/A'd-8	S = MB; A' = DPBF.	84A167
8.65	Piperidine, 2,2,6,6-tetramethyl-4,4'-[1,6-hexanediaminyl]bis-, polymer						
	C ₆ H ₆ / MeOII (80:20)	3.6×10^7			PL/A'd-8	S = MB; A' = DPBF; k calcd. for repeating unit.	84A167
8.66	1-Piperidineethanol, diester with [[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]butylpropanedioic acid						
	C ₆ H ₆ / MeOH (80:20)	4.3×10^6			PL/A'd-8	S = MB; A' = DPBF.	84A167
8.67	1-Piperidineethanol, 4-hydroxy-2,2,6,6-tetramethyl-, polymer with butanedioic acid (Tinuvin 622)						
	C ₆ H ₆ / MeOH (80:20)	$<1 \times 10^5$			PL/A'd-8	S = MB; A' = DPBF; k calcd. for repeating unit.	84A167
	CHCl ₃	2×10^5		293	CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91P158
8.68	1-Piperidineethanol, 2,2,6,6-tetramethyl-						
	CHCl ₃	$<2 \times 10^5$			CP/A'c-33	S = A' = Rub; No measurable effect.	777486
8.69	1-Piperidineethanol, 2,2,6,6-tetramethyl-, acetate (ester)						
	CHCl ₃	$<2 \times 10^5$			CP/A'c-33	S = A' = Rub; No measurable effect.	777486
8.70	Piperidin-3-ol, 1-methyl-						
	C ₆ H ₆ / MeOH (80:20)	2.5×10^7			PL/A'd-8	S = MB; A' = DPBF.	84A167

TABLE 8. Rate constants for the interaction of singlet oxygen with aliphatic and alicyclic amines. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
8.71	Piperidin-4-ol, 1-methyl-						
	C ₆ H ₆ / MeOH (80:20)	2.0×10^7			PL/A'd-8	S = MB; A' = DPBF.	84A167
8.72	Piperidin-4-ol, 1,2,2,6,6-pentamethyl-						
	C ₆ H ₆ / MeOH (80:20)	6.0×10^7			PL/A'd-8	S = MB; A' = DPBF.	84A167
	C ₆ H ₆ / EtOH (89:11)	$2.3 \times 10^3 (k_r)$		295	CP/Pa,A'c-17	S = RB; A' = Tetr; used $k_t^{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_t^{A'} = 3.3 \times 10^{-5}$; P = nitroxide radical.	757445
	CH ₂ Cl ₂	5×10^7		295	?	Method not given.	757445
	CH ₂ Cl ₂	5.2×10^7	1.4×10^{-4}		CP/A'c-19	S = A' = Tetr; used $k_d = 7.3 \times 10^3$ s ⁻¹ .	75F654
	CHCl ₃	3.3×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
	CHCl ₃	9.2×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486
8.73	Piperidin-4-ol, 1,2,2,6,6-pentamethyl-, diester with [[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]butylpropanedioic acid (Tinuvin 144)						
	C ₆ H ₆ / MeOH (80:20)	1.5×10^8			PL/A'd-8	S = MB; A' = DPBF.	84A167
8.74	Piperidin-4-ol, 1,2,2,6,6-pentamethyl-, diester with 1,10-decanedioic acid						
	CHCl ₃	1.8×10^8			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
8.75	Piperidin-4-ol, 1,2,2,6,6-pentamethyl-, triester with nitrilotriacetic acid						
	CHCl ₃	2.4×10^8			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
8.76	Piperidin-4-ol, 1,2,2,6-tetramethyl-, phenylacetate ester (Eucatropine)						
	CHCl ₃	2×10^6	5.3×10^{-3}		CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^4$ s ⁻¹ .	84F670
8.77	Piperidin-4-ol, 2,2,6,6-tetramethyl- (TEMP-4-OH)						
	C ₆ H ₆ / EtOH (50:50)	$<1.0 \times 10^5$			PL/A'd-8	S = MB; A' = DPBF.	84A167
	C ₆ H ₆ / EtOH (89:11)	$2.9 \times 10^3 (k_r)$		295	CP/Pa,A'c-17	S = RB; A' = Tetr; used $k_t^{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_t^{A'} = 4.1 \times 10^{-5}$; P = TEMPOL.	757445
	C ₆ H ₆ / EtOH (89:11)	5×10^5		295	CP/A'c-23	S = RB; A' = Tetr; used $k_d = 3 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	757445
	CHCl ₃	2.6×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
	CHCl ₃	$<2.0 \times 10^5$			CP/A'c-33	S = A' = Rub; No measurable effect.	777486
	DMF	5.0×10^7			CP/Pa-15	S = Pt(phen)(N ₃) ₂ ; P = 4-Hydroxy-2,2,6,6-tetramethylpiperidine N-oxyl; k_d not given.	90F196
	DMF	5.0×10^7			MP/Pa-15	S = HP; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; P = 4-Hydroxy-2,2,6,6-tetramethylpiperidine N-oxyl; esr detection.	89F311
	DMSO	5.2×10^6	0.01		CP/Pa-15	S = GV; used $k_d = 5.2 \times 10^4$ s ⁻¹ ; P = 4-Hydroxy-2,2,6,6-tetramethylpiperidine N-oxyl; esr detection.	89D112
	EtOH	8×10^5			CP/Pa-15	S = RB; used $k_d = 8 \times 10^4$ s ⁻¹ ; P = 4-Hydroxy-2,2,6,6-tetramethylpiperidine N-oxyl; detected by esr.	80D076
	H ₂ O pH = 8	4×10^7			CP/Pa-19	S = RB; Q = N ₃ ⁻ ; used $k_Q = 2 \times 10^9$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 0.025$; P = 4-Hydroxy-2,2,6,6-tetramethylpiperidine N-oxyl; detected by esr.	80D076

TABLE 8. Rate constants for the interaction of singlet oxygen with aliphatic and alicyclic amines. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
8.78 Piperidin-4-ol, 2,2,6,6-tetramethyl-, diester with 1,10-decanedioic acid (Tinuvin 770)							
	C ₆ H ₆ / MeOH (80:20)	<1 × 10 ⁵			PL/A'd-8	S = MB; A' = DPBF.	84A167
	CHCl ₃	2 × 10 ⁵		293	CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91P158
	CHCl ₃	1.5 × 10 ⁵			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
8.79 Piperidin-4-ol, 2,2,6,6-tetramethyl-, ester with copolymer of styrene + methacrylic acid							
	CHCl ₃	8.2 × 10 ⁵		293	CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91P158
8.80 Piperidin-4-ol, 2,2,6,6-tetramethyl-, triester with nitrilotriacetic acid							
	CHCl ₃	1.9 × 10 ⁶			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
8.81 Propanenitrile, 3-(diethylamino)-							
	CHCl ₃	2.7 × 10 ⁷			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486
8.82 1-Propanol, 3-(dimethylamino)-							
	C ₆ H ₆	6.8 × 10 ⁷			PL/Ld-2	S = Ac.	89E324
	CF ₃ CH ₂ OH	1.8 × 10 ⁵			PL/Ld-2	S = RB.	89E324
	CH ₃ CN	1.4 × 10 ⁸			PL/Ld-2	S = RB.	89E324
	CH ₃ COCH ₃	1.9 × 10 ⁸			PL/Ld-2	S = RB.	89E324
8.83 1-Propen-1-amine, N,N,2-trimethyl-							
	C ₆ H ₆	4.8 × 10 ⁸ (k_r)			CP/Ac,A'c-17	S = ZnTPP; A' = TME; used $k_r^{A'} = 4.8 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 10$.	75F656
	C ₆ H ₆ / CH ₃ CN (5:95)				CP/Ac,A'c-17	S = ZnTPP; A' = TME; meas. $k_r/k_r^{A'} = 9.4$.	75F656
	C ₆ H ₆ / CH ₃ CN (80:20)				CP/Ac,A'c-17	S = ZnTPP; A' = TME; meas. $k_r/k_r^{A'} = 7.8$.	75F656
	C ₆ H ₆ / DMSO (20:80)				CP/Ac,A'c-17	S = ZnTPP; A' = TME; meas. $k_r/k_r^{A'} = 9.1$.	75F656
	C ₆ H ₆ / DMSO (60:40)				CP/Ac,A'c-17	S = ZnTPP; A' = TME; meas. $k_r/k_r^{A'} = 7.6$.	75F656
	C ₆ H ₆ / DMSO (80:20)				CP/Ac,A'c-17	S = ZnTPP; A' = TME; meas. $k_r/k_r^{A'} = 11$.	75F656
	C ₆ H ₆ / MeOH (30:70)				CP/Ac,A'c-17	S = ZnTPP; A' = TME; meas. $k_r/k_r^{A'} = 2.9$.	75F656
	C ₆ H ₆ / MeOH (5:95)				CP/Ac,A'c-17	S = ZnTPP; A' = TME; meas. $k_r/k_r^{A'} = 2.2$.	75F656
	C ₆ H ₆ / n-C ₅ H ₁₂ (5:95)				CP/Ac,A'c-17	S = ZnTPP; A' = TME; meas. $k_r/k_r^{A'} = 7.1$.	75F656
8.84 Propylamine							
	CHCl ₃	2.3 × 10 ⁵			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486
8.85 Propylamine, 2-methyl-							
	CHCl ₃	4.1 × 10 ⁵			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777486
8.86 Propylamine, N-propyl-							
	c-C ₆ H ₁₂	9.7 × 10 ⁶			PL/Ld-2	S = 2-ACN.	90N078

TABLE 8. Rate constants for the interaction of singlet oxygen with aliphatic and alicyclic amines. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
8.86 Propylamine, <i>N</i>-propyl- — Continued							
	CHCl ₃	1.8×10^7			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	777486
8.87 Propyn-1-amine, <i>N,N</i>-diethyl-							
	CHCl ₃	2×10^8			CR/A'c-33	$A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$; ${}^1\text{O}_2^*$ from DMNO ₂ .	81E003
8.88 Pyrazolo[1,2-<i>a</i>][1,2]diazepine, hexahydro-							
	CH ₃ CN	7.1×10^8			PL/Ld-2	$S = \text{RB}$.	90E297
8.89 Pyrazolo[1,2-<i>a</i>]pyrazole, tetrahydro-							
	CH ₃ CN	7.7×10^8			PL/Ld-2	$S = \text{RB}$.	90E297
8.90 Pyrazolo[1,2-<i>a</i>]pyrazole, tetrahydro-2,2,6,6-tetramethyl-							
	C ₆ H ₆	1.8×10^9			PL/Ld-2	$S = \text{Ac}$.	90E297
	CH ₃ CN	1.3×10^9			PL/Ld-2	$S = \text{RB}$.	90E297
8.91 Pyridazine, hexahydro-1,2-dimethyl-							
	CH ₃ CN	4.6×10^8			PL/Ld-2	$S = \text{RB}$.	90E297
8.92 Pyridazino[1,2-<i>a</i>]pyridazine, octahydro-							
	C ₆ H ₆	2.8×10^8			PL/Ld-2	$S = \text{Ac}$.	90E297
	CF ₃ CH ₂ OH	2.5×10^6			PL/Ld-2	$S = \text{RB}$.	90E297
	CH ₃ CN	2.9×10^8			PL/Ld-2	$S = \text{RB}$.	90E297
	2-PrOH	7.4×10^7			PL/Ld-2	$S = \text{RB}$.	90E297
8.93 Pyridine, 3-(1-methyl-2-pyrrolidinyl)- (Nicotine)							
	CH ₃ CN/D ₂ O (80:20)	5.9×10^7 (k_q)			PL/A'd-5	$S = 2\text{-ACN}$; $A' = \text{DPBF}$; $k_q = 0$ estd. from lack of formation of O_2^- , ded. from buildup of radical anion in soln. contg. (2-20) $\times 10^{-5}$ mol L ⁻¹ 1,4-benzoquinone in CH ₃ CN/D ₂ O (1:4).	81A191
	CHCl ₃	3.8×10^7	2.6×10^{-4}		CP/A'c-16	$S = A' = \text{Rub}$; used $k_d = 1 \times 10^4 \text{ s}^{-1}$.	84F670
	MeOH		0.23	288	CP/Oc-?	$S = \text{RB}$; $A' = \alpha\text{-Terpinene}$; meas. $k_q/k_r = 5.4$.	587004
8.94 Pyridine, 1,2,3,6-tetrahydro-1-methyl-4-phenyl-							
	D ₂ O	3.1×10^4			PL/Ld-2	$S = \text{MD}$; P = <i>N</i> -Methyl-4-phenyl-2,3-dihydropyridinium.	91R251
8.95 1-Pyrrolidinamine, <i>N,N</i>-dimethyl-							
	CH ₃ CN	5.6×10^8			PL/Ld-2	$S = \text{RB}$.	90E297
8.96 Pyrrolidine							
	c-C ₆ H ₁₂ (mic)	1.9×10^8		313	CP/A'c-16	$S = \text{Fl}^2$; $A' = \text{DPBF}$; used $k_d = 5.9 \times 10^4 \text{ s}^{-1}$; DAP reverse micelles.	80N021
	CF ₃ CH ₂ OH	1.5×10^4			PL/Ld-2	$S = \text{RB}$.	89E324
	CH ₃ CN	1.4×10^8			PL/Ld-2	$S = \text{RB}$.	89E324
	MeOH	6.5×10^7		313	CP/A'c-16	$S = \text{Fl}^2$; $A' = \text{DPBF}$; used $k_d = 1.4 \times 10^5 \text{ s}^{-1}$.	80N021
	MeOH	2.1×10^6	0.042		CP/A'c-16	$S = \text{RB}$; $A' = \text{DPBF}$; used $k_d = 9.0 \times 10^4 \text{ s}^{-1}$.	736061
	2-PrOH	2.2×10^6			PL/Ld-2	$S = \text{RB}$.	89E324
8.97 Quinuclidine							
	MeOH	1.8×10^6	0.051		CP/A'c-16	$S = \text{RB}$; $A' = \text{DPBF}$; used $k_d = 9.0 \times 10^4 \text{ s}^{-1}$.	736061
8.98 Reserpine							
	CHCl ₃	2.7×10^7	3.8×10^{-4}		CP/A'c-16	$S = A' = \text{Rub}$; used $k_d = 1 \times 10^4 \text{ s}^{-1}$.	84F670

TABLE 8. Rate constants for the interaction of singlet oxygen with aliphatic and alicyclic amines. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
8.99 Strychnine							
	C ₆ D ₆	8.8×10^8			PR/Ld-2	S = 2-ACN.	89E113
	C ₆ H ₅ CH ₃	9.3×10^8			PL/Ld-2	S = 2-ACN; $\Delta H^\ddagger = 4 \text{ kJ mol}^{-1}$; $\Delta S^\ddagger = -77 \text{ J K}^{-1}$ mol ⁻¹ ; $E_a = 7 \text{ kJ mol}^{-1}$; studied at 183-363 K.	88A427 84E066
	C ₆ H ₆	9.0×10^8			PL/Ld-2	S = 2-ACN.	93E248
	C ₆ H ₆	1.0×10^9 (k_q)			PL/Ld-2	S = 2-ACN; comparison with TME shows strychnine 10 times less reactive, therefore k is at least 99% k_q .	84E066
	CH ₃ CN	6.4×10^8			PL/Ld-2	S = 2-ACN.	84E066
	CH ₃ COCl ₂	9.4×10^8			PL/Ld-2	S = 2-ACN.	84E066
	CHCl ₃	2.4×10^8	4.3×10^{-5}		CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^4 \text{ s}^{-1}$.	84F670
	EtOH	1.1×10^8			PL/Ld-2	S = 2-ACN.	84E066
8.100 Vincamine							
	CHCl ₃	4.3×10^7	2.3×10^{-4}		CP/A'c-16	S = A' = Rub; used $k_d = 1 \times 10^4 \text{ s}^{-1}$.	84F670
8.101 Vobasan-17-oic acid, 19,20-dihydro-3-oxo-, methyl ester, (20α) (Dregamine)							
	C ₆ H ₆		1.5×10^{-3}		CP/Ac-15	S = ZnTPP.	78F474
	MeOH/ C ₆ H ₆ (67:33)		2.0×10^{-3}		CP/Ac-15	S = ZnTPP.	78F474

TABLE 9. Rate constants for the interaction of singlet oxygen with aromatic amines.

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
9.1 Aniline							
	EtOH				CP/A'c-20	S = Eos,Ery, or RB; A' = TME; meas. $k_A/k_{A'} = 3.0$.	717299
	EtOH				CP/A'c-20	S = A; A' = TME; meas. $k_A/k_{A'} = 2.7$.	717299
	c-C ₆ H ₁₂ (mic)	3.9×10^7			CP/A'c-16	S = Fl ²⁻ ; A' = DPBF; used $k_d = 5.9 \times 10^4$ s ⁻¹ ; DAP reverse micelles.	80N021
	MeOH	2.0×10^9			CP/A'c-16	S = Fl ²⁻ ; A' = DPBF; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	80N021
	MeOH/ C ₆ H ₆ (67:33)	1.5×10^{-4}		298	CR/A'c-17	A' = Rub; used $\beta_{A'} = 1.4 \times 10^{-3}$ mol L ⁻¹ ; ¹ O ₂ * from (PhO) ₃ PO ₃ .	80M378
9.2 Aniline, 4-bromo-<i>N,N</i>-dimethyl-							
	MeOH	2×10^7			PL/A'd-8	S = MB; A' = DPBF.	74F640
	MeOH		3×10^{-3}		CP/A'c-16	S = RB; A' = DPBF.	736061
9.3 Aniline, 3-chloro-<i>N,N</i>-dimethyl-							
	MeOH	1.1×10^7			PL/A'd-8	S = MB; A' = DPBF.	74F640
	MeOH		7.1×10^{-3}		CP/A'c-16	S = RB; A' = DPBF.	736061
9.4 Aniline, 4-chloro-<i>N,N</i>-dimethyl-							
	MeOH/ H ₂ O (50:50)	3.6×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	83A094
9.5 Aniline, 3,4-dimethoxy-<i>N,N</i>-dimethyl-							
	MeOH/ H ₂ O (50:50)	8.8×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	83A094
9.6 Aniline, <i>N,N</i>-dimethyl-							
	C ₅ H ₅ N	1.0×10^8	5.9×10^{-4}		CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ .	757166
	c-C ₆ H ₁₂ (mic)	3.3×10^7			CP/A'c-16	S = Fl ²⁻ ; A' = DPBF; used $k_d = 5.9 \times 10^4$ s ⁻¹ ; DAP reverse micelles.	80N021
	CH ₃ CN	4.8×10^8			PL/Ld-2	S = RB.	89E324
	MeOH	1.2×10^9			CP/A'c-16	S = Fl ²⁻ ; A' = DPBF; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	80N021
	MeOH		0.029		CP/A'c-?	S = RB; A' = 2,5-DMF; solvent ?.	777055
	MeOH	1.2×10^8	7.6×10^{-4}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	736061
	MeOH	7.3×10^7			PL/A'd-8	S = MB; A' = DPBF.	737014
	MeOH/ C ₆ H ₆ (67:33)		2.2×10^{-3}	298	CR/A'c-17	A' = Rub; used $\beta_{A'} = 1.4 \times 10^{-3}$ mol L ⁻¹ ; ¹ O ₂ * from (PhO) ₃ PO ₃ .	80M378
	MeOH/ H ₂ O (50:50)	3.6×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	83A094
	2-PrOH	1.7×10^7			PL/Ld-2	S = RB.	89E324
9.7 Aniline, <i>N,N</i>-dimethyl-4-nitroso-							
	H ₂ O	6.8×10^7 (k_r)		298	CP/Ac-14	S = H ₂ TPPS ⁴⁻ ; used $k_d = 2.5 \times 10^5$ s ⁻¹ , $E_a = 15.9$ kJ mol ⁻¹ ; log(A) = 10.6; studied at 288-318 K; used $\phi_A(S) = 0.7$, phosphate buffer contg. 1.6% NaCl.	91R053
9.8 Aniline, <i>N,N</i>-diphenyl- (Triphenylamine)							
	C ₆ H ₅ CH ₃	1.6×10^5		293	PL/Ld-2	S = RBEE(Bu ₃ N); Pz also used as S.	92E220
9.9 Aniline, 4-(ethoxycarbonyl)- (Ethyl 4-aminobenzoate)							
	CHCl ₃	8×10^4			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80A041

TABLE 9. Rate constants for the interaction of singlet oxygen with aromatic amines. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
9.10 Aniline, 4-formyl-<i>N,N</i>-dimethyl- (4-(Dimethylamino)benzaldehyde)							
	CHCl ₃	1.1×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80A041
	MeOH	1×10^6			PL/A'd-8	S = MB; A' = DPBF.	74F640
	MeOH		0.037		CP/A'c-16	S = RB; A' = DPBF.	736061
9.11 Aniline, 2-methoxy-<i>N,N</i>-dimethyl-							
	MeOH/ H ₂ O (50:50)	3.2×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	83A094
9.12 Aniline, 3-methoxy-<i>N,N</i>-dimethyl-							
	MeOH	4.8×10^7			PL/A'd-8	S = MB; A' = DPBF.	74F640
	MeOH		2×10^{-3}		CP/A'c-16	S = RB; A' = DPBF.	736061
	MeOH/ H ₂ O (50:50)	8.4×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	83A094
9.13 Aniline, 4-methoxy-<i>N,N</i>-dimethyl-							
	MeOH	2×10^8			PL/A'd-8	S = MB; A' = DPBF.	74F640
	MeOH		5×10^{-4}		CP/A'c-16	S = RB; A' = DPBF.	736061
	MeOH/ H ₂ O (50:50)	4.4×10^8	6.2×10^{-4}	308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	83A094
9.14 Aniline, <i>N</i>-methyl-							
	c-C ₆ H ₁₂ (mic)	5.5×10^7			CP/A'c-16	S = Fl ²⁻ ; A' = DPBF; used $k_d = 5.9 \times 10^4$ s ⁻¹ ; DAP reverse micelles.	80N021
	MeOH	8.0×10^8			CP/A'c-16	S = Fl ²⁻ ; A' = DPBF; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	80N021
	MeOH	2.7×10^7	3.4×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	736061
	MeOH/ C ₆ H ₆ (67:33)		5.3×10^{-4}	298	CR/A'c-17	A' = Rub; used $\beta_{A'} = 1.4 \times 10^{-3}$ mol L ⁻¹ ; $^1\text{O}_2^*$ from (PhO) ₃ PO ₃ .	80M378
9.15 Aniline, <i>N,N</i>,2,4,6-pentamethyl-							
	MeOH/ H ₂ O (50:50)	1.5×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	83A094
9.16 Aniline, <i>N</i>-phenyl- (Diphenylamine)							
	MeOH	6.1×10^6	0.015		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	736061
	MeOH/ C ₆ H ₆ (67:33)		5.4×10^{-5}	298	CR/A'c-17	A' = Rub; used $\beta_{A'} = 1.4 \times 10^{-3}$ mol L ⁻¹ ; $^1\text{O}_2^*$ from (PhO) ₃ PO ₃ .	80M378
9.17 Aniline, <i>N,N</i>,2,4-tetramethyl-							
	MeOH/ H ₂ O (50:50)	1.3×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	83A094
9.18 Aniline, 4-(1,1,3,3-tetramethylbutyl)-<i>N</i>-[4-(1,1,3,3-tetramethylbutyl)phenyl]-							
	C ₆ H ₆ / EtOH (89:11)	2.6×10^4 (k_r)		295	CP/Pa,A'c-17	S = RB; A' = Tetr; used $k_r^{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 3.7 \times 10^{-4}$; P = nitroxide radicals.	757445
	C ₆ H ₆ / EtOH (89:11)	7.3×10^6		295	CP/A'c-23	S = RB; A' = Tetr; used $k_d = 3 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; A'' = Ni(II) dibutylthiocarbamate. used $k_{A''} = 1.6 \times 10^9$ L mol ⁻¹ s ⁻¹ .	757445
9.19 Aniline, <i>N,N</i>,4-trimethyl-							
	MeOH	1×10^8			PL/A'd-8	S = MB; A' = DPBF.	74F640
	MeOH		5.5×10^{-4}		CP/A'c-16	S = RB; A' = DPBF.	736061
	MeOH/ H ₂ O (50:50)	1.1×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	83A094

TABLE 9. Rate constants for the interaction of singlet oxygen with aromatic amines. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
9.20	Anthracenamine, N-methyl-9,10-diphenyl-						
	C ₆ H ₆	3.5×10^8	1.2×10^{-4}	298	CP/Ac-16	S = A; A' = 1,2-Dithiolane-3-pentanoic acid; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	757558
9.21	Benzoic acid, 4-(dimethylamino)-, ethyl ester						
	CHCl ₃	6×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80A041
9.22	Benzonitrile, 4-(dimethylamino)-						
	MeOH	5.7×10^5			PL/A'd-8	S = MB; A' = DPBF.	74F640
	MeOH		0.050		CP/A'c-16	S = RB; A' = DPBF.	736061
9.23	Biphenyl, 4,4'-diamino-						
	EtOH	3.0×10^9		273	CP/Oc-23	S = MB; A' = 2,5-DMF; used $k_d = 7.9 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^8$ L mol ⁻¹ s ⁻¹ ; k calcd. in [81Z251].	72F518
9.24	Naphthalene, 2-amino-						
	EtOH	9.6×10^8		273	CP/Oc-23	S = MB; A' = 2,5-DMF; used $k_d = 7.9 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^8$ L mol ⁻¹ s ⁻¹ ; k calcd. in [81Z251].	72F518
9.25	<i>o</i>-Phenylenediamine						
	EtOH	3.4×10^9		273	CP/Oc-23	S = MB; A' = 2,5-DMF; used $k_d = 7.9 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^8$ L mol ⁻¹ s ⁻¹ ; k calcd. in [81Z251].	72F518
9.26	<i>o</i>-Phenylenediamine, N,N,N',N'-tetramethyl-						
	MeOH/ H ₂ O (50:50)	1.1×10^9		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	83A094
9.27	<i>m</i>-Phenylenediamine, N,N,N',N'-tetramethyl-						
	MeOH/ H ₂ O (50:50)	7.4×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	83A094
9.28	<i>p</i>-Phenylenediamine, N-cyclohexyl-N'-phenyl-						
	EtOH	5.9×10^9		273	CP/Oc-20	S = MB; A' = 2,5-DMF; used $k_d = 7.9 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^8$ L mol ⁻¹ s ⁻¹ ; k calcd. in [81Z251].	72F518
	EtOH	1.1×10^{10}		273	CP/Oc-20	S = MB; A' = 2,5-DMF; used $k_{A'} = 2.3 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 47$; k calcd. in [81Z251].	72F518
9.29	<i>p</i>-Phenylenediamine, N,N'-diphenyl-						
	EtOH/ CHCl ₃ (50:50)	2.2×10^8		310	CR/LI-12	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	90A184
9.30	<i>p</i>-Phenylenediamine, N-(1-methylethyl)-N'-phenyl-						
	n-C ₁₆ H ₃₄	4.0×10^8		298	MD/A'c-33	A' = Rub; used $k_d = 9.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	EtOH	6.7×10^9		273	CP/Oc-23	S = MB; A' = 2,5-DMF; used $k_d = 7.9 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^8$ L mol ⁻¹ s ⁻¹ ; k calcd. in [81Z251].	72F518
	<i>i</i> -octane	2.1×10^8		298	CP/A'c-23	S = A' = Rub; used $k_d = 4.7 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	<i>i</i> -octane	3.3×10^8		CP/A'c-23	S = A' = Rub; used $k_d = 4 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	732066	
9.31	<i>p</i>-Phenylenediamine, N,N,N',N'-tetramethyl-						
	H ₂ O	$\sim 3 \times 10^9$ (k _r)			PL/Pa-14	S = Ery; used $k_d = 3.3 \times 10^5$ s ⁻¹ ; P = [TMPD] ⁺ ; used $\phi_{\Delta}(\text{Ery}) = 0.68$.	82A080
	MeOH	1×10^9			PL/A'd-8	S = MB; A' = DPBF.	74F640
	MeOH	1.5×10^{-4}			CP/A'c-16	S = RB; A' = DPBF.	736061

TABLE 9. Rate constants for the interaction of singlet oxygen with aromatic amines. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
9.31 <i>p</i>-Phenylenediamine, <i>N,N,N',N'</i>-tetramethyl- — Continued							
	MeOH/ H ₂ O (50:50)	1.8×10^9	1.5×10^{-4}	308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	83A094
9.32 Phenylhydrazine							
	C ₅ H ₅ N	1.4×10^8 (k_r)		PL/Ld-2	S = TPP, MPDEE or PdMP; meas. $\phi_{ox} = \phi_\Delta$.	80A398	
	CCl ₄	8.6×10^6		PL/Ld-2	S = MPDEE.	80A398	
	CH ₃ COCH ₃	9×10^7 (k_r)		PL/Ld-2	S = TPP, MPDEE or PdMP; meas. $\phi_{ox} = \phi_\Delta$.	80A398	
	CHCl ₃	7.5×10^6		PL/Ld-2	S = TPP, MPDEE or PdMP.	80A398	
9.33 Phthalazine-1,4-dione, 5-amino-2,3-dihydro-							
	H ₂ O pH = 10.1	9.3×10^8		CL/LI-12	S = RB; used $k_d = 2.4 \times 10^5$ s ⁻¹ .	90F502	
	H ₂ O pH = 7.1	1.4×10^9		CL/LI-12	S = RB; used $k_d = 2.4 \times 10^5$ s ⁻¹ .	90F502	

TABLE 10. Rate constants for the interaction of singlet oxygen with amino acids, peptides and proteins.

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
10.1 Alanine							
	D ₂ O pD = 8.4	2×10^6 (k_t)		293	CL/Ad-35	used $k_d = 2.9 \times 10^4$ s ⁻¹ ; high pressure O ₂ (0.195 mol L ⁻¹).	79A112
	D ₂ O pD = 8.1	1.3×10^6		295	CL/A'd-22	A' = BR ²⁻ ; used $k_d = 2.9 \times 10^4$ s ⁻¹ ; high pressure O ₂ ; recalc. [79A112].	757147
	H ₂ O pH = 10.6	3.0×10^7		310	CR/LI-12	used $k_d = 3.2 \times 10^5$ s ⁻¹ ; soln. cont. 5×10^{-4} mol L ⁻¹ CoCl ₂ , ¹ O ₂ * from autoxidation of oxytetracycline.	92M228
	H ₂ O/ MeOH (50:50)	$\leq 1 \times 10^7$			PL/A'd-5	S = MB; A' = DPBF.	72F516
10.2 β-Alanine							
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_t/k_t^{A'} = 0.82$; sensitizer immobilized on glass beads.	88R064
10.3 β-Alanine, L-histidyl-							
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_t/k_t^{A'} = 0.77$; sensitizer immobilized on glass beads.	88R064
10.4 Albumin							
	D ₂ O	2×10^8			PL/Ld-2	S = RF, water-sol. TPP derivs., or Chl a in micelles contg. Triton X-100; average for bovine or human serum albumin.	86F149
	D ₂ O pH = 7	5×10^8			PL/Ld-2	S = SnTPPSCl ₂ ⁴⁻ ; Soln. contg. 2×10^{-3} mol L ⁻¹ phosphate buffer and 1% NaCl wt/wt; human serum albumin.	90A022
10.5 Aposuperoxide dismutase							
	D ₂ O pD = 8.1	1.1×10^9		295	CP/A'd-22	S = MB; A' = BR ²⁻ ; used $k_d = 5.0 \times 10^4$ s ⁻¹ .	757147
	D ₂ O pD = 8.1	2.5×10^9		295	CL/A'd-22	A' = BR ²⁻ ; used $k_d = 5.0 \times 10^4$ s ⁻¹ ; high pressure O ₂ .	757147
10.6 Arginine							
	H ₂ O pH = 7.1	$\leq 1 \times 10^6$		298	CP/Oc-19	S = Phenosafranine; No measurable effect.	78A360
10.7 Carbonic anhydrase							
	D ₂ O pD = 8.1	8.0×10^8		295	CL/A'd-22	A' = BR ²⁻ ; used $k_d = 5.0 \times 10^4$ s ⁻¹ ; high pressure O ₂ .	757147
	D ₂ O pD = 8.1	6.5×10^8		295	CP/A'd-22	S = MB; A' = BR ²⁻ ; used $k_d = 5.0 \times 10^4$ s ⁻¹ .	757147
10.8 Cytochrome b							
	H ₂ O pH = 7.4	1.4×10^8			CP/Ac,Oc-18	S = RB; A' = His; used $k_d = 4.5 \times 10^5$ s ⁻¹ , $k_{A'} = 1 \times 10^8$ L mol ⁻¹ s ⁻¹ ; in submitochondria particles.	90R212
10.9 Cytochrome C (ferro)							
	CH ₃ CN/ D ₂ O (80:20)	4.9×10^8			PL/A'd-5	S = 2-ACN; A' = DPBF.	80R072
	H ₂ O pH = 7.4	5.6×10^8			CP/Ac,Oc-18	S = RB; A' = His; used $k_d = 4.5 \times 10^5$ s ⁻¹ , $k_{A'} = 1 \times 10^8$ L mol ⁻¹ s ⁻¹ ; in submitochondria particles.	90R212
10.10 Cytochrome C oxidase							
	H ₂ O pH = 7.4	1.0×10^8			CP/Ac,Oc-18	S = RB; A' = His; used $k_d = 4.5 \times 10^5$ s ⁻¹ , $k_{A'} = 1 \times 10^8$ L mol ⁻¹ s ⁻¹ ; in submitochondria particles.	90R212
10.11 Dehydrogenase, 3-hydroxybutyrate							
	H ₂ O pH = 7.4	6.1×10^8			CP/Ac,Oc-18	S = RB; A' = His; used $k_d = 4.5 \times 10^5$ s ⁻¹ , $k_{A'} = 1 \times 10^8$ L mol ⁻¹ s ⁻¹ ; in submitochondria particles.	90R212

TABLE 10. Rate constants for the interaction of singlet oxygen with amino acids, peptides and proteins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
10.11	Dehydrogenase, 3-hydroxybutyrate — Continued						
	H ₂ O pH = 7.2	1.2×10^9		295	CP/Ac,A'c-16	S = AlCl(tspc); A' = DPBF; estd. [¹ O ₂ *] from DPBF consumption in MeOH; k_d not given; enzyme inactivation rate in suspension of submitochondria particles.	87R074
10.12	Dehydrogenase, nicotinamide adenine dinucleotide (reduced)						
	H ₂ O pH = 7.4	1.7×10^8			CP/Ac,Oc-18	S = RB; A' = His; used $k_d = 4.5 \times 10^5$ s ⁻¹ , $k_{A'} = 1 \times 10^8$ L mol ⁻¹ s ⁻¹ ; in submitochondria particles.	90R212
10.13	Dehydrogenase, succinate						
	H ₂ O pH = 7.4	1.4×10^9			CP/Ac,Oc-18	S = RB; A' = His; used $k_d = 4.5 \times 10^5$ s ⁻¹ , $k_{A'} = 1 \times 10^8$ L mol ⁻¹ s ⁻¹ ; in submitochondria particles.	90R212
10.14	Dismutase, superoxide						
	D ₂ O pD = 8.1	8.2×10^8		295	CP/A'd-22	S = MR; A' = BR ²⁻ ; used $k_d = 5.0 \times 10^4$ s ⁻¹ .	757147
	D ₂ O pD = 8.1	2.6×10^9		295	CL/A'd-22	A' = BR ²⁻ ; used $k_d = 5.0 \times 10^4$ s ⁻¹ ; high pressure O ₂ .	757147
	H ₂ O pH = 7.1	2.7×10^9			CL/LI-12	S = RB; used $k_d = 2.4 \times 10^5$ s ⁻¹ .	90F502
10.15	Glycine						
	D ₂ O/ EtOH (75:25)	$<1 \times 10^5$		295	PL/Ld-2	S = RB.	94A113
10.16	Glycine, glycyl-						
	D ₂ O/ EtOH (75:25)	$<1 \times 10^5$		295	PL/Ld-2	S = RB.	94A113
10.17	Glycine, glycylglycyl-						
	D ₂ O/ EtOH (75:25)	$<1 \times 10^5$		295	PL/Ld-2	S = RB.	94A113
10.18	Glycine, glycylglycylglycyl-						
	D ₂ O/ EtOH (75:25)	$<1 \times 10^5$		295	PL/Ld-2	S = RB.	94A113
10.19	Glycine, glycyl-L-histidyl-						
	D ₂ O/ EtOH (75:25)	4.9×10^7		295	PL/Ld-2	S = RB.	94A113
10.20	Glycine, histidyl-						
	CH ₃ CN/ H ₂ O (50:50)	1.4×10^7			PL/Ld-2	S = RB or Eos.	93R059
	CH ₃ CN/ H ₂ O (50:50)	7.4×10^6 (k _r)			CP/Ac,A'c-17	S = RB; A' = DMA; $k_{A'}^{A'}$ not given.	93R059
	D ₂ O/ EtOH (75:25)	5.2×10^7		295	PL/Ld-2	S = RB.	94A113
	H ₂ O pH = 7	6.5×10^7			CP/A'c-18	S = RB; A' = TrpH; used $k_d = 2.5 \times 10^5$ s ⁻¹ , $k_{A'} = 6 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93R059
	H ₂ O pH = 7	6.6×10^7 (k _r)			CP/Oc-17	S = RB; A' = FFA; used $k_{A'}^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93R059
10.21	Glycine, tryptophyl-						
	EtOH/ CH ₃ CN (80:20)	1.0×10^7			PL/Ld-2	S = ZnTPP.	91A252
	EtOH/ CH ₃ CN (80:20)	7.4×10^5 (k _r)			CP/A'c-18	S = ZnTPP; A' = DMA; used $k_d = 5.3 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; used $k_A = 1.0 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A252
	H ₂ O pH = 7	3.3×10^7 (k _r)			CP/Oc-17	S = RB; A' = Met; used $k_{A'}^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A252

TABLE 10. Rate constants for the interaction of singlet oxygen with amino acids, peptides and proteins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
10.22 Glycine,L-tryptophylglycyl-							
	D ₂ O/ EtOH (75:25)	1.7 × 10 ⁷ (k_r)		295	CP/Ac-14,28	S = RB; used $k_d = 4.8 \times 10^4$ s ⁻¹ ; DMA as actinometer, k_r derived using $k_A = 3.1 \times 10^7$ L mol ⁻¹ s ⁻¹ .	94A113
	D ₂ O/ EtOH (75:25)	3.1 × 10 ⁷		295	PL/Ld-2	S = RB.	94A113
10.23 Histidine							
	D ₂ O/ EtOH (75:25)	4.6 × 10 ⁷		295	PL/Ld-2	S = RB.	94A113
	D ₂ O/ EtOH (75:25)	3.4 × 10 ⁷ (k_r)		295	CP/Ac-14,28	S = RB; used $k_d = 4.8 \times 10^4$ s ⁻¹ ; DMA as actinometer, k_r derived using $k_A = 4.6 \times 10^7$ L mol ⁻¹ s ⁻¹ .	94A113
	CH ₃ CN/ H ₂ O (50:50)	1.6 × 10 ⁷			PL/Ld-2	S = RB or Eos.	93R059
	CH ₃ CN/ H ₂ O (50:50)	8.1 × 10 ⁶ (k_r)			CP/Ac,A'c-17	S = RB; A' = DMA; $k_r^{A'}$ not given.	93R059
	D ₂ O pD = 7	4 × 10 ⁷			PL/Ld-2	S = Carboxyanthracene.	92E225
	D ₂ O pH = ~7	4.4 × 10 ⁷		293	PL/Ld-2	S = H ₂ TPPS ⁴ .	86A198
	D ₂ O pD = 7.4	6.1 × 10 ⁷			PL/A'd-5	S = MB; A' = ADPA.	81N048 80A205
	D ₂ O pD = 8.4	1.0 × 10 ⁸ (k_r)		293	CL/Ad-35	used $k_d = 2.9 \times 10^4$ s ⁻¹ ; high pressure O ₂ (0.195 mol L ⁻¹); unreactive at pH < 7.	79A112
	D ₂ O pD = 8.1	1.0 × 10 ⁸		295	CL/A'd-22	A' = BR ² ; used $k_d = 2.9 \times 10^4$ s ⁻¹ ; high pressure O ₂ ; recal. [79A112].	757147
	D ₂ O pD = 11.2	2.8 × 10 ⁸		310	CR/A'c-32	A' = DPBF; used $k_d = 1.5 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from DOPA/H ₂ O ₂ .	89M038
	D ₂ O pD = 11.2	9.3 × 10 ⁸		310	CR/A'c-32	A' = DPBF; used $k_d = 1.5 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from Dopamine/H ₂ O ₂ .	89M038
	D ₂ O (mic) pD = 9.3	6.7 × 10 ⁷			PL/A'd-5	S = 2-ACN; A' = DPBF; in vesicles (4.0 × 10 ⁻² mol L ⁻¹ DDAB).	82N027
	D ₂ O (mic) pD = 7.4	5.9 × 10 ⁷			PL/A'd-5	S = 2-ACN; A' = DPBF; micellar soln. 0.1 mol L ⁻¹ SDS or CTAB.	81N048
	H ₂ O pH = 10.6	5.7 × 10 ⁷		310	CR/LI-12	used $k_d = 3.2 \times 10^5$ s ⁻¹ ; soln. cont. 5 × 10 ⁻⁴ mol L ⁻¹ CoCl ₂ , ¹ O ₂ * from autoxidation of oxytetracycline.	92M228
	H ₂ O pH = 7	9.0 × 10 ⁷			CP/A'c-18	S = RB; A' = TrpH; used $k_d = 2.5 \times 10^5$ s ⁻¹ , $k_{A'} = 6 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93R059
	H ₂ O pH = 7	9.0 × 10 ⁷ (k_r)			CP/Oc-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93R059
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_r/k_r^{A'} = 0.94$; sensitizer immobilized on glass beads.	88R064
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_r/k_r^{A'} = 0.92$.	88R064
	H ₂ O pH = 7.1	3.2 × 10 ⁷		298	CP/Oc-19	S = Phenosafranine; Q = NaN ₃ ; used $k_Q = 2.0 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A360
	H ₂ O	5.4 × 10 ⁹	9.2 × 10 ⁻⁵	298	CP/Ac-15	S = MB; used $k_d = 5.0 \times 10^5$ s ⁻¹ .	78F061
	H ₂ O pH = 7.0		3.4 × 10 ⁻³	283	CP/Oc-15	S = MB; The mechanism of oxidation is not clear.	65F029
	H ₂ O pH = 8.0		2.9 × 10 ⁻³		CP/Oc-15	S = PF.	617008
	H ₂ O (mic)	6.7 × 10 ⁷			CP/A'c-16	S = 2-ACN; A' = DPBF; DDDAB liposomes.	85R008
	H ₂ O/ MeOH (50:50)	7 × 10 ⁶ (k_r)			CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_r^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; $k_r^{A'}$ in MeOH; k_r derived using $k_{TQ}^{A'} = 1 \times 10^6$ L mol ⁻¹ s ⁻¹ and $k_A = 5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	72F516

TABLE 10. Rate constants for the interaction of singlet oxygen with amino acids, peptides and proteins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
10.23	Histidine — Continued						
	H ₂ O/ MeOH (50:50)	5×10^7			PL/A'd-5	S = MB; A' = DPBF.	72F516
	MeOH	1.1×10^7		297	CR/P'a-16	A' = TEMP; k_d not given; formn. of TEMPO monitored by esr; soln. cont. MeONa and CoCl ₂ , ¹ O ₂ * from autoxidation of adrenaline.	92D227
10.24	L-Histidine, β -alanyl- (Carnosine)						
	D ₂ O pD = 7	3×10^7			PL/Ld-2	S = Carboxyanthracene.	92E225
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_r/k_r^{A'}$ = 0.72.	88R064
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_r/k_r^{A'}$ = 0.76; sensitizer immobilized on glass beads.	88R064
10.25	Histidine, β -alanyl-1-methyl- (Anserine)						
	D ₂ O pD = 7	3×10^7			PL/Ld-2	S = Carboxyanthracene.	92E225
10.26	L-Histidine, N-(4-amino-1-oxobutyl)- (Homocarnosine)						
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_r/k_r^{A'}$ = 0.91.	88R064
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_r/k_r^{A'}$ = 1.0; sensitizer immobilized on glass beads.	88R064
10.27	Histidine, glycyl-						
	CH ₃ CN/ H ₂ O (50:50)	9.2×10^6 (k_r)			CP/Ac,A'c-17	S = RB; A' = DMA; $k_r^{A'}$ not given.	93R059
	CH ₃ CN/ H ₂ O (50:50)	3.8×10^7			PL/Ld-2	S = RB or Eos.	93R059
	H ₂ O pH = 7	2.3×10^8			CP/A'c-18	S = RB; A' = TrpH; used $k_d = 1.5 \times 10^5$ s ⁻¹ , $k_{A'} = 6 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93R059
	H ₂ O pH = 7	8.3×10^7 (k_r)			CP/Oc-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93R059
10.28	L-Histidine, glycylglycyl-						
	D ₂ O/ EtOH (75:25)	5.6×10^7		295	PL/Ld-2	S = RB.	94A113
10.29	Luciferase						
	H ₂ O pH = 8.0	1.9×10^9 (k_r)		277	CP/Ac-17	S = RB; A' = His; used $k_r^{A'} = 2.8 \times 10^7$ L mol ⁻¹ s ⁻¹ ; sensitizer immobilized on glass beads.	86R129
10.30	Lysozyme						
	D ₂ O pD = 5.9	4.7×10^7 (k_r) 5.9×10^8 (k_q)		293	CP/Ac-14,28	S = Acridine Orange; used $k_d = 5.0 \times 10^4$ s ⁻¹ .	767105
	D ₂ O pD = 8.1	1.5×10^9		295	CL/A'd-22	A' = BR ²⁻ ; used $k_d = 5.0 \times 10^4$ s ⁻¹ ; high pressure O ₂ .	757147
	D ₂ O	1.3×10^9	4.0×10^{-5}	291	CP/Ac-15	S = 8-MOP; used $k_d = 5.0 \times 10^4$ s ⁻¹ .	757485
	D ₂ O	2.5×10^8 (k_r)			CP/Ac-28	S = Eos; used $k_d = 5.0 \times 10^4$ s ⁻¹ .	733045
	H ₂ O pH = 5.9	2.9×10^7 (k_r) 4.1×10^8 (k_q)		293	CP/Ac-14,28	S = Acridine Orange; used $k_d = 5.0 \times 10^5$ s ⁻¹ .	767105
	H ₂ O	1.3×10^8 (k_r)			CP/Ac-28	S = Eos; used $k_d = 5.0 \times 10^5$ s ⁻¹ .	733045
10.31	Phenylalanine						
	D ₂ O/ EtOH (75:25)	7×10^5		295	PL/Ld-2	S = RB.	94A113

TABLE 10. Rate constants for the interaction of singlet oxygen with amino acids, peptides and proteins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
10.32	Subtilisin BPN'						
	D ₂ O pH = 7.0	$3 \times 10^7 (k_r)$			CP/Ac-14,27	S = HP; used $k_d = 1.5 \times 10^4$ s ⁻¹ ; k_r estimated using $\phi_A(HP) = 0.75$.	83R008
10.33	Trypsin						
	H ₂ O pH = 7.0	8×10^9		298	CP/LI-12	S = Ret; used $k_d = 3.5 \times 10^5$ s ⁻¹ .	80R116
	H ₂ O pH = 8.0	7.1×10^9	7.0×10^{-5}	288	CP/Pa-15	S = MB and FMN; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; β calculated from data reported in [66F197].	73F659
10.34	Tryptamine						
	D ₂ O/ EtOH (75:25)	$2.7 \times 10^7 (k_r)$		295	CP/Ac-14,28	S = RB; used $k_d = 4.8 \times 10^4$ s ⁻¹ ; DMA as actinometer, k_r derived using $k_A = 5.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	94A113
	D ₂ O/ EtOH (75:25)	5.2×10^7		295	PL/Ld-2	S = RB.	94A113
	H ₂ O pH = 7	$7.8 \times 10^9 (k_r)$			CP/Oc-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93F191
	MeOH/ H ₂ O (50:50)	$9 \times 10^6 (k_r)$			CP/A'c-17	S = RB; A' = DPBF; used $k_r^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; derived $k_q = 4.6 \times 10^7$ L mol ⁻¹ s ⁻¹ using $k_A = 5.5 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A291
	MeOH/ H ₂ O (50:50)	5.5×10^7			CP/Ac-15	S = RB; used $k_d = 2.8 \times 10^5$ s ⁻¹ .	82A291
10.35	Tryptamine, N-methoxycarbonyl-						
	MeOH/ H ₂ O (50:50)	3.2×10^7		308	CR/Ac-31	used $k_q = 2.8 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	82A291
	MeOH/ H ₂ O (50:50)	3.2×10^7			CP/Ac-15	S = RB; used $k_d = 2.8 \times 10^5$ s ⁻¹ .	82A291
	MeOH/ H ₂ O (50:50)	$1.3 \times 10^7 (k_r)$		308	CR/A'c-17,31	A' = DPBF; used $k_r^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ ; derived $k_q = 1.9 \times 10^7$ L mol ⁻¹ s ⁻¹ using $k_A = 3.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A291
	MeOH/ H ₂ O (50:50)	$7 \times 10^6 (k_r)$			CP/A'c-17	S = RB; A' = DPBF; used $k_r^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; derived $k_q = 2.5 \times 10^7$ L mol ⁻¹ s ⁻¹ using $k_A = 3.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A291
10.36	Tryptophan						
	n-C ₇ H ₁₆ /H ₂ O (mic)	-3×10^7			PL/Ld-2	S = RB; reverse micelles contg. 0.25 mol L ⁻¹ AOT.	84N114
	MeOH/ D ₂ O (90:10)	1.3×10^7		293	PL/Ld-2	S = HP.	84F327
	MeOH/ H ₂ O (50:50)	3.8×10^7			CP/Ac,Oc-14,29	S = RB; A' = 2,5-DMF; used $k_d = 2.9 \times 10^5$ s ⁻¹ ; meas. $k_q/k_r = 2.8$.	83A397
	D ₂ O pH = ~7	$1.0 \times 10^7 (k_r)$			PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	86A198
	D ₂ O	$2.8 \times 10^7 (k_q)$			PL/Ld-2	S = RF, H ₂ TPPS ⁴⁻ , or Chl in Triton X-100 micelles.	86F149
	D ₂ O	5.6×10^7			PL/Ld-2	S = RF, H ₂ TPPS ⁴⁻ , or Chl in Triton X-100 micelles.	86F149
	D ₂ O	$~5 \times 10^7$			PL/Ld-2	S = RF, H ₂ TPPS ⁴⁻ , or Chl in Triton X-100 micelles.	86F149
	D ₂ O	$3.6 \times 10^7 (k_r)$		293	CP/Ac-14	S = HP; k_d not given; used $\phi_A(HP) = 0.41$.	84F327
	D ₂ O pD = 7.4	7.2×10^7		293	PL/Ld-2	S = HP.	84F327
	D ₂ O pD = 7.4	6.1×10^7			PL/A'd-5	S = MB; A' = ADPA.	81N048
	D ₂ O pD = 8.4	$3 \times 10^7 (k_r)$		293	CL/Ad-35	used $k_q = 2.9 \times 10^4$ s ⁻¹ ; high pressure O ₂ (0.195 mol L ⁻¹).	79A112
	D ₂ O pD = 8.1	5.1×10^7		295	CL/A'd-22	A' = BR ²⁻ ; used $k_d = 2.9 \times 10^4$ s ⁻¹ ; high pressure O ₂ ; recalc. [79A112].	75T147

TABLE 10. Rate constants for the interaction of singlet oxygen with amino acids, peptides and proteins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
10.36 Tryptophan — Continued							
	D ₂ O/ MeOH (70:30)	6.0×10^7		293	PL/Ld-2	S = HP.	84F327
	D ₂ O/ HCONH ₂ (70:30)	5.2×10^7		293	PL/Ld-2	S = HP.	84F327
	D ₂ O (mic) pD = 7.4	4.2×10^7			PL/A'd-5	S = 2-ACN; A' = DPBF; 0.1 mol L ⁻¹ CTAB.	81N048
	D ₂ O (mic) pD = 7.4	4.0×10^7			PL/A'd-5	S = 2-ACN; A' = DPBF; 0.1 mol L ⁻¹ SDS.	81N048
	D ₂ O/ EtOH (75:25)	1.3×10^7 (k_r)		295	CP/Ac-14,28	S = RB; used $k_d = 4.8 \times 10^4$ s ⁻¹ ; DMA as actinometer, k_r derived using $k_A = 3.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	94A113
	D ₂ O/ EtOH (75:25)	3.2×10^7		295	PL/Ld-2	S = RB.	94A113
	EtOH	$\leq 5.0 \times 10^6$			PL/A'd-5	S = MB; A' = DPBF.	78A338
	EtOH/ CH ₃ CN (80:20)	6.3×10^6			PL/Ld-2	S = ZnTPP.	91A252
	EtOH/ CH ₃ CN (80:20)	4.8×10^5 (k_r)			CP/Ac,A'c-14,28	S = ZnTPP; A' = DMA; used $k_d = 5.3 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; k_r derived using $k_A = 6.3 \times 10^6$ L mol ⁻¹ s ⁻¹ .	91A252
	H ₂ O pH = 10.6	2.6×10^7		310	CR/LJ-12	used $k_d = 3.2 \times 10^5$ s ⁻¹ ; soln. cont. 5×10^{-4} mol L ⁻¹ CoCl ₂ , ¹ O ₂ * from autoxidation of oxytetracycline.	92M228
	H ₂ O pH = 7	3.2×10^7 (k_r)			CP/Oc-17	S = RB; A' = Met; used $k_r^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A252
	H ₂ O	5.9×10^7 (k_r)		298	CP/Ac-14	S = H ₂ TPPS ⁴⁻ ; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; $E_a = 15.9$ kJ mol ⁻¹ ; log(A) = 10.6; studied at 288-318 K; used $\phi_A(S) = 0.7$, phosphate buffer contg. 1.6% NaCl.	91R053
	H ₂ O pH = 7	6.0×10^7 5×10^6 (k_r)			CP/Pa-14	S = GaCl(tspc); used $k_d = 2.4 \times 10^5$ s ⁻¹ ; P = Hydroperoxytryptophan; k_r estd. assuming $\phi_A(S) = 0.5$.	87F104
	H ₂ O pH = 7.4	6.6×10^7 (k_r)		293	CP/Ac-14	S = HP; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; k_d not given; used $\phi_A(HP) = 0.41$.	84F327
	H ₂ O pH = 7.0		1.8×10^{-3}	283	CP/Oc-15	S = MB; The mechanism of oxidation is not clear.	65F029
	H ₂ O/ MeOH (50:50)	9.0×10^7			CP/Pa-14	S = RB or GaCl(tspc); used $k_d = 2.8 \times 10^5$ s ⁻¹ ; P = Hydroperoxytryptophan.	87F104
	H ₂ O/ MeOH (50:50) pH = 10				CP/A'c-17	S = RB; A' = His; meas. $k_r/k_r^{A'} = 0.6$; sensitizer immobilized on glass beads.	83F166
	H ₂ O/ MeOH (50:50)	7.1×10^7			CP/Ac-15	S = RB; used $k_d = 2.8 \times 10^5$ s ⁻¹ .	82A291
	H ₂ O/ MeOH (50:50)	3×10^6 (k_r)			CP/A'c-17	S = RB; A' = DPBF; used $k_r^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; derived $k_q = 6.8 \times 10^7$ L mol ⁻¹ s ⁻¹ using $k_A = 7.1 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A291
	H ₂ O/ MeOH (50:50)	2.1×10^7		308	CR/Ac-31	used $k_d = 2.8 \times 10^5$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	82A291
	H ₂ O/ MeOH (50:50)	5×10^6 (k_r)		308	CR/A'c-17,31	A' = DPBF; used $k_r^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ ; derived $k_q = 1.6 \times 10^7$ L mol ⁻¹ s ⁻¹ , using $k_A = 2.1 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A291
	H ₂ O/ MeOH (50:50)	3.0×10^7			PL/A'd-5	S = MB; A' = DPBF.	78A338
	H ₂ O/ MeOH (50:50)	4×10^7			PL/A'd-5	S = MB; A' = DPBF.	72F516

TABLE 10. Rate constants for the interaction of singlet oxygen with amino acids, peptides and proteins. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
10.36	Tryptophan — Continued						
	H ₂ O/ MeOH (50:50)	4×10^6 (k_t)			CP/Ac,A'c-17	$S = MB; A' = DPBF$; used $k_t^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; k_t in MeOH; k_t derived using $k_{TQ}^{A'} = 2.0 \times 10^9$ L mol ⁻¹ s ⁻¹ and $k_A = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	72F516
	HCONH ₂ / D ₂ O (90:10)	8.7×10^7		293	PL/Ld-2	S = HP.	84F327
	HCONHCH ₃	1.3×10^8			PL/A'd-5	S = MB; A' = DPBF.	78A338
	MeOH	6×10^6			PL/A'd-5	S = MB; A' = DPBF.	78A338
10.37	Tryptophan, N-acetyl-						
	D ₂ O/ EtOH (75:25)	4.2×10^7		295	PL/Ld-2	S = RB.	94A113
10.38	Tryptophan, N-acetyl-, methyl ester						
	D ₂ O/ EtOH (75:25)	3.2×10^7		295	PL/Ld-2	S = RB.	94A113
	MeOH/ H ₂ O (50:50)	6×10^6 (k_t)		308	CR/A'c-17,31	A' = DPBF; used $k_t^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; ${}^1O_2^*$ from MNPO ₂ ; derived $k_q = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ , using $k_A = 2.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A291
	MeOH/ H ₂ O (50:50)	4×10^6 (k_t)			CP/A'c-17	S = RB; A' = DPBF; used $k_t^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; derived $k_q = 5.9 \times 10^7$ L mol ⁻¹ s ⁻¹ using $k_A = 6.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A291
	MeOH/ H ₂ O (50:50)	2.4×10^7		308	CR/Ac-31	used $k_d = 2.8 \times 10^5$ s ⁻¹ ; ${}^1O_2^*$ from MNPO ₂ .	82A291
	MeOH/ H ₂ O (50:50)	6.3×10^7			CP/Ac-15	S = RB; used $k_d = 2.8 \times 10^5$ s ⁻¹ .	82A291
10.39	L-Tryptophan, L-alanyl-						
	D ₂ O/ EtOH (75:25)	2.7×10^7 (k_t)		295	CP/Ac-14,28	S = RB; used $k_d = 4.8 \times 10^4$ s ⁻¹ ; DMA as actinometer, k_t derived using $k_A = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ .	94A113
	D ₂ O/ EtOH (75:25)	4.0×10^7		295	PL/Ld-2	S = RB.	94A113
10.40	L-Tryptophan, ethyl ester						
	D ₂ O/ EtOH (75:25)	3.1×10^7		295	PL/Ld-2	S = RB.	94A113
10.41	Tryptophan, glycyl-						
	EtOH/ CH ₃ CN (80:20)	1.2×10^7			PL/Ld-2	S = ZnTPP.	91A252
	EtOH/ CH ₃ CN (80:20)	9.8×10^5 (k_t)			CP/Ac,A'c-14,28	S = ZnTPP; A' = DMA; used $k_d = 5.3 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; k_t derived using $k_A = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A252
	H ₂ O pH = 7	5.1×10^7 (k_t)			CP/Oc-17	S = RB; A' = Met; used $k_t^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91A252
	H ₂ O pH = 7	$\sim 2 \times 10^8$	2×10^{-3}		CP/LJ-12	S = Ery; used $k_d = 3 \times 10^5$ s ⁻¹ .	90F511
10.42	Tryptophan, methyl ester						
	MeOH/ H ₂ O (50:50)	4×10^6 (k_t)			CP/A'c-17	S = RB; A' = DPBF; used $k_t^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; derived $k_q = 3.5 \times 10^7$ using $k_A = 3.9 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A291
	MeOH/ H ₂ O (50:50)	3.9×10^7			CP/Ac-15	S = RB; used $k_d = 2.8 \times 10^5$ s ⁻¹ .	82A291
10.43	Tryptophanamide						
	D ₂ O/ EtOH (75:25)	3.1×10^7		295	PL/Ld-2	S = RB.	94A113
10.44	L-Tryptophanamide, N-acetyl-						
	D ₂ O/ EtOH (75:25)	3.1×10^7		295	PL/Ld-2	S = RB.	94A113

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes.

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.1	Acridine, 3,6-diamino- (Proflavine)						
	D ₂ O pH = 5.9	3×10^8			CP/A'c-25	S = A; A' = Lysozyme; used $k_d = 5 \times 10^4$ s ⁻¹ , $k_{A'} = 6.4 \times 10^8$ L mol ⁻¹ s ⁻¹ ; Authors suggest that values are high due to quenching of sensitizer triplet.	81R011
	H ₂ O pH = 5.9	7×10^8			CP/A'c-25	S = A; A' = Lysozyme; used $k_d = 5 \times 10^5$ s ⁻¹ , $k_{A'} = 4.4 \times 10^8$ L mol ⁻¹ s ⁻¹ .	81R011
11.2	Acridinium, 3,6-diamino-10-methyl- (Acriflavine)						
	D ₂ O pH = 5.9	3×10^8			CP/A'c-25	S = A; A' = Lysozyme; used $k_d = 5 \times 10^4$ s ⁻¹ , $k_{A'} = 6.4 \times 10^8$ L mol ⁻¹ s ⁻¹ .	81R011
	H ₂ O pH = 5.9	9×10^8			CP/A'c-25	S = A; A' = Lysozyme; used $k_d = 5 \times 10^5$ s ⁻¹ , $k_{A'} = 4.4 \times 10^8$ L mol ⁻¹ s ⁻¹ .	81R011
11.3	9,10-Anthraquinone, 1-amino-						
	2-PrOH/ H ₂ O (83:17)			295	CR/Ac-32	A' = 1,2-AQ(NH ₂) ₂ ; meas. $k_A/k_{A'} = 0.02$; ${}^1\text{O}_2^*$ from NaOCl/H ₂ O ₂ .	88F582
11.4	9,10-Anthraquinone, 2-amino-						
	2-PrOH/ H ₂ O (83:17)			295	CR/Ac-32	A' = 1,2-AQ(NH ₂) ₂ ; meas. $k_A/k_{A'} = 0.15$; ${}^1\text{O}_2^*$ from NaOCl/H ₂ O ₂ .	88F582
11.5	9,10-Anthraquinone, 1-amino-4-hydroxy-						
	CCl ₄	4×10^6 $<10^5$ (k_r)			MP/LI-12,27	S = TPP; used $k_d = 36$ s ⁻¹ .	82A384
	CH ₃ COCH ₃			294	CP/Ac-19	S = A; A' = [(CH ₃) ₂ NCS ₂] ₂ Ni; meas. $k_A/k_{A'} = 0.061$.	80F634
	2-PrOH/ H ₂ O (83:17)			295	CR/Ac-32	A' = 1,2-AQ(NH ₂) ₂ ; meas. $k_A/k_{A'} = 0.55$; ${}^1\text{O}_2^*$ from NaOCl/H ₂ O ₂ .	88F582
11.6	9,10-Anthraquinone, 1-amino-2-methyl-						
	CCl ₄	1×10^5 $<10^5$ (k_r)			MP/LI-12,27	S = TPP; used $k_d = 36$ s ⁻¹ .	82A384
11.7	9,10-Anthraquinone, 1,4-bis(methylamino)-						
	CH ₃ COCH ₃			294	CP/Ac-19	S = A; A' = [(CH ₃) ₂ NCS ₂] ₂ Ni; meas. $k_A/k_{A'} = 0.12$.	80F634
11.8	9,10-Anthraquinone, 1,4-diamino-						
	C ₅ H ₅ N	1.4×10^9			PL/Ld-2	S = An.	87E959
	C ₅ H ₅ N	2×10^9			CP/A'c-16	S = TPBC; A' = Tetr; used $k_d = 1.6 \times 10^5$ s ⁻¹ .	85F675
	C ₆ H ₆	2.0×10^8			PL/Ld-2	S = An.	87E959
	C ₆ H ₆	3×10^8			CP/A'c-16	S = TPBC; A' = Tetr; used $k_d = 3.0 \times 10^5$ s ⁻¹ .	85F675
	CCl ₄	1×10^8 $<10^5$ (k_r)			MP/LI-12,27	S = TPP; used $k_d = 36$ s ⁻¹ .	82A384
	CH ₃ COCH ₃	6.2×10^8			PL/Ld-2	S = An.	87E959
	CH ₃ COCH ₃	9×10^8			CP/A'c-16	S = TPBC; A' = Tetr; used $k_d = 5.1 \times 10^5$ s ⁻¹ .	85F675
	2-PrOH/ H ₂ O (83:17)			295	CR/Ac-32	A' = 1,2-AQ(NH ₂) ₂ ; meas. $k_A/k_{A'} = 0.04$; ${}^1\text{O}_2^*$ from NaOCl/H ₂ O ₂ .	88F582
11.9	9,10-Anthraquinone, 1,5-diamino-						
	2-PrOH/ H ₂ O (83:17)			295	CR/Ac-32	A' = 1,2-AQ(NH ₂) ₂ ; meas. $k_A/k_{A'} = 0.15$; ${}^1\text{O}_2^*$ from NaOCl/H ₂ O ₂ .	88F582
11.10	9,10-Anthraquinone, 1,8-diamino-						
	2-PrOH/ H ₂ O (83:17)			295	CR/Ac-32	A' = 1,2-AQ(NH ₂) ₂ ; meas. $k_A/k_{A'} = 0.13$; ${}^1\text{O}_2^*$ from NaOCl/H ₂ O ₂ .	88F582

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.11	9,10-Anthraquinone, 2,6-diamino-						
	2-PrOH/H ₂ O (83:17)			295	CR/Ac-32	A' = 1,2-AQ(NH ₂) ₂ ; meas. $k_A/k_{A'} = 0.33$; $^1\text{O}_2^*$ from NaOCl/H ₂ O ₂ .	88F582
11.12	9,10-Anthraquinone, 1,5-diaminobromo-4,8-dihydroxy-						
	CCl ₄	1×10^7 $<10^5$ (k_t)			MP/LI-12,27	S = TPP; used $k_d = 36$ s ⁻¹ .	82A384
11.13	9,10-Anthraquinone, 1,4-diamino-2-methoxy-						
	CCl ₄	1×10^8 $<10^5$ (k_t)			MP/LI-12,27	S = TPP; used $k_d = 36$ s ⁻¹ .	82A384
	2-PrOH/H ₂ O (83:17)			295	CR/Ac-32	A' = 1,2-AQ(NH ₂) ₂ ; meas. $k_A/k_{A'} = 0.05$; $^1\text{O}_2^*$ from NaOCl/H ₂ O ₂ .	88F582
11.14	9,10-Anthraquinone, 1,4,5,8-tetraamino-						
	2-PrOH/H ₂ O (83:17)			295	CR/Ac-32	A' = 1,2-AQ(NH ₂) ₂ ; meas. $k_A/k_{A'} = 0.14$; $^1\text{O}_2^*$ from NaOCl/H ₂ O ₂ .	88F582
11.15	Benzene, 1-bromo-4-(diazophenylmethyl)-						
	CH ₃ CN	9.1×10^8 (k_t)			CP/Ac,A'c-17	S = MB; A' = DDM; used $k_t^{A'} = 1.4 \times 10^9$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 0.65$.	777113
11.16	Benzene, 1,1'-(diazomethylene)bis- (Diazodiphenylmethane, DDM)						
	CH ₃ CN	$\sim 6 \times 10^8$		300	PL/Pb-5	S = MB; P = Benzophenone oxide.	84A339
	CH ₃ CN	1.1×10^9	2.9×10^{-5}		CP/Ac-15	S = MB; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	777113
	CH ₃ CN	1.4×10^9 (k_t)			CP/Ac,A'c-17	S = MB; A' = DMA; used $k_t^{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 8.0$.	777113
	CH ₃ CN	2.4×10^9 (k_t)			CR/Ac,A'c-17	A' = DMA; used $k_t^{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 14$; $^1\text{O}_2^*$ from (PhO) ₃ PO ₃ .	777113
	CHCl ₃	1.1×10^9 (k_t)			CR/Ac,A'c-17	A' = DMA; used $k_t^{A'} = 9.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 12$; $^1\text{O}_2^*$ from (PhO) ₃ PO ₃ .	777113
	CHCl ₃	8.3×10^8	2×10^{-5}		CP/Ac-15	S = MB; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	777113
	MeOH	6.7×10^8	2.1×10^{-4}		CP/Ac-15	S = MB; used $k_d = 1.4 \times 10^4$ s ⁻¹ .	777113
11.17	Benzene, 1,1'-(diazomethylene)bis[4-chloro-						
	CH ₃ CN	7.3×10^8 (k_t)			CP/Ac,A'c-17	S = MB; A' = DMA; used $k_t^{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 4.3$.	777113
	CH ₃ CN	6.6×10^8	5×10^{-5}		CP/Ac-15	S = MB; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	777113
	CH ₃ CN	7.5×10^8 (k_t)			CP/Ac,A'c-17	S = MB; A' = DDM; used $k_t^{A'} = 1.4 \times 10^9$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 0.54$.	777113
	MeOH	3.8×10^8 (k_t)			CP/Ac,A'c-17	S = MB; A' = DDM; used $k_t^{A'} = 6.7 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 0.58$.	777113
11.18	Benzene, 1,1'-(diazomethylene)bis[4-methyl-						
	CH ₃ CN	2.3×10^9 (k_t)			CP/Ac,A'c-17	S = MB; A' = DDM; used $k_t^{A'} = 1.4 \times 10^9$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 1.6$.	777113
	MeOH	9.4×10^8 (k_t)			CP/Ac,A'c-17	S = MB; A' = DDM; used $k_t^{A'} = 6.7 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 1.4$.	777113
11.19	Benzene, 1,1'-(diazomethylene)bis[2,4,6-trimethyl-						
	CH ₃ CN	1.2×10^7			PL/Pb-5	S = MB; P = 2,4,6,2',4',6'- Hexamethylbenzophenone oxide.	89A105
11.20	Benzene, 1-methoxy-4-(diazophenylmethyl)-						
	CH ₃ CN	2.0×10^9 (k_t)			CP/Ac,A'c-17	S = MB; A' = DDM; used $k_t^{A'} = 1.4 \times 10^9$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 1.4$.	777113

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.21	1,4-Benzenediamine, N^4 -(6-amino-3-phenylpyrazolo[5,1-c]-1,2,4-triazol-7-ylidene)- N^1,N^1 -diethyl-3-methyl-						
	EtOH	$4.3 \times 10^5 (k_r)$			CP/Ac-17	S = AlCl(tspc); A' = Azo dye 10; used $k_r^{A'} = 2.4 \times 10^6$ L mol ⁻¹ s ⁻¹ ; values from graph.	91A448
11.22	1,4-Benzenediamine, N^1,N^1 -diethyl-3-methyl- N^4 -(6-methyl-3-phenylpyrazolo[5,1-c]-1,2,4-triazol-7-ylidene)-				CP/Ac-17	S = AlCl(tspc); A' = Azo dye 10; used $k_r^{A'} = 2.4 \times 10^6$ L mol ⁻¹ s ⁻¹ ; values from graph.	91A448
	EtOH	$3.5 \times 10^5 (k_r)$					
11.23	1,4-Benzenediamine, N^1,N^1 -diethyl-3-methyl- N^4 -(6-(phenylamino)-3-phenylpyrazolo[5,1-c]-1,2,4-triazol-7-ylidene)-				CP/Ac-17	S = AlCl(tspc); A' = Azo dye 10; used $k_r^{A'} = 2.4 \times 10^6$ L mol ⁻¹ s ⁻¹ ; values from graph.	91A448
	EtOH	$8.5 \times 10^5 (k_r)$					
11.24	1,4-Benzenediamine, N^1,N^1 -diethyl-3-methyl- N^4 -(3-phenylpyrazolo[5,1-c]-1,2,4-triazol-7-ylidene)-				CP/Ac-17	S = AlCl(tspc); A' = Azo dye 10; used $k_r^{A'} = 2.4 \times 10^6$ L mol ⁻¹ s ⁻¹ ; values from graph.	91A448
	EtOH	$2.7 \times 10^5 (k_r)$					
11.25	1,4-Benzenediamine, N^1,N^1 -diethyl-3-methyl- N^4 -(3-phenyl-6-(2-thienyl)pyrazolo[5,1-c]-1,2,4-triazol-7-ylidene)-				CP/Ac-17	S = AlCl(tspc); A' = Azo dye 10; used $k_r^{A'} = 2.4 \times 10^6$ L mol ⁻¹ s ⁻¹ ; values from graph.	91A448
	EtOH	$3.2 \times 10^5 (k_r)$					
11.26	Benzene propanamide, α -[(4-diethylamino)-2,6-dimethylphenyl]imino]- β -oxo- <i>N</i> -phenyl-				CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
	C ₅ H ₅ N	6.3×10^7					
11.27	Benzene propanamide, α -[(4-diethylamino)-2-methylphenyl]imino]- β -oxo- <i>N</i> -phenyl-				CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
	C ₅ H ₅ N	7.9×10^6					
11.28	Benzene propanamide, α -[(4-diethylamino)phenyl]imino]- β -oxo- <i>N</i> -phenyl-				CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
	C ₅ H ₅ N	4.0×10^6					
	C ₅ H ₅ N	1.6×10^7			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
11.29	Benz[e]indolium, 2-[7-(1,3-dihydro-1,1-dimethyl-3-(sulfobutyl)benz[e]indol-2-ylidene)-1,3,5-heptatrienyl]-1,1-dimethyl-3-(sulfobutyl)-, hydroxide, inner salt, Na salt						
	DMSO	$2.5 \times 10^5 (k_r)$	0.061	295	CL/Ac-15	S = Th; used $k_d = 1.3 \times 10^4$ s ⁻¹ ; k_r derived using $\phi_{\Delta}(S) = 0.87$.	82F531
	MeOH	$1.5 \times 10^6 (k_r)$	0.11	295	CL/Ac-15	S = Th; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_{\Delta}(S) = 0.87$.	82F531
11.30	Benz[e]indolium, 2-[2-[3-[[1,3-dihydro-1,1-dimethyl-3-(3-sulfopropyl)benz[e]indol-2-ylidene]ethylidene]-2-[4-(ethoxycarbonyl)-1-piperidinyl]-1-cyclopenten-1-yl]ethenyl]-[-1,1-dimethyl-3-(3-sulfopropyl)-, hydroxide, inner salt, compound with triethyl				CL/Ac-15	S = Th; used $k_d = 1.3 \times 10^4$ s ⁻¹ ; k_r derived using $\phi_{\Delta}(S) = 0.87$.	82F531
	DMSO	$1.7 \times 10^6 (k_r)$	9×10^{-3}	295	CL/Ac-15	S = Th; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_{\Delta}(S) = 0.87$.	82F531
	MeOH	$1.0 \times 10^7 (k_r)$	0.016	295	CL/Ac-15	S = Th; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_{\Delta}(S) = 0.87$.	82F531
11.31	1-Benzopyrylium, 8-[5-(6,7-dihydro-2,4-diphenyl-1-benzopyran-8-yl)-2,4-pentadienyliidene]-5,6,7,8-tetrahydro-2,4-diphenyl-, perchlorate				PL/A'd-5	S = MB; A' = DPBF.	727260
	CH ₃ CN	3×10^{10}					
11.32	Benzothiazolium, 2-[[2-butoxy-3-[(3-ethyl-2-benzothiazolylidene)methyl]-4-oxo-2-cyclobuten-1-ylidene]methyl]-3-ethyl-				CP/Pa-20	S = A; Q = Ni[S ₂ C ₂ (C ₆ H ₅) ₂]; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	90F554
	CHCl ₃	1.4×10^9		293			

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.33 Benzothiazolium, 5-chloro-2-[2-[3-[(5-chloro-3-ethyl-1-benzothiazolylidene)ethyldiene]-2-(diphenylamino)-1-cyclopenten-1-yl]ethenyl]-3-ethyl-, perchlorate							
	Propylene carbonate	$9.8 \times 10^9 (k_q)$		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690
	Propylene carbonate	$1.6 \times 10^8 (k_r)$		293	PL/A'd-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
	DMSO	$3.8 \times 10^6 (k_p)$	3.9×10^{-3}	295	CL/Ac-15	S = Th; used $k_d = 1.3 \times 10^4$ s ⁻¹ ; k_r derived using $\phi_{\Delta}(S) = 0.87$.	82F531
	MeOH	$1.4 \times 10^7 (k_r)$	0.011	295	CL/Ac-15	S = Th; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_{\Delta}(S) = 0.87$.	82F531
11.34 Benzothiazolium, 2-[7-(5-chloro-1,3,3-trimethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)-3-methyl-, iodide							
	Propylene carbonate	$<1 \times 10^7 (k_q)$		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690
	Propylene carbonate	$3 \times 10^7 (k_r)$		293	PL/A'd-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
11.35 Benzothiazolium, 2-[7-(5-fluoro-1,3,3-trimethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)-3-methyl-, iodide							
	Propylene carbonate	$7 \times 10^7 (k_r)$		293	PL/A'd-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
	Propylene carbonate	$2.0 \times 10^8 (k_q)$		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690
11.36 Benzothiazolium, 2-[7-(5-methoxy-1,3,3-trimethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)-3-methyl-, iodide							
	Propylene carbonate	$1.2 \times 10^9 (k_q)$		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690
	Propylene carbonate	$1.7 \times 10^8 (k_r)$		293	PL/A'd-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
11.37 Benzothiazolium, 3-methyl-2-[7-(1,3,3-trimethyl-2-indol-2-ylidene)-1,3,5-heptatrienyl]-, iodide							
	Propylene carbonate	$1.7 \times 10^8 (k_r)$		293	PL/A'd-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
	Propylene carbonate	$2.2 \times 10^6 (k_q)$		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690
11.38 Benzothiazolium, 2-[7-(1,3,3,5-tetramethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)-3-methyl-, iodide							
	Propylene carbonate	$2.5 \times 10^8 (k_q)$		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690
	Propylene carbonate	$6 \times 10^7 (k_r)$		293	PL/A'd-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
11.39 3-Benzoxazolepropanesulfonic acid, 2-[4-(1,3-dibutyltetrahydro-4,6-dioxo-2-thioxo-5-pyrimidinylidene)-2-butenyldiene]-, sodium salt (Merocyanine 540)							
	CD ₃ OD	2.1×10^9		294	PL/Ld-2	S = An.	91R017
	D ₂ O (ves)	5.6×10^9			CP/Ac-15	S = A; used $k_d = 4.0 \times 10^4$ s ⁻¹ ; soln. contg. DLPC.	92A123
	H ₂ O/ MeOH (50:50)	$3.0 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = A; A' = DPBF; $k_r^{A'}$ not given.	92A123
	H ₂ O/ MeOH (50:50)	4.5×10^9			CP/Ac-15	S = A; used $k_d = 9.1 \times 10^4$ s ⁻¹ .	92A123
	H ₂ O (ves)	$9.3 \times 10^7 (k_r)$			CP/Ac,A'c-17	S = A; A' = DPBF; $k_r^{A'}$ not given; soln. contg. DLPC.	92A123
	H ₂ O (ves)	5.8×10^9			CP/Ac-15	S = A; used $k_d = 4.0 \times 10^4$ s ⁻¹ ; soln. contg. DLPC.	92A123
11.40 2,3'-Biindolium, 1,1,3-trimethyl-2'-(2-(1,3,3-trimethylindolylidene)methyl)-, iodide							
	MeOH	$\leq 3.0 \times 10^4 (k_r)$			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_{\Delta}(S) = 0.75$.	767071

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.41	2,2'-Carbocyanine, 1,1'-diethyl-, chloride						
	MeOH	$3.4 \times 10^7 (k_r)$			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_{\Delta}(S) = 0.75$.	767071
11.42	4,4'-Carbocyanine, 1,1'-diethyl-						
	MeOH	2.0×10^9			PL/Ld-2	S = T(m-HOP)P.	90R164
11.43	4,4'-Carbocyanine, 1,1'-diethyl-, toluenesulfonate						
	CD ₃ OD	$1.6 \times 10^8 (k_r)$			CP/Ac-15	S = 2-ACN; used $k_d = 1.3 \times 10^4$ s ⁻¹ ; k_r derived using $\phi_{\Delta}(S) = 0.75$.	767071
	MeOH	$1.1 \times 10^8 (k_r)$			CP/Ac-15	S = 2-ACN, RB, Py, and fluorenone; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using ϕ_{Δ} for each S, k_r is an average for the 4 sensitizers.	767071
11.44	Crystal Violet						
	CH ₃ CN		5.8		CP/Ac-15	S = A.	82F525
11.45	2,2'-Cyanine, 1,1'-diethyl-, iodide						
	MeOH	$\sim 4.0 \times 10^5 (k_r)$			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_{\Delta}(S) = 0.75$.	767071
11.46	Cyclobutendiylium, 1,3-bis[4-(diethylamino)-2-hydroxyphenyl]-2,4-dihydroxy-						
	CHCl ₃	1.2×10^9	1.4×10^{-5}	293	CP/Pa-20	S = A; A' = Ni[S ₂ C ₂ (C ₆ H ₅) ₂] ₂ ; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	90F554
11.47	Cyclobutendiylium, 1,3-bis[4-(dimethylamino)-2-methylphenyl]-2,4-dihydroxy-						
	CHCl ₃	2.2×10^9		293	CP/Pa-20	S = A; A' = Ni[S ₂ C ₂ (C ₆ H ₅) ₂] ₂ ; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	90F554
11.48	Cyclobutendiylium, 1,3-bis[4-(N-methyl-N-octadecylamino)-2-methylphenyl]-2,4-dihydroxy-						
	CHCl ₃	1.4×10^9		293	CP/Pa-20	S = A; A' = Ni[S ₂ C ₂ (C ₆ H ₅) ₂] ₂ ; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	90F554
11.49	2,5-Cyclohexadien-1-one, 4-[[4-(diethylamino)-2,5-dimethylphenyl]imino]-3,5-dimethyl-						
	C ₅ H ₅ N	1.0×10^9			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
11.50	Diazene, 1-[4-N-(2-cyanoethyl)-N-ethylaminophenyl]-2-(4-methoxyphenyl)-						
	CH ₂ Cl ₂	$2.7 \times 10^4 (k_r)$			CP/Ac,A'c-17	S = MB; A' = DMA; used $k_r^{A'} = 2.1 \times 10^7$ L mol ⁻¹ s ⁻¹ in benzene [747312].	89F024
	CH ₂ Cl ₂	2×10^6			PL/Ld-2	S = TPP.	89F024
11.51	Diazene, 1-[4-N-(2-cyanoethyl)-N-ethylaminophenyl]-2-phenyl-						
	CH ₂ Cl ₂	2×10^6			PL/Ld-2	S = TPP.	89F024
	CH ₂ Cl ₂	$2.9 \times 10^4 (k_r)$			CP/Ac,A'c-17	S = MB; A' = DMA; used $k_r^{A'} = 2.1 \times 10^7$ L mol ⁻¹ s ⁻¹ in benzene [747312].	89F024
11.52	Diazene, 1-(4-diethylaminophenyl)-2-(2,4-dinitrophenyl)-						
	CH ₂ Cl ₂	1×10^6			PL/Ld-2	S = MB.	89F024
11.53	Diazene, 1-(4-diethylaminophenyl)-2-(4-methoxyphenyl)-						
	CD ₃ OD				CP/Ac-17	S = MB; A' = 4-(Diethylamino)azobenzene; meas. $k_r/k_r^{A'} = 14$; Rel. to A' in MeOH.	84F350
	CD ₃ OD/ D ₂ O (80:20)				CP/Ac-17	S = MB; A' = 4-(Diethylamino)azobenzene; meas. $k_r/k_r^{A'} = 74$; Rel. to A' in MeOH.	84F350
	CH ₂ Cl ₂	$1.3 \times 10^4 (k_r)$			CP/Ac,A'c-17	S = MB; A' = DMA; used $k_r^{A'} = 2.1 \times 10^7$ L mol ⁻¹ s ⁻¹ in benzene [747312].	89F024
	CH ₂ Cl ₂	1.0×10^7			PL/Ld-2	S = TPP.	89F024

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.53 Diazene, 1-(4-diethylaminophenyl)-2-(4-methoxyphenyl). — Continued							
	MeOH/ H ₂ O (80:20)				CP/Ac-17	S = MB; A' = 4-(Diethylamino)azobenzene; meas. $k_t/k_t^{A'}$ = 6; Rel. to A' in MeOH.	84F350
	MeOH/ H ₂ O (50:50)				CP/Ac-17	S = MB; A' = 4-(Diethylamino)azobenzene; meas. $k_t/k_t^{A'}$ = 22; Rel. to A' in MeOH.	84F350
	MeOH				CP/Ac-17	S = MB; A' = 4-(Diethylamino)azobenzene; meas. $k_t/k_t^{A'}$ = 1.	84F350
11.54 Diazene, 1-(4-diethylaminophenyl)-2-(3-nitrophenyl).							
	CH ₂ Cl ₂	3×10^6			PL/Ld-2	S = TPP.	89F024
11.55 Diazene, 1-(4-diethylaminophenyl)-2-(4-nitrophenyl).							
	CH ₂ Cl ₂	4×10^6			PL/Ld-2	S = MB.	89F024
	CH ₂ Cl ₂	1.7×10^4 (k_t)			CP/Ac,A'c-17	S = MB; A' = DMA; used $k_t^{A'} = 2.1 \times 10^7$ L mol ⁻¹ s ⁻¹ in benzene [747312].	89F024
11.56 Diazene, 1-(4-diethylaminophenyl)-2-phenyl-							
	CD ₃ OD				CP/Ac-17	S = MB; $k_t^{A'}/k_t^A$ (MeOH) = 11.	84F350
	CD ₃ OD/ D ₂ O (80:20)				CP/Ac-17	S = MB; $k_t^{A'}/k_t^A$ (MeOH) = 44.	84F350
	CH ₂ Cl ₂	1.1×10^4 (k_t)			CP/Ac,A'c-17	S = MB; A' = DMA; used $k_t^{A'} = 2.1 \times 10^7$ L mol ⁻¹ s ⁻¹ in benzene [747312].	89F024
	CH ₂ Cl ₂	6.5×10^6			PL/Ld-2	S = TPP.	89F024
	MeOH/ H ₂ O (50:50)				CP/Ac-17	S = MB; $k_t^{A'}/k_t^A$ (MeOH) = 24.	84F350
	MeOH/ H ₂ O (80:20)				CP/Ac-17	S = MB; $k_t^{A'}/k_t^A$ (MeOH) = 7.	84F350
11.57 Diazene, diphenyl-							
	CH ₂ Cl ₂	$<1 \times 10^5$			PL/Ld-2	S = TPP.	89F024
11.58 Diazene, 1-[4-N-ethyl-N-(2-hydroxyethyl)aminophenyl]-2-(4-methoxyphenyl).							
	CH ₂ Cl ₂	9×10^6			PL/Ld-2	S = TPP.	89F024
	CH ₂ Cl ₂	1.3×10^4 (k_t)			CP/Ac,A'c-17	S = MB; A' = DMA; used $k_t^{A'} = 2.1 \times 10^7$ L mol ⁻¹ s ⁻¹ in benzene [747312].	89F024
11.59 Diazene, 1-(4-methoxy-1-naphthyl)-2-phenyl-							
	CH ₂ Cl ₂	1.5×10^6			PL/Ld-2	S = TPP.	89F024
	CH ₂ Cl ₂	1.4×10^4 (k_t)			CP/Ac,A'c-17	S = MB; A' = DMA; used $k_t^{A'} = 2.1 \times 10^7$ L mol ⁻¹ s ⁻¹ in benzene [747312].	89F024
11.60 Diazene, 1-(4-methylphenyl)-2-[1-(phenylaminocarbonyl)-2-oxopropyl]-							
	DMAA	4.0×10^6 (k_t)			CP/A'c-17	S = RB; A' = DMA; used $k_t^{A'} = 1.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86F622
	DMF	3.0×10^6 (k_t)			CP/A'c-17	S = RB; A' = DMA; used $k_t^{A'} = 1.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86F622
11.61 Diazene, 1-(4-nitrophenyl)-2-phenyl-							
	CH ₂ Cl ₂	$<1 \times 10^5$			PL/Ld-2	S = TPP.	89F024
11.62 Diazene, 1-phenyl-2-[1-(phenylaminocarbonyl)-2-oxopropyl]-							
	DMAA	3×10^5 (k_t)			CP/A'c-17	S = RB; A' = DMA; used $k_t^{A'} = 1.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86F622
	DMF	1×10^5 (k_t)			CP/A'c-17	S = RB; A' = DMA; used $k_t^{A'} = 1.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86F622
	2-PrOH	3.2×10^6 (k_t)			CP/A'c-17	S = RB; A' = DMA; used $k_t^{A'} = 1.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86F622

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.63	Fluorene, 9-diazo-						
	CH ₃ CN	6.6×10^7		300	PL/Pb-5	S = MB; P = 9-Fluorenone oxide.	85A206
	CH ₃ CN	1.0×10^8 (k_t)			CP/Ac,A'c-17	S = MB; A' = DMA; used $k_t^{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 0.59$.	777113
	CH ₃ CN	1.0×10^8	3.3×10^{-4}		CP/Ac-15	S = MB; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	777113
	CHCl ₃	5.1×10^7	3×10^{-4}		CP/Ac-15	S = MB; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	777113
	CHCl ₃	6.8×10^7 (k_t)			CR/Ac,A'c-17	A' = DMA; used $k_t^{A'} = 9.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_t/k_t^{A'} = 0.74$; ${}^1O_2^*$ from (PhO) ₃ PO ₃ .	777113
	MeOH	3.3×10^7	4.2×10^{-3}		CP/Ac-15	S = MB; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	777113
11.64	Fluorescein, 2',7'-dibromo-4'-(hydroxymercuri)-, disodium salt (Merbromin)						
	CH ₃ CN/H ₂ O (50:50)	2.0×10^8			PL/Ld-2	S = A.	93F069
11.65	Fluorescein dianion, 2',4',5',7'-tetrabromo- (Eosin)						
	CH ₃ CN	7.2×10^7			CP/Oc-19	S = A; A' = 2M2P; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	84F191
	CH ₃ CN/H ₂ O (50:50)	2.7×10^7			PL/Ld-2	S = A.	93F069
	MeOH	2.4×10^7			CP/Oc-19	S = A; A' = 2M2P; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	84F191
11.66	Fluorescein dianion, 3,4,5,6-tetrachloro-2',4',5',7'-tetraiodo- (Rose Bengal, RB)						
	CD ₃ COCD ₃	9.9×10^6			PL/Ld-2	S = A.	89E324
	CH ₃ CN	5.0×10^7			CP/Oc-19	S = A; A' = 2M2P; used $k_d = 1.7 \times 10^4$ s ⁻¹ .	84F191
	CH ₃ CN/H ₂ O (50:50)	1.5×10^7			PL/Ld-2	S = A.	93F069
	MeOH	2.0×10^7			CP/Oc-19	S = A; A' = 2M2P; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	84F191
11.67	3,5-Heptanedione, 4-[[2-ethyl-4-(diethylamino)phenyl]imino]-2,2,6,6-tetramethyl-						
	C ₅ H ₅ N	$\sim 1.6 \times 10^7$			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
11.68	Indene-1,3-dione, 2-(2-quinolylidene)- (Quinophthalone)						
	CH ₃ COCH ₃			294	CP/Ac-19	S = A; A' = [(CH ₃) ₂ NCS ₂] ₂ Ni; meas. $k_A/k_{A'} = 0.068$.	80F634
11.69	Indigo						
	CHCl ₃				CP/Ac-17	S = A; A' = [(CH ₃) ₂ NCS ₂] ₂ Ni; meas. $k_t/k_t^{A'} = -0.025$.	79F582
11.70	Indolium, 2-[7-[4-bromo-1-(1,3,3-trimethyl-2-indol-2-ylidene)]-1,3,5-heptatrienyl]-1,3,3-trimetetrafluoroborate						
	Propylene carbonate	2.7×10^8 (k_q)		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_t/k_t^{A'}$.	87E690
	Propylene carbonate	3.0×10^8 (k_t)		293	PL/A'd-17	S = MB; A' = DPBF; used $k_t^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
11.71	Indolium, 2-[7-[4-chloro-1-(1,3,3-trimethyl-2-indol-2-ylidene)]-1,3,5-heptatrienyl]-1,3,3-trimethyl-, tetrafluoroborate						
	Propylene carbonate	3.6×10^8 (k_q)		293	PL/A'd-17	S = MB; A' = DPBF; used $k_t^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
	Propylene carbonate	2.6×10^8 (k_t)		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_t/k_t^{A'}$.	87E690
11.72	Indolium, 2-[7-[4-iodo-1-(1,3,3-trimethyl-2-indol-2-ylidene)]-1,3,5-heptatrienyl]-1,3,3-trimethyl-, iodide						
	Propylene carbonate	1.5×10^9 (k_q)		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_t/k_t^{A'}$.	87E690
	Propylene carbonate	2.6×10^8 (k_t)		293	PL/A'd-17	S = MB; A' = DPBF; used $k_t^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.73	Indolium, 2-[7-[4-iodo-1-(1,3,3-trimethyl-2-ylidene)-1,3,5-heptatrienyl]-1,3,3-trimethyl-, tetrafluoroborate						
	Propylene carbonate	$1.1 \times 10^8 (k_r)$		293	PL/A'd-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
11.74	Indolium, 2-[7-(1,3,3-trimethyl-2-ylidene)-1-[4-(2,2-dimethoxyethyl)-1,3,5-heptatrienyl]-1,3,3-trimethyl-, tetrafluoroborate						
	Propylene carbonate	$8 \times 10^7 (k_r)$		293	PL/A'd-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
	Propylene carbonate	$8.9 \times 10^8 (k_q)$		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690
11.75	Indolium, 2-[7-(1,3,3-trimethyl-2-ylidene)-1-[4-(1-piperidinio)-1,3,5-heptatrienyl]-1,3,3-trimethyl-, bis(tetrafluoroborate)						
	Propylene carbonate	$1.4 \times 10^9 (k_r)$		293	PL/A'd-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
	Propylene carbonate	$1.9 \times 10^9 (k_q)$		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690
11.76	Indolium, 2-[7-(1,3,3-trimethyl-2-ylidene)-1-[4-[3-(1,3,3-trimethyl-2-indolylidene)-2-propenyl]-1,3,5-heptatrienyl]-1,3,3-trimethyl-, tetrafluoroborate						
	Propylene carbonate	$2.0 \times 10^9 (k_q)$		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690
	Propylene carbonate	$3.8 \times 10^8 (k_r)$		293	PL/A'd-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
11.77	Leucomalachite Green						
	C ₆ H ₅ CH ₃	$<1 \times 10^5 (k_r)$		298	CP/Ac,A'c-17	S = A' = 9-Phenylanthracene.	79F148
	C ₆ H ₅ CH ₃	2×10^8	3×10^{-4}	298	CP/Ac-18	S = A' = DMA; used $k_d = 5.3 \times 10^4$ s ⁻¹ , $\beta_{A'} = 1.5 \times 10^{-3}$ mol L ⁻¹ ; meas. $\beta_A/\beta_{A'} = 0.2$.	79F148
11.78	Methanesulfonamide, N-[2-[(4-(1,5-dihydro-3-methyl-5-oxo-1-phenylpyrazol-4-ylidene)amino)methylphenyl]ethylamino]ethyl-						
	MeOH	$3 \times 10^7 (k_r)$		298	CP/Ac,A'c-	S = MB; A' = DPA; used $k_r^{A'} = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87F542
		$5 \times 10^6 (k_q)$		17,19			
11.79	2-Naphthalenecarboxamide, 4-[(4-(aminocarbonyl)phenyl]azo]-3-hydroxy-N-(2-methoxyphenyl)-						
	DMAA	7.0×10^6		CP/A'c-17	S = RB; A' = DMA; used $k_r^{A'} = 1.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86F622	
	DMF	3.6×10^6		CP/A'c-17	S = RB; A' = DMA; used $k_r^{A'} = 1.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86F622	
11.80	2-Naphthalenecarboxamide, 3-hydroxy-4-[(4-methylphenyl)azo]-N-phenyl-						
	DMAA	2.0×10^6		CP/A'c-17	S = RB; A' = DMA; used $k_r^{A'} = 1.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86F622	
	DMF	1.9×10^6		CP/A'c-17	S = RB; A' = DMA; used $k_r^{A'} = 1.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86F622	
11.81	1-Naphthalenone, 4-[(2,6-dimethyl-4-(diethylamino)phenyl]imino]-2-(2-methoxy-5-fluorosulfonylphenyl)aminocarbonyl-						
	C ₅ H ₅ N	1.4×10^9		CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166	
	C ₆ H ₆	4.7×10^9		CP/Oc-20	S = azine; A' = 2M2P; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.053$ mol L ⁻¹ ; No evidence of chemical reaction.	757166	
	CH ₃ CN	3.9×10^{10}		CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.3 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.018$ mol L ⁻¹ ; No evidence of chemical reaction.	757166	

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k ($\text{L mol}^{-1} \text{s}^{-1}$)	$\beta (k_d/k)$ (mol L^{-1})	T (K)	Method	Comment	Ref.
11.82	1-Naphthalenone, 4-[[4-(diphenylamino)phenyl]imino]-						
	$\text{C}_5\text{H}_5\text{N}$	$\leq 3.2 \times 10^6$			CP/Oc-20	$S = RB; A' = 2M2P$; used $k_d = 3.1 \times 10^4 \text{ s}^{-1}$, $\beta_{A'} = 0.043 \text{ mol L}^{-1}$; No evidence of chemical reaction.	757166
11.83	1-Naphthalenone, 4-[(<i>N</i> -methyl- <i>N</i> -phenyl)amino]imino-						
	CH_2Cl_2	8.9×10^7			PL/Ld-2	$S = MB$.	89F024
	CH_2Cl_2	$2 \times 10^3 (k_r)$			CP/Ac,A'c-17	$S = MB; A' = DMA$; used $k_r^{A'} = 2.1 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$ in benzene [747312].	89F024
11.84	1-Naphthalenone, 4-[[4-(phenylamino)phenyl]imino]-						
	$\text{C}_5\text{H}_5\text{N}$	$\leq 4.0 \times 10^7$			CP/Oc-20	$S = RB; A' = 2M2P$; used $k_d = 3.1 \times 10^4 \text{ s}^{-1}$, $\beta_{A'} = 0.043 \text{ mol L}^{-1}$; No evidence of chemical reaction.	757166
11.85	1-Naphthol, 5-[[3-(aminosulfonyl)phenyl]sulfonylamino]-4-[2-(methylsulfonyl)-4-nitrophenylazo]-, conjugate base						
	DMF	$9.3 \times 10^6 (k_r)$ $5.6 \times 10^7 (k_q)$	7.7×10^{-3}		CP/Ac-14	$S = RB$; used $k_d = 7.1 \times 10^4 \text{ s}^{-1}$; k_q includes possible back energy transfer; k_r derived using $\phi_{\Delta}(S) = 0.4$; soln. contains NH_4OH .	79F412
	H_2O	$1.3 \times 10^8 (k_r)$	$2.5 \times 10^{-3} (\beta_r)$		CP/Ac-14	$S = A$; used $k_d = 3.3 \times 10^5 \text{ s}^{-1}$; k_r derived using $\phi_{\Delta}(S) = 1.3 \times 10^{-4}$; soln. contains NH_4OH .	79F412
	MeOH	$4.8 \times 10^6 (k_r)$ $7.2 \times 10^7 (k_q)$			CP/Ac-14	$S = A$; used $k_d = 1.3 \times 10^5 \text{ s}^{-1}$; k_q includes possible back energy transfer; k_r derived using $\phi_{\Delta}(S) = 1.6 \times 10^{-3}$; soln. contains NH_4OH .	79F412
11.86	1-Naphthol, 5-[[3-(aminosulfonyl)phenyl]sulfonylamino]-4-[2-(methylsulfonyl)-4-nitrophenylazo]-, conjugate dibase						
	DMF		2.6×10^{-3}		CP/Ac-14	$S = RB$; soln. contains NH_4OH .	79F412
	DMF	$2.7 \times 10^7 (k_r)$			CP/Ac-16	$S = A; A' = DPBF$; used $k_d = 7.1 \times 10^4 \text{ s}^{-1}$; soln. contains NH_4OH .	79F412
	H_2O	$5 \times 10^8 (k_r)$	$7 \times 10^{-4} (\beta_r)$		CP/Ac-16	$S = A; A' = 2,5\text{-DMF}$; used $k_d = 3.3 \times 10^5 \text{ s}^{-1}$; soln. contains NH_4OH .	79F412
	MeOH		1.3×10^{-3}		CP/Ac-14	$S = RB$; soln. contains NH_4OH .	79F412
	MeOH	$1.0 \times 10^7 (k_r)$			CP/Ac-16	$S = A; A' = DPBF$; used $k_d = 1.3 \times 10^5 \text{ s}^{-1}$; soln. contains NH_4OH .	79F412
11.87	1-Naphthol, 4-(4-chlorophenylazo)-						
	MeOH			293	CP/Ac-17	$S = MB; A' = 4\text{-Phenylazo-1-naphthol}$; meas. $k_r/k_r^{A'} = 0.40$.	81A403
11.88	1-Naphthol, 5-methoxy-4-[2-(methylsulfonyl)-4-nitrophenyl]azo-, conjugate base						
	DMF	$4.3 \times 10^7 (k_r)$			CP/Ac-14	$S = RB$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$; k_r derived using $\phi_{\Delta}(S) = 0.4$; soln. contains NH_4OH .	79F412
	DMF	$4.6 \times 10^7 (k_r)$ $2.6 \times 10^8 (k_q)$			CP/Ac-14	$S = A$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$; k_q includes possible back energy transfer; k_r derived using $\phi_{\Delta}(S) = 2.7 \times 10^{-4}$; soln. contains NH_4OH .	79F412
11.89	1-Naphthol, 4-(4-methoxyphenylazo)-						
	MeOH			293	CP/Ac-17	$S = MB; A' = 4\text{-Phenylazo-1-naphthol}$; meas. $k_r/k_r^{A'} = 1.9$.	81A403
11.90	1-Naphthol, 4-(4-methylphenylazo)-						
	CD_3OD				CP/Ac-17	$S = MB; A' = 4\text{-}(Diethylamino)azobenzene$; meas. $k_r/k_r^{A'} = 2.5 \times 10^2$; Rel. to A' in MeOH.	84F350
	$\text{CD}_3\text{OD}/\text{D}_2\text{O}$ (80:20)				CP/Ac-17	$S = MB; A' = 4\text{-}(Diethylamino)azobenzene$; meas. $k_r/k_r^{A'} = 5.3 \times 10^2$; Rel. to A' in MeOH.	84F350
	CH_2Cl_2	$8.1 \times 10^4 (k_r)$			CL/Ac,A'c-17	$S = MB; A' = DMA$; used $k_r^{A'} = 2 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	89F024

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.90 1-Naphthol, 4-(4-methylphenylazo)- — Continued							
	CH ₂ Cl ₂	1.3×10^7			PL/Ld-2	S = TPP; charge-transfer; $k = 2.6 \times 10^7$ L mol ⁻¹ s ⁻¹ normalized to 100% hydrazone form.	89F024
11.91 1-Naphthol, 4-(4-nitrophenylazo)-							
	MeOH/H ₂ O (80:20)				CP/Ac-17	S = MB; A' = 4-(Diethylamino)azobenzene; meas. $k_t/k_t^{A'} = 50$; Rel. to A' in MeOH.	84F350
	MeOH/H ₂ O (50:50)				CP/Ac-17	S = MB; A' = 4-(Diethylamino)azobenzene; meas. $k_t/k_t^{A'} = 1.6 \times 10^2$; Rel. to A' in MeOH.	84F350
	MeOH				CP/Ac-17	S = MB; A' = 4-(Diethylamino)azobenzene; meas. $k_t/k_t^{A'} = 26$.	84F350
	MeOH			293	CP/Ac-17	S = MB; A' = 4-Phenylazo-1-naphthol; meas. $k_t/k_t^{A'} = 1.8$.	81A403
11.92 1-Naphthol, 4-phenylazo-							
	CH ₂ Cl ₂	2.1×10^7			PL/Ld-2	S = MB; charge-transfer; $k = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ normalized to 100% hydrazone form.	89F024
	CH ₂ Cl ₂	2.2×10^4 (k_t)			CL/Ac,A'c-17	S = MB; A' = DMA; used $k_t^{A'} = 2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	89F024
11.93 2-Naphthol, 1-(4-aminophenylazo)-							
	CH ₃ COCH ₃			294	CP/Ac-19	S = A; A' = [(CH ₃) ₂ NCS ₂] ₂ Ni; meas. $k_A/k_{A'} = 0.67$.	80F634
11.94 2-Naphthol, 1-(4-bromophenylazo)-							
	MeOH			293	CP/Ac-17	S = MB; A' = 1-Phenylazo-2-naphthol; meas. $k_t/k_t^{A'} = 0.58$.	81A403
11.95 2-Naphthol, 1-(4-chlorophenylazo)-							
	MeOH			293	CP/Ac-17	S = MB; A' = 1-Phenylazo-2-naphthol; meas. $k_t/k_t^{A'} = 0.54$.	81A403
11.96 2-Naphthol, 1-[4-(dimethylamino)phenylazo]-							
	MeOH			293	CP/Ac-17	S = MB; A' = 1-Phenylazo-2-naphthol; meas. $k_t/k_t^{A'} = 3.8$.	81A403
11.97 2-Naphthol, 1-(4-fluorophenylazo)-							
	MeOH			293	CP/Ac-17	S = MB; A' = 1-Phenylazo-2-naphthol; meas. $k_t/k_t^{A'} = 0.69$.	81A403
11.98 2-Naphthol, 1-(4-hydroxyphenylazo)-							
	MeOH			293	CP/Ac-17	S = MB; A' = 1-Phenylazo-2-naphthol; meas. $k_t/k_t^{A'} = 1.6$.	81A403
11.99 2-Naphthol, 1-(4-iodophenylazo)-							
	MeOH			293	CP/Ac-17	S = MB; A' = 1-Phenylazo-2-naphthol; meas. $k_t/k_t^{A'} = 0.50$.	81A403
11.100 2-Naphthol, 1-(4-methoxyphenylazo)-							
	CH ₂ Cl ₂	3.5×10^6			PL/Ld-2	S = MB; charge-transfer; $k = 4.5 \times 10^6$ L mol ⁻¹ s ⁻¹ normalized to 100% hydrazone form.	89F024
	MeOH			293	CP/Ac-17	S = MB; A' = 1-Phenylazo-2-naphthol; meas. $k_t/k_t^{A'} = 1.6$.	81A403
11.101 2-Naphthol, 1-(4-methyl-2-nitrophenylazo)-							
	DMAA	6×10^5			CP/A'c-17	S = RB; A' = DMA; used $k_t^{A'} = 1.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86F622

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k ($\text{L mol}^{-1} \text{s}^{-1}$)	$\beta (k_d/k)$ (mol L^{-1})	T (K)	Method	Comment	Ref.
11.101 2-Naphthol, 1-(4-methyl-2-nitrophenylazo)- — Continued							
	DMF	$\leq 2 \times 10^5$			CP/A'c-17	$S = \text{RB}; A' = \text{DMA}; \text{used } k_t^{A'} = 1.4 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	86F622
	2-PrOH	$\leq 7 \times 10^5$			CP/A'c-17	$S = \text{RB}; A' = \text{DMA}; \text{used } k_t^{A'} = 1.4 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	86F622
11.102 2-Naphthol, 1-(4-methylphenylazo)-							
	CD ₃ OD				CP/Ac-17	$S = \text{MB}; A' = 4\text{-}(Diethylamino)azobenzene; \text{meas. } k_t/k_t^{A'} = 2.7 \times 10^2; \text{ Rel. to } A' \text{ in MeOH.}$	84F350
	CD ₃ OD/ D ₂ O (80:20)				CP/Ac-17	$S = \text{MB}; A' = 4\text{-}(Diethylamino)azobenzene; \text{meas. } k_t/k_t^{A'} = 3.5 \times 10^2; \text{ Rel. to } A' \text{ in MeOH.}$	84F350
	CH ₂ Cl ₂	3×10^6			PL/Ld-2	$S = \text{MB}; \text{charge-transfer}; k = 4 \times 10^6 \text{ L mol}^{-1} \text{ s}^{-1}$ normalized to 100% hydrazone form.	89F024
	MeOH/ H ₂ O (50:50)				CP/Ac-17	$S = \text{MB}; A' = 4\text{-}(Diethylamino)azobenzene; \text{meas. } k_t/k_t^{A'} = 42; \text{ Rel. to } A' \text{ in MeOH.}$	84F350
	MeOH/ H ₂ O (80:20)				CP/Ac-17	$S = \text{MB}; A' = 4\text{-}(Diethylamino)azobenzene; \text{meas. } k_t/k_t^{A'} = 16; \text{ Rel. to } A' \text{ in MeOH.}$	84F350
	MeOH			293	CP/Ac-17	$S = \text{MB}; A' = 1\text{-Phenylazo-2-naphthol}; \text{meas. } k_t/k_t^{A'} = 1.3.$	81A403
	MeOH				CP/Ac-17	$S = \text{MB}; A' = 4\text{-}(Diethylamino)azobenzene; \text{meas. } k_t/k_t^{A'} = 6.$	84F350
11.103 2-Naphthol, 1-(4-nitrophenylazo)-							
	MeOH			293	CP/Ac-17	$S = \text{MB}; A' = 1\text{-Phenylazo-2-naphthol}; \text{meas. } k_t/k_t^{A'} = 0.15.$	81A403
	DMAA	1.1×10^6			CP/A'c-17	$S = \text{RB}; A' = \text{DMA}; \text{used } k_t^{A'} = 1.4 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	86F622
	DMF	1×10^7			CP/A'c-17	$S = \text{RB}; A' = \text{DMA}; \text{used } k_t^{A'} = 1.4 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	86F622
	2-PrOH	5.5×10^6			CP/A'c-17	$S = \text{RB}; A' = \text{DMA}; \text{used } k_t^{A'} = 1.4 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	86F622
11.104 2-Naphthol, 1-phenylazo-							
	CH ₂ Cl ₂	$2.0 \times 10^4 (k_t)$			CL/Ac,A'c-17	$S = \text{MB}; A' = \text{DMA}; \text{used } k_t^{A'} = 2 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	89F024
	CH ₂ Cl ₂	2×10^6			PL/Ld-2	$S = \text{MB}; \text{charge-transfer}; k = 2.5 \times 10^6 \text{ L mol}^{-1} \text{ s}^{-1}$ normalized to 100% hydrazone form.	89F024
11.105 Naphtho[1,2-d]thiazolium, 1-ethyl-2-[(1-ethyl)naphtho[1,2-d]thiazol-2-ylidene)methyl]-, chloride							
	MeOH	$4.0 \times 10^5 (k_t)$			CP/Ac-15	$S = 2\text{-ACN}; \text{used } k_d = 1.9 \times 10^5 \text{ s}^{-1}; k_t \text{ derived using } \phi_{\Delta}(S) = 0.75.$	767071
11.106 Naphtho[1,2-d]thiazolium, 1-ethyl-2-[(1-ethyl)naphtho[1,2-d]thiazol-2-ylidene)methyl]-1-butenyl-, bromide							
	MeOH	$7.3 \times 10^7 (k_t)$			CP/Ac-15	$S = 2\text{-ACN}; \text{used } k_d = 1.9 \times 10^5 \text{ s}^{-1}; k_t \text{ derived using } \phi_{\Delta}(S) = 0.75.$	767071
11.107 Naphtho[1,2-d]thiazolium, 1-ethyl-2-[(1-ethyl)naphtho[1,2-d]thiazol-2-ylidene)-1-propenyl]-, toluenesulfonate							
	MeOH	$1.6 \times 10^7 (k_t)$			CP/Ac-15	$S = 2\text{-ACN}; \text{used } k_d = 1.9 \times 10^5 \text{ s}^{-1}; k_t \text{ derived using } \phi_{\Delta}(S) = 0.75.$	767071
11.108 Naphtho[2,3-d]thiazolium, 2-[2-[2-(diphenylamino)-3-[3-(4-methoxy-4-oxobutyl)naphtho[2,3-d]thiazol-2-ylidene]ethylidene]1-cyclopenten-1-yl]ethenyl]-3-(4-methoxy-4-oxobutyl)-, perchlorate							
	DMSO	$2.3 \times 10^6 (k_t)$	6×10^{-3}		CL/Ac-15	$S = \text{Th}; \text{used } k_d = 1.3 \times 10^4 \text{ s}^{-1}; k_t \text{ derived using } \phi_{\Delta}(S) = 0.87.$	82F531
	MeOH	$1.4 \times 10^7 (k_t)$	0.012		CL/Ac-15	$S = \text{Th}; \text{used } k_d = 1.4 \times 10^5 \text{ s}^{-1}; k_t \text{ derived using } \phi_{\Delta}(S) = 0.87.$	82F531

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.109	2,2'-Oxatricarbocyanine, 3,3'-diethyl-, iodide						
	Propylene carbonate	$3.2 \times 10^8 (k_r)$		293	PL/A'd-17	$S = MB; A' = DPBF; \text{used } k_t^{A'} = 1.0 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	87E690
	Propylene carbonate	$9 \times 10^8 (k_q)$		293	PL/A'd-5	$S = MB; A' = DPBF; \text{Measured } k_q + k_r, k_q \text{ calcd. knowing } k_t \text{ from } k_t/k_t^{A'}$	87E690
11.110	Pentadienylum, 1,1,5,5-tetrakis[4-(diethylamino)phenyl]-						
	CH ₃ CN/ C ₆ H ₆ (80:20)	1×10^8 $1 \times 10^5 (k_r)$ $6.6 \times 10^7 (k_q)$		298	CP/Ac,A'c-23	$S = A' = \text{Rub}; \text{used } k_d = 3.4 \times 10^4 \text{ s}^{-1}, k_{A'} = 4.2 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$	90F360
11.111	Pantanamide, 2-[(4-diethylamino)-2-methylphenyl]imino]-4,4-dimethyl-3-oxo-N-phenyl-	C ₅ H ₅ N	2.5×10^7		CP/Oc-20	$S = RB; A' = 2M2P; \text{used } k_d = 3.1 \times 10^4 \text{ s}^{-1}, \beta_{A'} = 0.043 \text{ mol L}^{-1}; \text{No evidence of chemical reaction.}$	757166
11.112	Pantanamide, 2-[(4-diethylamino)phenyl]imino]-4,4-dimethyl-3-oxo-N-phenyl-	C ₅ H ₅ N	7.9×10^6		CP/Oc-20	$S = RB; A' = 2M2P; \text{used } k_d = 3.1 \times 10^4 \text{ s}^{-1}, \beta_{A'} = 0.043 \text{ mol L}^{-1}; \text{No evidence of chemical reaction.}$	757166
11.113	Phenothiazinium, 3,7-bis(dimethylamino)- (Methylene Blue, MB)	CCl ₄ / MeOH (98:2)	$<5 \times 10^7$	293	PL/A'd-5	$S = A; A' = DPBF; \text{no effect of [MB] on decay rate up to } 1.9 \times 10^{-5} \text{ mol L}^{-1}$	83A371
	CHCl ₃		6.8×10^{-4}		CP/Oc-19	$S = A; A' = 2M2P$	84F673
	D ₂ O pH = 5.9	4×10^8			CP/A'c-25	$S = A; A' = \text{Lysozyme}; \text{used } k_d = 5 \times 10^4 \text{ s}^{-1}, k_{A'} = 6.4 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$	81R011
	H ₂ O pH = 5.9	3×10^8			CP/A'c-25	$S = A; A' = \text{Lysozyme}; \text{used } k_d = 5 \times 10^5 \text{ s}^{-1}, k_{A'} = 4.4 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$	81R011
	MeOD	1.6×10^7			PL/Ld-2	$S = A$	88A165
	MeOH	2.3×10^8	4.8×10^{-4}		CP/Ac-14	$S = A; \text{used } k_d = 1.1 \times 10^5 \text{ s}^{-1}$	87F479
11.114	Phenothiazinium, 3,7-diamino- (Thionine)	D ₂ O	6.8×10^7		PL/Ld-2	$S = MPDME$	83E235
11.115	Propanediamide, 2-[(4-diethylamino)-2-ethylphenyl]imino]-N,N'-diphenyl-	C ₅ H ₅ N	-1.6×10^7		CP/Oc-20	$S = RB; A' = 2M2P; \text{used } k_d = 3.1 \times 10^4 \text{ s}^{-1}, \beta_{A'} = 0.043 \text{ mol L}^{-1}; \text{No evidence of chemical reaction.}$	757166
11.116	1,3-Propanedione, 2-[[2-ethyl-4-(diethylamino)phenyl]imino]-1,3-diphenyl-	C ₅ H ₅ N	-1.6×10^7		CP/Oc-20	$S = RB; A' = 2M2P; \text{used } k_d = 3.1 \times 10^4 \text{ s}^{-1}, \beta_{A'} = 0.043 \text{ mol L}^{-1}; \text{No evidence of chemical reaction.}$	757166
11.117	Pyrazole, 4,5-dihydro-5-[4-(diethylamino)phenyl]-3-[2-[4-(diethylamino)phenyl]ethenyl]-1-phenyl-	MeOH/ C ₆ H ₆ (68:32)	5.8×10^9		CP/P'a-23	$S = RB; A' = DPBF; \text{used } k_d = 1.8 \times 10^5 \text{ s}^{-1}, k_{A'} = 6.3 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$	83A063
11.118	Pyrazole, 4,5-dihydro-5-[4-(dimethylamino)phenyl]-3-[2-[4-(dimethylamino)phenyl]ethenyl]-1-phenyl-	MeOH/ C ₆ H ₆ (80:20)	1.9×10^9		CP/P'a-23	$S = RB; A' = 2M2P; \text{used } k_d = 1.8 \times 10^5 \text{ s}^{-1}, k_{A'} = 8.1 \times 10^5 \text{ L mol}^{-1} \text{ s}^{-1}$	83A063
11.119	Pyrazole, 4,5-dihydro-1,5-diphenyl-3-(2-phenylethenyl)-	MeOH/ C ₆ H ₆ (80:20)	7.2×10^8		CP/P'a-23	$S = RB; A' = 2M2P; \text{used } k_d = 1.8 \times 10^5 \text{ s}^{-1}, k_{A'} = 8.1 \times 10^5 \text{ L mol}^{-1} \text{ s}^{-1}$	83A063
	MeOH/ C ₆ H ₆ (68:32)	1.7×10^8			CP/P'a-23	$S = RB; A' = DPBF; \text{used } k_d = 1.8 \times 10^5 \text{ s}^{-1}, k_{A'} = 6.3 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$	83A063

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.120	Pyrazole, 4,5-dihydro-5-(4-methoxyphenyl)-3-[2-(4-methoxyphenyl)ethenyl]-1-phenyl-						
	MeOH/ C ₆ H ₆ (80:20)	6.0×10^8			CP/P'a-23	S = RB; A' = 2M2P; used $k_d = 1.8 \times 10^5$ s ⁻¹ , $k_{A'} = 8.1 \times 10^5$ L mol ⁻¹ s ⁻¹ .	83A063
	MeOH/ C ₆ H ₆ (68:32)	1.3×10^9			CP/P'a-23	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ , $k_{A'} = 6.3 \times 10^8$ L mol ⁻¹ s ⁻¹ .	83A063
11.121	2-Pyrazoline, 3-(4-chlorophenyl)-1,5-diphenyl-				CP/Ac-17	S = MB; A' = 1,3,5-Triphenyl-2-pyrazoline; meas. $k_r/k_{r'}^{A'} = 2.1$.	86F225
	CH ₂ Cl ₂						
11.122	2-Pyrazoline, 5-(4-chlorophenyl)-1,3-diphenyl-				CP/Ac-17	S = MB; A' = 1,3,5-Triphenyl-2-pyrazoline; meas. $k_r/k_{r'}^{A'} = 1.5$.	86F225
	CH ₂ Cl ₂						
11.123	2-Pyrazoline, 5-deutero-1,3,5-triphenyl-				CP/Ac-17	S = MB; A' = 1,3,5-Triphenyl-2-pyrazoline; meas. $k_r/k_{r'}^{A'} = 0.48$; electron transfer.	86F225
	CH ₂ Cl ₂						
11.124	2-Pyrazoline, 3-(4-methoxyphenyl)-1,5-diphenyl-				CP/Ac-17	S = MB; A' = 1,3,5-Triphenyl-2-pyrazoline; meas. $k_r/k_{r'}^{A'} = 0.3$.	86F225
	CH ₂ Cl ₂						
11.125	2-Pyrazoline, 5-(4-methoxyphenyl)-1,3-diphenyl-				CP/Ac-17	S = MB; A' = 1,3,5-Triphenyl-2-pyrazoline; meas. $k_r/k_{r'}^{A'} = 0.8$.	86F225
	CH ₂ Cl ₂						
11.126	2-Pyrazoline, 3-(4-methylphenyl)-1,5-diphenyl-				CP/Ac-17	S = MB; A' = 1,3,5-Triphenyl-2-pyrazoline; meas. $k_r/k_{r'}^{A'} = 0.7$.	86F225
	CH ₂ Cl ₂						
11.127	2-Pyrazoline, 5-(4-methylphenyl)-1,3-diphenyl-				CP/Ac-17	S = MB; A' = 1,3,5-Triphenyl-2-pyrazoline; meas. $k_r/k_{r'}^{A'} = 1.1$.	86F225
	CH ₂ Cl ₂						
11.128	3-Pyrazolinone, 4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2-phenyl-				298 CP/Ac,A'c- 17,19	S = MB; A' = DPA; used $k_r^{A'} = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87F542
	MeOH	$4 \times 10^7 (k_r)$ $4 \times 10^6 (k_q)$					
11.129	3-Pyrazolinone, 4-[4-[(ethyl(2-hydroxyethyl)amino)-2-methylphenyl]imino-5-methyl-2-phenyl-				298 CP/Ac,A'c- 17,19	S = MB; A' = DPA; used $k_r^{A'} = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87F542
	MeOH	$4 \times 10^7 (k_r)$ $5 \times 10^6 (k_q)$					
11.130	3-Pyrazolinone, 4-[4-[(2-hydroxyethyl)ethylamino]phenyl]imino-5-methyl-2-phenyl-				298 CP/Ac,A'c- 17,19	S = MB; A' = DPA; used $k_r^{A'} = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87F542
	MeOH	$2 \times 10^7 (k_r)$ $5 \times 10^6 (k_q)$					
11.131	Pyrazolo[1,2-a]benzotriazole				CP/Ac-15	S = RB; used $k_d = 1.6 \times 10^4$ s ⁻¹ .	83F136
	CH ₃ CN	5×10^6	3×10^{-3}				
11.132	Pyrazolo[1,2-a]benzotriazole, 1,3-dimethyl-				CP/Ac-15	S = RB; used $k_d = 1.6 \times 10^4$ s ⁻¹ .	83F136
	CH ₃ CN	2×10^7	8×10^{-4}				
11.133	Pyrazol-3-one, 4-[(4-aminophenyl)imino]-2,4-dihydro-5-methyl-2-phenyl-				CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.3 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.018$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
	CH ₃ CN	5.0×10^7					
	MeOH	$4 \times 10^6 (k_q)$		298	CP/Ac,A'c-19	S = MB; A' = DPA.	87F542

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.134	Pyrazol-3-one, 4-[(4-amino-2,3,5,6-tetramethylphenyl)imino]-2,4-dihydro-5-methyl-2-phenyl-						
	CH ₃ CN	5.0×10^8			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.3 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.018$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
11.135	Pyrazol-3-one, 2-(4-bromophenyl)-4-(4-diethylamino-2-methylphenyl)-2,4-dihydro-imino-5-methylcarbamyl-						
	MeOH	$4 \times 10^7 (k_r)$		298	CP/Ac,A'c-17	S = MB; A' = DPA; used $k_r^{A'} = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87F542
11.136	Pyrazol-3-one, 2-(3-chlorophenyl)-4-(4-diethylamino-2-methylphenyl)-2,4-dihydro-imino-5-methylcarbamyl-						
	MeOH	$4 \times 10^7 (k_r)$		298	CP/Ac,A'c-17	S = MB; A' = DPA; used $k_r^{A'} = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87F542
11.137	Pyrazol-3-one, 2-(4-chlorophenyl)-4-(4-diethylamino-2-methylphenyl)-2,4-dihydro-imino-5-methylcarbamyl-						
	MeOH	$4 \times 10^7 (k_r)$		298	CP/Ac,A'c-17,19	S = MB; A' = DPA; used $k_r^{A'} = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87F542
$4 \times 10^6 (k_q)$							
11.138	Pyrazol-3-one, 4-[[4-(diethylamino)-2,6-dimethylphenyl]imino]-2,4-dihydro-5-(benzoylamino)-2-phenyl-						
	C ₅ H ₅ N	1.0×10^9			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
11.139	Pyrazol-3-one, 4-[[4-(diethylamino)-2,6-dimethylphenyl]imino]-2,4-dihydro-5-(1,1-dimethylethyl)-2-phenyl-						
	C ₅ H ₅ N	1.0×10^9			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
11.140	Pyrazol-3-one, 4-[[4-(diethylamino)-2,6-dimethylphenyl]imino]-2,4-dihydro-5-methyl-2-phenyl-						
	C ₅ H ₅ N	6.3×10^8			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
	C ₆ H ₆	1.1×10^9			CP/Oc-20	S = azine; A' = 2M2P; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.053$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
	CH ₃ CN	4.0×10^9			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.3 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.018$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
11.141	Pyrazol-3-one, 4-[[4-(diethylamino)-2-methylphenyl]imino]-2,4-dihydro-5-(1,1-dimethylethyl)-2-phenyl-						
	C ₅ H ₅ N	1.7×10^8			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
	C ₆ H ₆	1.3×10^8			CP/Oc-20	S = azine; A' = 2M2P; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.053$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
	CH ₃ CN	1.3×10^9			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.3 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.018$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
11.142	Pyrazol-3-one, 4-(4-diethylamino-2-methylphenyl)imino-2,4-dihydro-5-methylcarbamyl-2-(3-methoxyphenyl)-						
	MeOH	$5 \times 10^7 (k_r)$		298	CP/Ac,A'c-17	S = MB; A' = DPA; used $k_r^{A'} = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87F542
11.143	Pyrazol-3-one, 4-(4-diethylamino-2-methylphenyl)imino-2,4-dihydro-5-methylcarbamyl-2-(3-methylphenyl)-						
	MeOH	$4 \times 10^7 (k_r)$		298	CP/Ac,A'c-17,19	S = MB; A' = DPA; used $k_r^{A'} = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87F542
		$4 \times 10^6 (k_q)$					
11.144	Pyrazol-3-one, 4-[[4-(diethylamino)-2-methylphenyl]imino]-2,4-dihydro-5-methyl-2-phenyl-						
	C ₅ H ₅ N	4.0×10^7			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.144	Pyrazol-3-one, 4-[[4-(diethylamino)-2-methylphenyl]imino]-2,4-dihydro-5-methyl-2-phenyl-	— Continued					
	MeOH	$4 \times 10^7 (k_r)$ $5 \times 10^6 (k_q)$		298 17,19	CP/Ac,A'c-	S = MB; A' = DPA; used $k_r^{A'} = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87F542
11.145	Pyrazol-3-one, 4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2,4-dihydro-2-(4-nitrophenyl)-						
	MeOH	$4 \times 10^7 (k_r)$		298	CP/Ac,A'c-17	S = MB; A' = DPA; used $k_r^{A'} = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87F542
11.146	Pyrazol-3-one, 4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2,4-dihydro-2-(2,4,6-trichlorophenyl)-						
	MeOH	$4 \times 10^7 (k_r)$		298	CP/Ac,A'c-17	S = MB; A' = DPA; used $k_r^{A'} = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87F542
11.147	Pyrazol-3-one, 4-[[4-(diethylamino)phenyl]imino]-2,4-dihydro-5-methyl-2-phenyl-						
	C ₅ H ₅ N	4.0×10^7			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
	C ₆ H ₆	1.2×10^7			CP/Oc-20	S = azine; A' = 2M2P; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.053$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
	CH ₃ CN	4.0×10^8			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.3 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.018$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
	MeOH	$3 \times 10^7 (k_r)$ $4 \times 10^6 (k_q)$		298 17,19	CP/Ac,A'c-	S = MB; A' = DPA; used $k_r^{A'} = 1.8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	87F542
11.148	Pyrazol-3-one, 2,4-dihydro-4-[(4-methoxyphenyl)imino]-5-methyl-2-phenyl-						
	C ₅ H ₅ N				CP/Oc-20	S = RB; A' = 2M2P; No measurable effect.	757166
	CH ₃ CN				CP/Oc-20	S = RB; A' = 2M2P; No measurable effect.	757166
11.149	Pyrazol-3-one, 4-[[4-(dimethylamino)-3,5-dimethylphenyl]imino]-2,4-dihydro-5-methyl-2-phenyl-						
	CH ₃ CN	6.3×10^6			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.3 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.018$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
11.150	Pyrazol-3-one, 4-[[4-(dimethylamino)phenyl]imino]-2,4-dihydro-5-methyl-2-phenyl-						
	C ₅ H ₅ N	4.0×10^7			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
	MeOH	$4 \times 10^6 (k_q)$		298	CP/Ac,A'c-19	S = MB; A' = DPA.	87F542
11.151	Pyrazol-3-one, 4-[[4-(dimethylamino)-2,3,5,6-tetramethylphenyl]imino]-2,4-dihydro-5-methyl-2-phenyl-						
	CH ₃ CN	4.0×10^6			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.3 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.018$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
11.152	Pyrazol-3-one, 4-[[4-(diphenylamino)phenyl]imino]-2,4-dihydro-5-methyl-2-phenyl-						
	C ₅ H ₅ N	5.0×10^7			CP/Oc-20	S = RB; A' = 2M2P; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $\beta_{A'} = 0.043$ mol L ⁻¹ ; No evidence of chemical reaction.	757166
11.153	Pyrazol-3-one, 4,4'-[(2,3,5,6-tetramethyl-1,4-phenylene)dinitrilo]bis[2,4-dihydro-5-methyl-2-phenyl-						
	C ₅ H ₅ N				CP/Oc-20	S = MB; A' = 2M2P; No measurable effect.	757166
	CH ₃ CN				CP/Oc-20	S = RB; A' = 2M2P; No measurable effect.	757166
11.154	Pyrazolo[5,1-c]-1,2,4-triazole, 7-[4-(N-ethyl-[N-(2-methylsulfonylamino)ethyl]amino-2-methylphenylimino]-6-methyl-3-[4-[3-(3-butyl-4-hydroxyphenoxy)dodecyl]methoxy]carbonylamino]phenyl]propyl]-						
	EtOH	$<6 \times 10^7$			PL/A'd-8	S = An; A' = DPBF.	91A448
11.155	Pyrazolo[5,1-c]-1,2,4-triazole-6-carboxylic acid, 7-[[4-(diethylamino)-2-methylphenyl]imino]-3-phenyl-, ethyl ester						
	EtOH	2×10^8			PL/A'd-8	S = An; A' = DPBF.	91A448

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.155	Pyrazolo[5,1- <i>c</i>]-1,2,4-triazole-6-carboxylic acid, 7-[[4-(diethylamino)-2-methylphenyl]imino]-3-phenyl-, ethyl ester	—	—	—	Continued		
	EtOH	1.4×10^6 (k_r)			CP/Ac-17	S = AlCl(tspc); A' = Azo dye 10; used $k_r^{A'} = 2.4 \times 10^6$ L mol ⁻¹ s ⁻¹ ; values from graph.	91A448
	EtOH	9×10^7			CP/Ac-16	S = AlCl(tspc); A' = DPBF; used $k_d = 8 \times 10^4$ s ⁻¹ .	91A448
11.156	Pyrido[2,1- <i>b</i>]benzothiazolium, 4-[(2,3-dihydro)pyrido[2,1- <i>b</i>]benzothiazol-4-yl)methylene]-1,2,3,4-tetrahydro-, iodide						
	MeOH	2.7×10^7 (k_r)			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(S) = 0.75$.	767071
11.157	Pyrylium, 2,6-bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)selenopyran-4-ylidene]-3-propenyl]-						
	MeOH/ H ₂ O (50:50)	4×10^6 (k_r)			CP/Ac-14	S = RB; used $k_d = 1.7 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(RB) = 0.77$.	90F157
	MeOH/ H ₂ O (50:50)	3×10^6 (k_r)			CP/Ac-14	S = MB; used $k_d = 1.7 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(MB) = 0.50$.	90F157
11.158	Pyrylium, 2,6-bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)telluropyran-4-ylidene]-3-propenyl]-						
	MeOH/ H ₂ O (50:50)	8.5×10^7 (k_r)			CP/Ac-14	S = MB; used $k_d = 1.7 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(MB) = 0.50$.	90F157
	MeOH/ H ₂ O (50:50)	9.4×10^7 (k_r)			CP/Ac-14	S = RB; used $k_d = 1.7 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(RB) = 0.77$.	90F157
11.159	Pyrylium, 4,4'--(1,3-propenyl)bis[2,6-di(1,1-dimethylethyl)-						
	MeOH/ H ₂ O (50:50)	3×10^5 (k_r)			CP/Ac-14	S = RB; used $k_d = 1.7 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(RB) = 0.77$.	90F157
11.160	Selenopyrylium, 2,6-bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)selenopyran-4-ylidene]-3-propenyl]-						
	MeOH/ H ₂ O (50:50)	5×10^6 (k_r)			CP/Ac-14	S = RB; used $k_d = 1.7 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(RB) = 0.77$.	90F157
11.161	Telluropyrylium, 2,6-bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)telluropyran-4-ylidene]-3-propenyl]-						
	MeOH/ H ₂ O (99:1)	9×10^6 (k_r)			CP/Ac-14	S = MB; used $k_d = 1 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(MB) = 0.50$.	90F157
	MeOH/ H ₂ O (50:50)	1.8×10^8 (k_r)			CP/Ac-14	S = MB; used $k_d = 1.7 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(MB) = 0.50$.	90F157
	MeOH/ H ₂ O (50:50)	1.8×10^8 (k_r)			CP/Ac-14	S = RB; used $k_d = 1.7 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(RB) = 0.77$.	90F157
	H ₂ O	8×10^8 (k_r)			CP/Ac-14	S = MB; used $k_d = 2.4 \times 10^5$ s ⁻¹ , k_r derived using $\phi_\Delta(MB) = 0.50$.	90F157
							92F189
11.162	Telluropyrylium, 2,6-diphenyl-4-(2,6-diphenyltelluropyran-4-ylidene)methyl-						
	MeOH/ H ₂ O (99:1)	1×10^7 (k_r)	298	CP/Ac-14	S = RB; used $k_d = 1 \times 10^5$ s ⁻¹ .		92F189
11.163	2,2'-Thiacarbocyanine, 5,5'-dichloro-3,3-diethyl-, bromide						
	MeOH	1.3×10^6 (k_r)			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(S) = 0.75$.	767071
11.164	2,2'-Thiacarbocyanine, 5,5'-dichloro-3,9,3'-triethyl-, bromide						
	MeOH	7.1×10^6 (k_r)			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(S) = 0.75$.	767071
11.165	2,2'-Thiacarbocyanine, 5,5'-dicyano-3,9,3'-triethyl-, tetrafluoroborate						
	MeOH	3.3×10^6 (k_r)			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(S) = 0.75$.	767071
11.166	2,2'-Thiacarbocyanine, 3,3'-diethyl-, toluenesulfonate						
	MeOH	3.5×10^6 (k_r)			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(S) = 0.75$.	767071

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.167	2,2'-Thiacarbocyanine, 3,3'-diethyl-5,5'-dimethoxy-, toluenesulfonate						
	MeOH	$6.7 \times 10^6 (k_r)$			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(S) = 0.75$.	767071
11.168	2,2'-Thiacarbocyanine, 3,3'-diethyl-8,10-dimethyl-, toluenesulfonate						
	MeOH	$2.9 \times 10^8 (k_r)$			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(S) = 0.75$.	767071
11.169	2,2'-Thiacarbocyanine, 3,3'-diethyl-8,10-ethanediyl-, toluenesulfonate						
	MeOH	$1.2 \times 10^8 (k_r)$			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(S) = 0.75$.	767071
11.170	2,2'-Thiacarbocyanine, 3,3'-diethyl-8,9-(1,3-propanediyl)-, iodide						
	MeOH	$1.5 \times 10^8 (k_r)$			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(S) = 0.75$.	767071
11.171	2,2'-Thiacarbocyanine, 3,9,3'-triethyl-, bromide						
	MeOH	$1.6 \times 10^7 (k_r)$			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(S) = 0.75$.	767071
11.172	2,2'-Thiacarbocyanine, 3,9,3'-triethyl-5,5'-dimethoxy-, toluenesulfonate						
	MeOH	$3.1 \times 10^7 (k_r)$			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(S) = 0.75$.	767071
11.173	2,2'-Thiadicarbocyanine, 3,3'-diethyl-, iodide						
	MeOH	$1.3 \times 10^7 (k_r)$			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(S) = 0.75$.	767071
11.174	2,2'-Thiatricarbocyanine, 3,3'-diethyl-, iodide						
	Propylene carbonate	$2.7 \times 10^9 (k_q)$		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_q/k_r^{A'}$.	87E690
	Propylene carbonate	$3.8 \times 10^8 (k_r)$		293	PL/A'd-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
	MeOH	$5.1 \times 10^7 (k_r)$			CP/Ac-15	S = 2-ACN; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(S) = 0.75$.	767071
11.175	Thiopyrylium, 2,6-bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)selenopyran-4-ylidene]-3-propenyl]-						
	MeOH/ H ₂ O (50:50)	$9.0 \times 10^6 (k_r)$		298	CP/Ac-14	S = RB; used $k_d = 1.7 \times 10^5$ s ⁻¹ .	92F189
11.176	Thiopyrylium, 2,6-bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)thiopyran-4-ylidene]-3-propenyl]-						
	MeOH/ H ₂ O (50:50)	$1.3 \times 10^7 (k_r)$			CP/Ac-14	S = RB; used $k_d = 1.7 \times 10^5$ s ⁻¹ ; k_r derived using $\phi_\Delta(RB) = 0.77$.	90F157
11.177	1,2,3-Triazolo[1,2-a]benzotriazole						
	CH ₃ CN	4×10^4	0.4		CP/Ac-15	S = RB; used $k_d = 1.6 \times 10^4$ s ⁻¹ .	83F136
11.178	Tricarbocyanine, 5,5'-dichloro-1,1',3,3,3',3'-hexamethyl-, iodide						
	Propylene carbonate	$4.0 \times 10^9 (k_q)$		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_q/k_r^{A'}$.	87E690
	Propylene carbonate	$4 \times 10^7 (k_r)$		293	PL/A'd-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
11.179	Tricarbocyanine, 5,5'-difluoro-1,1',3,3,3',3'-hexamethyl-, iodide						
	Propylene carbonate	$7 \times 10^7 (k_r)$		293	PL/A'd-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.0 \times 10^9$ L mol ⁻¹ s ⁻¹ .	87E690
	Propylene carbonate	$1.0 \times 10^9 (k_q)$		293	PL/A'd-5	S = MB; A' = DPBF; Measured $k_q + k_r$, k_q calcd. knowing k_r from $k_q/k_r^{A'}$.	87E690

TABLE 11. Rate constants for the interaction of singlet oxygen with diazo compounds and dyes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.180 Tricarbocyanine, 5,5'-dimethoxy-1,1',3,3,3',3'-hexamethyl-, iodide							
	Propylene carbonate	$1.5 \times 10^8 (k_r)$		293	PL/A'd-17	$S = MB; A' = DPBF$; used $k_r^{A'} = 1.0 \times 10^9 L$ mol ⁻¹ s ⁻¹ .	87E690
	Propylene carbonate	$2.6 \times 10^9 (k_q)$		293	PL/A'd-5	$S = MB; A' = DPBF$; Measured $k_q + k_r, k_q$ calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690
11.181 Tricarbocyanine, 1,1',3,3,3',3'-hexamethyl-, fluoride							
	Propylene carbonate	$8 \times 10^7 (k_r)$		293	PL/A'd-17	$S = MB; A' = DPBF$; used $k_r^{A'} = 1.0 \times 10^9 L$ mol ⁻¹ s ⁻¹ .	87E690
	Propylene carbonate	$5.4 \times 10^8 (k_q)$		293	PL/A'd-5	$S = MB; A' = DPBF$; Measured $k_q + k_r, k_q$ calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690
11.182 Tricarbocyanine, 1,1',3,3,3',3'-hexamethyl-, iodide							
	Propylene carbonate	$7 \times 10^7 (k_r)$		293	PL/A'd-17	$S = MB; A' = DPBF$; used $k_r^{A'} = 1.0 \times 10^9 L$ mol ⁻¹ s ⁻¹ .	87E690
	Propylene carbonate	$1.8 \times 10^9 (k_q)$		293	PL/A'd-5	$S = MB; A' = DPBF$; Measured $k_q + k_r, k_q$ calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690
	DMSO	$1.9 \times 10^5 (k_r)$	0.078		CL/Ac-15	$S = Th$; used $k_d = 1.3 \times 10^4 s^{-1}$; k_r derived using $\phi_{\Delta}(S) = 0.87$.	82F531
	MeOH	$1.2 \times 10^6 (k_r)$	0.14		CL/Ac-15	$S = Th$; used $k_d = 1.4 \times 10^5 s^{-1}$; k_r derived using $\phi_{\Delta}(S) = 0.87$.	82F531
11.183 Tricarbocyanine, 1,1',3,3,3',3'-hexamethyl-, perchlorate							
	Propylene carbonate	$1.0 \times 10^8 (k_r)$		293	PL/A'd-17	$S = MB; A' = DPBF$; used $k_r^{A'} = 1.0 \times 10^9 L$ mol ⁻¹ s ⁻¹ .	87E690
	Propylene carbonate	$7.7 \times 10^8 (k_q)$		293	PL/A'd-5	$S = MB; A' = DPBF$; Measured $k_q + k_r, k_q$ calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690
	CH ₃ CN/C ₆ H ₆ (80:20)	8×10^7 $1.6 \times 10^6 (k_r)$ $4 \times 10^7 (k_q)$		298	CP/Ac,A'c-23	$S = A' = Rub$; used $k_d = 3.4 \times 10^4 s^{-1}$, $k_{A'} = 4.2 \times 10^7 L$ mol ⁻¹ s ⁻¹ .	90F360
11.184 Tricarbocyanine, 1,1',3,3,3',3',5,5'-octamethyl-, iodide							
	Propylene carbonate	$5 \times 10^7 (k_r)$		293	PL/A'd-17	$S = MB; A' = DPBF$; used $k_r^{A'} = 1.0 \times 10^9 L$ mol ⁻¹ s ⁻¹ .	87E690
	Propylene carbonate	$5.8 \times 10^8 (k_q)$		293	PL/A'd-5	$S = MB; A' = DPBF$; Measured $k_q + k_r, k_q$ calcd. knowing k_r from $k_r/k_r^{A'}$.	87E690

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes.

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.1	Acetatobis(acetylacetonato)manganese(III)						
	CH ₃ COCH ₃ / H ₂ O (95:5)	2.2 × 10 ⁷			PL/Ld-2	S = PdMP.	82A412
12.2	Ammine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	4.8 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.3	Aniline[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	1.9 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.4	Aqua[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II)						
	C ₆ H ₅ CH ₃	1.2 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.5	Aqua[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	1.4 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.6	Bis(acetylacetonato)chloroacetatomanganese(III)						
	CH ₃ COCH ₃ / H ₂ O (95:5)	2.7 × 10 ⁷			PL/Ld-2	S = PdMP.	82A412
12.7	Bis(acetylacetonato)cobalt(III)						
	C ₆ H ₅ CH ₃	1 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
	CCl ₄ / CHCl ₃ (90:10)	1.5 × 10 ⁸			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_A/(k_A[A'] + k_d) = 8.2 \times 10^4$ L mol ⁻¹ at [A'] = 5 × 10 ⁻⁶ mol L ⁻¹ .	74F645
12.8	Bis(acetylacetonato)copper(II)						
	C ₆ H ₅ CH ₃	1 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
	CCl ₄ / CHCl ₃ (90:10)	3.7 × 10 ⁶			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_A/(k_A[A'] + k_d) = 2.1 \times 10^3$ L mol ⁻¹ at [A'] = 5 × 10 ⁻⁶ mol L ⁻¹ .	74F645
12.9	Bis(acetylacetonato)manganese(III)						
	CH ₃ COCH ₃ / H ₂ O (95:5)	2.5 × 10 ⁷			PL/Ld-2	S = PdMP.	82A412
12.10	Bis(acetylacetonato)nickel(II)						
	BuOCH ₂ CH ₂ OH	7.5 × 10 ⁷		273	MD/A'c-33	A' = Rub; used $k_d = 3.8 \times 10^5$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319
	C ₆ H ₅ Br	6.5 × 10 ⁷		273	MD/A'c-33	A' = Rub; used $k_d = 1.3 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k/[k_d/(k_d[A']) + k_{A'}] = 0.5$ at [A'] = 1.5 × 10 ⁻⁴ mol L ⁻¹ . A exists as a trimer in solution.	737333
	C ₆ H ₅ CH ₃	3 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
	CCl ₄ / CHCl ₃ (90:10)	8.2 × 10 ⁷			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_A/(k_A[A'] + k_d) = 4.6 \times 10^4$ L mol ⁻¹ at [A'] = 5 × 10 ⁻⁶ mol L ⁻¹ .	74F645
	CH ₂ Cl ₂	1.5 × 10 ⁸		298	CP/A'c-23	S = A' = Rub; used $k_d = 8 \times 10^3$ s ⁻¹ , $k_{A'} = 8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	CH ₂ Cl ₂	7.5 × 10 ⁷			CP/A'c-23	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	732066

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.10	Bis(acetylacetonato)nickel(II) — Continued						
	CH ₃ COCH ₃ / H ₂ O (95:5)	1.5×10^8			PL/Ld-2	S = PdMP.	82A412
	CH ₂ Cl ₂	1.8×10^8			CP/A'c-19	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88F059
12.11	Bis(acetylacetonato)nickel(II) dihydrate						
	CCl ₄ / MeOH (98:2)	6.6×10^7			FP/A'd-5	S = MB; A' = DPBF.	78E238
12.12	Bis(acetylacetonato)trichloroacetatomanganese(III)						
	CH ₃ COCH ₃ / H ₂ O (95:5)	2.2×10^7			PL/Ld-2	S = PdMP.	82A412
12.13	Bis(acetylacetonato)zinc(II)						
	CCl ₄ / CHCl ₃ (90:10)	1.1×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_A/(k_A[A'] + k_d) = 6.1 \times 10^3$ L mol ⁻¹ at [A'] = 5×10^{-6} mol L ⁻¹ .	74F645
12.14	Bis(5-bromo-2-hydroxybenzaldehydato)nickel(II) dihydrate						
	CHCl ₃	5.3×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.15	Bis(2-butene-2,3-dithiolato)nickel(II)						
	C ₆ H ₅ CH ₃	2.8×10^{10}			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
	CH ₃ CN/ C ₆ H ₆ (80:20)	1.3×10^{10}		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
		$9 \times 10^5 (k_p)$					
		$1.4 \times 10^{10} (k_q)$					
12.16	Bis[O-butyl-3,5-di-(1,1-dimethylethyl)-4-hydroxybenzylphosphonato]nickel(II)						
	BuOCH ₂ CH ₂ OH	3.4×10^7		273	MD/A'c-33	A' = Rub; used $k_d = 3.8 \times 10^5$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319
	C ₆ H ₅ Br	1.3×10^7		273	MD/A'c-33	A' = Rub; used $k_d = 1.3 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k/[k_d/(A') + k_{A'}] = 0.1$ at [A'] = 1.5×10^{-4} mol L ⁻¹ . A exists as a trimer in solution.	737333
	C ₆ H ₅ CH ₃	2.0×10^8			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
	C ₆ H ₆	1.4×10^7			PL/A'd-8	S = MB; A' = DPBF.	76F902
	CCl ₄ / CHCl ₃ (90:10)	9.0×10^6			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_A/(k_A[A'] + k_d) = 5.2 \times 10^3$ L mol ⁻¹ at [A'] = 5×10^{-6} mol L ⁻¹ .	74F645
	CH ₂ Cl ₂	1.6×10^7		298	CP/A'c-23	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	74F642
	CH ₂ Cl ₂	1.4×10^7		298	CP/A'c-23	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319
	CH ₂ Cl ₂ / MeOH/ C ₅ H ₅ N (94:3:3)	1.0×10^7		298	CR/A'c-32	A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; O ₂ * from (PhO) ₃ PO ₃ .	727319
12.17	Bis[2-[(butylimino)methyl]-4-bromophenolato]nickel(II)						
	CHCl ₃	3.7×10^9			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.18	Bis[2-[(butylimino)methyl]-4-methoxyphenolato]nickel(II)						
	n-C ₁₆ H ₃₄	4.0×10^8		298	MD/A'c-33	A' = Rub; used $k_d = 9.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	n-C ₁₆ H ₃₄	2.0×10^8		298	MD/A'c-33	A' = Rub; used $k_d = 9.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.18 Bis[2-[(butylimino)methyl]-4-methoxyphenolato]nickel(II) — Continued							
	CH ₂ Cl ₂	2.0×10^9		298	CP/A'c-23	S = A' = Rub; used $k_d = 8 \times 10^3$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	CHCl ₃	3.4×10^9			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
	i-octane/ MeOH/ C ₅ H ₅ N (94:3:3)	3.5×10^9		298	CR/A'c-32	A' = Rub; used $k_d = 5.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from (PhO) ₃ PO ₃ .	727319
	i-octane	2.6×10^9		298	CP/A'c-23	S = A' = Rub; used $k_d = 4.7 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
12.19 Bis[2-[(butylimino)methyl]phenolato]nickel(II)							
	CHCl ₃	2.8×10^9			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.20 Bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenethionato)cobalt(II)							
	C ₆ H ₅ CH ₃	2.7×10^9			PL/Ld-2	S = TPP.	88A507
	C ₆ H ₅ CH ₃	3.5×10^7 (k_r)			CP/Ac,A'c-17	S = PP; A' = DPBF; used $k_r^{A'} = 6.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	88A507
	CH ₃ CN	1.9×10^9			PL/Ld-2	S = TPP.	88A507
	CH ₃ CN	8.5×10^6 (k_r)			CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.1 \times 10^9$ L mol ⁻¹ s ⁻¹ .	88A507
12.21 Bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenethionato]nickel(II)							
	C ₆ H ₅ CH ₃	4.7×10^9			PL/Ld-2	S = TPP.	88A507
	C ₆ H ₅ CH ₃	5.0×10^7 (k_r)			CP/Ac,A'c-17	S = PP; A' = DPBF; used $k_r^{A'} = 6.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	88A507
	CH ₃ CN	1.3×10^7 (k_r)			CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.1 \times 10^9$ L mol ⁻¹ s ⁻¹ .	88A507
	CH ₃ CN	8.4×10^9			PL/Ld-2	S = TPP.	88A507
12.22 Bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenonato]copper(II)							
	C ₆ H ₅ CH ₃	5.0×10^7 (k_r)			CP/Ac,A'c-17	S = PP; A' = DPBF; used $k_r^{A'} = 6.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	88A507
	C ₆ H ₅ CH ₃	1.5×10^9			PL/Ld-2	S = TPP.	88A507
12.23 Bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenethionato]zinc(II)							
	C ₆ H ₅ CH ₃	5.0×10^7			PL/Ld-2	S = TPP.	88A507
	C ₆ H ₅ CH ₃	1.3×10^7 (k_r)			CP/Ac,A'c-17	S = PP; A' = DPBF; used $k_r^{A'} = 6.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	88A507
	CH ₃ CN	9.0×10^6 (k_r)			CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.1 \times 10^9$ L mol ⁻¹ s ⁻¹ .	88A507
	CH ₃ CN	1.9×10^8			PL/Ld-2	S = TPP.	88A507
	CHCl ₃	1.2×10^8			PL/Ld-2	S = TPP.	88A507
12.24 Bis[2-[(cyclohexylimino)methyl]phenolato]nickel(II)							
	C ₆ H ₅ CH ₃	4.7×10^9			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.25 Bis[1,2-di(2-bromophenyl)-1,2-ethenedithiolato]nickel(II)							
	CH ₃ CN/ C ₆ H ₆ (80:20)	1.1×10^{10} 9×10^5 (k_r) 1.1×10^{10} (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.26 Bis(dibutylthiocarbamato)cobalt(II)							
	CCl ₄ / MeOH (98:2)	1.2×10^9			FP/A'd-5	S = MB; A' = DPBF.	78E238

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.27	Bis(dibutylthiocarbamato)copper(II)						
	CCl ₄ / MeOH (98:2)	<5 × 10 ⁸			FP/A'd-5	S = MB; A' = DPBF; no effect for [A] = 5 × 10 ⁻⁶ mol L ⁻¹ .	78E238
12.28	Bis(dibutylthiocarbamato)nickel(II)						
	n-C ₁₆ H ₃₄	9.7 × 10 ⁸		298	MD/A'c-33	A' = Rub; used $k_d = 9 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88F415
	n-C ₁₆ H ₃₄	1.0 × 10 ⁹		298	MD/A'c-33	A' = Rub; used $k_d = 9.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	n-C ₁₆ H ₃₄	9.0 × 10 ⁸		298	MD/A'c-33	A' = Rub; used $k_d = 9.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319
	C ₆ H ₅ Br	2.6 × 10 ⁸		273	MD/A'c-33	A' = Rub; used $k_d = 1.3 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k/[k_d/[A] + k_{A'}]$ = 2.0 at [A'] = 1.5 × 10 ⁻⁴ mol L ⁻¹ .	737333
	C ₆ H ₅ CH ₃	4.3 × 10 ⁹			CP/A'c-25	S = A' = Rub; uscd $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
	C ₆ H ₆	6 × 10 ⁹		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
	C ₆ H ₆ / EtOH (89:11)	5.2 × 10 ⁹ (k_d)		295	CP/P'a-19	S = RB; A' = TEMP-4-OH; used $k_d = 3 \times 10^4$ s ⁻¹ ; P' = TEMPOL.	757445
	CCl ₄ / MeOH (98:2)	4.1 × 10 ⁹			FP/A'd-5	S = MB; A' = DPBF.	78E238
	CH ₂ Cl ₂	1.2 × 10 ⁹		293	CP/Oc-22	S = Chl a; A' = Soybean oil; used $k_d = 1.0 \times 10^4$ s ⁻¹ .	91U026
	CH ₂ Cl ₂	3.5 × 10 ⁹	2.1 × 10 ⁻⁶		CP/A'c-19	S = A' = Tetr; used $k_d = 7.3 \times 10^3$ s ⁻¹ .	75F654
	CH ₂ Cl ₂	9.0 × 10 ⁹		298	CP/A'c-23	S = A' = Rub; used $k_d = 8 \times 10^3$ s ⁻¹ , $k_{A'} = 8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	CH ₂ Cl ₂	8.0 × 10 ⁹		298	CP/A'c-23	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	74F642
	CH ₂ Cl ₂ / MeOH/ C ₅ H ₅ N (94:3:3)	≥1.0 × 10 ⁹		298	CR/A'c-32	A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from (PhO) ₃ PO ₃ . Interference by chelate ozonide reaction to give colored products.	727319
	CHCl ₃	7.7 × 10 ⁹			CP/P'a-17	S = 1,5-Diamino-4,8-dihydroxyanthraquinone; A' = DMA; used $k_d = 3.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	80E446
	CHCl ₃	8.1 × 10 ⁹			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
	CS ₂ / MeOH (98:2)	4 × 10 ⁹			FP/A'd-5	S = MB; A' = DPBF.	737334
	DMF	2.0 × 10 ⁹			CP/A'c-25	S = TTMPP; A' = DPBF; used $k_d = 1.0 \times 10^5$ s ⁻¹ , $k_{A'} = 1.1 \times 10^9$ L mol ⁻¹ s ⁻¹ .	92F505
	DMF	3.6 × 10 ¹⁰		301	MP/P'a-19	S = Pt(phen)(BCAT); A' = 2,2,6,6-Tetramethyl-4-piperidinol; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; P' = 2,2,6,6-Tetramethyl-4-hydroxypiperidine-1-oxy; detected by esr.	89F181
	DMF	1.1 × 10 ¹⁰			MP/P'a-19	S = HP; A' = TEMP-4-OH; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; P' = TEMPOL; esr detection.	89F311
	DMF	~2 × 10 ¹⁰		301	MP/P'a-19	S = Pt(phen)(N ₃) ₂ ; A' = TEMPO-4-OH; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; P' = TEMPOL; detected by esr.	88A276
	i-octane	1.7 × 10 ¹⁰		298	CP/A'c-23	S = A' = Rub; used $k_d = 4.7 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	i-octane	7.0 × 10 ⁹		298	CP/A'c-23	S = A' = Rub; used $k_d = 4.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319
12.29	Bis(dibutylthiocarbamato)zinc(II)						
	C ₆ H ₅ Br	≤1 × 10 ⁶		273	MD/A'c-33	A' = Rub; No measurable effect.	737333
	CS ₂ / MeOH (98:2)	2 × 10 ⁷			FP/A'd-5	S = MB; A' = DPBF.	737334

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.29	Bis(dibutylthiocarbamato)zinc(II) — Continued						
	i-octane	$\leq 1 \times 10^6$			CP/A'c-23	S = A' = Rub; used $k_d = 4.7 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	74F341
12.30	Bis[O,O'-di-(4- <i>tert</i> -butylphenyl)phosphorodithiolato]cobalt(II)	C_6H_6	1.2×10^9		CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
12.31	Bis[O,O'-di-(4- <i>tert</i> -butylphenyl)phosphorodithiolato]copper(II)	C_6H_6	7.0×10^6		CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
12.32	Bis[O,O'-di-(4- <i>tert</i> -butylphenyl)phosphorodithiolato]nickel(II)	C_6H_6	2.3×10^9		CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
12.33	Bis[1,2-di(2-chlorophenyl)-1,2-ethenedithiolato]nickel(II)	CH_3CN/C_6H_6 (80:20)	1.1×10^{10} $1 \times 10^6 (k_p)$ $1.2 \times 10^{10} (k_q)$	298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.34	Bis[1,2-di(4-chlorophenyl)-1,2-ethenedithiolato]nickel(II)	CH_3CN/C_6H_6 (80:20)	1.2×10^{10} $1.1 \times 10^6 (k_p)$ $1.2 \times 10^{10} (k_q)$	298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.35	Bis(dicyclohexylphosphinodithiolato)nickel(II)	$CHCl_3$	5.7×10^9		CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.36	Bis(O,O'-dicyclohexylphosphorodithiolato)cobalt(II)	$C_6H_5CH_3$	2.7×10^9		CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.37	Bis[1,2-di(2,4-dichlorophenyl)-1,2-ethenedithiolato]nickel(II)	CH_3CN/C_6H_6 (80:20)	1.2×10^{10} $3.1 \times 10^6 (k_p)$ $1.3 \times 10^{10} (k_q)$	298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.38	Bis[1,2-di(3,4-dichlorophenyl)-1,2-ethenedithiolato]nickel(II)	CH_3CN/C_6H_6 (80:20)	1.2×10^{10} $2.4 \times 10^6 (k_p)$ $1.2 \times 10^{10} (k_q)$	298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.39	Bis[1,2-di(dimethylaminophenyl)-1,2-ethenedithiolato]nickel(II)	CH_3CN/C_6H_6 (80:20)	9.8×10^9 $3 \times 10^5 (k_p)$ $9.8 \times 10^9 (k_q)$	298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.40	Bis(diethylthiocarbamato)nickel(II)	$CCl_4/MeOH$ (98:2)	6.6×10^9		FP/A'd-5	S = MB; A' = DPBF.	78E238
	DMF	3×10^{10}		301	MP/P'a-19	S = Pt(DHBA)(DPA); A' = TEMP-4-OH; used $k_d = 1.4 \times 10^5$ s ⁻¹ , $k_{A'} = 6.7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; P' = TEMPOL; detected by esr.	94F004
	DMF	1.5×10^{10}		301	CP/P'a-19	S = Pt(phen)(DMT); A' = 2,2,6,6-Tetramethyl-4-piperidinol; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; P' = TEMPOL obs. by esr.	92F210
	DMF	1.3×10^{11}			CP/A'c-22	S = A' = Pt(phen)(DMT); used $k_d = 1.4 \times 10^5$ s ⁻¹ .	91F098

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.41 Bis[1,2-di(4-ethylphenyl)-1,2-ethenedithiolato]cobalt(II)							
	C ₆ H ₆	5.5 × 10 ⁹ 1.6 × 10 ⁶ (k_r) 5.4 × 10 ⁹ (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.42 Bis[1,2-di(4-ethylphenyl)-1,2-ethenedithiolato]nickel(II)							
	C ₆ H ₆	7.8 × 10 ⁹ 1 × 10 ⁵ (k_r) 7.1 × 10 ⁹ (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.43 Bis[1,2-di(4-ethylphenyl)-1,2-ethenedithiolato]platinum(II)							
	C ₆ H ₆	8.5 × 10 ⁹ 2 × 10 ⁵ (k_r) 8.0 × 8.0 × 10 ⁹ (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.44 Bis(O,O'-diethylphosphorodithiolato)nickel(II)							
	CH ₃ CN/C ₆ H ₆ (80:20)	6.8 × 10 ⁹ 3.6 × 10 ⁶ (k_r) 8.2 × 10 ⁹ (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
	CHCl ₃	9.5 × 10 ⁹			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.45 Bis[2,3-dihydro-N-2-pyridinyl-3-(2-pyridinylimino)isoindol-1-amine]nickel(II)							
	CHCl ₃	1.6 × 10 ⁸			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.46 Bis(diisopropylthiocarbamato)cobalt(II)							
	n-C ₁₆ H ₃₄	9.0 × 10 ⁸		298	MD/A'c-33	A' = Rub; used $k_d = 9.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	CH ₂ Cl ₂	1.9 × 10 ⁹			CP/A'c-23	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	732066
	i-octane	1.9 × 10 ⁹		298	CP/A'c-23	S = A' = Rub; used $k_d = 4.7 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
12.47 Bis(diisopropylthiocarbamato)copper(II)							
	CH ₂ Cl ₂	<1.0 × 10 ⁷			CP/A'c-23	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	732066
12.48 Bis(diisopropylthiocarbamato)iron(III)							
	n-C ₁₆ H ₃₄	1.2 × 10 ⁹		298	MD/A'c-33	A' = Rub; used $k_d = 9.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	CH ₂ Cl ₂	4.3 × 10 ⁹		298	CP/A'c-23	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 8 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	CH ₂ Cl ₂	3.9 × 10 ⁹		298	CP/A'c-23	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	74F642
	CH ₂ Cl ₂	3.8 × 10 ⁹			CP/A'c-23	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	732066
	i-octane	3.8 × 10 ⁹		298	CP/A'c-23	S = A' = Rub; used $k_d = 4.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	74F642
12.49 Bis(diisopropylthiocarbamato)manganese(I)							
	CH ₂ Cl ₂	<1.0 × 10 ⁷			CP/A'c-23	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	732066
12.50 Bis(diisopropylthiocarbamato)nickel(II)							
	CH ₂ Cl ₂	3.4 × 10 ⁹			CP/A'c-23	S = A' = Rub; used $k_d = 4.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	732066

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.51	Bis(diisopropylthiocarbamato)zinc(II)						
	CH ₂ Cl ₂	<1.0 × 10 ⁷			CP/A'c-23	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	732066
12.52	Bis(<i>O,O'</i> -diisopropylphosphorodithiolato)nickel(II)						
	C ₂ H ₅ CO ₂ CH ₃	6.4 × 10 ⁹		293	PL/Ld-2	S = Pz and 2-ACN.	92E220
	C ₆ H ₅ CH ₃	6.5 × 10 ⁹		293	PL/Ld-2	S = Pz and 2-ACN.	92E220
	CCl ₄ / MeOH (98:2)	7.6 × 10 ⁹			FP/A'd-5	S = MB; A' = DPBF.	78E238
	CH ₂ Cl ₂	5.4 × 10 ⁹			CP/A'c-23	S = A' = Rub; used $k_d = 4.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	732066
12.53	Bis[1,2-di(4-methoxyphenyl)-1,2-ethanedithionato]nickelate(I), tetrabutylammonium salt						
	CH ₃ CN/ C ₆ H ₆ (80:20)	8.6 × 10 ⁹ 8.4 × 10 ⁶ (k_p) 8.9 × 10 ⁹ (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.54	Bis[1,2-di(4-methoxyphenyl)-1,2-ethenedithiolato]nickel(II)						
	CH ₃ CN/ C ₆ H ₆ (80:20)	1.0 × 10 ¹⁰ 4 × 10 ⁵ (k_p) 1.0 × 10 ¹⁰ (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.55	Bis(dimethylthiocarbamato)bismuth(II)						
	C ₆ H ₆ / EtOH (89:11)			295	CP/P'a-19	S = RB; A' = TEMP-4-OH; P' = TEMPOL; no effect for [A] = 1.8 × 10 ⁻⁵ mol L ⁻¹ .	757445
	CH ₂ Cl ₂	9.2 × 10 ⁴	0.067		CP/A'c-19	S = A' = Tetr; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $\beta_{A'} = 4.0 \times 10^{-4}$ mol L ⁻¹ .	75F654
12.56	Bis(dimethylthiocarbamato)nickel(II)						
	DMF	3 × 10 ³			CP/A'c-19	S = RB; A' = DMA; used $k_d = 1.4 \times 10^5$ s ⁻¹ , $k_{A'} = 1.4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86F622
12.57	Bis[2-[(1,1-dimethylethyl)imino)methyl]phenolato]nickel(II)						
	C ₆ H ₆	2.6 × 10 ⁸			PL/A'd-8	S = MB; A' = DPBF.	76F902
12.58	Bis[4-(1,1-dimethylethyl)phenylsalicylato]nickel(II)						
	CCl ₄ / CHCl ₃ (90:10)	1.5 × 10 ⁷			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_{A'}/(k_A[A'] + k_d) = 8.5 \times 10^3$ L mol ⁻¹ at [A'] = 5 × 10 ⁻⁶ mol L ⁻¹ .	74F645
12.59	Bis[3,5-di(1-methylethyl)salicylato]cobalt(II)						
	C ₆ H ₅ CH ₃	3.0 × 10 ⁷			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.60	Bis[3,5-di(1-methylethyl)salicylato]nickel(II)						
	C ₆ H ₅ CH ₃	5.0 × 10 ⁷			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.61	Bis(<i>O,O'</i> -di(4-methylphenyl)phosphorodithiolato)cobalt(II)						
	C ₆ H ₆	1.1 × 10 ⁹			CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
12.62	Bis(<i>O,O'</i> -di(4-methylphenyl)phosphorodithiolato)copper(II)						
	C ₆ H ₆	2.0 × 10 ⁶			CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
12.63	Bis(<i>O,O'</i> -di(4-methylphenyl)phosphorodithiolato)nickel(II)						
	C ₆ H ₆	2.3 × 10 ⁹			CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.64 Bis(diphenyldithiocarbamato)nickel(II)							
	CHCl ₃	6.3×10^9			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
	<i>i</i> -octane	$<1.0 \times 10^6$		298	CP/A'c-23	$S = A' = \text{Rub}$; used $k_d = 4.7 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
12.65 Bis(1,2-diphenyl-1,2-ethanedithionato)nickelate(I), tetrabutylammonium salt							
	CH ₃ CN/ C ₆ H ₆ (80:20)	9.2×10^9 $4.3 \times 10^6 (k_p)$ $9.7 \times 10^9 (k_q)$		298	CP/Ac,A'c-23	$S = A' = \text{Rub}$; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.66 Bis[1,2-diphenyl-1,2-ethenedithiolato]nickel(II)							
	C ₆ H ₅ CH ₃	2.2×10^{10}			CP/A'c-25	$S = A' = \text{Rub}$; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
	CH ₃ CN/ C ₆ H ₆ (80:20)	1.3×10^{10} $9 \times 10^5 (k_p)$ $1.2 \times 10^{10} (k_q)$		298	CP/Ac,A'c-23	$S = A' = \text{Rub}$; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
	CHCl ₃	1.6×10^{10}		293	CP/P'a-23	$S = A' = 2,4\text{-Di}[2\text{-hydroxy-4-}\text{(diethylamino)phenyl}squarylium$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 1.2 \times 10^9$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 13$.	90F554
	CHCl ₃	1.1×10^{10}			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.67 Bis(O,O'-diphenylphosphorodithiolato)chromium(II)							
	C ₆ H ₆	3.2×10^6			CP/A'c-23	$S = A' = \text{Rub}$; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
12.68 Bis(O,O'-diphenylphosphorodithiolato)cobalt(II)							
	C ₆ H ₆	1.2×10^9			CP/A'c-23	$S = A' = \text{Rub}$; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
12.69 Bis(O,O'-diphenylphosphorodithiolato)copper(II)							
	C ₆ H ₆	1.7×10^6			CP/A'c-23	$S = A' = \text{Rub}$; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
12.70 Bis(O,O'-diphenylphosphorodithiolato)lead(II)							
	C ₆ H ₆	3.4×10^7			CP/A'c-23	$S = A' = \text{Rub}$; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
12.71 Bis(O,O'-diphenylphosphorodithiolato)nickel(II)							
	C ₆ H ₆	2.5×10^9			CP/A'c-23	$S = A' = \text{Rub}$; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
	CHCl ₃	1.1×10^{10}			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.72 Bis(O,O'-diphenylphosphorodithiolato)zinc(II)							
	C ₆ H ₆	5.0×10^6			CP/A'c-23	$S = A' = \text{Rub}$; used $k_d = 3 \times 10^7$ s ⁻¹ .	78F497
12.73 Bis(dithioacetylacetone)cobalt(II)							
	C ₆ H ₅ CH ₃	9.4×10^8		293	PL/Ld-2	$S = \text{An or PPDME}$; $E_a = 4.2$ kJ mol ⁻¹ ; $\log(A) = 9.7$.	86A540
	CH ₂ Cl ₂	1.8×10^8		293	PL/Ld-2	$S = \text{An or PPDME}$.	86A540
	CH ₃ CN	7.2×10^8		293	PL/Ld-2	$S = \text{An or PPDME}$.	86A540
	CH ₃ COCH ₃	9.0×10^8		293	PL/Ld-2	$S = \text{An or PPDME}$.	86A540
	CHCl ₃	7.4×10^8		293	PL/Ld-2	$S = \text{An or PPDME}$.	86A540

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.74	Bis(dithioacetylacetonato)nickel(II)						
	C ₆ H ₅ Br	2.3 × 10 ⁸		273	MD/A'c-23	A' = Rub; used $k_d = 1.3 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k/[k_d/(k_d/[A']) + k_{A'}] = 1.8$ at [A'] = 1.5 × 10 ⁻⁵ mol L ⁻¹ .	737331
12.75	Bis[1,2-di(4-trifluoromethylphenyl)-1,2-ethanedithionato]nickelate(I), tetrabutylammonium salt						
	CH ₃ CN/C ₆ H ₆ (80:20)	1.1 × 10 ¹⁰ 2.1 × 10 ⁶ (k_p) 1.0 × 10 ¹⁰ (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.76	Bis[1,2-di(4-trifluoromethylphenyl)-1,2-ethenedithiolato]nickel(II)						
	CH ₃ CN/C ₆ H ₆ (80:20)	1.4 × 10 ¹⁰ 2.7 × 10 ⁶ (k_p) 1.4 × 10 ¹⁰ (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.77	Bis[1,2-di(3,4,5-trimethoxyphenyl)-1,2-ethanedithionato]nickelate(I), tetrabutylammonium salt						
	CH ₃ CN/C ₆ H ₆ (80:20)	7.0 × 10 ⁹ 4.9 × 10 ⁶ (k_p) 7.6 × 10 ⁹ (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.78	Bis[1,2-di(3,4,5-trimethoxyphenyl)-1,2-ethanedithionato]nickelate(I), 4-[1,5,5-tri(4-diethylaminophenyl)pentadienylidene]anilinium salt						
	CH ₃ CN/C ₆ H ₆ (80:20)	5.6 × 10 ⁹ 5 × 10 ⁵ (k_p) 5.1 × 10 ⁹ (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.79	Bis[1,2-di(3,4,5-trimethoxyphenyl)-1,2-ethenedithiolato]nickel(II)						
	CH ₃ CN/C ₆ H ₆ (80:20)	9.7 × 10 ⁹ 4 × 10 ⁵ (k_p) 9.6 × 10 ⁹ (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.80	Bis[2-[(dodecylimino)methyl]phenolato]cobalt(II)						
	C ₆ H ₅ CH ₃	2.4 × 10 ⁹			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.81	Bis[2-[(dodecylimino)methyl]phenolato]copper(II)						
	C ₆ H ₅ CH ₃	5.0 × 10 ⁷			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.82	Bis[2-[(dodecylimino)methyl]phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	7.0 × 10 ⁹			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.83	Bis[O-ethyl-3,5-di-(1,1-dimethylethyl)-4-hydroxybenzylphosphonato]nickel(II)						
	CH ₂ Cl ₂	3.6 × 10 ⁶	2.0 × 10 ⁻³		CP/A'c-19	S = A' = Tetr; used $k_d = 7.3 \times 10^3$ s ⁻¹ .	75F654
	CHCl ₃	2.2 × 10 ⁷			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.84	Bis[5-ethyl-3-[(3-pyridinylimino)methyl]-2-thiophenethionato]cobalt(II)						
	C ₆ H ₅ CH ₃	5.0 × 10 ⁷ (k_p)			CP/Ac,A'c-17	S = PP; A' = DPBF; used $k_r^{A'} = 6.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	88A507
	C ₆ H ₅ CH ₃	1.5 × 10 ⁹			PL/Ld-2	S = TPP.	88A507
	CH ₃ CN	4.0 × 10 ⁶ (k_p)			CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_r^{A'} = 1.1 \times 10^9$ L mol ⁻¹ s ⁻¹ .	88A507
	CH ₃ CN	2.0 × 10 ⁹			PL/Ld-2	S = TPP.	88A507
12.85	Bis[1,1,1,4,4-hexafluoro-2-butene-2,3-dithiolato]nickel(II)						
	CHCl ₃	8.1 × 10 ⁹			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.86	Bis[hydrotris(1-pyrazolyl)borato]nickel(II)						
	CHCl ₃	2.1×10^6			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	79A050
12.87	Bis(2'-hydroxyacetophenone oximato)nickel(II)				CP/A'c-25	$S = A' = \text{Rub}$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$, $k_{A'} = 1.7 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	752063
12.88	Bis(2-hydroxybenzaldehydato)nickel(II) dihydrate				CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	79A050
12.89	Bis(2-hydroxybenzaldehyde oximato)nickel(II)				CP/A'c-25	$S = A' = \text{Rub}$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$, $k_{A'} = 1.7 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	752063
12.90	Bis[2-hydroxybenzaldehyde phenylhydrazone]nickel(II)				CP/A'c-25	$S = A' = \text{Rub}$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$, $k_{A'} = 1.7 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	752063
12.91	Bis[2'-hydroxy-4'- <i>tert</i> -butyloctadecanophenone oximato]palladium(II)			298	PL/A'd-8	$S = \text{An}; A' = \text{DPBF}$.	737438
12.92	Bis(2-hydroxy-5-methoxybenzaldehydato)nickel(II) dihydrate				CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	79A050
12.93	Bis[2'-hydroxy-4'-methylacetophenone oximato]nickel(II)			298	PL/A'd-8	$S = \text{An}; A' = \text{DPBF}$.	737438
12.94	Bis[2-hydroxy-5-methylbenzophenonato]nickel(II)				CP/A'c-25	$S = A' = \text{Rub}$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$, $k_{A'} = 1.7 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	752063
12.95	Bis[2'-hydroxy-4'-methyldodecanophenone oximato]nickel(II)				CP/A'c-25	$S = A' = \text{Rub}$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$, $k_{A'} = 1.7 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	752063
	C ₆ H ₆	2.7×10^9		298	PL/A'd-8	$S = \text{An}; A' = \text{DPBF}$.	737438
12.96	Bis[2-(iminomethyl)phenolato]nickel(II)				CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	79A050
12.97	Bis(4-imino-2-pentanonato)nickel(II)				CP/A'c-25	$S = A' = \text{Rub}$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$, $k_{A'} = 1.7 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	752063
12.98	Bis[<i>O</i> -(1-methylethyl)carbonodithionato]nickel(II)			298	CP/A'c-23	$S = A' = \text{Rub}$; used $k_d = 4.0 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 7.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	74F642
12.99	Bis[4-methyl-1,2-benzenedithiolato]cobalt(II) tetrabutylammonium salt				CP/A'c-25	$S = A' = \text{Rub}$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$, $k_{A'} = 1.7 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	752063
12.100	Bis[4-methyl-1,2-benzenedithiolato]nickel(II) tetrabutylammonium salt				CP/A'c-25	$S = A' = \text{Rub}$; used $k_d = 1 \times 10^5 \text{ s}^{-1}$, $k_{A'} = 1.7 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	752063

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.101	Bis[2-[(1-methylethylimino)methyl]phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	5.9×10^9			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
	CHCl ₃	2.6×10^9			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.102	Bis[N-methyl-7-(methylimino)-1,3,5-cycloheptatrien-1-aminato]nickel(II)						
	CHCl ₃	6.1×10^9			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.103	Bis(4-methylphenyldithiocarbamato)nickel(II)						
	CHCl ₃	1.1×10^{10}			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.104	Bis[N-(4-methylphenyl)-7-[(4-methylphenyl)imino]-1,3,5-cycloheptatrien-1-aminato]nickel(II)						
	CHCl ₃	5.6×10^9			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.105	Bis[2-[(1-methylpropylimino)methyl]phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	4.0×10^9			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.106	Bis[2-[(4-phenylamino)phenyl]imino]methyl]phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	1.7×10^9			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.107	Bis(phenyldithiocarbamato)nickel(II)						
	C ₆ H ₆	4.4×10^9		303	CP/P'a-16	S = A' = Rub; k_d not given.	87A107
	CHCl ₃	1.1×10^{10}			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
	ClCF ₂ CCl ₂ F	7.2×10^9		303	CR/P'a-32	A' = Rub; $^1O_2^*$ from DMNO ₂ ; $k_{A'}$ and k_d not given.	87A107
12.108	Bis[2-[(phenylimino)methyl]phenolato]cobalt(II)						
	C ₆ H ₅ CH ₃	3.2×10^9			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.109	Bis[2-[(phenylimino)methyl]phenolato]copper(II)						
	C ₆ H ₅ CH ₃	4.0×10^8			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.110	Bis[2-[(phenylimino)methyl]phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	7.8×10^9			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.111	Bis[2,2'-thiobis[O,O'-di(4- <i>tert</i> -butylphenyl)phosphorodithiolato]cadmium(II)						
	C ₆ H ₆	8.4×10^6			CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
12.112	Bis[2,2'-thiobis[O,O'-di(4- <i>tert</i> -butylphenyl)phosphorodithiolato]cobalt(II)						
	C ₆ H ₆	8.7×10^8			CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
12.113	Bis[2,2'-thiobis[O,O'-di(4- <i>tert</i> -butylphenyl)phosphorodithiolato]lead(II)						
	C ₆ H ₆	1.0×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
12.114	Bis[2,2'-thiobis[O,O'-di(4- <i>tert</i> -butylphenyl)phosphorodithiolato]nickel(II)						
	C ₆ H ₆	2.2×10^9			CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.115	Bis[2,2'-thiobis[O,O'-di(4- <i>tert</i> -butylphenyl)phosphorodithiolato]zinc(II)						
	C ₆ H ₆	4.4 × 10 ⁶			CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	78F497
12.116	Bis(2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)phenolato]nickel(II))						
	n-C ₁₆ H ₃₄	1.3 × 10 ⁸		298	MD/A'c-33	A' = Rub; used $k_d = 9.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319
	C ₆ H ₅ Br	3.9 × 10 ⁷		273	MD/A'c-33	A' = Rub; used $k_d = 1.3 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_A/[(k_d/[A']) + k_{A'}] = 0.3$ at [A'] = 1.5 × 10 ⁻⁴ mol L ⁻¹ .	737333
	C ₆ H ₅ CH ₃	2.7 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
	C ₆ H ₆	1.1 × 10 ⁸			PL/A'd-8	S = MB; A' = DPBF.	76F902
	CCl ₄ /CHCl ₃ (90:10)	5.7 × 10 ⁷			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_A/[(k_d/[A']) + k_{A'}] = 3.2 \times 10^4$ L mol ⁻¹ at [A'] = 5 × 10 ⁻⁶ mol L ⁻¹ .	74F645
	CS ₂ /MeOH (98:2)	5 × 10 ⁸			FP/A'd-5	S = MB; A' = DPBF.	737334
	<i>i</i> -octane/ MeOH/ C ₅ H ₅ N (94:3:3)	2.0 × 10 ⁸		298	CR/A'c-32	A' = Rub; used $k_d = 5.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from (PhO) ₃ P(O ₃) ₂ .	727319
	<i>i</i> -octane	9.6 × 10 ⁷		298	CP/A'c-23	S = A' = Rub; used $k_d = 4.7 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	<i>i</i> -octane	1.3 × 10 ⁸		298	CP/A'c-23	S = A' = Rub; used $k_d = 4.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319
12.117	Bis[3,4,6-trichloro-1,2-benzenedithiolato]nickelate(I), tetrabutylammonium salt						
	CH ₃ CN/ C ₆ H ₆ (80:20)	6.5 × 10 ⁹ 1 × 10 ⁶ (k_d) 6.7 × 10 ⁹ (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 3.4 \times 10^4$ s ⁻¹ , $k_{A'} = 4.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.118	Bis(2,4,6-trimethylbenzenesulfonato)nickel(II)						
	C ₆ H ₆ /EtOH (50:50)	1 × 10 ⁷ 3 × 10 ⁷ (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 5.6 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
12.119	Bromocarbonylbis(triphenylphosphine)iridium				PL/Ld-2	S = TPP.	94A126
12.120	Butylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II)						
	C ₆ H ₅ CH ₃	2.5 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.121	Butylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)phenolato]cobalt(II)]						
	C ₆ H ₅ CH ₃	3.7 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.122	Butylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)phenolato]nickel(II)]						
	n-C ₁₆ H ₃₄	1.0 × 10 ⁸		298	MD/A'c-33	A' = Rub; used $k_d = 9.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	n-C ₁₆ H ₃₄	8.0 × 10 ⁷		298	MD/A'c-33	A' = Rub; used $k_d = 9.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319
	BuOCH ₂ CH ₂ OH	2.8 × 10 ⁸		273	MD/A'c-33	A' = Rub; used $k_d = 3.8 \times 10^5$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319
	C ₆ H ₅ Br	1.3 × 10 ⁸		273	MD/A'c-33	A' = Rub; used $k_d = 1.3 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_A/[(k_d/[A']) + k_{A'}] = 1.0$ at [A'] = 1.5 × 10 ⁻⁴ mol L ⁻¹ . A exists as a tetramer in solution.	737333
	C ₆ H ₅ CH ₃	4.0 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
	C ₆ H ₆	1.1 × 10 ⁸			PL/A'd-8	S = MB; A' = DPBF.	76F902

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.122	Butylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) — Continued						
	CCl ₄ / CHCl ₃ (90:10)	1.1×10^8			CP/A'c-23	$S = A' = Rub$; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_A/(k_A[A'] + k_d) = 6.1 \times 10^4$ L mol ⁻¹ at $[A'] = 5 \times 10^{-6}$ mol L ⁻¹ .	74F645
	CH ₂ Cl ₂	3.7×10^7		293	CP/Oc-22	$S = Chl\; a$; $A' = Soybean$ oil; used $k_d = 1.0 \times 10^4$ s ⁻¹ .	91U026
	CHCl ₃	1.7×10^8			CP/A'c-33	$S = A' = Rub$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
	CS ₂ / MeOH (98:2)	2×10^8			FP/A'd-5	$S = MB$; $A' = DPBF$.	737334
	<i>i</i> -octane/ MeOH/ C ₅ H ₅ N (94:3:3)	2.7×10^8		298	CR/A'c-32	$A' = Rub$; used $k_d = 5.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; $^1O_2^*$ from (PhO) ₃ PO ₃ .	727319
	<i>i</i> -octane	2.0×10^8		298	CP/A'c-23	$S = A' = Rub$; used $k_d = 4.7 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
	<i>i</i> -octane	1.4×10^8		298	CP/A'c-23	$S = A' = Rub$; used $k_d = 4.0 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	74F642
	<i>i</i> -octane	1.8×10^8		298	CP/A'c-23	$S = A' = Rub$; used $k_d = 4 \times 10^4$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319
12.123	Carbonylchlorobis(triphenylphosphine)iridium						
	C ₆ D ₆	3.9×10^8 $2 \times 10^7 (k_r)$	3.5×10^{-6}		PL/Ac,A'c- 15,27	$S = MB$; $A' = TME$; used $k_d = 1.1 \times 10^2$ s ⁻¹ , assumed $f_r^{A'} = 1$.	93A050
	C ₆ D ₆	4.3×10^8			PL/Ld-2	$S = MB$.	93A050
	CDCl ₃	2.6×10^8			PL/Ld-2	$S = MB$.	93A050
	CH ₂ Cl ₂	2.6×10^8			PL/Ld-2	$S = TPP$.	94A126
	CHCl ₃	$2 \times 10^7 (k_r)$			CP/Ac,A'c-17	$S = MB$; $A' = DMA$; used $k_r^{A'} = 2.9 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.67$.	93A050
12.124	Carbonylchlorobis(triphenylphosphine)rhodium						
	CDCl ₃	3.1×10^8			PL/Ld-2	$S = MB$ or C ₆₀ .	93E539
	CDCl ₃	$1 \times 10^8 (k_r)$			PL/Ac,A'c- 15,27	$S = MB$ or C ₆₀ ; $A' = TME$; assumed $f_r^{A'} = 1$; used $k_A = 3.1 \times 10^6$ L mol ⁻¹ s ⁻¹ .	93E539
12.125	Carbonyliodobis(triphenylphosphine)iridium						
	CH ₂ Cl ₂	3×10^8			PL/Ld-2	$S = TPP$.	94A126
12.126	Cobalt(II) acetate						
	CH ₃ COCH ₃ / H ₂ O (95:5)	4×10^7			PL/Ld-2	$S = PdMP$.	82A412
12.127	Cobalt(II) chloride hexahydrate						
	BuOCH ₂ CH ₂ OH	4.8×10^7		273	MD/A'c-33	$A' = Rub$; used $k_d = 3.8 \times 10^5$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319
12.128	Cobalt(II) ion						
	CH ₃ COCH ₃ / H ₂ O (95:5)	2×10^7			PL/Ld-2	$S = PdMP$.	82A412
12.129	Copper(II) ion						
	D ₂ O	6.4×10^7			PL/Ld-2	$S = RB$.	84N150
12.130	Cyclohexylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II)						
	C ₆ H ₅ CH ₃	3.4×10^8			CP/A'c-25	$S = A' = Rub$; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.131	Cyclohexylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II)				CP/A'c-25	$S = A' = Rub$; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.132	Dodecylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II)						
	C ₆ H ₅ CH ₃	3.2 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.133	1,1'-(7,13-Dimethyl-1,4,8,12-tetraazacyclopentadeca-4,7,12,15-tetraene-6,14-diyl)bis[ethanoato]nickel(II)						
	CHCl ₃	1.6 × 10 ⁹			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A050
12.134	Dodecylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II)						
	C ₆ H ₅ CH ₃	3.9 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.135	Dodecylamine[2,2'-thiobis(4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	2.6 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.136	2,2'-(1,2-Ethanediyl)bis(nitrilodecylidyne)bis[4-methylphenolato]nickel(II)						
	C ₆ H ₅ CH ₃	4.8 × 10 ⁹			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.137	2,2'-(1,2-Ethanediyl)bis(nitriloethylidyne)bis[phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	3.4 × 10 ⁹			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.138	[3,3]-[1,2-Ethanediyl]bis(nitrilomethylidyne)bis[5-ethyl-2-thiophenethionato]copper(II)						
	C ₆ H ₅ CH ₃	2.9 × 10 ⁸			PL/Ld-2	S = TPP.	88A507
	C ₆ H ₅ CH ₃	4.3 × 10 ⁵ (k_f)			CP/Ac,A'c-17	S = PP; A' = DPBF; used $k_f^{A'} = 6.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	88A507
	CH ₃ CN	3.4 × 10 ⁸			PL/Ld-2	S = TPP.	88A507
	CH ₃ CN	4.0 × 10 ⁵ (k_f)			CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_f^{A'} = 1.1 \times 10^9$ L mol ⁻¹ s ⁻¹ .	88A507
12.139	2,2'-(1,2-Ethanediyl)bis(nitrilomethylidyne)bis[phenolato]cobalt(II)						
	C ₆ H ₅ CH ₃	1.0 × 10 ¹⁰			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.140	2,2'-(1,2-Ethanediyl)bis(nitrilomethylidyne)bis[phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	5.3 × 10 ⁹			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.141	Ethylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II)						
	C ₆ H ₅ CH ₃	4.1 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.142	Ethylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	3.1 × 10 ⁸			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
12.143	[N,N'-Ethylenebis(5-sulfosalicylideneiminato)]nickelate(II) disodium salt						
	H ₂ O	1.6 × 10 ⁹			CR/A'c-32	A' = AES; used $k_{A'} = 4.7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_A/k_{A'} = 34$; ${}^1\text{O}_2^*$ from H ₂ O ₂ /OCl ⁻ .	85A177
12.144	Ferrocene						
	C ₆ H ₆	9 × 10 ⁶		198	PL/A'd-8	S = An; A' = DPBF.	737438
	CCl ₄ /CHCl ₃ (90:10)	3.0 × 10 ⁶			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_A/(k_A[A'] + k_d) = 1.7 \times 10^3$ L mol ⁻¹ at [A'] = 5×10^{-6} mol L ⁻¹ .	74F645
12.145	Hexaamminecobalt(III) ion						
	D ₂ O	1.2 × 10 ⁶			PL/Ld-2	S = RB.	84N150

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.146	Hydroxybis(dimethylglyoximato)pyridinecobalt(II)						
	CH ₂ Cl ₂ / MeOH (90:10)	4.5×10^7			PL/A'd-5	S = MB; A' = DPBF.	87F655
12.147	Hydroxybis(dimethylglyoximato)triethylaminecobalt(II)						
	CH ₂ Cl ₂ / MeOH (90:10)	1.5×10^8			PL/A'd-5	S = MB; A' = DPBF.	87F655
12.148	Hydroxybis(dimethylglyoximato)triphenylphosphinecobalt(II)						
	CH ₂ Cl ₂ / MeOH (90:10)	$<5 \times 10^6$			PL/A'd-5	S = MB; A' = DPBF.	87F655
12.149	Hydroxybis(diphenylglyoximato)pyridinecobalt(II)						
	CH ₂ Cl ₂ / MeOH (90:10)	2×10^7			PL/A'd-5	S = MB; A' = DPBF.	87F655
12.150	Hydroxybis(diphenylglyoximato)triphenylphosphinecobalt(II)						
	CH ₂ Cl ₂ / MeOH (90:10)	$<5 \times 10^6$			PL/A'd-5	S = MB; A' = DPBF.	87F655
12.151	Manganese(II) chloride hexahydrate						
	BuOCH ₂ CH ₂ OH	$<1.0 \times 10^6$		273	MD/A'c-33	A' = Rub; used $k_d = 3.8 \times 10^5$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319
12.152	Manganese(II) chloride tetrahydrate						
	<i>i</i> -octane	$\leq 1.0 \times 10^6$		298	CP/A'c-23	S = A' = Rub; used $k_d = 4.7 \times 10^4$ s ⁻¹ , $k_{A'} = 7.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	747341
12.153	[4-Methyl-1,2-benzenedithiolato](2,2'-bipyridine)nickel(II) [Ni(bpy)(DMT)]						
	DMF	7×10^9			MP/P'a-19	S = HP; A' = TEMP-4-OH; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; P' = TEMPOL; esr detection.	89F311
12.154	[4-Methyl-1,2-benzenedithiolato](1,10-phenanthroline)nickel(II)						
	DMF	1.1×10^{10}			MP/P'a-19	S = HP; A' = TEMP-4-OH; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; P' = TEMPOL; esr detection.	89F311
12.155	[4-Methyl-1,2-benzenedithiolato](1,10-phenanthroline)platinum(II) [Pt(phen)(DMT)]						
	DMF	4.7×10^9			CP/Ac-19	S = A; Q = Bis(diethylthiocarbamato)nickel(II); used $k_d = 1.4 \times 10^5$ s ⁻¹ .	91F098
12.156	2,2'-Methylenebis[<i>O,O'</i> -di(4- <i>tert</i> -butylphenyl)dithiophosphate]cobalt(II)						
	C ₆ H ₆	1.8×10^8			CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A434
12.157	2,2'-Methylenebis[<i>O,O'</i> -di(4- <i>tert</i> -butylphenyl)dithiophosphate]lead(II)						
	C ₆ H ₆	4.3×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A434
12.158	2,2'-Methylenebis[<i>O,O'</i> -di(4- <i>tert</i> -butylphenyl)dithiophosphate]nickel(II)						
	C ₆ H ₆	2.7×10^8			CP/A'c-23	S = A' = Rub; used $k_d = 2.8 \times 10^4$ s ⁻¹ , $k_{A'} = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A434
12.159	Methylthiolato(triethylphosphino)gold(I)						
	C ₆ H ₆	3.7×10^7		303	CP/P'a-19	S = A' = Rub; k_d not given.	87A107
	ClCF ₂ CCl ₂ F	4.5×10^7		303	CR/P'a-32	A' = Rub; $^1\text{O}_2^*$ from DMNO ₂ ; k_d and $k_{A'}$ not given.	87A107
12.160	2,2'-[1,8-Naphthylenebis(nitrilomethylidyne)]bis[phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	1.2×10^{10}			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
11.161	Nickel(II) acetate						
	CH ₃ COCH ₃ / H ₂ O (95:5)	2×10^7			PL/Ld-2	S = PdMP.	82A412
11.162	Nickel(II) dichloride						
	C ₅ H ₅ N	1.6×10^7			PL/Ld-2	S = PdMP.	82A412
	CH ₃ COCH ₃ / H ₂ O (95:5)	3×10^7			PL/Ld-2	S = PdMP.	82A412
11.163	Nickel(II) dichloride hexahydrate						
	BuOCH ₂ CH ₂ OH	3.1×10^8		273	MD/A'c-33	A' = Rub; used $k_d = 3.8 \times 10^5$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	727319
	C ₆ H ₆ / EtOH (50:50)	9.2×10^7 8.9×10^7 (k_q)		298	CP/Ac,A'c-23	S = A' = Rub; used $k_d = 5.6 \times 10^4$ s ⁻¹ , $k_{A'} = 3.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	90F360
11.164	Nickel(I) ion						
	D ₂ O pD = 10.6	3.3×10^7		295	PL/A'd-5	S = 2-ACN; A' = BR ²⁻	82N021
11.165	Nickelocene						
	tert-BuOH	2.8×10^9		295	PL/A'd-5	S = DMTBP; A' = DPBF.	87E181
	c-C ₆ H ₁₂	4.5×10^9		295	PL/A'd-5	S = DMTBP; A' = DPBF.	87E181
	C ₆ H ₆	5.4×10^9		295	PL/A'd-5	S = DMTBP; A' = DPBF.	87E181
	CCl ₄	2.7×10^9		295	PL/A'd-5	S = DMTBP; A' = DPBF.	87E181
	CH ₃ CN	6.4×10^9		295	PL/A'd-5	S = DMTBP; A' = DPBF.	87E181
	CH ₃ COCH ₃	4.8×10^9		295	PL/A'd-5	S = DMTBP; A' = DPBF.	87E181
	CHCl ₃	4.4×10^9		295	PL/A'd-5	S = DMTBP; A' = DPBF.	87E181
	MeOH	6.8×10^9		295	PL/A'd-5	S = DMTBP; A' = DPBF.	87E181
	2-ProOH	2.9×10^9		295	PL/A'd-5	S = DMTBP; A' = DPBF.	87E181
11.166	2,2'-[1,2-Phenylenebis(nitrilomethylidyne)]bis[phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	3.7×10^9			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
11.167	[N,N'-o-Phenylenebis(5-sulfosalicylideneiminato)]nickelate(II) disodium salt						
	H ₂ O	-10^8			CR/A'c-32	A' = AES; used $k_{A'} = 4.7 \times 10^7$ L mol ⁻¹ s ⁻¹ , meas. $k_A/k_{A'} = 10$; $^1\text{O}_2^*$ from H ₂ O ₂ /OCl ⁻ ; k is concn. dependent.	85A177
11.168	Propylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	3.0×10^8			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
11.169	[N,N'-Propylenebis(5-sulfosalicylideneiminato)]nickelate(II) disodium salt						
	H ₂ O	1.3×10^9			CR/A'c-32	A' = AES; used $k_{A'} = 4.7 \times 10^7$ L mol ⁻¹ s ⁻¹ , meas. $k_A/k_{A'} = 28$; $^1\text{O}_2^*$ from H ₂ O ₂ /OCl ⁻ .	85A177
11.170	Tetra-O-acetylglucose-1-thiolato(triethylphosphino)gold(I) (Auranofin)						
	C ₆ H ₆	2×10^6		303	CP/P'a-19	S = A' = Rub; k_d not given.	87A107
	ClCF ₂ CCl ₂ F	7.5×10^6		303	CR/P'a-32	A' = Rub; $^1\text{O}_2^*$ from DMNO ₂ ; $k_{A'}$ and k_d not given.	87A107
11.171	Triethanolamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II)						
	C ₆ H ₅ CH ₃	1.1×10^8			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063

TABLE 12. Rate constants for the interaction of singlet oxygen with metal complexes. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
12.172	Tris(acetylacetonato)chromate(III)						
	CCl ₄ / CHCl ₃ (90:10)	5.0×10^5			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_A/(k_A[A'] + k_d) = 3.0 \times 10^2$ L mol ⁻¹ at [A'] = 5×10^{-6} mol L ⁻¹ .	74F645
12.173	Tris(acetylacetonato)cobalt(III)						
	CCl ₄ / CHCl ₃ (90:10)	9.2×10^8			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_A/(k_A[A'] + k_d) = 5.2 \times 10^5$ L mol ⁻¹ at [A'] = 5×10^{-6} mol L ⁻¹ .	74F645
	CH ₂ Cl ₂ / MeOH (90:10)	6.5×10^8			PL/A'd-5	S = MB; A' = DPBF.	87F655
12.174	Tris(acetylacetonato)iron(III)						
	CCl ₄ / CHCl ₃ (90:10)	8.7×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 1.4 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Measured $k_A/(k_A[A'] + k_d) = 4.9 \times 10^4$ L mol ⁻¹ at [A'] = 5×10^{-6} mol L ⁻¹ .	74F645
12.175	Tris(acetylacetonato)manganese(III)						
	C ₆ H ₅ CH ₃	5×10^8			CP/A'c-25	S = A' = Rub; used $k_d = 1 \times 10^5$ s ⁻¹ , $k_{A'} = 1.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	752063
	CH ₃ COCH ₃ / H ₂ O (95:5)	1.6×10^8			PL/Ld-2	S = PdMP.	82A412
12.176	Tris(2,2'-bipyridine)cobalt(II) ion						
	CH ₃ COCH ₃ / H ₂ O (95:5)	$<1 \times 10^6$			PL/Ld-2	S = PdMP.	82A412
12.177	Tris(2,2'-bipyridine)ruthenium(II) dichloride						
	CCl ₄ / MeOH (98:2)	$\leq 1.0 \times 10^9$			FP/A'd-5	S = MB; A' = DPBF; no effect for [A] = 1×10^{-6} mol L ⁻¹ .	78E238
12.178	Tris(1,10-phenanthroline)cobalt(II) ion						
	CH ₃ COCH ₃ / H ₂ O (95:5)	$<1 \times 10^6$			PL/Ld-2	S = PdMP.	82A412

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds.

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.1	Azepine-1-carbodithioate ion, hexahydro- (Hexamethylenedithiocarbamate)						
	EtOH	1.5×10^8			CP/Oc-23	$S = RB; A' = Car$; used $k_d = 1.0 \times 10^4$ s ⁻¹ , $k_{A'} = 5.0 \times 10^5$ L mol ⁻¹ s ⁻¹ ; Measured $k_{A'}/(k_d + k_A [A]) = 7 \times 10^4$ L mol ⁻¹ at $[A] = 3.35 \times 10^{-4}$ mol L ⁻¹ .	727116
13.2	Benzamide, 2-(methylseleno)-N-phenyl-	9.2×10^6			PL/Ld-2	$S = H_2TPPS^4$.	91R262
13.3	Benzamide, 2-(methylthio)-N-phenyl-	7.8×10^5			PL/Ld-2	$S = H_2TPPS^4$.	91R262
13.4	Benzamide, N-phenyl-2-(phenylmethyl)seleno-				PL/Ld-2	$S = H_2TPPS^4$.	91R262
	CD ₃ OD	5.2×10^6			PL/Ld-2	$S = H_2TPPS^4$.	91R262
13.5	Benzamide, N-phenyl-2-(phenylmethyl)thio-	3.4×10^5			PL/Ld-2	$S = H_2TPPS^4$.	91R262
13.6	Benzanamine, N,N-dimethyl-4-(methylthio)-						
	C ₆ H ₆	3.0×10^7		308	CR/A'c-16	$A' = DPBF$; used $k_d = 5.4 \times 10^4$ s ⁻¹ ; $^1O_2^*$ from MNPO ₂ .	85M098
	MeOH/H ₂ O (50:50)	3.6×10^8		308	CR/A'c-16	$A' = DPBF$; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; $^1O_2^*$ from MNPO ₂ .	85M098
13.7	Benzene, 1-bromo-4-(methylthio)-						
	CHCl ₃	1.1×10^6			CP/A'c-33	$S = A' = Rub$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
13.8	Benzene, (butylthio)-						
	CH ₃ COCH ₃	1.6×10^6			PL/Ld-2	$S = RB$.	91F286 92F225
	CH ₃ COCH ₃	1.1×10^4 (k_t)			CP/Ac,A'c-17	$S = RB$; $A' = Limonene$; used $k_t^{A'} = 1.7 \times 10^5$ L mol ⁻¹ s ⁻¹ .	92F225
13.9	Benzene, 1-(butylthio)-3-chloro-						
	CH ₃ COCH ₃	5.3×10^5			PL/Ld-2	$S = RB$.	91F286 92F225
13.10	Benzene, 1-(butylthio)-4-chloro-						
	CH ₃ COCH ₃	7.7×10^5			PL/Ld-2	$S = RB$.	91F286 92F225
13.11	Benzene, 1-(butylthio)-4-fluoro-						
	CH ₃ COCH ₃	1.3×10^6			PL/Ld-2	$S = RB$.	91F286 92F225
13.12	Benzene, 1-(butylthio)-4-methoxy-						
	CH ₃ COCH ₃	4.3×10^6			PL/Ld-2	$S = RB$.	91F286 92F225
13.13	Benzene, 1-(butylthio)-4-methyl-						
	CH ₃ COCH ₃	2.8×10^6			PL/Ld-2	$S = RB$.	91F286 92F225
13.14	Benzene, 1-chloro-3-(methylthio)-						
	CHCl ₃	5.5×10^5			CP/A'c-33	$S = A' = Rub$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
13.15	Benzene, 1-chloro-4-(methylthio)-						
	CHCl ₃	1.0×10^6			CP/A'c-33	$S = A' = Rub$; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.	
13.15	Benzene, 1-chloro-4-(methylthio)-	— Continued						
	MeOH	8.3×10^5			CP/A'c-16	S = RB; A' = DPF; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	79A085	
13.16	Benzene, 1-chloro-4-(phenylthio)-				CP/Ac,Pa-17	S = RB; A' = (C ₆ H ₅) ₂ S; meas. $k_r/k_r^{A'} = 0.93$.	80F104	
13.17	Benzene, [(2,3-dimethyl-2-butenyl)thio]-				PL/Ad-5	S = RB.	89F298	
	CH ₃ COCH ₃	2.3×10^6						
13.18	Benzene, [(2,3-dimethyl-2-butenyl)thio]-4-methoxy-				PL/Ad-5	S = RB.	89F298	
	CH ₃ COCH ₃	7.6×10^6						
13.19	Benzene, [(2,3-dimethyl-2-butenyl)thio]-4-methyl-				PL/Ad-5	S = RB.	89F298	
	CH ₃ COCH ₃	3.1×10^6						
13.20	Benzene, 1-(1,1-dimethylethyl)-4-(methylthio)-				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085	
	CHCl ₃	4.7×10^6						
13.21	Benzene, 1-fluoro-4-(methylthio)-				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085	
13.22	Benzene, 1-methoxy-4-(methylthio)-							
	C ₆ H ₆	2×10^6		308	CR/A'c-16	A' = DPBF; used $k_d = 5.4 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	85M098	
	CHCl ₃	7.6×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085	
	MeOH	5.3×10^6			CP/A'c-16	S = RB; A' = DPF; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	79A086	
	MeOH/ H ₂ O (50:50)	4.6×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	85M098	
13.23	Benzene, 1-methoxy-4-(phenylthio)-				CP/Ac,Pa-17	S = RB; A' = (C ₆ H ₅) ₂ S; meas. $k_r/k_r^{A'} = 2.5$.	80F104	
	MeOH							
13.24	Benzene, 1-methyl-3-(methylthio)-				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085	
	CHCl ₃	3.1×10^6						
13.25	Benzene, 1-methyl-4-(methylthio)-							
	C ₆ H ₆	1.5×10^6		308	CR/A'c-16	A' = DPBF; used $k_d = 5.4 \times 10^4$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	85M098	
	CHCl ₃	4.6×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085	
	MeOH	3.1×10^6			CP/A'c-16	S = RB; A' = DPF; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	79A086	
	MeOH/ H ₂ O (50:50)	3.2×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; ¹ O ₂ * from MNPO ₂ .	85M098	
13.26	Benzene, 1-methyl-4-(phenylthio)-				CP/Ac,Pa-17	S = RB; A' = (C ₆ H ₅) ₂ S; meas. $k_r/k_r^{A'} = 1.9$.	80F104	
	MeOH							
13.27	Benzene, methylseleno-				296	CR/LI-12	used $k_d = 1.7 \times 10^3$ s ⁻¹ ; ¹ O ₂ * from DMNO ₂ .	90F069
	CCl ₄	1.4×10^8						
13.28	Benzene, methyltelluro-				296	CR/LI-12	used $k_d = 1.7 \times 10^3$ s ⁻¹ ; ¹ O ₂ * from DMNO ₂ .	90F069
	CCl ₄	3.8×10^9						

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.29	Benzene, (methylthio)-						
	C ₆ H ₆	8×10^5		308	CR/A'c-16	A' = DPBF; used $k_d = 5.4 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	85M098
	CCl ₄	7.1×10^6		296	CR/LI-12	used $k_d = 1.7 \times 10^3$ s ⁻¹ ; ${}^1\text{O}_2^*$ from DMNO ₂ .	90F069
	CHCl ₃	2.3×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
	MeOH	2.0×10^6			CP/A'c-16	S = RB; A' = DPF; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	79A086
	MeOH/H ₂ O (50:50)	2.3×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from MNPO ₂ .	85M098
13.30	Benzene, [(methylthio)methyl]-						
	CHCl ₃	1.2×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
13.31	Benzene, 1-nitro-4-(phenylthio)-						
	MeOH				CP/Ac,Pa-17	S = RB; A' = (C ₆ H ₅) ₂ S; meas. $k_r/k_r^{A'}$ = 0.30.	80F104
13.32	Benzene, [(phenylmethyl)seleno]-						
	CHCl ₃	6×10^6		253	CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82N064
13.33	Benzene, [(phenylimethyl)thio]-						
	CHCl ₃	8.5×10^5		253	CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82N064
13.34	Benzene, 1,1'-thiobis-						
	CHCl ₃	8.0×10^4			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
	MeOH	$\sim 1.0 \times 10^5$			CP/A'c-16	S = RB; A' = DPF; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; Theoretical intercept of 1.0 used in calculation.	79A086
13.35	Benzene methanesulfenamide, N,N-diethyl-						
	C ₆ H ₆	2.0×10^6			PL/Ld-2	S = TPP.	94F024
13.36	1,2-Benzisoselenazol-3-one, 7-chloro-2-phenyl-						
	CD ₃ OD	3.0×10^5			PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	91R262
13.37	1,2-Benzisoselenazol-3-one, 7-fluoro-2-phenyl-						
	CD ₃ OD	4.5×10^5			PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	91R262
13.38	1,2-Benzisoselenazol-3-one, 7-methoxy-2-phenyl-						
	CD ₃ OD	1.3×10^6			PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	91R262
13.39	1,2-Benzisoselenazol-3-one, 7-nitro-2-phenyl-						
	CD ₃ OD	1.1×10^5			PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	91R262
13.40	1,2-Benzisoselenazol-3-one, 2-phenyl-						
	CD ₃ OD	2.5×10^6			PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	91R262
13.41	1,2-Benzothiazol-3-one, 2-phenyl-						
	CD ₃ OD	2.3×10^5			PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	91R262
13.42	Benzyl mercaptan						
	MeOH	1.2×10^5	0.73		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
13.43	Bicyclo[2.2.1]heptane-2-thione						
	CHCl ₃	1.8×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A059 83F028

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.44	Bicyclo[2.2.1]heptane-2-thione, 3,3-dimethyl-						
	CHCl ₃	1×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A093 83F028
13.45	Bicyclo[2.2.1]heptane-2-thione, 1,3,3,7,7-pentamethyl-				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A093 83F028
13.46	Bicyclo[2.2.1]heptane-2-thione, 1,3,3-trimethyl- (Thiofenchone)				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A093 83F028
13.47	Bicyclo[2.2.1]heptane-2-thione, 1,7,7-trimethyl- (Thiocamphor)				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A093 83F028
13.48	Bicyclo[3.1.1]hept-3-ene-2-thione, 4,6,6-trimethyl-				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	85F151
13.49	2,3-Butanediol, 1,4-dimercapto-, <i>erythro</i> - (Dithioerythritol)						
	CHCl ₃ / EtOH (50:50)	7.4×10^8 2.9×10^6 (k_t)		310	CR/LI,Ac-12,14	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2$ * from NDPO ₂ .	90E622
13.50	2,3-Butanediol, 1,4-dimercapto-, <i>threo</i> - (Dithiothreitol)						
	CHCl ₃ / EtOH (50:50)	9.8×10^8 2.8×10^6 (k_t)		310	CR/LI,Ac-12,14	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2$ * from NDPO ₂ .	90E622
	D ₂ O pD = 7.4	1.5×10^6 (k_t)		310	CR/Ac-17	A' = CysSH; used $k_t^{A'} = 8.3 \times 10^6$ L mol ⁻¹ s ⁻¹ ; ${}^1\text{O}_2$ * from NDPO ₂ .	91R055
	D ₂ O pD = 7.4	1.9×10^6		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2$ * from NDPO ₂ .	91R055
13.51	1-Butanethiol						
	MeOH	5.3×10^4	1.7		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
13.52	1-Butanol, 4-(4-methylphenyl)thio-						
	CH ₃ COCH ₃	4.1×10^4 (k_t)			CP/Ac-17	S = RB; A' = Limonene; used $k_t^{A'} = 1.7 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_t = 64.9$.	91F286 92F225
	CH ₃ COCH ₃	2.7×10^6			PL/Ld-2	S = RB.	91F286 92F225
13.53	2-Butene, [(2-methyl-3-(phenylsulfinyl))-						
	C ₆ H ₆		11 (β_r)		CP/Ac-17	S = TPP; A' = Linalool; used $\beta_r^{A'} = 0.18$ mol L ⁻¹ .	89F400
13.54	2-Butene, 2-(phenylsulfinyl)- (E)						
	C ₆ H ₆		26 (β_r)		CP/Ac-17	S = TPP; A' = Linalool; used $\beta_r^{A'} = 0.18$ mol L ⁻¹ .	89F400
13.55	2-Butene, 2-(phenylsulfinyl)- (Z)						
	C ₆ H ₆		$>1 \times 10^2$ (β_r)		CP/Ac-17	S = TPP; A' = Linalool; used $\beta_r^{A'} = 0.18$ mol L ⁻¹ .	89F400
13.56	<i>tert</i> -Butyl ethyl sulfide						
	CHCl ₃	6.4×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
13.57	Butyl methyl sulfide						
	CHCl ₃	2.9×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.58 <i>tert</i> -Butyl propyl sulfide							
	CHCl ₃	7.2×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
13.59 <i>sec</i> -Butyl propyl sulfide							
	CHCl ₃	1.1×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
13.60 1,3-Cyclobutanedithione, 2,2,4,4-tetramethyl-							
	CHCl ₃	2.2×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A093
13.61 Cyclobutanethione, 2,2,4,4-tetramethyl-							
	CHCl ₃	3.1×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A093
13.62 Cyclobutanone, 2,2,4,4-tetramethyl-3-thioxo-							
	CHCl ₃	8.5×10^4			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A093
13.63 Cyclohexanamine, <i>N,N'</i> -[dithiobis[(5-ethyl-2,3-thiophenediy) methylidyne]]bis-							
	C ₆ H ₅ CH ₃	2.9×10^8			PL/Ld-2	S = TPP.	88A507
	C ₆ H ₅ CH ₃	3.2×10^8 (k_t)			CP/Ac,A'c-17	S = PP; A' = DPBF; used $k_t^{A'} = 6.7 \times 10^8$ L mol ⁻¹ s ⁻¹ .	88A507
	CH ₃ CN	1.1×10^8 (k_t)			CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_t^{A'} = 1.1 \times 10^9$ L mol ⁻¹ s ⁻¹ .	88A507
	CH ₃ CN	3.7×10^8			PL/Ld-2	S = TPP.	88A507
	CHCl ₃	5.6×10^7			PL/Ld-2	S = TPP.	88A507
13.64 Cyclohexanamine, <i>N</i> -[(5-methyl-2-(methylthio)-3-thienyl)methylene]-							
	C ₆ H ₅ CH ₃	2.5×10^5			PL/Ld-2	S = TPP.	88A507
	CH ₃ CN	2.0×10^5 (k_t)			CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_t^{A'} = 1.1 \times 10^9$ L mol ⁻¹ s ⁻¹ .	88A507
	CH ₃ CN	2.2×10^6			PL/Ld-2	S = TPP.	88A507
	CHCl ₃	6.0×10^5			PL/Ld-2	S = TPP.	88A507
	2-ProOH	3.4×10^5			PL/Ld-2	S = TPP.	88A507
13.65 2-Cyclohexene-1-thione, 3,5,5-trimethyl-							
	CHCl ₃	6.1×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	85F151
13.66 2-Cyclopentene-1-thione, 3-(4-methoxyphenyl)-							
	CHCl ₃	2.3×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	85F151
13.67 Cysteine							
	CHCl ₃ /EtOH (50:50)	2.4×10^8 7×10^7 (k_t)		310	CR/LI,Ac-12,14	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	90E622
	D ₂ O pD = 7.4	8.3×10^6		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	91R055
	D ₂ O	$\leq 4 \times 10^4$			PL/Ld-2	S = H ₂ TPPS ⁴⁻ or H ₂ TMpyP ⁴⁺ ; calcd. from $k_{obs} = 8.9 \times 10^6$ at pH 7 and $pK_a(\text{D}_2\text{O}) = 8.6$.	88A164
	D ₂ O	1×10^5			PL/Ld-2	S = RF, water-sol. TPP derivs., or Chl a in micelles contg. Triton X-100.	86F149
	D ₂ O/EtOH (75:25)	5×10^7		295	PL/Ld-2	S = RB.	94A113
	EtOH	8×10^6			CP/Pa-15	S = Ret; used $k_d = 3.5 \times 10^5$ s ⁻¹ ; CysS [•] obs. by esr.	85A443

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.68	Cysteine, negative ion						
	D ₂ O	1.5×10^8			PL/Ld-2	S = H ₂ TPPS ⁴⁻ or H ₂ TMpyP ⁴⁺ ; calcd. from k_{obs} = 8.9×10^6 at pH 7 and pK _a = 8.6.	88A164
13.69	L-Cysteine, N-acetyl-						
	D ₂ O pD = 7.4	$6 \times 10^5 (k_r)$		310	CR/Ac-17	A' = CysSH; used $k_r^{A'} = 8.3 \times 10^6$ L mol ⁻¹ s ⁻¹ ; $^1\text{O}_2^*$ from NDPO ₂ .	91R055
	D ₂ O pD = 7.4	6×10^5		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from NDPO ₂ .	91R055
	D ₂ O	$\leq 4 \times 10^4$			PL/Ld-2	S = H ₂ TPPS ⁴⁻ or H ₂ TMpyP ⁴⁺ ; calcd. from k_{obs} = 2.5×10^6 at pH 7 and pK _a (D ₂ O) = 10.0.	88A164
13.70	L-Cysteine, N-acetyl-, negative ion						
	D ₂ O	2.1×10^8			PL/Ld-2	S = H ₂ TPPS ⁴⁻ or H ₂ TMpyP ⁴⁺ ; calcd. from k_{obs} = 2.5×10^6 at pH 7 and pK _a = 8.6.	88A164
13.71	Dibenzyl sulfide						
	CHCl ₃	6.4×10^6		253	CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82N064
	MeOH	5.5×10^{-3}		293	CP/Oc-15	S = FICl ₄ ²⁻ ; $E_a = 1.7$ kJ mol ⁻¹ .	68F288
	MeOH	9.0×10^{-3}		293	CP/Oc-15	S = MB; $E_a = 7.1$ kJ mol ⁻¹ .	68F288
	MeOH	0.012		293	CP/Oc-15	S = RB; $E_a = 3.8$ kJ mol ⁻¹ .	68F288
	MeOH	6.7×10^{-3}		293	CP/Oc-15	S = FlBr ₄ Cl ₄ ²⁻ ; $E_a = 5.4$ kJ mol ⁻¹ .	68F288
	MeOH	8.0×10^{-3}		293	CP/Oc-15	S = DNT; $E_a = 6.7$ kJ mol ⁻¹ .	68F288
13.72	Diethyl sulfide						
	CHCl ₃	2.3×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
	EtOH			298	?	S = Chrysene, 2-bromochrysene, MB; meas. $k_q/k_r = \sim 0.7$.	74F648
	MeOH	2.5×10^7	7.2×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	85F480
	MeOH	8.2×10^6	0.011		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
13.73	Di-sec-butyl sulfide						
	CHCl ₃	1.8×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
13.74	Di-tert-butyl sulfide						
	CHCl ₃	1×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
	MeOH	$\sim 1.5 \times 10^5$			CP/A'c-16	S = RB; A' = DPF; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; Theoretical intercept of 1.0 used in calculation.	79A086
13.75	Diethyl disulfide						
	MeOH	1.6×10^7	5.5×10^{-3}		CP/A'c-16	S = MB; A' = DPBF; used $k_d = 9.0 \times 10^4$ s ⁻¹ .	72F519
13.76	Diethyl sulfide						
	(C ₂ H ₅) ₂ O	5×10^6	6.0×10^{-3}	296	CP/A'c-16	S = ZnTPP; A' = DPF; used $k_d = 3 \times 10^4$ s ⁻¹ .	82F357
			4.1×10^{-3}	195			
	C ₆ H ₅ CH ₃	1.4×10^7	2.9×10^{-3}	296	CP/A'c-16	S = ZnTPP; A' = DPF; used $k_d = 4.0 \times 10^4$ s ⁻¹ .	82F357
	C ₆ H ₆	2.0×10^7			CP/A'c-16	S = ZnTPP; A' = DPF; used $k_d = 4 \times 10^4$ s ⁻¹ .	79A086
	C ₆ H ₆	7.6×10^6			CP/Pa-20	S = ZnTPP; A' = Car; used $k_d = 4.0 \times 10^4$ s ⁻¹ , $k_{A'} = 1.3 \times 10^{10}$ L mol ⁻¹ s ⁻¹ ; Measured $k_{A'}/(k_d + k[A]) = 1.7 \times 10^4$ L mol ⁻¹ at [A] = 0.1 mol L ⁻¹ ; k calcd. in [81Z251].	71F580
	C ₆ H ₆				CP/Ac-20	S = ZnTPP; A' = DPA; meas. $k_A/k_{A'} = 6.0$.	71F580

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.76 Diethyl sulfide — Continued							
	C ₆ H ₆	0.032			CP/Pa-15	S = ZnTPP.	71F580
	C ₆ H ₆ / MeOH (80:20)	6×10^6 $3 \times 10^5 (k_r)$		298	CP/P'a-20	S = MB; A' = 2M2P; used $k_d = 1.0 \times 10^5$ s ⁻¹ , $\beta_{A'} = 0.04$ mol L ⁻¹ ; P = Diethyl sulfoxide.	70F734
	CH ₃ CN	2.1×10^7			PL/Ld-2	S = Ac.	89A099
	CH ₃ COCH ₃	2.7×10^7	8.9×10^{-4} 9.7×10^{-4}	296 195	CP/A'c-16	S = RB; A' = DPF; used $k_d = 2.4 \times 10^4$ s ⁻¹ .	82F357
	ClCF ₂ CCl ₂ F	3.2×10^6			PL/Ld-2	S = DNT.	87A072
	H ₂ O pH = 7.0	$2 \times 10^7 (k_r)$		292	MP/Ac,A'c-17	S = RB; A' = FFA; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	87A180
	MeOH	2.1×10^7	8.7×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	85F480
	MeOH	1.7×10^7	6.6×10^{-3} 0.011	296 195	CP/A'c-16	S = RB; A' = DPF; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	82F357
	MeOH	1.7×10^7			CP/A'c-16	S = MB; A' = DPF; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	79A086
	MeOH		0.02		CP/Pa-15	S = ZnTPP.	71F580
13.77 Diisobutyl sulfide							
	MeOH	2.9×10^7	6.1×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	85F480
13.78 Diisopropyl sulfide							
	CHCl ₃	2.2×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
	MeOH	2.5×10^6			CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.1 \times 10^5$ s ⁻¹ .	79A086
13.79 Dimethyldithiocarbamate ion							
	H ₂ O pH = 7.1	$\sim 8 \times 10^7$		298	CP/Oc-19	S = Phenosafranine; Q = NaN ₃ ; used $k_Q = 2.0 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A360
13.80 Dimethyl sulfide							
	C ₆ H ₆ / MeOH (50:50)			293	MP/Pa-17	S = TPP; rate is 19 times faster than in benzene.	81F311
	MeOH	5.8×10^7	3.1×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	85F480
13.81 Dimethyl sulfoxide							
	C ₆ H ₆ / MeOH (50:50)			293	MP/Pa-17	S = TPP; A' = CH ₃ SCH ₃ ; meas. $k_r/k_r^{A'} = 4.2 \times 10^{-3}$.	81F311
	C ₆ H ₆			293	MP/Pa-17	S = TPP; A' = CH ₃ SCH ₃ ; meas. $k_r/k_r^{A'} = 0.021$.	81F311
13.82 Dipropyl sulfide							
	MeOH	3.1×10^7	5.8×10^{-3}		CP/A'c-16	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ .	85F480
13.83 Diselenide, diphenyl-							
	CCl ₄	1.2×10^6		296	CR/LI-12	used $k_d = 1.7 \times 10^3$ s ⁻¹ ; ${}^1O_2^*$ from DMNO ₂ .	90F069
13.84 Disulfide, diphenyl-							
	CCl ₄	3.0×10^5		296	CR/LI-12	used $k_d = 1.7 \times 10^3$ s ⁻¹ ; ${}^1O_2^*$ from DMNO ₂ .	90F069
13.85 Ditelluride, diphenyl-							
	CCl ₄	1.1×10^8		296	CR/LI-12	used $k_d = 1.7 \times 10^3$ s ⁻¹ ; ${}^1O_2^*$ from DMNO ₂ .	90F069
13.86 1,5-Dithiacyclooctane							
	C ₆ H ₅ CH ₃	1.7×10^7		298	PL/Ld-2	S = TPP.	92F104
	CH ₃ COCH ₃	$3.7 \times 10^7 (k_r)$		298	CP/Pa-17	S = RB; A' = TME; used $k_r^{A'} = 2.7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	92F104
	CH ₃ COCH ₃	5.3×10^7		298	PL/Ld-2	S = RB.	92F104

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.86	1,5-Dithiacyclooctane	— Continued					
	CHCl ₃	8.2×10^6 (k_r)			CP/Pa-17	S = TPP; A' = TME; used $k_t^{A'} = 2.7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	92F103
	CHCl ₃	1.2×10^7			PL/Ld-2	S = TPP.	92F103
	2-PrOH	2.4×10^7		298	PL/Ld-2	S = RB.	92F104
13.87	1,5-Dithiacyclooctane 1-oxide						
	CH ₃ COCH ₃	2.4×10^5 (k_r)		298	CP/Pa-17	S = RB; A' = DMHD; used $k_t^{A'} = 7.4 \times 10^5$ L mol ⁻¹ s ⁻¹ .	92F104
	CH ₃ COCH ₃	3.3×10^6		298	PL/Ld-2	S = RB.	92F104
	CHCl ₃	8.7×10^4 (k_r)			CP/Pa-17	S = TPP; A' = 2M2P; used $k_t^{A'} = 7.6 \times 10^5$ L mol ⁻¹ s ⁻¹ .	92F103
	CHCl ₃	2.1×10^5			PL/Ld-2	S = TPP.	92F103
13.88	1,4-Dithiane						
	C ₆ H ₅ CH ₃	2.0×10^6		298	PL/Ld-2	S = TPP.	92F104
	CH ₃ CN		~0.050		CP/Pa-15	S = RB; Slope estimated as tangent to curve.	71F580
	CH ₃ CN		0.017		CP/Pa-15	S = RB; Solvent contains 0.52% H ₂ O by molarity.	71F580
	CH ₃ CN		9.8×10^{-3}		CP/Pa-15	S = RB; Solvent contains 0.05% H ₂ O by molarity.	71F580
	CH ₃ CN/H ₂ O (95:5)		6.7×10^{-3}		CP/Pa-15	S = RB.	71F580
	CH ₃ CN/H ₂ O (61:39)		0.017		CP/Pa-15	S = RB.	71F580
	CH ₃ COCH ₃	4.8×10^6		298	PL/Ld-2	S = RB.	92F104
	CH ₃ COCH ₃	2.4×10^5 (k_r)		293	CP/Pa-17	S = RB; A' = DMHD; used $k_t^{A'} = 7.4 \times 10^5$ L mol ⁻¹ s ⁻¹ ; $\Delta H^\ddagger = -26$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -226$ J K ⁻¹ mol ⁻¹ ; studied at 202–293 K; P = sulfoxide + sulfone.	91F170
	2-PrOH	2.1×10^6		298	PL/Ld-2	S = RB.	92F104
13.89	1,3-Dithiane-2-thione, 4,5-diphenyl-						
	C ₆ H ₆ / MeOH (83:17)		1.4×10^{-3}	293	CP/Oc-15	S = MB.	77F794
13.90	1,2-Dithiolane-3-pentanoic acid (Lipoic acid)						
	C ₆ H ₆	1.0×10^8	3.5×10^{-4}	298	CP/A'c-23	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	74F641
	CHCl ₃ / EtOH (50:50)	1.3×10^8		310	CR/LI-12	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	90E622
13.91	Ethanesulfonate ion, 2,2'-dithiobis- (Dimesna)						
	CHCl ₃ / EtOH (50:50)	6×10^6		310	CR/LI-12	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	90E622
13.92	Ethanesulfonate ion, 2-mercaptop-, (Mesna)						
	CHCl ₃ / EtOH (50:50)	2.4×10^7		310	CR/LI,Ac-12,14	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	90E622
		2.8×10^6 (k_r)					
	D ₂ O	2.7×10^6		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	91R055
	pD = 7.4						
13.93	Ethanethiol, 2-amino- (Cysteamine)						
	CHCl ₃ / EtOH (50:50)	4.1×10^7		310	CR/LI,Ac-12,14	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	90E622
		1.1×10^7 (k_r)					
	H ₂ O	8.3×10^6			PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	91R020

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_q/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.94	Ethanethiol, 2-[(3-aminopropyl)amino]-						
	CHCl ₃ / EtOH (50:50)	5.3×10^7		310	CR/LI, Ac- 12,14	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	90E622
		7.0×10^6 (k_t)					
	H ₂ O	7.4×10^6			PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	91R020
13.95	Ethanethiol, 2-hydroxy-,						
	D ₂ O	$\leq 4 \times 10^4$			PL/Ld-2	S = H ₂ TPPS ⁴⁻ or H ₂ TMpyP ⁴⁺ ; calcd. from $k_{obs} = 3.0 \times 10^5$ at pH 7 and $pK_a(D_2\text{O}) = 9.9$.	88A164
13.96	Ethanethiol, 2-hydroxy-, negative ion						
	D ₂ O	3.0×10^8			PL/Ld-2	S = H ₂ TPPS ⁴⁻ or H ₂ TMpyP ⁴⁺ ; calcd. from $k_{obs} = 3.0 \times 10^5$ at pH 7.	88A164
13.97	Ethanol, 2-(4-methylphenyl)thio-						
	CH ₃ COCH ₃	4.2×10^4 (k_t)			CP/Ac-17	S = RB; A' = Limonene; used $k_t^{A'} = 1.7 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_t = 34.8$.	91F286 92F225
	CH ₃ COCH ₃	1.5×10^6			PL/Ld-2	S = RB.	91F286 92F225
13.98	Fluorene, 9-(phenylsulfonyl)-, anion						
	tert-BuOH	4.6×10^{-4}		303	CP/Pa-15	S = RB; P = 9-Fluorenone.	707250
	tert-BuOH	3.8×10^{-4}		303	CP/Pa-15	S = A; P = 9-Fluorenone.	707250
13.99	Glutathione						
	CHCl ₃ / EtOH (50:50)	5.9×10^7		310	CR/LI, Ac- 12,14	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	90E622
		1.7×10^6 (k_t)					
	D ₂ O $pD = 7.4$	2.4×10^6		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	91R055
	D ₂ O	8.7×10^5			PL/Ld-2	S = MB ⁺ .	91R251
	D ₂ O	$\leq 4 \times 10^4$			PL/Ld-2	S = H ₂ TPPS ⁴⁻ or H ₂ TMpyP ⁴⁺ ; calcd. from $k_{obs} = 2.9 \times 10^6$ at pH 7 and $pK_a(D_2\text{O}) = 9.2$.	88A164
13.100	Glutathione, negative ion						
	D ₂ O	1.9×10^8			PL/Ld-2	S = H ₂ TPPS ⁴⁻ or H ₂ TMpyP ⁴⁺ ; calcd. from $k_{obs} = 2.9 \times 10^6$ at pH 7 and pK_a .	88A164
13.101	Glycine, N-(2-mercapto-1-oxopropyl)-						
	D ₂ O $pD = 7.4$	4.1×10^6		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	91R055
	D ₂ O	$\leq 4 \times 10^4$			PL/Ld-2	S = H ₂ TPPS ⁴⁻ or H ₂ TMpyP ⁴⁺ ; calcd. from $k_{obs} = 2.7 \times 10^6$ at pH 7 and $pK_a(D_2\text{O}) = 10.4$.	88A164
13.102	Glycine, N-(2-mercaptop-1-oxopropyl)-, negative ion						
	D ₂ O	1.8×10^8			PL/Ld-2	S = H ₂ TPPS ⁴⁻ or H ₂ TMpyP ⁴⁺ ; calcd. from $k_{obs} = 2.7 \times 10^6$ at pH 7 and pK_a .	88A164
13.103	1-Hexanol, 6-(4-methylphenyl)thio-						
	CH ₃ COCH ₃	6.1×10^3 (k_t)			CP/Ac-17	S = RB; A' = α -Pinene; used $k_t^{A'} = 4.3 \times 10^4$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_t = 359$.	91F286 92F225
	CH ₃ COCH ₃	2.2×10^6			PL/Ld-2	S = RB.	91F286 92F225
13.104	Imidazole-4-ethanaminium, α -carboxy-2,3-dihydro-N,N,N-trimethyl-2-thioxo-, S, (Ergothioneine)						
	D ₂ O $pD = 7$	2×10^7			PL/Ld-2	S = Carboxyanthracene.	92E225
	D ₂ O $pD = 7.4$	3.1×10^7		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	91R055

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.104	Imidazole-4-ethanaminium, α -carboxy-2,3-dihydro- <i>N,N,N</i> -trimethyl-2-thioxo-, <i>S</i> , (Ergothioneine) — Continued						
	D ₂ O pH = 7	2.3×10^7			PL/Ld-2	S = H ₂ TPPS ⁴⁻ or H ₂ TMpyP ⁴⁺ ; observed value; pK _a (D ₂ O) = 8.4.	88A164
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_t/k_t^{A'} = 1.1$; sensitizer immobilized on glass beads.	88R064
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_t/k_t^{A'} = 0.20$.	88R064
13.105	Imidazole-4-ethanaminium, α -carboxy-2,3-dihydro- <i>N,N,N</i> -trimethyl-2-thioxo-, ion	D ₂ O	5.0×10^8		PL/Ld-2	S = H ₂ TPPS ⁴⁻ or H ₂ TMpyP ⁴⁺ ; calcd. from $k_{obs} = 2.3 \times 10^7$ at pH 7 and pK _a (D ₂ O) = 8.4.	88A164
13.106	Imidazole-4-propionic acid, α -amino-2,3-dihydro-2-thioxo-, (<i>S</i>)	H ₂ O pH = 7.0		283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_t/k_t^{A'} = 1.1$; sensitizer immobilized on glass beads.	88R064
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_t/k_t^{A'} = 0.26$.	88R064
13.107	L-Leucine, L-methionyl-	D ₂ O/ EtOH (75:25)	2.1×10^7	295	PL/Ld-2	S = RB.	94A113
13.108	Methionine	CH ₃ CN/ H ₂ O (50:50)	$3 \times 10^6 (k_t)$		CP/Ac,A'c-17	S = RB; A' = DMA; $k_t^{A'}$ not given.	93R059
	CH ₃ CN/ H ₂ O (50:50)	9×10^6			PL/Ld-2	S = RB or Eos.	93R059
	D ₂ O pD = 7.4	1.3×10^7		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from NDPO ₂ .	91R055
	D ₂ O pD = 11.2	4.1×10^8		310	CR/A'c-32	A' = DPBF; used $k_d = 1.5 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from Dopamine/H ₂ O ₂ .	89M038
	D ₂ O pD = 11.2	2.1×10^8		310	CR/A'c-32	A' = DPBF; used $k_d = 1.5 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from DOPA/H ₂ O ₂ .	89M038
	D ₂ O pD = 7.4	1.5×10^7			PL/A'd-5	S = MB; A' = ADPA.	81N048
	D ₂ O pD = 8.4	$1.6 \times 10^7 (k_t)$		293	CL/Ad-35	used $k_d = 2.9 \times 10^4$ s ⁻¹ ; high pressure O ₂ (0.195 mol L ⁻¹); statistical error of 30% and systematic error of the same order.	79A112
	D ₂ O pD = 11	1.6×10^7			CP/A'c-16	S = RB; A' = DPF; k_d not given; plots are nonlinear, used slope as [A] → 0.	777433
	D ₂ O pD = 7	3.3×10^7			CP/A'c-16	S = RB; A' = DPF; k_d not given; plots are nonlinear, used slope as [A] → 0.	777433
	D ₂ O pD = 8.1	1.7×10^7		295	CL/A'c-22	A' = BRH ₂ ; used $k_d = 2.9 \times 10^4$ s ⁻¹ ; high pressure O ₂ ; recal. [79A112].	757147
	D ₂ O (mic) pD = 7.4	1.8×10^7			PL/A'd-5	S = 2-ACN; A' = DPBF; 0.1 mol L ⁻¹ SDS.	81N048
	D ₂ O (mic) pD = 7.4	1.6×10^7			PL/A'd-5	S = 2-ACN; A' = DPBF; 0.1 mol L ⁻¹ CTAB.	81N048
	D ₂ O/ EtOH (75:25)	1.3×10^7		295	PL/Ld-2	S = RB.	94A113
	H ₂ O pH = 10.6	2.8×10^7		310	CR/LI-12	used $k_d = 3.2 \times 10^5$ s ⁻¹ ; soln. cont. 5×10^{-4} mol L ⁻¹ CoCl ₂ ; $^1\text{O}_2^*$ from autoxidation of oxytetracycline.	92M228
	H ₂ O pH = 7	$2.1 \times 10^7 (k_t)$			CP/Oc-17	S = RB; A' = FFA; used $k_t^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93R059
	H ₂ O pH = 7	2.1×10^7			CP/A'c-18	S = RB; A' = TrpH; used $k_d = 2.5 \times 10^5$ s ⁻¹ , $k_{A'} = 6 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93R059

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.108	Methionine — Continued						
	H ₂ O pH = 10	4×10^7			CP/Ac-16	S = Eos; A' = 2,4-Cl ₂ C ₆ H ₃ O'; used $k_d = 5 \times 10^5$ s ⁻¹ .	87F537
	H ₂ O pH = 7.1	8.6×10^6		298	CP/Oc-19	S = Phenosafranine; Q = NaN ₃ ; used $k_Q = 2.0 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A360
	H ₂ O pH = 11	$\leq 5.0 \times 10^7$			CP/A'c-16	S = RB; A' = DPF; used $k_d = 1 \times 10^6$ s ⁻¹ ; Plots are nonlinear. Used slope as [A] → 0.	777433
	H ₂ O pH = 7	1.2×10^7			CP/A'c-16	S = RB; A' = DPF; used $k_d = 1 \times 10^6$ s ⁻¹ ; Plots are nonlinear. Used slope as [A] → 0.	777433
	H ₂ O pH = 6		0.052	298	CP/Ac-14	S = PF; β derived using $\phi_{isc}(PF) = 0.73$.	70F732
	H ₂ O		0.020	283	CP/Oc-15	S = MB; The mechanism of oxidation is not clear.	65F029
	H ₂ O/MeOH (50:50)	3×10^7			PL/A'd-5	S = MB; A' = DPBF.	72F516
	H ₂ O/MeOH (50:50)	5×10^6 (k_t)			CP/Ac,A'c-17	S = MB; A' = DPBF; used $k_t^{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; k_t derived using $k_{TQ}^{A'} = 3 \times 10^8$ L mol ⁻¹ s ⁻¹ and $k_A = 3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	72F516
13.109	Methionine, glycyl-						
	CH ₃ CN/H ₂ O (50:50)	1.5×10^6 (k_t)			CP/Ac,A'c-17	S = RB; A' = DMA; $k_t^{A'}$ not given.	93R059
	CH ₃ CN/H ₂ O (50:50)	1.5×10^7			PL/Ld-2	S = RB or Eos.	93R059
	D ₂ O/EtOH (75:25)	2.1×10^7		295	PL/Ld-2	S = RB.	94A113
	H ₂ O pH = 7	1.3×10^7 (k_t)			CP/Oc-17	S = RB; A' = FFA; used $k_t^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93R059
	H ₂ O pH = 7	9.0×10^7			CP/A'c-18	S = RB; A' = TrpH; used $k_d = 1.5 \times 10^5$ s ⁻¹ , $k_{A'} = 6 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93R059
13.110	Methionine, S-methyl-N-(phenylmethyl)carbonyl-, methyl ester						
	CHCl ₃	1.4×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
13.111	Methionine methyl ester						
	CH ₃ CN/H ₂ O (50:50)	1.2×10^6 (k_t)			CP/Ac,A'c-17	S = RB; A' = DMA; $k_t^{A'}$ not given.	93R059
	CH ₃ CN/H ₂ O (50:50)	1.4×10^7			PL/Ld-2	S = RB or Eos.	93R059
	H ₂ O pH = 7	1.0×10^8			CP/A'c-18	S = RB; A' = TrpH; used $k_d = 1.5 \times 10^5$ s ⁻¹ , $k_{A'} = 6 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93R059
	H ₂ O pH = 7	7×10^6 (k_t)			CP/Oc-17	S = RB; A' = FFA; used $k_t^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93R059
13.112	Morpholine, 4-[(phenylmethyl)thio]-						
	C ₆ H ₆	1.5×10^6 (k_t)			CP/Pa,P'a-17	S = TPP; A' = Octalin; used $k_t^{A'} = 1.8 \times 10^6$ L mol ⁻¹ s ⁻¹ ; P = 4-[(Phenylmethyl)sulfinyl]morpholine; $k_{obs} = 2.9 \times 10^6$ L mol ⁻¹ s ⁻¹ , $f_f^{A'} = 2$.	94F024
	C ₆ H ₆	1.5×10^6 (k_t)			CP/Pa,P'a-17	S = TPP; A' = Adamantylideneadamantane; used $k_t^{A'} = 3.5 \times 10^5$ L mol ⁻¹ s ⁻¹ ; P = 4-[(Phenylmethyl)sulfinyl]morpholine; $k_{obs} = 2.9 \times 10^6$ L mol ⁻¹ s ⁻¹ , $f_f^{A'} = 2$.	94F024
	C ₆ H ₆	1.3×10^6			PL/Ld-2	S = TPP; product formation rate studies suggest no physical quenching.	94F024

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.113	2-Naphthalenethione, 1,2-dihydro-1,1-dimethyl-						
	CHCl ₃	4.3×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	85F151
13.114	2-Naphthalenethione, 1,2-dihydro-1,1,3-trimethyl-				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	85F151
13.115	Octanoate ion, 6,8-dimercapto- (Dihydrolipoate ion)						
	CHCl ₃ / EtOH (50:50)	5×10^6		310	CR/LI,Ac-12,14	used $k_d = 1 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from NDPO ₂ .	90E622
		$5 \times 10^6 (k_r)$					
13.116	Penicillamine						
	D ₂ O pD = 7.4	4.6×10^6		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from NDPO ₂ .	91R055
	H ₂ O	2.4×10^6			PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	91R020
13.117	3-Pentanethione, 2,2,4,4-tetramethyl-				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A093
	CHCl ₃	1.0×10^4					
13.118	1-Pentanol, 5-(4-methylphenyl)thio-				CP/Ac-17	S = RB; A' = α -Pinene; used $k_r^{A'} = 4.3 \times 10^4$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_r = 192$.	91F286 92F225
	CH ₃ COCH ₃	$1.5 \times 10^4 (k_r)$			PL/Ld-2	S = RB.	91F286 92F225
13.119	3-Pentanol, 3,4-dimethyl-1-[(4-methylphenyl)thio]-				CP/Ac-17	S = RB; A' = Limonene; used $k_r^{A'} = 1.7 \times 10^5$ L mol ⁻¹ s ⁻¹ ; $\Delta H^\ddagger = -22$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -226$ J K ⁻¹ mol ⁻¹ ; studied at 195-293 K; $k_r = 9.7 \times 10^5$ L mol ⁻¹ s ⁻¹ at 223 K when A' = Cyclopentene and $k_r^{A'} = 3.9 \times 10^4$ L mol ⁻¹ s ⁻¹ .	91F170
	CH ₃ COCH ₃	$6.5 \times 10^4 (k_r)$		293			
	CH ₃ COCH ₃	3.7×10^6			PL/Ld-2	S = RB.	91F286 92F225
	CH ₃ COCH ₃	$7.1 \times 10^4 (k_r)$			CP/Ac-17	S = RB; A' = Limonene; used $k_r^{A'} = 1.7 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_r = 51.1$.	91F286 92F225
13.120	Phenol, 4-(methylthio)-						
	C ₆ H ₆	1.1×10^7		308	CR/A'c-16	A' = DPBF; used $k_d = 5.4 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	85M098
	MeOH/ H ₂ O (50:50)	$1.3 \times 10^8 (k_r)$		308	CR/A'c-17	A' = DPBF; $^1\text{O}_2^*$ from MNPO ₂ . $k_r^{A'}$ not given.	85M098
	MeOH/ H ₂ O (50:50)	1.6×10^8		308	CR/A'c-16	A' = DPBF; used $k_d = 2.8 \times 10^5$ s ⁻¹ ; $^1\text{O}_2^*$ from MNPO ₂ .	85M098
13.121	Phenothiazine						
	C ₆ H ₅ Br/ MeOH (67:33)	4.2×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 4.9 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777240
	MeOH/ C ₆ H ₆ (90:10)			288	CP/Pa-17	S = MB; measured ($\beta_r/\beta_r(\text{MeOH})$) = 0.94.	757623
	MeOH/ C ₆ H ₆ (80:20)			288	CP/Pa-17	S = MB; measured ($\beta_r/\beta_r(\text{MeOH})$) = 1.0.	757623
	MeOH/ C ₆ H ₆ (20:80)			288	CP/Pa-17	S = MB; measured ($\beta_r/\beta_r(\text{MeOH})$) = 1.5.	757623
	MeOH/ HOCH ₂ CH ₂ OH (25:75)			288	CP/Pa-17	S = MB; measured ($\beta_r/\beta_r(\text{MeOH})$) = 0.34.	757623

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.121	Phenothiazine — Continued						
	MeOH/ HOCH ₂ CH ₂ OH (75:25)			288	CP/Pa-17	S = MB; measured (β_s/β_t (MeOH)) = 0.51.	757623
13.122	Phenothiazine, 2-chloro-10-dimethylaminopropyl- (Chlorpromazine)	3.5×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 4.9 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777240
13.123	Phenothiazine, 10-methyl-						
	C ₆ H ₅ Br/ MeOH (67:33)	$\leq 1.2 \times 10^6$			CP/A'c-23	S = A' = Rub; used $k_d = 4.9 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	777240
	CHCl ₃	2.9×10^6		253	CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82N064
	H ₂ O/1-BuOH/c- C ₆ H ₁₂ (72:16:4) (mic)	8.4×10^4 (k_r)		289	CP/LI,Pa- 12,27	S = RB; used $k_d = 1.6 \times 10^5$ s ⁻¹ ; P = 10-Methylphenothiazine-5-oxide; 8% SDS; used $k_q = 5.0 \times 10^6$ L mol ⁻¹ s ⁻¹ , $\phi_\Delta(RB) = 0.75$.	87N090
	MeOD	5.3×10^6 2.7×10^5 (k_r) 5.0×10^6 (k_q)		289	CP/LI,Pa- 12,27	S = RB; used $k_d = 4.4 \times 10^3$ s ⁻¹ ; P = 10-Methylphenothiazine-5-oxide; used $\phi_\Delta(RB) = 0.75$.	87N090
13.124	Phenothiazine, 10-(12-sulfonatododecyl)-						
	H ₂ O (mic)	4.9×10^5 (k_r) 1.1×10^6 (k_q)		289	CP/LI,Pa- 12,27	S = RB; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; P = 10-(12-Sulfonatododecyl)phenothiazine-9-oxide; [A] = 10^{-2} mol L ⁻¹ ; counter ion Na ⁺ ; used $\phi_\Delta(RB) = 0.75$.	87N090
13.125	Piperidine, 1-[(phenylimethyl)thio]-						
	C ₆ H ₆	3.6×10^6			PL/Ld-2	S = TPP.	94F024
13.126	Pivalothiophenone						
	CHCl ₃	4.3×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A093
13.127	Pivalothiophenone, 4'-chloro-				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A093
13.128	Pivalothiophenone, 4'-fluoro-				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A093
13.129	Pivalothiophenone, 4'-methoxy-				CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82A093
13.130	L-Proline, 1-(3-mercaptop-2-methyl-1-oxopropyl)-, (S)- (Captopril)						
	CHCl ₃ / EtOH (50:50)	4×10^6		310	CR/LI-12	used $k_d = 1 \times 10^5$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	90E622
	D ₂ O pD = 7.4	$<1 \times 10^5$		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	91R055
13.131	1-Propanethiol, 3-amino-2-hydroxy-						
	H ₂ O	8×10^5			PL/Ld-2	S = H ₂ TPPS ⁺ .	91R020
13.132	1-Propanol, 3-(3-chlorophenyl)thio-				PL/Ld-2	S = RB.	91F286 92F225

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.133	1-Propanol, 3-(4-chlorophenyl)thio-						
	CH ₃ COCH ₃	8.6×10^5			PL/Ld-2	S = RB.	91F286 92F225
13.134	1-Propanol, 3-(4-fluorophenyl)thio-						
	CH ₃ COCH ₃	1.3×10^6			PL/Ld-2	S = RB.	91F286 92F225
13.135	1-Propanol, 3-(4-methoxyphenyl)thio-						
	CH ₃ COCH ₃	4.1×10^6			PL/Ld-2	S = RB.	91F286 92F225
13.136	1-Propanol, 3-(4-methylphenyl)thio-						
	CH ₃ COCH ₃	7.1×10^4 (k_r)			CP/Ac-17	S = RB; A' = Limonene; used $k_r^{A'} = 1.7 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_q/k_r = 31.3$.	91F286 92F225
	CH ₃ COCH ₃	2.3×10^6			PL/Ld-2	S = RB.	91F286 92F225
13.137	1-Propanol, 3-(phenylthio)-						
	CH ₃ COCH ₃	3.3×10^4 (k_r)			CP/Ac-17	S = RB; A' = α -Pinene; used $k_r^{A'} = 4.3 \times 10^4$ L mol ⁻¹ s ⁻¹ .	92F225
	CH ₃ COCH ₃	1.5×10^6			PL/Ld-2	S = RB.	91F286 92F225
13.138	Pyran-4-thione						
	CH ₂ Cl ₂	2×10^{-3}		298	CP/Pa-15	S = MB; P = Pyran-4-one.	717403
13.139	Pyran-4-thione, 2,6-dimethyl-						
	CH ₂ Cl ₂	3×10^{-4}		298	CP/Pa-15	S = MB; P = 2,6-Dimethylpyran-4-one.	717403
13.140	Pyran-4-thione, 2,6-diphenyl-						
	CH ₂ Cl ₂	1.4×10^{-3}		298	CP/Ac-15	S = MB.	717403
13.141	Pyrazole-3-selone, 4-(aminomethylene)-2,4-dihydro-5-methyl-2-phenyl-, (Z)-						
	1-BuOH	1.8×10^8 (k_r)	6.1×10^{-4} (β_r)	295	CP/Ac-14	S = A; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	c-C ₆ H ₁₂	1.2×10^9 (k_r)	4.8×10^{-5} (β_r)	295	CP/Ac-14	S = A; used $k_d = 5.9 \times 10^4$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	CCl ₄	1.3×10^6 (k_r)	3×10^{-5} (β_r)	295	CP/Ac-14	S = A; used $k_d = 38$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	CDCl ₃	2.6×10^7 (k_r)	4.5×10^{-5} (β_r)	295	CP/Ac-14	S = A; used $k_d = 1.2 \times 10^3$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	CH ₃ CN	1.3×10^8 (k_r)	1.3×10^{-4} (β_r)	295	CP/Ac-14	S = A; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	CHCl ₃	2.0×10^7 (k_r)	2.0×10^{-4} (β_r)	295	CP/Ac-14	S = A; used $k_d = 4.0 \times 10^3$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	EtOH	2.8×10^8 (k_r)	3.5×10^{-4} (β_r)	295	CP/Ac-14	S = A; used $k_d = 1 \times 10^5$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	EtOH	1.3×10^8 (k_r)	7.4×10^{-4} (β_r)	295	CP/Ac-14	S = Ery; used $k_d = 1 \times 10^5$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	EtOH	2.2×10^8 (k_r)	4.5×10^{-4} (β_r)	295	CP/Ac-14	S = Eos; used $k_d = 1 \times 10^5$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	EtOH	7×10^7 (k_r)	1.4×10^{-3} (β_r)	295	CP/Ac-14	S = Rhodamine S; used $k_d = 1 \times 10^5$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	EtOH	8×10^7 (k_r)	1.2×10^{-3} (β_r)	295	CP/Ac-14	S = MB; used $k_d = 1 \times 10^5$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	MeOH	2.6×10^8 (k_r)	5.5×10^{-4} (β_r)	295	CP/Ac-14	S = A; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	MeOH-d ₃		3.1×10^{-4} (β_r)	295	CP/Ac-14	S = A; used $\phi_\Delta(S)$.	85F646
	2-PrOH		3.1×10^{-4} (β_r)	295	CP/Ac-14	S = A; used $\phi_\Delta(S)$.	85F646
	dioxane	3.6×10^8 (k_r)	9.1×10^{-5} (β_r)	295	CP/Ac-14	S = A; used $k_d = 3.3 \times 10^4$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
13.142	Pyrazole-3-thione, 4-(aminomethylene)-2,4-dihydro-5-methyl-2-phenyl-						
	1-BuOH	9.2×10^7 (k_r)	1.2×10^{-3} (β_r)	295	CP/Ac-14	S = A; used $k_d = 1.1 \times 10^5$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	c-C ₆ H ₁₂	5×10^8 (k_r)	1.1×10^{-4} (β_r)	295	CP/Ac-14	S = A; used $k_d = 5.9 \times 10^4$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	C ₆ H ₅ CH ₃	1.4×10^8 (k_r)	2.7×10^{-4} (β_r)		CP/Ac-14	S = A; used $k_d = 4.2 \times 10^4$ s ⁻¹ ; used $\phi_\Delta(S)$.	84F244

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.142 Pyrazole-3-thione, 4-(aminomethylene)-2,4-dihydro-5-methyl-2-phenyl- — Continued							
	<i>n</i> -C ₇ H ₁₆	$7.0 \times 10^8 (k_r)$	$8.4 \times 10^{-5} (\beta_r)$		CP/Ac-14	S = A; used $k_d = 5.9 \times 10^4$ s ⁻¹ ; used $\phi_\Delta(S)$.	84F244
	CCl ₄	$1.9 \times 10^6 (k_r)$	$2 \times 10^{-5} (\beta_r)$	295	CP/Ac-14	S = A; used $k_d = 38$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	CDCl ₃	$2.4 \times 10^7 (k_r)$	$5.0 \times 10^{-5} (\beta_r)$	295	CP/Ac-14	S = A; used $k_d = 1.2 \times 10^3$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	CH ₃ CN	$1.3 \times 10^8 (k_r)$	$1.3 \times 10^{-4} (\beta_r)$	295	CP/Ac-14	S = A; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	CH ₃ CN	$2.8 \times 10^8 (k_r)$	$6.7 \times 10^{-5} (\beta_r)$		CP/Ac-14	S = A; used $k_d = 3.3 \times 10^4$ s ⁻¹ ; used $\phi_\Delta(S)$.	84F244
	CHCl ₃	$1.9 \times 10^7 (k_r)$	$2.1 \times 10^{-4} (\beta_r)$	295	CP/Ac-14	S = A; used $k_d = 4.0 \times 10^3$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	CHCl ₃	$7.4 \times 10^7 (k_r)$	$2.3 \times 10^{-4} (\beta_r)$		CP/Ac-14	S = A; used $k_d = 1.7 \times 10^4$ s ⁻¹ ; used $\phi_\Delta(S)$.	84F244
	EtOH	$8.3 \times 10^7 (k_r)$	$1.2 \times 10^{-3} (\beta_r)$	295	CP/Ac-14	S = A; used $k_d = 1 \times 10^5$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	EtOH/H ₂ O (96:4)	$8.5 \times 10^7 (k_r)$	$9.7 \times 10^{-4} (\beta_r)$		CP/Ac-14	S = A; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; used $\phi_\Delta(S)$.	84F244
	MeOH	$8.4 \times 10^7 (k_r)$	$1.7 \times 10^{-3} (\beta_r)$	295	CP/Ac-14	S = A; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	MeOH	$8.7 \times 10^7 (k_r)$	$1.6 \times 10^{-3} (\beta_r)$		CP/Ac-14	S = A; used $k_d = 1.4 \times 10^5$ s ⁻¹ ; used $\phi_\Delta(S)$.	84F244
	MeOH- <i>d</i> ₃		$8.1 \times 10^{-4} (\beta_r)$	295	CP/Ac-14	S = A; used $\phi_\Delta(S)$.	85F646
	2-PrOH		$7.3 \times 10^{-4} (\beta_r)$	295	CP/Ac-14	S = A; used $\phi_\Delta(S)$.	85F646
	dioxane	$1.0 \times 10^8 (k_r)$	$3.3 \times 10^{-4} (\beta_r)$	295	CP/Ac-14	S = A; used $k_d = 3.3 \times 10^4$ s ⁻¹ ; used $\phi_\Delta(S)$.	85F646
	dioxane	$2.0 \times 10^8 (k_r)$	$1.5 \times 10^{-4} (\beta_r)$		CP/Ac-14	S = A; used $k_d = 3.1 \times 10^4$ s ⁻¹ ; used $\phi_\Delta(S)$.	84F244
13.143 Thiacyclohex-2-ene, 2-methyl-, 1-oxide							
	C ₆ H ₆		$>1 \times 10^2 (\beta_r)$		CP/Ac-17	S = TPP; A' = Linalool; used $\beta_r^{A'} = 0.18$ mol L ⁻¹ .	89F400
13.144 Thiacyclopent-2-ene, 2-methyl-, 1-oxide							
	C ₆ H ₆		$>1 \times 10^2 (\beta_r)$		CP/Ac-17	S = TPP; A' = Linalool; used $\beta_r^{A'} = 0.18$ mol L ⁻¹ .	89F400
13.145 Thiane							
	C ₆ H ₅ CH ₃	1.3×10^7		298	PL/Ld-2	S = TPP.	92F104
	C ₆ H ₅ CH ₃	$2 \times 10^7 (k_r)$		198	CP/Pa-17	S = TPP; A' = TME; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; $\Delta H^\ddagger = -21$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -205$ J K ⁻¹ mol ⁻¹ ; studied at 198-293 K; P = sulfoxide + sulfone.	91F170
	CH ₃ COCH ₃	1.6×10^7		298	PL/Ld-2	S = RB.	92F104
	CH ₃ COCH ₃	$4.7 \times 10^5 (k_r)$		298	CP/Pa-17	S = RB; A' = TME; used $k_r^{A'} = 2.7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	92F104
	CH ₃ COCH ₃	$9.4 \times 10^6 (k_r)$		198	CP/Pa-17	S = RB; A' = TME; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; $\Delta H^\ddagger = -17$ kJ mol ⁻¹ ; $\Delta S^\ddagger = -192$ J K ⁻¹ mol ⁻¹ ; studied at 198-293 K; P = sulfoxide + sulfone.	91F170
	CHCl ₃	1.3×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
	2-PrOH	1.4×10^7		298	PL/Ld-2	S = RB.	92F104
13.146 Thiepane							
	CHCl ₃	1.3×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085
13.147 Thiepin, 4,5-didehydro-2,3,6,7-tetrahydro-3,3,6,6-tetramethyl-							
	CHCl ₃	3.5×10^6			CR/A'c-33	A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from DMNO ₂ .	81E003
13.148 Thiepin, 4,5-didehydro-2,3,6,7-tetrahydro-3,3,6,6-tetramethyl-, 1,1-dioxide							
	CHCl ₃	3×10^5			CR/A'c-33	A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from DMNO ₂ .	81E003
13.149 Thiepin, 4,5-didehydro-2,3,6,7-tetrahydro-3,3,6,6-tetramethyl-, 1-oxide							
	CHCl ₃	7.2×10^4			CR/A'c-33	A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from DMNO ₂ .	81E003

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.150	Thiepin, 2,3,6,7-tetrahydro-3,3,6,6-tetramethyl-						
	CHCl ₃	3.5×10^6			CR/A'c-33	$A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$; ${}^1\text{O}_2^*$ from DMNO ₂ .	81E003
13.151	Thiirane, 2,3-diphenyl-, (E)						
	ClCF ₂ CCl ₂ F	4.3×10^3			PL/Ld-2	S = DNT.	87A072
13.152	Thiirane, methyl-						
	ClCF ₂ CCl ₂ F	6.6×10^4			PL/Ld-2	S = DNT.	87A072
13.153	Thiobenzamide, 4-chloro-						
	CH ₂ Cl ₂			273	CP/Ac,A'c-17	S = Poly-RB; $A' = \text{C}_6\text{H}_5\text{CSNH}_2$; meas. $k_t/k_t^{A'} = 0.80$.	84F196
13.154	Thiobenzamide, 4-methoxy-						
	CH ₂ Cl ₂			273	CP/Ac,A'c-17	S = Poly-RB; $A' = \text{C}_6\text{H}_5\text{CSNH}_2$; meas. $k_t/k_t^{A'} = 1.1$.	84F196
13.155	Thiobenzamide, 4-methyl-						
	CH ₂ Cl ₂			273	CP/Ac,A'c-17	S = Poly-RB; $A' = \text{C}_6\text{H}_5\text{CSNH}_2$; meas. $k_t/k_t^{A'} = 1.0$.	84F196
13.156	Thiobenzamide, 4-nitro-						
	CH ₂ Cl ₂			273	CP/Ac,A'c-17	S = Poly-RB; $A' = \text{C}_6\text{H}_5\text{CSNH}_2$; meas. $k_t/k_t^{A'} = 0.45$.	84F196
13.157	Thiobenzamide, 4-(trifluoromethyl)-						
	CH ₂ Cl ₂			273	CP/Ac,A'c-17	S = Poly-RB; $A' = \text{C}_6\text{H}_5\text{CSNH}_2$; meas. $k_t/k_t^{A'} = 0.52$.	84F196
13.158	Thiobenzophenone, 4-chloro-						
	CHCl ₃	1.1×10^6			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	82A093
13.159	Thiobenzophenone, 4,4'-dichloro-						
	CHCl ₃	2×10^5			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	82A093
13.160	Thiobenzophenone, 4,4'-dimethoxy-						
	CHCl ₃	9.6×10^6			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	82A093
13.161	Thiobenzophenone, 4,4'-dimethyl-						
	CHCl ₃	3.7×10^6			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	82A093
13.162	Thiobenzophenone, 4-methoxy-						
	CHCl ₃	5.4×10^6			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	82A093
13.163	Thiobenzophenone, 4-phenyl-						
	CHCl ₃	3.1×10^6			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	82A093
13.164	2,2'-Thiodiethanol						
	MeOH	0.016		293	CP/Oc-15	$S = \text{FeCl}_4^{2-}$; $E_a = 6.3 \text{ kJ mol}^{-1}$.	68F288
	MeOH	0.013		293	CP/Oc-15	$S = \text{FeBr}_4\text{Cl}_4^{2-}$; $E_a = 3.8 \text{ kJ mol}^{-1}$.	68F288
	MeOH	0.010		293	CP/Oc-15	$S = \text{DNT}$; $E_a = 8.4 \text{ kJ mol}^{-1}$.	68F288
	MeOH	0.022		293	CP/Oc-15	$S = \text{MB}$; $E_a = 7.5 \text{ kJ mol}^{-1}$.	68F288
	MeOH	0.025		293	CP/Oc-15	$S = \text{RB}$; $E_a = 5.4 \text{ kJ mol}^{-1}$.	68F288

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.165	Thioketene, di- <i>tert</i> -butyl-						
	CHCl ₃	1.2×10^7			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	84A070
13.166	Thioketene, 2,2,6,6-tetramethylcyclohexyl-				CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	84A070
13.167	Thiophene						
	MeOH	2.0×10^5	0.45		CP/A'c-16	$S = \text{MB}$; $A' = \text{DPBF}$; used $k_d = 9.0 \times 10^4 \text{ s}^{-1}$.	72F519
	MeOH		$\geq 6.0 \times 10^2$	293	CP/Oc-15	$S = \text{RB}$; $E_a = 25 \text{ kJ mol}^{-1}$.	68F288
13.168	Thiophene, tetrahydro-						
	CHCl ₃	4.2×10^7			CP/A'c-33	$S = A' = \text{Rub}$; used $k_d = 1.7 \times 10^4 \text{ s}^{-1}$, $k_{A'} = 5.3 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$.	79A085
	MeOH	2.8×10^6	0.032		CP/A'c-16	$S = \text{MB}$; $A' = \text{DPBF}$; used $k_d = 9.0 \times 10^4 \text{ s}^{-1}$.	72F519
13.169	2,5-Thiophenedithione, 3,4-bis[(cyclohexylamino)methylene]dihydro-, (Z,Z)						
	C ₆ H ₅ CH ₃	1.0×10^7 (k_r)			CP/Ac,A'c-17	$S = \text{PP}$; $A' = \text{DPBF}$; used $k_r^{A'} = 6.7 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	88A507
	C ₆ H ₅ CH ₃	9.0×10^7			PL/Ld-2	$S = \text{TPP}$.	88A507
	CH ₃ CN	4.0×10^6 (k_r)			CP/Ac,A'c-17	$S = \text{MB}$; $A' = \text{DPBF}$; used $k_r^{A'} = 1.1 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$.	88A507
	CH ₃ CN	1.6×10^8			PL/Ld-2	$S = \text{TPP}$.	88A507
	CHCl ₃	4.8×10^7			PL/Ld-2	$S = \text{TPP}$.	88A507
13.170	2-Thiophenethione, 3-[(cyclohexylamino)methylene]-5-ethyl-, (Z)-						
	C ₆ H ₅ CH ₃	1.8×10^8 (k_r)			CP/Ac,A'c-17	$S = \text{PP}$; $A' = \text{DPBF}$; used $k_r^{A'} = 6.7 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$.	88A507
	C ₆ H ₅ CH ₃	1.6×10^8			PL/Ld-2	$S = \text{TPP}$.	88A507
	CH ₃ CN	7.7×10^8 (k_r)			CP/Ac,A'c-17	$S = \text{MB}$; $A' = \text{DPBF}$; used $k_r^{A'} = 1.1 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$.	88A507
	CH ₃ CN	9.0×10^8			PL/Ld-2	$S = \text{TPP}$.	88A507
	CHCl ₃	2.5×10^8			PL/Ld-2	$S = \text{TPP}$.	88A507
	2-PrOH	5.5×10^8			PL/Ld-2	$S = \text{TPP}$.	88A507
13.171	2-Thiophenethione, 3,3'-[1,2-ethanediylbis(iminomethylidyne)]bis[5-ethyl-, (Z,Z)]						
	C ₆ H ₅ Cl	2.1×10^8			PL/Ld-2	$S = \text{TPP}$.	88A507
	CH ₃ CN	5.8×10^8			PL/Ld-2	$S = \text{TPP}$.	88A507
	CH ₃ CN	2.7×10^8 (k_r)			CP/Ac,A'c-17	$S = \text{MB}$; $A' = \text{DPBF}$; used $k_r^{A'} = 1.1 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$.	88A507
	CHCl ₃	2.0×10^8			PL/Ld-2	$S = \text{TPP}$.	88A507
13.172	Thiopyran-4-thione						
	CH ₂ Cl ₂	$<1.0 \times 10^{-4}$		298	CP/Pa-15	$S = \text{MB}$; P = Thiopyran-4-one.	717403
13.173	Thiopyran-4-thione, 2,6-diphenyl-						
	CH ₂ Cl ₂	2.6×10^{-3}		298	CP/Ac-15	$S = \text{MB}$.	717403
13.174	Thiourea						
	C ₅ H ₅ N	6×10^7 (k_r)			PL/Ld-2	$S = \text{PdMP}$; meas. $\phi_{\text{ox}} = \phi_{\Delta}$.	80A398
	C ₅ H ₅ N	9×10^7		273	CP/LI-12	$S = \text{MP}$; used $k_d = 5.9 \times 10^4 \text{ s}^{-1}$.	78F549
	CH ₃ COCH ₃	2.5×10^7 (k_r)			PL/Ld-2	$S = \text{PdMP}$; meas. $\phi_{\text{ox}} = \phi_{\Delta}$.	80A398
	CH ₃ COCH ₃	1.3×10^8		273	CP/LI-12	$S = \text{MP}$; used $k_d = 5.0 \times 10^4 \text{ s}^{-1}$.	78F549
	EtOH/H ₂ O (96:4)	$\sim 1 \times 10^8$		273	CP/LI-12	$S = \text{MP}$; k_d not given.	78F549

TABLE 13. Rate constants for the interaction of singlet oxygen with sulfur, selenium, and tellurium compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.174 Thiourea — Continued							
	H ₂ O pH = 7.1	4.4 × 10 ⁶		298	CP/Oc-19	S = Phenosafranine; Q = NaN ₃ ; used $k_Q = 2.0 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A360
	MeOH		0.024		CL/Oc-15	S = Ru(bpy) ₃ ²⁺ .	777221
	MeOH		0.040	293	CP/Oc-15	S = MB; $E_a = 6.3$ kJ mol ⁻¹ .	68F288
13.175 Thiourea, allyl-							
	<i>i</i> -C ₅ H ₁₁ OH		0.023	293	CP/Oc-14	S = Chl a.	56F007
	C ₅ H ₅ N	~2.7 × 10 ⁷		285	CP/P'a-20	S = A' = DMA; meas. $k_A/k_{A'} = 0.7$; k derived using $(k(TME)/k_{A'}) = 1.1$ and $k(TME) = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ .	737202
	<i>c</i> -C ₆ H ₁₁ OH		≤1 × 10 ⁻³	293	CP/Oc-14	S = Chl a.	56F007
	C ₆ H ₅ CH ₂ OH		9.2 × 10 ⁻³	293	CP/Oc-14	S = Chl a.	56F007
	H ₂ O pH = 7.1	4 × 10 ⁶		298	CP/Oc-19	S = Phenosafranine; Q = NaN ₃ ; used $k_Q = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78F020
	H ₂ O pH = 5.9		0.01	310	CP/Oc-15	S = PF.	617008
	H ₂ O (mic) pH = 7.0	4.5 × 10 ⁶			CP/Oc-19	S = Chl a; Q = N ₃ ⁻ ; used $k_Q = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; S solubilized in Triton X-100 micelles (1.0 % by volume).	78A278
	MeOH		0.10	293	CP/Oc-15	S = Chl a.	56F007
13.176 Thiourea, methyl-							
	H ₂ O pH = 7.1	2 × 10 ⁶		298	CP/Oc-19	S = Phenosafranine; Q = NaN ₃ ; used $k_Q = 2 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78A360
13.177 1,4-Thioxane							
	CHCl ₃	1.5 × 10 ⁶			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	79A085

TABLE 14. Rate constants for the interaction of singlet oxygen with inorganic compounds.

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
14.1 Ammonia							
	MeOH/C ₆ H ₆ (67:33)	3.3×10^{-4} (β_r)		298	CR/A'c-17	A' = Rub; used $\beta_r^{A'} = 1.4 \times 10^{-3}$ mol L ⁻¹ ; $^1\text{O}_2^*$ from (PhO) ₃ PO ₃ .	80M378
14.2 Azide ion							
	n-C ₇ H ₁₆ /H ₂ O (mic)	3.3×10^8			PL/Ld-2	S = RB; reverse micelles contg. 0.25 mol L ⁻¹ AOT and 2% H ₂ O.	84N114
	CH ₃ CN	4.8×10^9			PL/Ld-2	S = MB.	89A346
	DMF	1.2×10^8			CP/Ac-22	S = A' = Pt(phen)(DMT); used $k_d = 1.4 \times 10^5$ s ⁻¹ .	89F462
	DMSO	6×10^8			CP/P'a-25	S = GV; A' = TEMP-4-OH; used $k_d = 5.2 \times 10^4$ s ⁻¹ , $k_{A'} = 5.2 \times 10^6$ L mol ⁻¹ s ⁻¹ ; P = 4-Hydroxy-2,2,6,6-tetramethylpiperidine N-oxyl; counter ion = Na ⁺ , measured by esr.	89D112
	D ₂ O	3×10^8			PL/Ld-2	S = MB; k for I → 0; at I = 3 (NaCl) $k = 7.1 \times 10^8$ L mol ⁻¹ s ⁻¹ .	92F181
	D ₂ O pD = 11.2	9×10^7		310	CR/A'c-32	A' = DPBF; used $k_d = 1.5 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from DOPA/H ₂ O ₂ .	89M038
	D ₂ O pD = 11.2	5×10^7		310	CR/A'c-32	A' = DPBF; used $k_d = 1.5 \times 10^4$ s ⁻¹ ; $^1\text{O}_2^*$ from Dopamine/H ₂ O ₂ .	89M038
	D ₂ O pH = 7.0	6.3×10^8	4.0×10^{-5}		CP/LI-12	S = RB; used $k_d = 2.5 \times 10^4$ s ⁻¹ .	87E150
	D ₂ O pH = 7.0	6.9×10^8	3.1×10^{-5}		CP/LI-12	S = Eos; used $k_d = 2.2 \times 10^4$ s ⁻¹ .	87E150
	D ₂ O	4.4×10^8			PL/Ld-2	S = H ₂ TPPS ⁴⁻ ; counter ion Na ⁺ .	86F149
	D ₂ O pD = 7.4	5.1×10^8			PL/A'd-5	S = MB; A' = ADPA.	81N048
	D ₂ O (mic) pD = 7.4	3.9×10^8			PL/A'd-5	S = 2-ACN; A' = DPBF; 0.1 mol L ⁻¹ SDS.	81N048
	D ₂ O (mic) pD = 7.4	8.2×10^8			PL/A'd-5	S = 2-ACN; A' = DPBF; 0.1 mol L ⁻¹ CTAB.	81N048
	D ₂ O (ves)	1.1×10^8			CP/A'c-16	S = RB; A' = DMA; used $k_d = 1.8 \times 10^4$ s ⁻¹ ; 0.05% egg yolk lecithin.	86N104
	EtOH	2.2×10^8			PL/Ld-2	S = MB; k for I → 0; at I = 3 (tetramethylammonium chloride) $k = 5.5 \times 10^8$ L mol ⁻¹ s ⁻¹ .	92F181
	EtOH	2.0×10^8			CP/A'c-16	S = MC 540; A' = DPBF; used $k_d = 6.5 \times 10^4$ s ⁻¹ .	88F151
	EtOH/H ₂ O (96:4)	3.9×10^8			CP/P'a-18	S = HA; A' = TEMP; used $k_d = 6.5 \times 10^4$ s ⁻¹ ; P = 2,2,6,6-Tetramethylpiperidine-N-oxyl; monitored by esr.	90R162
	H ₂ O pH = 10.6	2.9×10^7		310	CR/LI-12	used $k_d = 3.2 \times 10^5$ s ⁻¹ ; soln. cont. 5×10^{-4} mol L ⁻¹ CoCl ₂ ; $^1\text{O}_2^*$ from autoxidation of oxytetracycline.	92M228
	H ₂ O pH = 7	4.5×10^8			CP/A'c-18	S = RB; A' = TrpH; used $k_d = 2.5 \times 10^5$ s ⁻¹ , $k_{A'} = 6 \times 10^8$ L mol ⁻¹ s ⁻¹ .	93R059
	H ₂ O pH = 5.4	3.8×10^8			CL/LI-12	S = RB; used $k_d = 2.4 \times 10^5$ s ⁻¹ .	91E570 89E798
	H ₂ O pH = 4.7-8.1	$<7 \times 10^5$ (k_r) 5.0×10^8 (k_q)		298	CP/A'c-16,17	S = RB, Humic acids; A' = FFA; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; meas. $k_r/k_r^{A'} \leq 0.0061$; used $k_r^{A'} = 1.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; counter ion Na ⁺ .	87A063
	H ₂ O pH = 7.0	5.8×10^8	4.1×10^{-4}		CP/LI-12	S = Eos or RB; used $k_d = 2.4 \times 10^5$ s ⁻¹ .	87E150
	H ₂ O pH = 10	1×10^9			CP/Oc-16	S = Eos; A' = 2,4-Cl ₂ C ₆ H ₃ O ⁻ ; used $k_d = 5 \times 10^5$ s ⁻¹ .	87F537

TABLE 14. Rate constants for the interaction of singlet oxygen with inorganic compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
14.2 Azide ion — Continued							
	H ₂ O pH = 7	6.4×10^8		298	CP/A'c-16	S = UP; A' = TrpH; used $k_d = 3 \times 10^5$ s ⁻¹ .	86F678
	H ₂ O	1.5×10^9			CP/Oc-22	S = MB; A' = PBN; used $k_d = 5 \times 10^5$ s ⁻¹ .	80A330
	H ₂ O pH = 8.4	1.7×10^9		298	CP/A'c-20	S = MB; A' = TrpH; used $k_d = 5.0 \times 10^5$ s ⁻¹ .	78F061
	H ₂ O	2.1×10^9			CP/P'a-23	S = A' = 1-AnS; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; A'' = KI, P'' = I ₃ ⁻ .	78F184
	H ₂ O (mic)	2.5×10^8		303	CP/A'c-20	S = HA; A' = DPBF; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; 2.9 $\times 10^{-4}$ mol L ⁻¹ SDS.	92N179
	H ₂ O (mic)	6.0×10^8		303	CP/A'c-20	S = HA; A' = DPBF; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; 2.2 $\times 10^{-4}$ mol L ⁻¹ Triton X-100.	92N179
	H ₂ O (mic)	7.6×10^8		303	CP/A'c-20	S = HA; A' = DPBF; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; 2.3 $\times 10^{-4}$ mol L ⁻¹ TDPB.	92N179
	H ₂ O (mic)	1.8×10^9		313	CP/A'c-23	S = Py; A' = DPBF; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; A' and S solubilized in DTAC micelles.	78A174
	H ₂ O (mic)	7.9×10^8		313	CP/A'c-23	S = Py; A' = DPBF; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; A' and S solubilized in SDS micelles.	78A174
	H ₂ O (mic) pH = 7.0	2.3×10^9		298	CP/A'c-20	S = MB; A' = DMA; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; A' solubilized in DTAC micelles.	78F061
	H ₂ O (mic) pH = 7.0	2.2×10^9		298	CP/A'c-20	S = MB; A' = 2,5-Diphenyloxazole; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; A' solubilized in DTAC micelles.	78F061
	H ₂ O (ves)	1.7×10^8			CP/A'c-16	S = A' = MC 540; used $k_d = 4.0 \times 10^4$ s ⁻¹ ; soln. contg. DLPC.	92A123
	H ₂ O (ves)	9.9×10^7			CP/A'c-16	S = RB; A' = DMA; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; 0.05% egg yolk lecithin.	86N104
	H ₂ O/ MeOH (80:20)	1.3×10^9			CP/P'a-23	S = A' = 1-AnS; used $k_d = 4.0 \times 10^5$ s ⁻¹ ; A'' = KI, P'' = I ₃ ⁻ .	78F183
	H ₂ O/ MeOH (60:40)	1.1×10^9			CP/P'a-23	S = A' = 1-AnS; used $k_d = 3.2 \times 10^5$ s ⁻¹ ; A'' = KI, P'' = I ₃ ⁻ .	78F183
	H ₂ O/ MeOH (40:60)	5.2×10^8			CP/P'a-23	S = A' = 1-AnS; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; A'' = KI, P'' = I ₃ ⁻ .	78F183
	H ₂ O/ MeOH (20:80)	3.0×10^8			CP/P'a-23	S = A' = 1-AnS; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; A'' = KI, P'' = I ₃ ⁻ .	78F183
	MeOH	1.6×10^8		297	CR/P'a-16	A' = TEMP; formn. of TEMPO monitored by esr; soln. cont. MeONa and CoCl ₂ , ¹ O ₂ * from autoxidation of adrenaline; k_d not given.	92D227
	MeOH	2.3×10^8			CP/A'c-16	S = MC 540; A' = DPBF; used $k_d = 1 \times 10^5$ s ⁻¹ .	88F151
	MeOH	2.3×10^8		303	PL/Tb-3	S = Ery; $E_a = 7.2$ kJ mol ⁻¹ ; studied at 303-323 K.	82A140
	MeOH	2.3×10^8			CP/A'c-23	S = Py; A' = DPBF; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	78A174
	MeOH	2.2×10^8			PL/A'd-5	S = MB; A' = DPBF.	72F515
	MeOH/ C ₆ H ₆ (67:33)		0.018 (β_f)	298	CR/A'c-17	A' = Rub; used $\beta_f^{A'} = 1.4 \times 10^{-3}$ mol L ⁻¹ ; ¹ O ₂ * from (PhO) ₃ PO ₃ .	80M378
	MeOH/ HOCH ₂ CH ₂ OH (90:10)	2.5×10^8		303	PL/Tb-3	S = Ery.	82A140
	MeOH/ HOCH ₂ CH ₂ OH (80:20)	3.0×10^8		303	PL/Tb-3	S = Ery.	82A140
	MeOH/ HOCH ₂ CH ₂ OH (70:30)	3.8×10^8		303	PL/Tb-3	S = Ery.	82A140

TABLE 14. Rate constants for the interaction of singlet oxygen with inorganic compounds. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
14.1 Azide ion — Continued							
	MeOH/ HOCH ₂ CH ₂ OH (60:40)	4.9×10^8		303	PL/Tb-3	S = Ery.	82A140
	MeOH/ HOCH ₂ CH ₂ OH (50:50)	6.9×10^8		303	PL/Tb-3	S = Ery.	82A140
14.2 Hydrogen azide							
	H ₂ O pH = 1.9	$<2 \times 10^6$		298	CP/A'c-16	S = MB; A' = FFA; used $k_d = 2.5 \times 10^5$ s ⁻¹ .	87A063
14.3 Bromide ion							
	C ₆ H ₅ Br/ CH ₃ COCH ₃ (67:33)	$\leq 10^6$			CP/A'c-23	S = A' = Rub; No measurable effect; LiBr or tetrabutylammonium bromide.	767126
	C ₆ H ₅ Br/ CH ₃ COCH ₃ (67:33)	$\sim 1 \times 10^6$			CP/A'c-23	S = A' = Rub; used $k_d = 3.2 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ ; dicyclohexano-18-crown-6-polyether, potassium bromide.	767126
	C ₆ H ₅ Br/ MeOH (67:33)	$\leq 10^6$			CP/A'c-23	S = A' = Rub; No measurable effect; LiBr or tetrabutylammonium bromide.	767126
14.4 Chloride ion							
	C ₆ H ₅ Br/ CH ₃ COCH ₃ (67:33)	$\leq 10^6$			CP/A'c-23	S = A' = Rub; No measurable effect; tetrabutylammonium chloride.	767126
	C ₆ H ₅ Br/ MeOH (67:33)	$\leq 10^6$			CP/A'c-23	S = A' = Rub; No measurable effect; tetrabutylammonium chloride.	767126
	CH ₃ CN	2.4×10^4			PL/Ld-2	S = MB; 0.1 mol L ⁻¹ tetraethylammonium chloride	89A346
	D ₂ O	10^3			PL/Ld-2	S = RF, H ₂ TPPS ⁴⁻ , or Chl in Triton X-100 micelles; counter ion Na ⁺ .	86F149
14.5 Deuterium peroxide							
	D ₂ O	8.8×10^2		293	PL/Ld-2	S = PTSA; soln. contg. 2-9 mol L ⁻¹ D ₂ O ₂ .	89A506
14.6 Hydrazine hydrate							
	C ₅ H ₅ N	8.7×10^7 (k_r)			PL/Ld-2	S = TPP, MPDEE or PdMP; meas. $\phi_{ox} = \phi_{\Delta^-}$	80A398
	CCl ₄	6.1×10^5			PL/Ld-2	S = MPDEE.	80A398
	CH ₃ COCH ₃	5.5×10^7 (k_r)			PL/Ld-2	S = TPP, MPDEE or PdMP; meas. $\phi_{ox} = \phi_{\Delta^-}$	80A398
	MeOH/ C ₆ H ₆ (67:33)		4.0×10^{-4} (β_r)	298	CR/A'c-17	A' = Rub; used $\beta_r^{A'} = 1.4 \times 10^{-3}$ mol L ⁻¹ ; ¹ O ₂ * from (PhO) ₃ PO ₃ .	80M378
14.7 Hydroxide-<i>d</i> ion							
	D ₂ O	5.5×10^3		293	PL/Ld-2	S = PTSA; soln. contg. 0.5-6 mol L ⁻¹ NaOD.	89A506
14.8 Iodide ion							
	C ₆ H ₅ Br/ CH ₃ COCH ₃ (67:33)	9.1×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 3.2 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ ; [CH ₃ (CH ₂) ₂ CH ₂] ₄ NI.	767126
	C ₆ H ₅ Br/ CH ₃ COCH ₃ (67:33)	8.1×10^7			CP/A'c-23	S = A' = Rub; used $k_d = 3.2 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ ; from LiI.	767126
	C ₆ H ₅ Br/ CH ₃ COCH ₃ (67:33)	2.8×10^8			CP/A'c-23	S = A' = Rub; used $k_d = 3.2 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ ; dicyclohexano-18-crown-6-polyether potassium iodide.	767126
	C ₆ H ₅ Br/ MeOH (67:33)	$\leq 10^6$			CP/A'c-23	S = A' = Rub; No measurable effect; LiI or KI or tetrabutylammonium iodide.	767126

TABLE 14. Rate constants for the interaction of singlet oxygen with inorganic compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
14.9 Iodide ion — Continued							
	D ₂ O pH = -7	8.7×10^5			PL/Ld-2	S = 2-AnS.	88A398
	H ₂ O pH = -7	8.7×10^5			PL/Ld-2	S = 2-AnS.	88A398
	H ₂ O	8.7×10^6	0.058		CP/Pa-23	S = A' = 1-AnS; Q = N ₃ ⁻ ; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; P = Triiodine ion.	78F183
	H ₂ O	7.2×10^6		301	CP/Pa-23	S = 1-AnS; Q = NaN ₃ ; used $k_d = 5.0 \times 10^5$ s ⁻¹ , $k_Q = 2.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; P = Triiodine ion; from KI.	777074
	H ₂ O	7.2×10^6		301	CP/Pa-23	S = 1,5-Anthracenedisulfonate ion; Q = NaN ₃ ; used $k_d = 5.0 \times 10^5$ s ⁻¹ , $k_Q = 2.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; P = Triiodine ion; from KI.	777074
	H ₂ O/ MeOH (60:40)	6.4×10^6		301	CP/Pa-23	S = 2-AnS; Q = NaN ₃ ; used $k_d = 5.0 \times 10^5$ s ⁻¹ , $k_Q = 2.2 \times 10^8$ L mol ⁻¹ s ⁻¹ ; P = Triiodine ion; from KI.	777074
	H ₂ O/ MeOH (20:80)	3.0×10^6	0.11		CP/Pa-23	S = A' = 1-AnS; Q = N ₃ ⁻ ; used $k_d = 3.2 \times 10^5$ s ⁻¹ ; P = Triiodine ion; from KI.	78F183
	H ₂ O/ MeOH (40:60)	3.4×10^5	0.56		CP/Pa-23	S = A' = 1-AnS; Q = N ₃ ⁻ ; used $k_d = 1.9 \times 10^5$ s ⁻¹ ; P = Triiodine ion; from KI.	78F183
	H ₂ O/ MeOH (80:20)	1.5×10^6	0.16		CP/Pa-23	S = A' = 1-AnS; Q = N ₃ ⁻ ; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; P = Triiodine ion; from KI.	78F183
	H ₂ O/ MeOH (80:20)	5.6×10^6	0.072		CP/Pa-23	S = A' = 1-AnS; Q = N ₃ ⁻ ; used $k_d = 4.0 \times 10^5$ s ⁻¹ ; P = Triiodine ion; from KI.	78F183
14.10 Iodine							
	C ₆ H ₆	1.4×10^9			PL/A'd-8	S = An; A' = DPBF; Quenching is non-linear.	84E291
14.11 Nitrite ion							
	H ₂ O pH = 8.3	3.1×10^6			CP/Oc-16	S = Eos; A' = N ₃ ⁻ ; used $k_d = 2.5 \times 10^5$ s ⁻¹ , $k = 7.4 \times 10^4$ L mol ⁻¹ sec ⁻¹ was also evaluated; $k = 4.4 \times 10^4$ L mol ⁻¹ sec ⁻¹ was also quoted [92D008].	88F485
14.12 Oxygen							
	Perfluorodecalin	4.1×10^3		295	PL/Ld-2	S = PHO.	91E427
	n-C ₆ F ₁₄	2.6×10^3		295	PL/Ld-2	S = PHO.	91E427
	C ₆ H ₆	$\sim 6 \times 10^4$			CP/A'c-?	S = A' = DMA; Reported as unpublished data.	73F659
	CCl ₄	3.9×10^3		295	PL/Ld-2	S = PHO.	91E427
	CS ₂	1.3×10^4		298	FP/Ld-2	S = Per.	82A322
	ClCF ₂ CCl ₂ F	3.2×10^3		295	PL/Ld-2	S = PHO.	91E427
	ClCF ₂ CCl ₂ F	9.2×10^2		298	FP/Ld-2	S = Per.	82A322
	ClCF ₂ CCl ₂ F	2.5×10^3		293	PL/A'd-10	A' = DPBF; high pressure O ₂ .	79A113
	ClCF ₂ CCl ₂ F	2.7×10^3			PL/A'd-10	A' = DPBF; high pressure O ₂ .	747102
	H ₂ O pH = 6.2	0.060			CP/A'c-19	S = Ac; A' = Leuco fluorescein.	68F287
	H ₂ O pH = 6.2	1.9×10^{-3}			CP/A'c-19	S = Th ⁺ ; A' = Leuco fluorescein.	68F287
	H ₂ O pH = 6.2	1.2×10^{-3}			CP/A'c-19	S = MB; A' = Leuco fluorescein.	68F287
	H ₂ O pH = 6.2	2.2×10^{-3}			CP/A'c-19	S = Eos; A' = Leuco fluorescein.	68F287
	O ₂	5.9×10^2		77	PL/Ld-2	Liquid oxygen, direct laser excitation.	82E575
	O ₂	6.0×10^2		77	PL/Ld-2	Liquid oxygen, direct laser excitation, 1064 nm.	80E616

TABLE 14. Rate constants for the interaction of singlet oxygen with inorganic compounds. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_q/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
14.12 Oxygen — Continued							
	O ₂	6.0×10^2		77	PL/Ld-2	Liquid oxygen, direct laser excitation, 1060 nm; $k = 1.3 \times 10^3 (T/300)^{0.5}$ [719112].	746110
14.13 Oxygen-18							
	Perfluorodecalin	60		295	PL/Ld-2	S = PHO.	91E427
	n-C ₆ F ₁₄	1×10^2		295	PL/Ld-2	S = PHO.	91E427
	ClCF ₂ CCl ₂ F	1×10^2		295	PL/Ld-2	S = PHO.	91E427
	O ₂	3		77	PL/Ld-2	Liquid oxygen, direct laser excitation.	82E575
14.14 Superoxide radical anion							
	C ₆ H ₅ Br/CH ₃ CN (67:33)	3.6×10^7		CP/A'c-23	S = A' = Rub; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ ; Showed $k_q \gg k_r$; tetrabutylammonium superoxide.	757578	
	CH ₃ CN	2×10^8		PL/Ld-2	S = 2-ACN; KO ₂ /18-crown-6.	91A394	
	CH ₃ CN	2×10^8		PL/Ld-2	S = Ru(bpy) ₃ ²⁺ ; KO ₂ /18-crown-6.	91A394	
	CH ₃ CN	7×10^9	5×10^{-6}	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 3.3 \times 10^4$ s ⁻¹ ; tetramethylammonium superoxide.	766072	
	DMSO	1.6×10^9	3.3×10^{-5}	CP/A'c-16	S = RB; A' = DPBF; used $k_d = 5.2 \times 10^4$ s ⁻¹ ; tetramethylammonium superoxide.	766072	

TABLE 15. Rate constants for the interaction of singlet oxygen with oximes, hydrazones, nitrones, nitroso compounds and *N*-oxides.

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
15.1	Acetaldehyde, (2,6-diphenylpyran-4-ylidene)-, dimethyl hydrazone						
	MeOH/ C ₆ H ₆ (80:20)	$3.0 \times 10^9 (k_r)$			CP/A'c-17	S = RB; A' = 2M2P; used $k_r^{A'} = 8.1 \times 10^5$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 3.7 \times 10^3$.	86A438
	MeOH/ C ₆ H ₆ (80:20)	$2.2 \times 10^{10} (k_r)$			CP/A'c-17	S = RB; A' = DPBF; used $k_r^{A'} = 6.3 \times 10^8$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 35$.	86A438
15.2	Acetone oxime						
	MeOH	$\leq 1.0 \times 10^4 (k_r)$			CP/A'c-17	S = RB; A' = 2M2B; used $k_r^{A'} = 1.3 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 7.7 \times 10^{-3}$.	767197
15.3	Acetone, (1-methylethylidene)hydrazone (Acetone azine)						
	CCl ₃ F		253	CP/Pa-17	S = TPP; A' = 3,4,4,5-Tetramethylpyrazole; meas. $k_r/k_r^{A'} = 3.1$.	79F278	
	CCl ₄		253	CP/Pa-17	S = TPP; A' = 3,4,4,5-Tetramethylpyrazole; meas. $k_r/k_r^{A'} = 3.2$; with 10^{-4} mol L ⁻¹ TPP.	79F278	
	CCl ₄		253	CP/Pa-17	S = TPP; A' = 3,4,4,5-Tetramethylpyrazole; meas. $k_r/k_r^{A'} = 3.8$; with 10^{-6} mol L ⁻¹ TPP.	79F278	
	CDCl ₃		253	CP/Pa-17	S = TPP; A' = 3,4,4,5-Tetramethylpyrazole; meas. $k_r/k_r^{A'} = 3.2$.	79F278	
	CH ₂ Cl ₂		253	CP/Pa-17	S = TPP; A' = 3,4,4,5-Tetramethylpyrazole; meas. $k_r/k_r^{A'} = 3.3$.	79F278	
	CHCl ₃		253	CP/Pa-17	S = TPP; A' = 3,4,4,5-Tetramethylpyrazole; meas. $k_r/k_r^{A'} = 3.0$.	79F278	
	CHCl ₃		253	CP/Pa-17	S = MB; A' = 3,4,4,5-Tetramethylpyrazole; meas. $k_r/k_r^{A'} = 3.8$.	79F278	
15.4	Benzaldehyde, 4-diethylamino-, diphenylhydrazone						
	MeOH/ C ₆ H ₆ (80:20)	1.8×10^9			CP/P'a-23	S = RB; A' = 2M2P; used $k_d = 1.8 \times 10^5$ s ⁻¹ , $k_{A'} = 8.1 \times 10^5$ L mol ⁻¹ s ⁻¹ .	83A063
	MeOH/ C ₆ H ₆ (68:32)	8.8×10^8			CP/P'a-23	S = RB; A' = DPBF; used $k_d = 1.8 \times 10^5$ s ⁻¹ , $k_{A'} = 6.3 \times 10^8$ L mol ⁻¹ s ⁻¹ .	83A063
15.5	Benzenamine, <i>N</i> -(diphenylmethylene)-, <i>N</i> -oxide (Triphenyl nitrone)						
	c-C ₆ H ₁₂	5×10^6	296	PL/A'd-8	S = ZnTPP; A' = DPBF.	91A297	
	C ₆ H ₆	3.9×10^7	296	PL/A'd-8	S = ZnTPP; A' = DPBF.	91A297	
	CH ₃ CN	3.3×10^7	296	PL/A'd-8	S = MB; A' = DPBF.	91A297	
	MeOH	1.0×10^7	296	PL/A'd-8	S = MB; A' = DPBF.	91A297	
15.6	Benzenamine, <i>N</i> -fluoren-9-ylidene, <i>N</i> -oxide (Phenyl fluorenyl nitrone)						
	C ₆ H ₆	6.2×10^7	296	PL/A'd-8	S = ZnTPP; A' = DPBF.	91A297	
15.7	Benzenamine, <i>N</i> -[(2-hydroxyphenyl)methylene]-, <i>N</i> -oxide (2-Hydroxyphenyl phenyl nitrone)						
	C ₆ H ₆	4.5×10^7	296	PL/A'd-8	S = ZnTPP; A' = DPBF.	91A297	
15.8	Benzenamine, 4-methyl- <i>N</i> -(phenylmethylene)-, <i>N</i> -oxide (4-Methylphenyl phenyl nitrone)						
	C ₆ H ₆	4.3×10^7	296	PL/A'd-8	S = ZnTPP; A' = DPBF.	91A297	
15.9	Benzenamine, <i>N</i> -[(4-methylphenyl)methylene]-, <i>N</i> -oxide (Phenyl 4-methylphenyl nitrone)						
	c-C ₆ H ₁₂	2.1×10^7	296	PL/A'd-8	S = ZnTPP; A' = DPBF.	91A297	
	C ₆ H ₆	5.5×10^7	296	PL/A'd-8	S = ZnTPP; A' = DPBF.	91A297	
	CH ₃ CN	4.6×10^7	296	PL/A'd-8	S = MB; A' = DPBF.	91A297	
	MeOH	7.7×10^7	296	PL/A'd-8	S = MB; A' = DPBF.	91A297	
15.10	Benzenamine, <i>N</i> -(phenylmethylene)-, <i>N</i> -oxide (Diphenyl nitrone)						
	c-C ₆ H ₁₂	8×10^6	296	PL/A'd-8	S = ZnTPP; A' = DPBF.	91A297	
	C ₆ H ₆	3.7×10^7	296	PL/A'd-8	S = ZnTPP; A' = DPBF.	91A297	

TABLE 15. Rate constants for the interaction of singlet oxygen with oximes, hydrazones, nitrones, nitroso compounds and *N*-oxides. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
15.10 Benzenamine, <i>N</i>-(phenylmethylene)-, <i>N</i>-oxide (Diphenyl nitrone) — Continued							
	CH ₃ CN	3.3×10^7		296	PL/A'd-8	S = MB; A' = DPBF.	91A297
	MeOH	5.1×10^7		296	PL/A'd-8	S = MB; A' = DPBF.	91A297
15.11 Benzenemethanamine, <i>N</i>-hydroxy-<i>N</i>-(phenylmethyl)- (Dibenzylhydroxylamine)							
	CCl ₄	1.2×10^4	2.2×10^{-3}		CP/Pa-14	S = DCA; used $k_d = 26$ s ⁻¹ ; P = α -Phenyl- <i>N</i> -benzylnitrene.	92F491
15.12 Benzenemethanamine, <i>N</i>-(phenylmethylene)-, <i>N</i>-oxide (Phenyl benzyl nitrone)							
	C ₆ H ₆	1.4×10^7		296	PL/A'd-8	S = ZnTPP; A' = DPBF.	91A297
15.13 Benzophenone oximate anion							
	MeOH	3.4×10^5 (k_r)			CP/A'c,Pa-17	S = RB; A' = 2M2B; used $k_r^{A'} = 1.3 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.26$; P = Benzophenone.	767197
15.14 Benzophenone oxime							
	MeOH	7.7×10^4 (k_r)			CP/A'c,Pa-17	S = RB; A' = 2M2B; used $k_r^{A'} = 1.3 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.059$; P = Benzophenone.	767197
15.15 Benzophenone oxime <i>O</i>-methyl ether							
	MeOH	2.0×10^5 (k_r)			CP/A'c,Pa-17	S = RB; A' = 2M2B; used $k_r^{A'} = 1.3 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.15$; P = Benzophenone.	767197
15.16 Cyclohexane, 1,4-dichloro-1,4-dinitroso-, (<i>E</i>)-							
	CH ₂ Cl ₂ / MeOH (69:31)	5.3×10^9			CR/A'c-32	A' = Rub; used $k_d = 1.9 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; $^1\text{O}_2^*$ from (PhO) ₃ PO ₃ .	76F900
15.17 Cyclohexane, 1,4-dichloro-1,4-dinitroso-, (<i>Z</i>)-							
	CH ₂ Cl ₂ / MeOH (69:31)	9.2×10^9			CP/A'c-32	S = Chl a; A' = DPBF; used $k_d = 2.1 \times 10^4$ s ⁻¹ , $k_{A'} = 8 \times 10^8$ L mol ⁻¹ s ⁻¹ ; Assumed that quenching is due to equilibrium mixture of this species with 1,4-dichloro-2,3-diazabicyclo[2.2.2]oct-2-ene 2,3-dioxide.	76F900
	CH ₂ Cl ₂ / MeOH (69:31)	1.2×10^{10}			CR/A'c-32	A' = Rub; used $k_d = 2.1 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; $^1\text{O}_2^*$ from (PhO) ₃ PO ₃ ; Assumed that quenching is due to this species in equilibrium with 1,4-dichloro-2,3-diazabicyclo[2.2.2]oct-2-ene 2,3-dioxide.	76F900
15.18 2,3-Diazabicyclo[2.2.2]oct-2-ene, 1,4-dichloro-, 2,3-dioxide							
	CH ₂ Cl ₂ / MeOH (69:31)	8.0×10^7			CR/A'c-32	A' = Rub; used $k_d = 2.1 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; $^1\text{O}_2^*$ from (PhO) ₃ PO ₃ ; Assumed that quenching is due to this species only.	76F900
15.19 Dimethylglyoxime							
	CH ₂ Cl ₂ / MeOH (90:10)	1×10^6			PL/A'd-5	S = MB; A' = DPBF.	87F655
15.20 3,7,4,6-Ethanediylidenepentalenzo[1,6-<i>cd</i>]pyridazine, 3,7a-dichloro-3a,4,5,5a,6,7,7a,7b-octahydro-, 1,2-dioxide							
	CH ₂ Cl ₂ / MeOH (69:31)	$<2 \times 10^6$			CR/A'c-32	A' = Rub; used $k_d = 2.1 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; $^1\text{O}_2^*$ from (PhO) ₃ PO ₃ .	76F900
15.21 Ethylamine, <i>N</i>-ethoxy-							
	n-C ₆ H ₁₄	3×10^4			CP/A'c-16	S = A' = Rub; k_d not given.	87F639
	CH ₃ CN	8×10^5			CP/A'c-16	S = RB; A' = DMA; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	87F639
15.22 Ethylamine, <i>N</i>-ethyl-<i>N</i>-ethoxy-							
	n-C ₆ H ₁₄	6×10^5			CP/A'c-16	S = A' = Rub; k_d not given.	87F639

TABLE 15. Rate constants for the interaction of singlet oxygen with oximes, hydrazones, nitrones, nitroso compounds and *N*-oxides. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.	
15.22	Ethylamine, <i>N</i> -ethyl- <i>N</i> -ethoxy-	— Continued						
	CH ₃ CN	2.1×10^6			CP/A'c-16	S = RB; A' = DMA; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	87F639	
15.23	Ethylamine, <i>N</i> -ethyl- <i>N</i> -hydroxy-				CP/A'c-16	S = A' = Rub; k_d not given.	87F639	
	n-C ₆ H ₁₄	2.5×10^7			CP/A'c-16	S = A' = Rub; used $k_d = 4.2 \times 10^4$ s ⁻¹ .	87F639	
	C ₆ H ₆	2.7×10^7			CP/A'c-16	S = RB; A' = DMA; used $k_d = 3.3 \times 10^4$ s ⁻¹ .	87F639	
	CH ₃ CN	1.8×10^7			CP/A'c-16	S = RB; A' = DMA; used $k_d = 3.8 \times 10^4$ s ⁻¹ .	87F639	
	CH ₃ COCH ₃	2.5×10^7			CP/A'c-16	S = RB; A' = DMA; k_d not given.	87F639	
	CHCl ₃	1.2×10^7			CP/A'c-16	S = RB; A' = DMA; k_d not given.	87F639	
	D ₂ O	$<10^5$			CP/LI-12	S = RB; used $k_d = 1.8 \times 10^4$ s ⁻¹ .	93D154	
	HCONH ₂	1.0×10^7			CP/A'c-16	S = RB; A' = DMA; k_d not given.	87F639	
	MeOH	3×10^6			CP/A'c-16	S = RB; A' = DMA; used $k_d = 2.0 \times 10^4$ s ⁻¹ .	87F639	
15.24	Ethylamine, <i>N</i> -ethyl- <i>N</i> -hydroxy-, conjugate acid				CP/LI-12	S = RB; used $k_d = 1.8 \times 10^4$ s ⁻¹ .	93D154	
	D ₂ O	$<5 \times 10^3$				pH = 8.5		
15.25	Isoindole, 1,1,3-triphenyl-, 2-oxide			296	PL/A'd-8	S = ZnTPP; A' = DPBF.	91A297	
	C ₆ H ₆	8×10^6						
15.26	Nitroxide, bis[4-(1,1-dimethylethyl)phenyl]-				293	PL/Ld-2	S = An and PP; $E_a = 1.0$ kJ mol ⁻¹ ; log(A) = 7.4.	86E108
	C ₆ H ₅ CH ₃	1.6×10^7			293	PL/Ld-2	S = An and PP.	86E108
	CH ₃ CN	1.7×10^7			293	PL/Ld-2	S = HP.	92F402
	CH ₃ CN/ MeOH (96:4)	7.5×10^6						
15.27	Nitroxide, bis[4-(1-methyl-1-phenylethyl)phenyl]-				293	PL/Ld-2	S = An and PP; $E_a = 5.0$ kJ mol ⁻¹ ; log(A) = 7.3.	86E108
	C ₆ H ₅ CH ₃	2.5×10^6						
15.28	Piperidin-1-ol, 2,2,6,6-tetramethyl- (TEMPOH)				CP/A'c-18	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88F059	
	CH ₂ Cl ₂	1.5×10^8						
15.29	1-Piperidinyloxy, 4-amino-2,2,6,6-tetramethyl-				PL/A'd-8	S = MB; A' = DPBF.	84A167	
	C ₆ H ₆ / MeOH (80:20)	8.5×10^5						
15.30	1-Piperidinyloxy, 4-azido-2,2,6,6-tetramethyl-				PL/Ld-2	S = HP.	92F402	
	CH ₃ CN/ MeOH (96:4)	8.3×10^6						
15.31	1-Piperidinyloxy, 4-hydroxy-4-(2-naphthyl)-2,2,6,6-tetramethyl-,				293	PL/Ld-2	S = An and PP; $E_a = 7.5$ kJ mol ⁻¹ ; log(A) = 6.3.	86E108
	CHCl ₃	1.0×10^5						
15.32	1-Piperidinyloxy, 4-hydroxy-4-(2-phenylethynyl)-2,2,6,6-tetramethyl-				293	PL/Ld-2	S = An and PP; $E_a = 1.3$ kJ mol ⁻¹ ; log(A) = 5.1.	86E108
	CHCl ₃	7.5×10^4			293	PL/Ld-2	S = An and PP.	86E108
	EtOH	1.7×10^5						
15.33	1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl- (TEMPOL)				293	PL/Ld-2	S = An and PP; $E_a = 5.4$ kJ mol ⁻¹ ; log(A) = 6.2.	86E108
	C ₆ H ₅ CH ₃	1.6×10^5			293	PL/Ld-2	S = TPP.	93E090
	CCl ₄	9.8×10^4			293	PL/Ld-2	S = TPP.	93E090
	CH ₂ Cl ₂	9.5×10^4			293	PL/Ld-2	S = An and PP.	86E108
	CH ₃ CN	1.4×10^5			293	PL/Ld-2	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
	CHCl ₃	1.2×10^6			293	PL/Ld-2	S = An and PP.	86E108
	CHCl ₃	1.1×10^5			293	PL/Ld-2	S = An and PP.	86E108

TABLE 15. Rate constants for the interaction of singlet oxygen with oximes, hydrazones, nitrones, nitroso compounds and *N*-oxides. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
15.33	1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl- (TEMPOL) — Continued						
	EtOH	1.2×10^5		293	PL/Ld-2	S = An and PP.	86E108
15.34	1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, benzoate						
	CHCl ₃	1.4×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
15.35	1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, butyrate						
	CHCl ₃	1.1×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
15.36	1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, crotonate						
	CHCl ₃	1.6×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
15.37	1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, diester with 1,10-decanedioic acid						
	CHCl ₃	3.1×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
15.38	1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, diester with 1,6-hexanedioic acid						
	CHCl ₃	3.1×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
15.39	1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, diester with terephthalic acid						
	CHCl ₃	3.4×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
15.40	1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, methacrylate						
	CHCl ₃	1.7×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
15.41	1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, pentanoate						
	CHCl ₃	1.7×10^6			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88P313
15.42	1-Piperidinyloxy, 4-iodo-2,2,6,6-tetramethyl-						
	CH ₃ CN/ MeOH (96:4)	3.7×10^5			PL/Ld-2	S = HP.	92F402
15.43	1-Piperidinyloxy, 2,2,6,6-tetramethyl- (TEMPO)						
	C ₆ H ₆ / MeOH (80:20)	$<1.0 \times 10^5$			PL/A'd-8	S = MB; A' = DPBF.	84A167
	C ₆ H ₅ CH ₃	5.5×10^5		293	PL/Ld-2	S = An and PP; $E_a = 8.4$ kJ mol ⁻¹ ; $\log(A) = 7.2$.	86E108
	CCl ₄	1.4×10^5			PL/Ld-2	S = TPP.	93E090
	CH ₂ Cl ₂	1.6×10^5			PL/Ld-2	S = TPP.	93E090
	CH ₂ Cl ₂	2.8×10^6			CP/A'c-18	S = A' = Rub; used $k_d = 7.3 \times 10^3$ s ⁻¹ , $k_{A'} = 7 \times 10^7$ L mol ⁻¹ s ⁻¹ .	88F059
	CH ₃ CN	4.3×10^5		293	PL/Ld-2	S = An and PP.	86E108
	EtOII	2.5×10^5		293	PL/Ld-2	S = An and PP.	86E108
15.44	4-Piperidone, 2,2,6,6-tetramethyl-1-oxyl- (TAN)						
	C ₆ H ₅ CH ₃	1.2×10^6		293	PL/Ld-2	S = An and PP; $E_a = 6.7$ kJ mol ⁻¹ ; $\log(A) = 7.3$.	86E108
	CCl ₄	6.2×10^5			PL/Ld-2	S = TPP.	93E090
	CH ₂ Cl ₂	6.3×10^5			PL/Ld-2	S = TPP.	93E090
	CH ₃ CN	1.3×10^6		293	PL/Ld-2	S = An and PP.	86E108
	EtOH	1.3×10^6		293	PL/Ld-2	S = An and PP.	86E108
	MeOH/ C ₆ H ₆ (67:33)	3.4×10^{-5}		298	CR/A'c-17	A' = Rub; used $\beta_{A'} = 1.4 \times 10^{-3}$ mol L ⁻¹ , ¹ O ₂ * from (PhO) ₃ PO ₃ .	80M378

TABLE 15. Rate constants for the interaction of singlet oxygen with oximes, hydrazones, nitrones, nitroso compounds and *N*-oxides. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
15.45	2-Propanamine, 2-methyl- <i>N</i> -(1-methyl-4-pyridylmethylene)-, <i>N</i> -oxyl						
	H ₂ O	8.0×10^7			CP/Ac-23	S = MB; Q = N ₃ ⁻ ; used $k_d = 5 \times 10^5$ s ⁻¹ , $k_Q = 1.5 \times 10^9$ L mol ⁻¹ s ⁻¹ .	80A339
15.46	2-Propanamine, 2-methyl- <i>N</i> -phenylmethylene-, <i>N</i> -oxyl (Phenyl <i>N</i> - <i>tert</i> -butyl nitrone, PBN)						
	C ₆ H ₅ CH ₃	9.0×10^6		293	PL/A'd-8	S = PP; A' = DPBF; $E_a = 0.4 \pm 0.4$ kJ mol ⁻¹ ; studied at 223-293 K.	88F462
	C ₆ H ₅ CH ₃	9.0×10^6		293	PL/Ld-2	S = PP; $E_a = 0.4 \pm 0.4$ kJ mol ⁻¹ ; studied at 223-293 K.	86A307
	C ₆ H ₅ CH ₃	$\leq 1 \times 10^5$ (k_r)		293	CP/A'c-17	S = PP; A' = Tetr; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86A307 88F462
	C ₆ H ₆	8.3×10^6		293	PL/A'd-8	S = PP; A' = DPBF.	84A247 85F538
	C ₆ H ₆	1.4×10^5 (k_r)		293	CL/Ac,A'c-17	S = PP; A' = Tetr; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.011$.	84A247 85F538
	CH ₃ CN	1.8×10^7		293	PL/A'd-8	S = PP; A' = DPBF.	88F462
	CH ₃ CN	1.8×10^7		293	PL/Ld-2	S = PP.	86A307
	CH ₃ CN	$\leq 1 \times 10^5$ (k_r)		293	CP/A'c-17	S = PP; A' = Tetr; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86A307
	CHCl ₃	4.4×10^6		293	PL/A'd-8	S = PP; A' = DPBF.	88F462
	CHCl ₃	4.4×10^6		293	PL/Ld-2	S = PP.	86A307
	CHCl ₃	$\leq 1 \times 10^5$ (k_r)		293	CP/A'c-17	S = PP; A' = Tetr; used $k_r^{A'} = 1.2 \times 10^7$ L mol ⁻¹ s ⁻¹ .	86A307
	H ₂ O	1.4×10^8			CP/Ac-23	S = MB; Q = N ₃ ⁻ ; used $k_d = 5 \times 10^5$ s ⁻¹ , $k_Q = 1.5 \times 10^9$ L mol ⁻¹ s ⁻¹ .	80A339
15.47	2-Propanamine, 2-methyl- <i>N</i> -(4-pyridylmethylene)-, <i>N,N</i> -dioxy- (4-POBN)						
	H ₂ O	1.2×10^8			CP/Ac-23	S = MB; Q = N ₃ ⁻ ; used $k_d = 5 \times 10^5$ s ⁻¹ , $k_Q = 1.5 \times 10^9$ L mol ⁻¹ s ⁻¹ .	80A339
15.48	2-Propanamine, 2-methyl- <i>N</i> -(4-pyridylmethylene)-, <i>N</i> -oxyl (4-PyBN)						
	H ₂ O	1.4×10^8			CP/Ac-23	S = MB; Q = N ₃ ⁻ ; used $k_d = 5 \times 10^5$ s ⁻¹ , $k_Q = 1.5 \times 10^9$ L mol ⁻¹ s ⁻¹ .	80A339
15.49	2-Propanamine, 2-methyl- <i>N</i> -(2-sulfonatophenyl)methylene-, <i>N</i> -oxyl (2-SSPB)						
	H ₂ O	9.4×10^7			CP/Ac-23	S = MB; Q = N ₃ ⁻ ; used $k_d = 5 \times 10^5$ s ⁻¹ , $k_Q = 1.5 \times 10^9$ L mol ⁻¹ s ⁻¹ .	80A339
15.50	Propane, 2-methyl-2-nitroso-						
	CH ₂ Cl ₂ / MeOH (94:6)	9.3×10^9			CR/A'c-32	A' = Rub; used $k_d = 1.1 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ¹ O ₂ * from (PhO) ₃ PO ₃ .	76F900
15.51	Pyrrole, 3,4-dihydro-2,2,3-trimethyl-, 1-oxide						
	CHCl ₃	5×10^7			CP/A'c-16	S = MB; A' = DPBF; k_d not given; $k = k_q$ (No reaction observed).	75F653
15.52	Pyrrole, 3,4-dihydro-3,5,5-trimethyl-, 1-oxide						
	CHCl ₃	2.1×10^7			CP/A'c-16	S = MB; A' = DPBF; k_d not given.	75F653
15.53	1-Pyrrolidinyloxy, 3-cyano-2,2,5,5-tetramethyl-4-nitro-						
	CH ₃ CN/ MeOH (96:4)	1.7×10^5			PL/Ld-2	S = HP.	92F402
15.54	1-Pyrrolidinyloxy, 2,5-di-(4-hydroxy-3-nitrophenyl)-2,5-dimethyl-						
	CH ₃ CN/ MeOH (96:4)	3.1×10^5			PL/Ld-2	S = HP.	92F402

TABLE 15. Rate constants for the interaction of singlet oxygen with oximes, hydrazones, nitrones, nitroso compounds and *N*-oxides. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
13.55	1-Pyrrolidinyloxy, 2-(4-fluoro-3-nitrophenyl)-2,5,5-trimethyl-						
	CH ₃ CN/ MeOH (96:4)	1.7×10^5			PL/Ld-2	S = HP.	92F402
13.56	1-Pyrrolinyl-1-oxy, 5,5-dimethyl-						
	H ₂ O	1.8×10^7			CP/Ac-23	S = MB; Q = N ₃ ⁻ ; used $k_d = 5 \times 10^5$ s ⁻¹ , $k_Q = 1.5 \times 10^9$ L mol ⁻¹ s ⁻¹ .	80A339

TABLE 16. Rate constants for the interaction of singlet oxygen with some compounds of biological interest.

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
16.1	Adenosine						
	D ₂ O pH = ~7	<2 × 10 ⁵		293	PL/Ld-2	S = H ₂ TPPS ⁴⁻ ; No reaction at [A] = 2 × 10 ⁻² mol L ⁻¹ .	86A198
	H ₂ O pH = 7.1	≤1 × 10 ⁶		298	CP/Oc-19	S = Phenosafranine; No measurable effect.	78A360
16.2	Adenosine 5'-monophosphate						
	D ₂ O pH = ~7	4.3 × 10 ⁴		293	PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	86A198
16.3	Adenosine triphosphate						
	D ₂ O	4 × 10 ⁴			PL/Ld-2	S = RF, H ₂ TPPS ⁴⁻ , or Chl in Triton X-100 micelles.	86F149
16.4	4-Androsten-3-one, 17-hydroxy-4-methyl-		>2 × 10 ²		CP/Ac-18	S = Poly-RB or RB or MB; A' = Linalool; used $\beta_{A'} = 0.18$ mol L ⁻¹ ; solvent is CH ₂ Cl ₂ or MeOH or C ₆ H ₆ .	80F111
16.5	Androst-5-en-17-one, 3-(acetoxy)- (3β)	C ₆ H ₆ /C ₅ H ₅ N (91:9)	6.3	293	CP/Oc-14	S = HP.	57F008
16.6	Ascorbate ion						
	C ₅ H ₅ N	1.1 × 10 ⁷		303	CR/A'c-23	A' = DMA; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $k_{A'} = 2.1 \times 10^7$ L mol ⁻¹ s ⁻¹ , ¹ O ₂ * from MNPO ₂ .	83F407
	C ₅ H ₅ N	8.4 × 10 ⁶		303	CP/A'c-23	S = TPP; A' = DMA; used $k_d = 3.1 \times 10^4$ s ⁻¹ , $k_{A'} = 2.1 \times 10^7$ L mol ⁻¹ s ⁻¹ .	83F407
	C ₅ H ₅ N	7 × 10 ⁷			PL/Ld-2	S = TPP, MPDEE or PdMP.	80A398
	CD ₃ OD	1.5 × 10 ⁸		293	PL/Ld-2	S = HP.	90F411
	CH ₃ COCH ₃	9 × 10 ⁵			PL/Ld-2	S = TPP, MPDEE or PdMP.	80A398
	D ₂ O pH = ~7	1.6 × 10 ⁸		293	PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	86A198
	D ₂ O	4 × 10 ⁷			PL/Ld-2	S = RF, water-sol. TPP derivs., or Chl a in micelles contg. Triton X-100; counter ion Na ⁺ .	86F149
	D ₂ O pD = 7.2	2.5 × 10 ⁶			CP/LI-12	S = Chrysenesulfonate ion; used $k_d = 1.8 \times 10^4$ s ⁻¹ .	83A205
	H ₂ O pH = 6.8	8.3 × 10 ⁶			CP/LI-12	S = Chrysenesulfonate ion; used $k_d = 2.6 \times 10^5$ s ⁻¹ .	83A205
	MeOH	0.012		293	CP/Oc-15	S = RB; $E_a = 6.3$ kJ mol ⁻¹ .	68F288
16.7	Azirino[2'3':3,4]pyrrolo[1,2-a]indole-4,7-dione, 6-amino-8-[(aminocarbonyl)oxy]methyl]-1,1a,2,8,8a,8b-hexahydro-8a-methoxy-5-methyl- (Mitomycin C)	H ₂ O	8.9 × 10 ⁹	297	CP/Ac-14	S = RF; used $k_d = 5.0 \times 10^5$ s ⁻¹ .	79F816
16.8	Benz[e]naphtho[2'3':5,6]fluoreno[1,9ab]oxepin-5,10,19-trione, 5c,8,8a,16-tetrahydro-1,8,11,15,18-pentahydroxy-13-methyl- (Rubellin A)	CD ₃ OD	≤8 × 10 ⁶		CP/LI-12	S = RB; used $k_d = 4.2 \times 10^3$ s ⁻¹ .	92F063
16.9	Benzo[ghi]perylene-4,11-dione, 1-acetyl-1,2-dihydro-5,10-dihydroxy-2-(1-hydroxyethyl)-3,7,8,12-tetramethoxy- (Elsinochrome B)	CD ₃ OD	≤5.5 × 10 ⁶		CP/LI-12	S = RB; used $k_d = 4.2 \times 10^3$ s ⁻¹ .	92F063
16.10	Benzo[ghi]perylene-4,11-dione, 1,2-diacetyl-1,2-dihydro-5,10-dihydroxy-3,7,8,12-tetramethoxy-, trans- (Elsinochrome A)	CD ₃ OD	≤5 × 10 ⁶		CP/LI-12	S = RB; used $k_d = 4.2 \times 10^3$ s ⁻¹ .	92F063
16.11	Benzopyran-6-ol, 3,4-dihydro-2,5,7,8-tetramethyl-2-(4,8,12-trimethyltridecyl)-, acetate (α-Tocopheryl acetate)	C ₅ H ₅ N	≤1.6 × 10 ⁶		CP/A'c-23	S = A' = Rub; used $k_d = 6.0 \times 10^4$ s ⁻¹ , $k_{A'} = 4 \times 10^7$ L mol ⁻¹ s ⁻¹ .	743112

TABLE 16. Rate constants for the interaction of singlet oxygen with some compounds of biological interest. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
16.12	1,4-Benzoquinone						
	C ₆ H ₆ / EtOH (67:33)	1.6 × 10 ⁷			CP/A'c-19	S = RB; A' = Chl a; used $k_d = 1 \times 10^5$ s ⁻¹ .	78F404
	EtOH	3.4 × 10 ⁷			CP/A'c-19	S = RB; A' = Chl a; used $k_d = 1 \times 10^5$ s ⁻¹ .	78F404
16.13	1,4-Benzoquinone, methyl-						
	C ₆ H ₅ CH ₃	<10 ⁴			PL/Ld-2	S = TPP.	93Y025
16.14	Cholesterol						
	C ₅ H ₅ N		0.89	293	CP/Oc-14	S = HP.	57F008
	C ₆ D ₆	5.7 × 10 ⁴			PL/Ld-2	S = TPP.	89A331
	MeOH/ H ₂ O (95:5)	2.5 × 10 ⁸		310	CR/LI-12	used $k_d = 1.8 \times 10^5$ s ⁻¹ ; soln. cont. 0.05 mol L ⁻¹ MeONa and 5 × 10 ⁻⁴ mol L ⁻¹ CoCl ₂ , ¹ O ₂ * from autoxidation of oxytetracycline.	92M228
16.15	Cholesteryl benzoate						
	C ₆ H ₆ / C ₅ H ₅ N (91:9)	4.0		293	CP/Oc-14	S = HP.	57F008
16.16	Cobrynic acid, bis(cyano)-7-de(carboxymethyl)-7,8-didehydro-, hexamethyl ester						
	C ₆ D ₆	8 × 10 ⁶		293	CP/Ac-14	S = A; k_d not given.	83F222
	CHCl ₃	5 × 10 ⁶		293	CP/Ac-14	S = A; k_d not given.	83F222
	MeOH	6 × 10 ⁶		293	CP/Ac-14	S = A; used $k_d = 1 \times 10^5$ s ⁻¹ .	83F222
16.17	Coenzyme A						
	CHCl ₃ / EtOH (50:50)	8.9 × 10 ⁷ 1.2 × 10 ⁶ (k_t)		310	CR/LI,Ac-12,14	used $k_d = 1 \times 10^5$ s ⁻¹ ; ¹ O ₂ * from NDPO ₂ .	90E622
16.18	Cytidine						
	H ₂ O pH = 7.1	≤1 × 10 ⁶		298	CP/Oc-19	S = Phenosafranine; No measurable effect.	78A360
16.19	Cytidine 5'-monophosphate						
	D ₂ O pH = ~7	2.8 × 10 ⁴		293	PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	86A198
16.20	Cytosine						
	D ₂ O pH = ~7	<8 × 10 ⁴		293	PL/Ld-2	S = H ₂ TPPS ⁴⁻ ; No reaction at [A] = 5 × 10 ⁻² mol L ⁻¹ .	86A198
	DMSO/ H ₂ O (75:25)	6.8 × 10 ⁴		298	CP/A'c-16	S = MB; A' = DPBF; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	86R210
16.21	2'-Deoxyadenosine 5'-monophosphate						
	H ₂ O pH = 7.0, 10.5	<10 ⁵			PL/Ld-2	S = RB.	87A043
	D ₂ O pH = ~7	7.5 × 10 ⁴		293	PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	86A198
16.22	2'-Deoxycytidine-5'-monophosphate						
	H ₂ O pH = 7.0, 10.5	<10 ⁵			PL/Ld-2	S = RB.	87A043
16.23	2'-Deoxyguanosine 5'-monophosphate						
	D ₂ O pH = ~7	5 × 10 ⁶		293	PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	86A198
	H ₂ O pH = 7.0 pH = 10.5	5.3 × 10 ⁶ 5.7 × 10 ⁷			PL/Ld-2	S = RB.	87A043

TABLE 16. Rate constants for the interaction of singlet oxygen with some compounds of biological interest. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
16.24	Deoxyribonucleic acid (DNA)						
	D ₂ O pH = 7	7×10^5			PL/Ld-2	S = AlCl(tspc).	89R092
	H ₂ O pH = 7.0 pH = 10.5	5.1×10^5 2.3×10^6			PL/Ld-2	S = RB.	87A043
16.25	3'-Deoxythymidine 5'-monophosphate						
	D ₂ O pH = ~7	1.4×10^5		293	PL/Ld-2	S = H ₂ TPPS ⁴⁺ .	86A198
16.26	Ergosterol						
	C ₅ H ₃ N		4.5×10^{-3}	293	CP/Oc-14	S = HP.	57F008
	C ₆ H ₆		2.1×10^7		PL/Ld-2	S = TPP.	87E055
16.27	Ergosterol acetate						
	CHCl ₃	2.7×10^7			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81F445
16.28	1,3,5(10),8-Estratetraen-17-one, 3-methoxy-						
	MeOH	1.6×10^7 (k_r)	9.2×10^{-3} (β_r)		CP/Ac-14	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	91F405
16.29	1,3,5(10),9(11)-Estratetraen-17-one, 3-(acetoxy)-						
	MeOH	5.5×10^5 (k_r)		273	CP/Ac-17	S = RB; A' = 3-Methoxy-1,3,5(10),9(11)-estratetraen-17-one; used $k_r^{A'} = 4.3 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.13$.	89A202
16.30	1,3,5(10),9(11)-Estratetraen-17-one, 3-(benzoyloxy)-						
	MeOH	7.5×10^5 (k_r)		273	CP/Ac-17	S = RB; A' = 3-Methoxy-1,3,5(10),9(11)-estratetraen-17-one; used $k_r^{A'} = 4.3 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.17$.	89A202
16.31	1,3,5(10),9(11)-Estratetraen-17-one, 3-hydroxy-						
	MeOH	6.8×10^6 (k_r)		273	CP/Ac-17	S = RB; A' = 3-Methoxy-1,3,5(10),9(11)-estratetraen-17-one; used $k_r^{A'} = 4.3 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 1.6$.	89A202
16.32	1,3,5(10),9(11)-Estratetraen-17-one, 3-methoxy-						
	MeOH	4.3×10^6 (k_r)	0.033 (β_r)	273	CP/Ac-14	S = RB; used $k_d = 1.4 \times 10^5$ s ⁻¹ .	89A202
16.33	1,3,5(10),9(11)-Estratetraen-17-one, 3-(toluenesulfonyloxy)-						
	MeOH	2.7×10^5 (k_r)		273	CP/Ac-17	S = RB; A' = 3-Methoxy-1,3,5(10),9(11)-estratetraen-17-one; used $k_r^{A'} = 4.3 \times 10^6$ L mol ⁻¹ s ⁻¹ ; meas. $k_r/k_r^{A'} = 0.06$.	89A202
16.34	Glucose						
	D ₂ O	1.4×10^4			PL/Ld-2	S = RF, water-sol. TPP derivs., or Chl a in micelles contg. Triton X-100.	86F149
16.35	Guanine						
	DMSO/H ₂ O (75:25)	5.0×10^5		298	CP/A'c-16	S = MB; A' = DPBF; used $k_d = 1.2 \times 10^5$ s ⁻¹ , $k_{A'} = 6.9 \times 10^8$ L mol ⁻¹ s ⁻¹ .	86R210
16.36	Guanosine						
	D ₂ O pD = 7.4	6.2×10^6		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	91R127
	H ₂ O pH = 7.1	$\leq 1 \times 10^6$		298	CP/Oc-19	S = Phenosafranine; No measurable effect.	78A360
	H ₂ O/MeOH (50:50)	$\leq 1 \times 10^7$			PL/A'd-5	S = MB; A' = DPBF.	72F516

TABLE 16. Rate constants for the interaction of singlet oxygen with some compounds of biological interest. — Continued

	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
16.47	Guanosine, 2'-deoxy-						
	D ₂ O pD = 7.4	5.2×10^6		310	CR/LI-12	used $k_d = 1.9 \times 10^4$ s ⁻¹ ; ${}^1\text{O}_2^*$ from NDPO ₂ .	91R127
	D ₂ O pH = ~7	5×10^6		293	PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	86A198
	H ₂ O pH = 10				CP/A'c-17	S = MB; A' = BHMF; meas. $k_t/k_{t'}^{A'} = 0.4$; sensitizer is immobilized on glass beads.	83F166
16.48	Guanosine 5'-monophosphate						
	D ₂ O pH = ~7	5×10^6		293	PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	86A198
16.49	Imidazo[1,2- <i>a</i>]pyrazin-3-one, 4-methyl-6-[4-[2-[3-carboxy-4-(6-hydroxy-3-xanthenon-9-yl)phenylthiocarbamylethoxy]phenyl]- (FCLA)						
	H ₂ O pH = 7.1	8.0×10^7			CL/LI-12	S = RB; used $k_d = 2.4 \times 10^5$ s ⁻¹ .	90F502
16.50	Imidazo[1,2- <i>a</i>]pyrazin-3-one, 2-methyl-6-(4-methoxyphenyl)- (MCLA)						
	H ₂ O pH = 5.4	2.9×10^9			CL/LI-12	S = RB; used $k_d = 2.4 \times 10^5$ s ⁻¹ .	91E570 89E798
16.51	Imidazo[1,2- <i>a</i>]pyrazin-3-one, 2-methyl-6-phenyl- (CLA)						
	H ₂ O pH = 7.1	6.3×10^8			CL/LI-12	S = RB; used $k_d = 2.4 \times 10^5$ s ⁻¹ .	90F502
16.52	Inosine diphosphate, ester with 1,4-dihydro-1-β-D-ribofuranosyl-3-pyridinecarboxamide						
	CH ₃ CN/ D ₂ O (80:20)	1.6×10^8			PL/A'd-5	S = 2-ACN; A' = DPBF.	81A191
	CH ₃ CN/ D ₂ O (20:80)	8.8×10^7 (k_t)			PL/A'd-5	S = 2-ACN; A' = DPBF; k_t calcd. from k_A ; $56 \pm 11\%$ electron transfer giving O ₂ ⁻ , detd. from buildup of radical anion in soln. contg. (2-20) × 10 ⁻⁵ mol L ⁻¹ 1,4-benzoquinone.	81A191
		7.9×10^7 (k_q)					
16.53	Methyl stearate						
	C ₅ H ₁₁ N	$\leq 5 \times 10^4$ (k_t)			CP/Ac,A'c-17	S = PP; A' = Cholesterol; No measurable effect.	743115
16.54	1,4-Naphthoquinone, 2-methyl-						
	EtOH	2.4×10^6			CP/A'c-19	S = RB; A' = Chl a; used $k_d = 1 \times 10^5$ s ⁻¹ .	78F404
16.55	1,8-Naphthyridine-3-carboxylic acid, 1,4-dihydro-1-ethyl-7-methyl-4-oxo- (Nalidixic acid)						
	D ₂ O pH = 4.4	$\sim 2.5 \times 10^8$			PL/Ld-12	S = A; used $k_d = 3 \times 10^4$ s ⁻¹ ; estd. from increased lifetime on decreasing concn.	88R070
16.56	1,8-Naphthyridine-3-carboxylic acid, 1,4-dihydro-1-ethyl-7-methyl-4-oxo-, anion (Nalidixic acid anion)						
	D ₂ O pH = 8.9	$\sim 1 \times 10^8$			PL/Ld-12	S = A; used $k_d = 3 \times 10^4$ s ⁻¹ ; estd. from increased lifetime on decreasing concn.	88R070
16.57	Nicotinamide adenine dinucleotide						
	D ₂ O	3×10^4			PL/Ld-2	S = RF, water-sol. TPP derivs., or Chl a in micelles contg. Triton X-100.	86F149
	D ₂ O	$< 5 \times 10^6$			PL/A'd-5	S = 2-ACN; A' = DPBF.	80R072 81A191
16.58	Nicotinamide adenine dinucleotide, reduced						
	CH ₃ CN/ MeOH (75:25)	7.5×10^7			PL/A'd-5	S = 2-ACN; A' = DPBF.	81A191
	CH ₃ CN/ D ₂ O (20:80)	4.3×10^7 (k_t)			PL/A'd-5	S = 2-ACN; A' = DPBF; k_t calcd. from k_A ; $54.5 \pm 15\%$ electron transfer giving O ₂ ⁻ , detd. from buildup of radical anion in soln. contg. (2-20) × 10 ⁻⁵ mol L ⁻¹ 1,4-benzoquinone.	81A191
		3.6×10^7 (k_q)					

TABLE 16. Rate constants for the interaction of singlet oxygen with some compounds of biological interest. — Continued

No.	Solvent	k ($\text{L mol}^{-1} \text{s}^{-1}$)	$\beta (k_d/k)$ (mol L^{-1})	T (K)	Method	Comment	Ref.
16.48	Nicotinamide adenine dinucleotide, reduced	— Continued					
	D ₂ O	2×10^7			PL/Ld-2	S = RF, water-sol. TPP derivs., or Chl a in micelles contg. Triton X-100.	86F149
	D ₂ O (mic)	7.9×10^7			PL/A'd-5	S = 2-ACN; A' = DPBF; 0.1 mol L ⁻¹ SDS.	80R072 81A191
	H ₂ O pH = 7.4	$\sim 2 \times 10^9$			CP/Oc-19	S = FMN; Q = N ₃ ⁻ ; used $k_Q = 2 \times 10^9 \text{ L mol}^{-1} \text{s}^{-1}$.	81A422
16.49	Nicotinamide adenine dinucleotide phosphate	D ₂ O	2×10^5		PL/Ld-2	S = RF, H ₂ TPPS ⁴⁻ , or Chl in Triton X-100 micelles.	86F149
16.50	Nicotinamide-adenine dinucleotide phosphate, reduced	CH ₃ CN/ D ₂ O (80:20)	1.3×10^8		PL/A'd-5	S = 2-ACN; A' = DPBF.	81A191
	CH ₃ CN/ D ₂ O (20:80)	$8.4 \times 10^7 (k_r)$ $5.0 \times 10^7 (k_q)$			PL/A'd-5	S = 2-ACN; A' = DPBF; k_r calcd. from k_A ; $63 \pm 9\%$ electron transfer giving O ₂ ⁻ , detd. from buildup of radical anion in soln. contg. (2-20) $\times 10^{-5}$ mol L ⁻¹ 1,4-benzoquinone.	81A191
16.51	Nicotinamide mononucleotide, reduced	CH ₃ CN/ D ₂ O (80:20)	8.0×10^7		PL/A'd-5	S = 2-ACN; A' = DPBF.	81A191
16.52	Ommochrome	D ₂ O pD = 7.5-8.1	3×10^6		PL/Ld-2	S = ?; rate is per monomer unit of MW = 600; ommochromes extracted from shrimp eyes contain pigments of the ommin group, e.g., phenothiazine-phenoxazine structures with 3-amino-3-carboxy-1-oxopropyl side chains.	87R227
16.53	3,10-Perylenedione, 4,9-dihydroxy-1,12-bis(2-hydroxypropyl)-2,6,7,11-tetramethoxy- (Phleichrome)	CD ₃ OD	$\leq 6 \times 10^6$		CP/LI-12	S = RB; used $k_d = 4.2 \times 10^3 \text{ s}^{-1}$.	92F063
16.54	Perylo[1,12-def]-1,3-dioxepin-5,11-dione, 6-amino-12-hydroxy-8,9-bis(2-hydroxypropyl)-7,10-dimethoxy- (Aminocercosporin)	CD ₃ OD	$\leq 3 \times 10^6$		CP/LI-12	S = RB; used $k_d = 4.2 \times 10^3 \text{ s}^{-1}$.	92F063
16.55	Perylo[1,12-def]-1,3-dioxepin-5,11-dione, 6,12-dihydroxy-8,9-bis(2-hydroxypropyl)-7,10-dimethoxy- (Cercosporin)	CD ₃ OD	3.5×10^6		CP/LI-12	S = RB; used $k_d = 4.2 \times 10^3 \text{ s}^{-1}$.	92F063
16.56	Perylo[1,12-def]-1,3-dioxepin-5,12-dione, 6,11-diamino-8,9-bis(2-hydroxypropyl)-7,10-dimethoxy-	CD ₃ OD	3.6×10^6		CP/LI-12	S = RB; used $k_d = 4.2 \times 10^3 \text{ s}^{-1}$.	92F063
16.57	Perylo[1,12-def]-1,3-dioxepin-6,11-dione, 8,9-bis(2-hydroxypropyl)-5,7,10,12-tetramethoxy-	CD ₃ OD	1.1×10^7		CP/LI-12	S = RB; used $k_d = 4.2 \times 10^3 \text{ s}^{-1}$.	92F063
16.58	Perylo[1,12-def]-1,3-dioxepin-6,11-dione, 5-hydroxy-8,9-bis(2-hydroxypropyl)-7,10,12-trimethoxy-	CD ₃ OD	$\leq 10^7$		CP/LI-12	S = RB; used $k_d = 4.2 \times 10^3 \text{ s}^{-1}$.	92F063
16.59	Phenanthro[1,10,9,8-opgra]perylene-7,14-dione, 1,3,4,6,8,13-hexahydroxy-10,11-dialkyl.	CH ₃ CN	2.0×10^{10}		PL/Ld-2	S = Riboflavin tetrabutyrate; Red pigment extracted from <i>Blepharisma japonicum</i> ; on irradiation red form converts to blue form which reacts with ${}^1\text{O}_2^*$ with $k = 4.4 \times 10^{10} \text{ L mol}^{-1} \text{s}^{-1}$; k probably overestimated due to impurities in the crude extracts.	92R107
16.60	Phenanthro[1,10,9,8-opgra]perylene-7,14-dione, 1,3,4,6,8,13-hexahydroxy-10,11-dimethyl- (Hypericin, HYP)	EtOH	5.0×10^9		PL/Ld-2	S = Riboflavin tetrabutyrate.	92R107

TABLE 16. Rate constants for the interaction of singlet oxygen with some compounds of biological interest. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
16.61	Pregna-5,16-dien-20-one, 3-(acetoxy)- (3 β)						
	C ₆ H ₆ / C ₅ H ₅ N (91:9)	4.7		293	CP/Oc-14	S = HP.	57F008
16.62	Pregn-5-en-20-one, 3-hydroxy-, (3 β)						
	C ₅ H ₅ N	1.7		293	CP/Oc-15	S = HP.	587002
	C ₆ H ₆ / C ₅ H ₅ N (60:40)	1.8		293	CP/Oc-15	S = HP.	587002
	C ₆ H ₆ / C ₅ H ₅ N (71:29)	1.8		293	CP/Oc-14	S = HP.	57F008
16.63	Purine, 6-amino- (Adenine)						
	D ₂ O pH = ~7	<4 × 10 ⁵		293	PL/Ld-2	S = H ₂ TPPS ⁴⁺ ; No reaction at [A] = 9 × 10 ⁻³ mol L ⁻¹ .	86A198
	DMSO/ H ₂ O (75:25)	7.2 × 10 ⁴		298	CP/A'c-16	S = MB; A' = DPBF; used $k_d = 1.2 \times 10^5$ s ⁻¹ , $k_{A'} = 6.9 \times 10^3$ L mol ⁻¹ s ⁻¹ .	86R210
	H ₂ O pH = 7.1	≤1 × 10 ⁶		298	CP/Oc-19	S = Phenosafranine; No measurable effect.	78A360
16.64	Purin-2,6,8-trione, 7,9-dihydro-, anion (Urate ion)						
	D ₂ O pD = 7.4	3.6 × 10 ⁸		298	PL/Ld-2	S = H ₂ TMpyP ⁴⁺ .	90R134
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_t/k_r^{A'} = 1.0$; sensitizer immobilized on glass beads.	88R064
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_t/k_r^{A'} = 0.50$.	88R064
16.65	2-Pyrimidinamine, N,N,4,5-tetramethyl-6-(phenylmethoxy)-						
	CHCl ₃	9.0 × 10 ⁶			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81A363
16.66	Pyrimidin-4-ol, 5-butyl-2-(dimethylamino)-6-methyl- (Dimethirimol)						
	CHCl ₃	5.5 × 10 ⁶			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81A363
16.67	Pyrimidin-4-ol, 5-butyl-2-(ethylamino)-6-methyl- (Ethirimol)						
	CHCl ₃	2.3 × 10 ⁶			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81A363
16.68	Pyrimidin-4-ol, 2-(diethylamino)-6-methyl-						
	CHCl ₃	7.0 × 10 ⁵			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81A363
16.69	Pyrimidin-4-ol, 2-(dimethylamino)-5,6-dimethyl-						
	CHCl ₃	1.2 × 10 ⁷			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81A363
	H ₂ O pH = 4.0	4.4 × 10 ⁷			CP/Oc-15	S = RB; used $k_d = 5 \times 10^5$ s ⁻¹ .	83A315
	pH = 7.0	8 × 10 ⁷					
	pH = 9.0	1.4 × 10 ⁸					
	H ₂ O	9.0 × 10 ⁷		296	CP/Oc-15	S = RB; used $k_d = 5 \times 10^5$ s ⁻¹ ; $\Delta S^\ddagger = 150.5$ J K ⁻¹ mol ⁻¹ ; $E_a = 71.9$ kJ mol ⁻¹ ; studied at 279–296 K.	83A315
16.70	Pyrimidin-4-ol, 2-(dimethylamino)-6-methyl-						
	CHCl ₃	1.0 × 10 ⁶			CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	81A363

TABLE 16. Rate constants for the interaction of singlet oxygen with some compounds of biological interest. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
16.71	Rhodopsin						
	H ₂ O pH = 8	1.1×10^9			CP/Pa-15	S = Ret; used $k_d = 3.5 \times 10^5$ s ⁻¹ ; membrane suspension; sulfur radicals obs. by esr.	85A44 ¹
16.72	Riboflavin						
	MeOD	6.0×10^7			PL/Ld-2	S = A.	88A165
16.73	Sitosteryl acetate						
	C ₆ H ₆ / C ₅ H ₅ N (91:9)		6.5	293	CP/Oc-14	S = HP.	57F008
16.74	Stigmasteryl acetate						
	C ₆ H ₆ / C ₅ H ₅ N (91:9)		4.1	293	CP/Oc-14	S = HP.	57F008
16.75	Sucrose						
	D ₂ O	3×10^4			PL/Ld-2	S = RF, water-sol. TPP derivs., or Chl a in micelles contg. Triton X-100.	86F149
16.76	4-Thiouridine						
	D ₂ O pH = ~7	1.2×10^6		293	PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	86A198
16.77	Thymidine						
	D ₂ O pH = ~7	3×10^4		293	PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	86A198
16.78	Thymidine 5'-monophosphate						
	H ₂ O pH = 7.0, 10.5	<10 ⁵			PL/Ld-2	S = RB.	87A043
16.79	Thymine						
	D ₂ O pH = ~7	<1.6 × 10 ⁵		293	PL/Ld-2	S = H ₂ TPPS ⁴⁻ ; No reaction at [A] = 2.5×10^{-2} mol L ⁻¹ .	86A198
	DMSO/ H ₂ O (75:25)	2.5×10^4		298	CP/A'c-16	S = MB; A' = DPBF; used $k_d = 1.2 \times 10^5$ s ⁻¹ .	86R210
	H ₂ O pH = 7.1	$\leq 1 \times 10^6$		298	CP/Oc-19	S = Phenosafranine; No measurable effect.	78A360
16.80	Uridine						
	D ₂ O pH = ~7	5×10^4		293	PL/Ld-2	S = H ₂ TPPS ⁴⁻ .	86A198

TABLE 17. Rate constants for the interaction of singlet oxygen with miscellaneous substrates.

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
17.1	Acetic acid						
	CCl ₄	2.3×10^3			MP/LI-12	S = TPP; used $k_d = 42$ s ⁻¹ .	83E425 83F151
17.2	Acetone						
	CCl ₄	1.9×10^3		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329
17.3	Acetone-d ₆						
	CCl ₄	95		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329
17.4	Acetonitrile						
	CCl ₄	9×10^2		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329
17.5	Acetonitrile-d ₃						
	CCl ₄	55		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329
17.6	Butanoic acid						
	C ₆ D ₆	3.7×10^3			PL/Ld-2	S = TPP.	89A331
17.7	Butanoic acid, 3,3-dimethyl-2-oxo-						
	MeOH	$\sim 1 \times 10^6$			PL/A'd-5	S = MB; A' = DPBF.	79A386
17.8	1-Butanol						
	CCl ₄	2×10^3			PL/A'd-8	S = MB; A' = DPBF.	76F903
17.9	Chloroform						
	CCl ₄	5×10^2		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329
17.10	Chloroform-d						
	CCl ₄	22		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329
17.11	2,5-Cyclohexadien-4-one, 1,2-ethanediylidenebis[3-[2-hydroxy-5-methyl-3-(1,1-dimethylethyl)benzyl]-5-(1,1-dimethylethyl)]-						
	CH ₂ Cl ₂	8.2×10^9		303	CP/Oc-23	S = MB; A' = TME; used $k_d = 1.2 \times 10^4$ s ⁻¹ , $k_{A'} = 3.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; k calcd. in [81Z251].	76F921
17.12	2,5-Cyclohexadien-4-one, 4,4'-(1,2-ethanediylidene)bis[2,6-bis(1,1-dimethylethyl)-						
	CH ₂ Cl ₂	5.4×10^9		303	CP/Oc-23	S = MB; A' = Rub; used $k_d = 1.2 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; k calcd. in [81Z251].	76F921
	CH ₂ Cl ₂	2.3×10^9		303	CP/Oc-23	S = MB; A' = CHD; used $k_d = 1.2 \times 10^4$ s ⁻¹ , $k_{A'} = 3.5 \times 10^6$ L mol ⁻¹ s ⁻¹ ; k calcd. in [81Z251].	76F921
	CH ₂ Cl ₂	4.3×10^9		303	CP/Oc-23	S = MB; A' = TME; used $k_d = 1.2 \times 10^4$ s ⁻¹ , $k_{A'} = 4.0 \times 10^7$ L mol ⁻¹ s ⁻¹ ; k calcd. in [81Z251].	76F921
17.13	Cyclohexane						
	CCl ₄	6.4×10^3		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329
17.14	Cyclohexane-d ₁₂						
	CCl ₄	3.4×10^2		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329
17.15	1,3-Cyclohexanedione, 5,5-dimethyl-						
	D ₂ O pD = 11.2	9×10^7		310	CR/A'c-32	A' = DPBF; used $k_d = 1.5 \times 10^4$ s ⁻¹ , ¹ O ₂ * from Dopamine/H ₂ O ₂ .	89M038

TABLE 17. Rate constants for the interaction of singlet oxygen with miscellaneous substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	β (k_d/k) (mol L ⁻¹)	T (K)	Method	Comment	Ref.
17.15 1,3-Cyclohexanedione, 5,5-dimethyl- — Continued							
	D ₂ O pD = 11.2	1.2×10^8		310	CR/A'c-32	A' = DPBF; used $k_d = 1.5 \times 10^4$ s ⁻¹ ; $^1\text{O}_2$ * from DOPA/H ₂ O ₂ .	89M038
17.16 Dichloromethane							
	CCl ₄	1.1×10^3		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329
17.17 Dichloromethane-<i>d</i>₂							
	CCl ₄	5.3×10^2		298	CP/LI-12	S = Benz[de]anthracen-7-one, phenalenone or fluorenone; used $k_d = 32$ s ⁻¹ .	82E329
17.18 1,1-Dimethylethyl methyl ether							
	CD ₃ OD	4.1×10^3			CP/LI-12	S = RB; used $k_d = 4.0 \times 10^3$ s ⁻¹ .	92A386
	(CH ₃) ₃ COCH ₃	4.1×10^3			PL/Ld-2	S = TPP.	92A386
17.19 2,2-Dimethylpropyl methyl ether							
	CD ₃ OD	4.9×10^3			CP/LI-12	S = RB; used $k_d = 4.0 \times 10^3$ s ⁻¹ .	92A386
	(CH ₃) ₃ CCH ₂ OCH ₃	2.9×10^3			PL/Ld-2	S = TPP.	92A386
17.20 1,4-Dioxane							
	CCl ₄	7×10^2			PL/A'd-8	S = MB; A' = DPBF.	76F903
17.21 Disilirane, 1,1,2,2-tetrakis(2,4,6-trimethylphenyl)-							
	C ₆ H ₆	6.4×10^4			CP/LI-12	S = TPP; k_d not given.	93F160
17.22 Docosanoic acid (Behenic acid)							
	CCl ₄	1.1×10^4			MP/LI-12	S = TPP; used $k_d = 42$ s ⁻¹ .	83E425 83F151
17.23 Dodecane							
	CCl ₄	1.1×10^4 (k_q)			MP/LI-12	S = TPP; used $k_d = 39$ s ⁻¹ .	83E813
17.24 Dodecanoic acid							
	CCl ₄	6.1×10^3			MP/LI-12	S = TPP; used $k_d = 42$ s ⁻¹ .	83E425 83F151
17.25 Dodecyl sulfate, sodium salt							
	H ₂ O (mic)	1.0×10^9		313	CP/A'c-23	S = Py; A' = DPBF; Q = NaN ₃ ; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; S and A' solubilized in SDS micelles.	78A174
17.26 Dodecytrimethylammonium chloride							
	H ₂ O (mic)	1.7×10^8		313	CP/A'c-23	S = Py; A' = DPBF; Q = NaN ₃ ; used $k_d = 5.0 \times 10^5$ s ⁻¹ ; S and A' solubilized in DTAC micelles.	78A174
17.27 Eicosanoic acid (Arachidic acid)							
	C ₆ D ₆	1.2×10^4			PL/Ld-2	S = TPP.	89A331
	CCl ₄	1.0×10^4			MP/LI-12	S = TPP; used $k_d = 42$ s ⁻¹ .	83E425 83F151
17.28 Ethane, iodo-							
	C ₆ H ₅ Br/ CH ₃ COCH ₃ (67:33)	$\leq 10^6$			CP/A'c-23	S = A' = Rub; No measurable effect.	767126
	C ₆ H ₅ Br/ MeOH (67:33)	$\leq 10^6$			CP/A'c-23	S = A' = Rub; No measurable effect.	767126
	C ₆ H ₆	2×10^4			PL/A'd-8	S = An; A' = DPBF.	76F902

TABLE 17. Rate constants for the interaction of singlet oxygen with miscellaneous substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
17.29	Ethanol						
	CCl ₄	2×10^3			PL/A'd-8	S = MB; A' = DPBF.	76F903
17.30	Ethylene-propylene-ethylideneborbornene terpolymer						
	c-C ₆ H ₁₂	$3.6 \times 10^4 (k_r)$		298	CP/Ac-17	S = A' = DPBF; used $k_t^{A'} = 5.4 \times 10^8$ L mol ⁻¹ s ⁻¹ ; [A] = 5 g/L, assuming that reaction is only at the ethylideneborbornene groups, $k = 8.8 \times 10^7$.	84P629
17.31	Formamide, N,N-dimethyl-						
	EtOH	$3.0 \times 10^2 (k_r)$			CP/Pa-19	S = RB; used $k_d = 8.3 \times 10^4$ s ⁻¹ ; used k_q/k_r (not reported); k_r derived assuming a singlet oxygen mechanism.	75F655
17.32	Fullerene-C ₆₀						
	C ₆ D ₆	5×10^5			PL/Ld-2	S = TPP.	91E003
	CCl ₄	6.1×10^4			PL/Ld-2	S = A.	93E091
17.33	Fullerene-C ₇₀						
	C ₆ F ₆	3×10^6			PL/Ld-2	S = A.	93E301
17.34	Heptane						
	CCl ₄ / MeOH (96:4)	4×10^2			PL/A'd-8	S = MB; A' = DPBF.	777162
17.35	Heptane, 3-methyl -						
	CCl ₄ / MeOH (96:4)	6×10^2			PL/A'd-8	S = MB; A' = DPBF.	777162
17.36	Heptane, 4-methyl -						
	CCl ₄ / MeOH (96:4)	5×10^2			PL/A'd-8	S = MB; A' = DPBF.	777162
17.37	Hexadecanoic acid						
	C ₆ D ₆	1.2×10^4			PL/Ld-2	S = TPP.	89A331
	CCl ₄	8.0×10^3			MP/LI-12	S = TPP; used $k_d = 42$ s ⁻¹ .	83E425 83F151
17.38	Hexanoic acid						
	C ₆ D ₆	5.6×10^3			PL/Ld-2	S = TPP.	89A331
	CCl ₄	3.9×10^3			MP/LI-12	S = TPP; used $k_d = 42$ s ⁻¹ .	83E425 83F151
17.39	Malonic acid						
	D ₂ O	4×10^4			PL/Ld-2	S = RF, water-sol. TPP derivs., or Chl a in micelles contg. Triton X-100.	.86F149
17.40	Methanol						
	CCl ₄	3×10^3			PL/A'd-8	S = MB; A' = DPBF.	76F903
17.41	5,8-Methano-[1,2,4]triazolo[1,2-a]pyridazine-1,3-dione, tetrahydro-2-methyl-						
	CH ₂ Cl ₂	4×10^4			PR/A'd-8	S = An; A' = DPBF.	79A457
17.42	Methyl esters of fatty acids from peanut oil						
	MeOH/ CHCl ₃ (80:20)	8×10^4			PL/Ld-2	S = ZnTPP.	91U181
17.43	Methyl esters of fatty acids from soybean oil						
	MeOH	1.7×10^5			CP/A'c-33	S = A' = Rub; used $k_d = 1.4 \times 10^5$ s ⁻¹ , $k_{A'} = 6 \times 10^7$ L mol ⁻¹ s ⁻¹ .	91U201
	MeOH/ CHCl ₃ (80:20)	3.6×10^5			PL/Ld-2	S = ZnTPP.	91U181

TABLE 17. Rate constants for the interaction of singlet oxygen with miscellaneous substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
17.44	Methyl esters of fatty acids from sunflower oil						
	MeOH/CHCl ₃ (80:20)	9×10^4			PL/Ld-2	S = ZnTPP.	91U181
17.45	4-Morpholineethanesulfonic acid (MES)						
	D ₂ O	1.6×10^4			PL/Ld-2	S = RF, water-sol. TPP derivs., or Chl a in micelles contg. Triton X-100.	86F149
17.46	Oxadisilirane, 2,2,3,3-tetrakis(2,6-diethylphenyl)-						
	C ₆ H ₆	8.4×10^4			CP/LI-12	S = TPP; k_d not given.	93F160 90F241
17.47	Oxadisilirane, 2,2,3,3-tetrakis(2,6-diisopropylphenyl)-						
	C ₆ H ₆	7.1×10^4			CP/LI-12	S = TPP; k_d not given.	93F160 90F241
17.48	Oxalate ion						
	D ₂ O	2×10^4			PL/Ld-2	S = RF, water-sol. TPP derivs., or Chl a in micelles contg. Triton X-100; counter ion K ⁺ .	86F149
17.49	Oxirane, 2,3-diphenyl-, (E)-						
	ClCF ₂ CCl ₂ F	5.0×10^3			PL/Ld-2	S = DNT.	87A072
17.50	Pentane						
	CCl ₄	5.9×10^3			MP/LI-12	S = TPP; used $k_d = 39$ s ⁻¹ .	83E813
17.51	Pentanedioic acid, 2-oxo- (α -Ketoglutaric acid)						
	MeOH	$\sim 3 \times 10^6$			PL/A'd-5	S = MB; A' = DPBF.	79A386
17.52	Phosphoric acid, mono(2-aminoethyl) mono(2,3-dihydroxypropyl) ester (Phosphatidylethanolamine)						
	CCl ₄	10^5			MP/LI-12	S = PP; used $k_d = 36$ s ⁻¹ .	79F463
17.53	Phosphoric acid, mono(2-dimethylammonioethyl) mono(2,3-dihydroxypropyl) ester (Phosphatidylcholine, Lecithin)						
	CCl ₄	10^5			MP/LI-12	S = PP; used $k_d = 36$ s ⁻¹ .	79F463
	D ₂ O (ves)	7.4×10^5			CP/Pa-15	S = RB; used $k_d = 1.8 \times 10^4$ s ⁻¹ ; P = Lecithin hydroperoxide; 0.05% egg yolk lecithin, ave. MW per monomer = 750.	86N104
	H ₂ O (ves)	7.5×10^5			CP/Pa-15	S = RB; used $k_d = 2.5 \times 10^5$ s ⁻¹ ; P = Lecithin hydroperoxide; 0.05% egg yolk lecithin.	86N104
17.54	Phycol-400						
	D ₂ O	2.4×10^7			PL/I.d-2	S = RF, water-sol. TPP derivs., or Chl a in micelles contg. Triton X-100.	86F149
17.55	Polycarbonate						
	ClI ₂ Cl ₂	3.9×10^3			PL/Ld-2	S = An or Ac.	86E884
17.56	Poly(ethylene glycol)						
	D ₂ O	4×10^4			PL/Ld-2	S = RF, water-sol. TPP derivs., or Chl a in micelles contg. Triton X-100.	86F149
17.57	Poly(methyl methacrylate)						
	CH ₂ Cl ₂	2.2×10^3			PL/Ld-2	S = An.	86E809
17.58	Polystyrene						
	CH ₂ Cl ₂	3.3×10^3			PL/Ld-2	S = An or Ac.	86E884
17.59	1,3-Propanediol, 2-amino-2-(hydroxymethyl)- (Tris)						
	D ₂ O	1.5×10^4			PL/Ld-2	S = RF, water-sol. TPP derivs., or Chl a in micelles contg. Triton X-100.	86F149

TABLE 17. Rate constants for the interaction of singlet oxygen with miscellaneous substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
17.60	Propanoic acid, 2-oxo-3-phenyl- (Phenylpyruvic acid)						
	MeOH	<7 × 10 ⁶			PL/A'd-5	S = MB; A' = DPBF.	79A386
17.61	2-Propanol						
	CCl ₄	2 × 10 ³			PL/A'd-8	S = MB; A' = DPBF.	76F903
17.62	2-Propanol, 2-methyl- (<i>tert</i> -BuOH)						
	CCl ₄	2 × 10 ³			PL/A'd-8	S = MB; A' = DPBF.	76F903
17.63	Quadracyclane						
	CHCl ₃	4.8 × 10 ⁴			CR/A'c-33	A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ ; ${}^1\text{O}_2^*$ from DMNO ₂ .	81E003
17.64	Singlet oxygen						
	C ₆ F ₆	2.6 × 10 ⁷			PL/Ld-2	S = Chr; self-annihilation from high intensity photolysis; M = ${}^1\text{O}_2^*$.	87F333
	c-C ₆ H ₁₂	<4 × 10 ⁷			PL/Ld-2	S = Chr; self-annihilation from high intensity photolysis; M = ${}^1\text{O}_2^*$.	87F333
	CCl ₄	1 × 10 ⁹			PL/Ld-2	S = PdMPDME; [${}^1\text{O}_2^*$] estd. from sensitizer triplet concn.	79E846
	O ₂	2 × 10 ⁵		77	PL/Ld-2	In liquid ¹⁸ O ₂ $k = (2 \pm 0.6) \times 10^6$. Emission is monitored at 635 nm.	82E575
17.65	Soybean oil						
	CH ₂ Cl ₂	1.4 × 10 ⁵	0.073	293	CP/Oc-15	S = Chl a; used $k_d = 1.0 \times 10^4$ s ⁻¹ .	91U026
	CH ₂ Cl ₂	1.0 × 10 ⁵	0.11	298	CP/Pa-15	S = Chl a; used $k_d = 1.1 \times 10^4$ s ⁻¹ ; measured peroxide formation.	91U180 91U128
17.66	Stearic acid						
	C ₆ D ₆	1.2 × 10 ⁴			PL/Ld-2	S = TPP.	89A331
	CCl ₄	9.0 × 10 ³			MP/LI-12	S = TPP; used $k_d = 42$ s ⁻¹ .	83E425 83F151
17.67	Tetradecanoic acid (Myristic acid)						
	C ₆ D ₆	9.9 × 10 ³			PL/Ld-2	S = TPP.	89A331
17.68	Tetrahydrofuran						
	CCl ₄	2 × 10 ³			PL/A'd-8	S = MB; A' = DPBF.	76F903
17.69	1,2,4-Triazolo[1,3]cyclopropa[1,2,3-cd]cyclopropa[gh]pentalene, 1,2,5,6,6a,6b,6c,6d-octahydro-1,6[1',2']-4-methyl-3,5-dioxo-						
	CH ₂ Cl ₂	1.0 × 10 ⁵			PR/A'd-8	S = An; A' = DPBF.	79A457
17.70	1,2,4-Triazolo[1,3]cyclopropa[1,2,3-cd]cyclopropa[gh]pentalene, 1,2,5,6,6a,6b,6c,6d-octahydro-1,6[1',2']-4-methyl-3,5-dioxo-						
	CH ₂ Cl ₂	3.4 × 10 ⁴			PR/A'd-8	S = An; A' = DPBF.	79A457
17.71	Tributyl phosphite						
	CH ₃ COCH ₃				CP/Pa,P'a-17	S = RB; A' = (C ₂ H ₅ O) ₃ P; meas. $k_r/k_r^{A'} = 0.78$.	747495
17.72	Tricyclohexyl phosphite						
	CH ₃ COCH ₃				CP/Pa,P'a-17	S = RB; A' = (C ₂ H ₅ O) ₃ P; meas. $k_r/k_r^{A'} = 0.60$.	747495
17.73	Triethyl phosphite						
	CH ₃ COCH ₃				CP/Pa,P'a-17	S = MB; A' = (CH ₃ O) ₃ P; meas. $k_r/k_r^{A'} = 1.5$.	747495
	C ₆ H ₆ / MeOH (80:20)	2.5 × 10 ⁷			CP/Pa-19	S = RB; A' = Car; k_d and $k_{A'}$ not given.	747495
17.74	Trimethyl phosphite						
	C ₆ H ₆	4.7 × 10 ⁴			PL/Ld-2	S = Ac.	89A099

TABLE 17. Rate constants for the interaction of singlet oxygen with miscellaneous substrates. — Continued

No.	Solvent	k (L mol ⁻¹ s ⁻¹)	$\beta (k_d/k)$ (mol L ⁻¹)	T (K)	Method	Comment	Ref.
17.74 Trimethyl phosphite — Continued							
	C ₆ H ₆ / MeOH (80:20)	1.5×10^7			CP/Pa-19	S = RB; A' = Car; k_d and $k_{A'}$ not given.	74F495
	C ₆ H ₆ / MeOH (80:20)	3.9×10^4			PL/Ld-2	S = Ac.	89A099
	CH ₃ CN	6.6×10^4			PL/Ld-2	S = Ac.	89A099
	CH ₃ COCH ₃	6.3×10^4			PL/Ld-2	S = Ac.	89A099
	MeOD	2.7×10^4			PL/Ld-2	S = Ac.	89A099
17.75 Triphenylphosphine							
	CH ₂ Cl ₂ / MeOH (90:10)	5×10^6			PL/A'd-5	S = MB; A' = DPBF.	87F655
	CHCl ₃	8.5×10^6		253	CP/A'c-33	S = A' = Rub; used $k_d = 1.7 \times 10^4$ s ⁻¹ , $k_{A'} = 5.3 \times 10^7$ L mol ⁻¹ s ⁻¹ .	82N064
	MeOH		6.0×10^{-3}	293	CP/Oc-15	S = MB; $E_a = 5.0$ kJ mol ⁻¹ .	68F288
	MeOH		5.0×10^{-3}	293	CP/Oc-15	S = RB; $E_a = 5.9$ kJ mol ⁻¹ .	68F288
17.76 Undecanoic acid							
	C ₆ D ₆	9.3×10^3			PL/Ld-2	S = TPP.	89A331
17.77 Urea							
	H ₂ O pH = 7.0			283	CP/A'c-17	S = RB; A' = BHMF; meas. $k_t/k_{t'}^{A'} = 1.1$.	88R064
17.78 Urea, allyl-							
	H ₂ O pH = 7.1	$<2 \times 10^5$		298	CP/Oc-19	S = Phenosafranine; Q = NaN ₃ ; used $k_Q = 2.0 \times 10^8$ L mol ⁻¹ s ⁻¹ .	78F020
17.79 Valeric acid							
	C ₆ D ₆	4.0×10^3			PL/Ld-2	S = TPP.	89A331

12. References to Tables

- 57F004 Reactions in the liquid phase photochemistry of anthracene. Part I. The photo-oxidation of anthracenes in solution. Bowen, E.J., *Discuss. Faraday Soc.* : 143-6 (1953)
- 57F004 The photochemistry of anthracenes. Part 3. Inter-relations between fluorescence quenching, dimerization, and photo-oxidation. Bowen, E.J.; Tanner, D.W., *Trans. Faraday Soc.* **51**: 475-81 (1955)
- 57F007 A diffusion-controlled step in chlorophyll-sensitized photochemical autoxidations. Livingston, R.; Owens, K.E., *J. Am. Chem. Soc.* **78**: 3301-5 (1956)
- 57F008 Zur photosensibilisierten Autoxydation der Steroide. Darstellung von Steroid-Hydroperoxyden Mittels Phototoxischer Photosensibilisatoren. Schenck, G.O.; Gollnick, K.; Neumueller, O.A., *Ann. Chem.* **603**: 46-59 (1957)
- 58F002 Zur photosensibilisierten Autoxydation der Steroide. Synthese tertiärer Steroid-Hydroperoxyde, insbesondere des Δ^6 -Allopregnend- β -ol-20-on-5 α -Hydroperoxyds. Schenck, G.O.; Neumueller, O.A., *Ann. Chem.* **618**: 194-201 (1958)
- 58F004 Cinétique et inhibition de réactions photosensibilisées en présence d'oxygène moléculaire. Schenck, G.O.; Gollnick, K., *J. Chim. Phys. Phys.-Chim. Biol.* **55**: 892-8 (1958)
- 59F003 A further study of the photochemical autoxidation of anthracenes. Livingston, R.; Rao, V.S., *J. Phys. Chem.* **63**: 794-9 (1959)
- 59F008 Photo-oxidation, sensitised by proflavine, of furfuryl alcohol, N-allyl thiourea and histidine. Sluyterman, L.A.E., *Recl. Trav. Chim. Pays-Bas* **80**: 989-1002 (1961)
- 62F005 Chemismus und Kinetik der durch Xanthenfarbstoffe photosensibilisierten O_2 -Übertragungen. (Analyse der Beteiligung angeregter Singulett- und Tripletzustände sowie der Natur der Terminationsprozesse). Gollnick, K., Ph.D. Thesis, Univ. Goettingen, 1962, 144p.
- 63F028 Reactivities in photosensitized olefin oxidations. Kopecky, K.R.; Reich, H.J., *Can. J. Chem.* **43**: 2265-70 (1965)
- 65F029 On the mechanism of the photo-oxidation of amino acids sensitized by methylene blue. Weil, L., *Arch. Biochem. Biophys.* **110**: 57-68 (1965)
- 66F041 Excited singlet molecular oxygen in photooxidation. Wilson, T., *J. Am. Chem. Soc.* **88**: 2898-902 (1966)
- 66F197 Quantum yield studies of the dye-sensitized photoinactivation of trypsin. Glad, B.W.; Spikes, J.D., *Radiat. Res.* **27**: 237-49 (1966)
- 68F285 Quenching of singlet oxygen by tertiary aliphatic amines. Effect of DABCO. Ouannes, C.; Wilson, T., *J. Am. Chem. Soc.* **90**: 6527-8 (1968)
- 68F287 Reactions by singlet oxygen and via transient complex (between dye and oxygen), produced simultaneously through the interaction between triplet eosine and oxygen. Koizumi, M.; Usui, Y., *Tetrahedron Lett.* : 6011-4 (1968)
- 68F288 Zur Photosensibilisierten Sauerstoffübertragung. Untersuchung der Terminationsstufen durch Belichtungen bei tiefen Temperaturen. Koch, E., *Tetrahedron* **24**: 6295-318 (1968)
- 68F289 Type II photosensitized oxygenation reactions. Gollnick, K., *Adv. Chem. Ser.* **77**: 78-101 (1968)
- 68F292 Chemistry of singlet oxygen. V. Reactivity and kinetic characterization. Higgins, R.; Foote, C.S.; Cheng, H., *Adv. Chem.* **77**: 102-17 (1968)
- 69F388 The photoperoxidation of unsaturated organic molecules. IV. The photosensitized reaction. Stevens, B.; Algar, B.E., *J. Phys. Chem.* **73**: 1711-5 (1969)
- 70F079 The photoperoxidation of unsaturated organic molecules. VI. The inhibited reaction. Algar, B.E.; Stevens, B., *J. Phys. Chem.* **74**: 3029-34 (1970)
- 70F188 Chemistry of singlet oxygen. X. Carotenoid quenching parallels biological protection. Foote, C.S.; Chang, Y.C.; Denny, R.W., *J. Am. Chem. Soc.* **92**: 5216-8 (1970)
- 70F250 Photo-oxidation of carbanions: The reactivity of singlet oxygen with 9-benzenesulphonylfluoren-9-yl anion. Bethell, D.; Wilkinson, R.G., *Chem. Commun.* : 1178-9 (1970)
- 70F200 1,2 Cycloaddition of singlet oxygen to 9,9'-bifluorenylidene. Richardson, W.H.; Hodge, V., *J. Org. Chem.* **35**: 1216-7 (1970)
- 70F454 Photoinduced reactions. XL. Addition of singlet oxygen to monocyclic aromatic ring. Saito, I.; Kato, S.; Matsuura, T., *Tetrahedron Lett.* : 239-42 (1970)
- 70F732 Proflavine-sensitized photooxidation of methionine. Jori, G.; Cauzzo, G., *Photochem. Photobiol.* **12**: 231-7 (1970)
- 70F733 Competitive modes of reaction of singlet oxygen. Bartlett, P.D.; Mendenhall, G.D.; Schaap, A.P., *Ann. New York Acad. Sci.* **171**: 79-87 (1970)
- 70F734 Quenching of singlet oxygen. Foote, C.S.; Denny, R.W.; Weaver, L.; Chang, Y.; Peters, J., *Ann. New York Acad. Sci.* **171**: 139-45 (1970)
- 70F735 Photosensitised oxygenation as a function of the triplet energy of sensitizers. Gollnick, K.; Franken, T.; Schade, G.; Doerhoefer, G., *Ann. N.Y. Acad. Sci.* **171**: 89-107 (1970)
- 71F299 Cinétique d'oxydation de l'aniline par l'oxygène singulet formé à partir d'aniline à l'état triplet. Pouyet, B.; Chapelon, R., *C. R. Hebd. Séances Acad. Sci., Ser. C* **272**: 1753-6 (1971)
- 71F356 Chemistry of singlet oxygen. XIII. Solvent effects on the reaction with olefins. Foote, C.S.; Denny, R.W., *J. Am. Chem. Soc.* **93**: 5168-71 (1971)
- 71F398 Solvent effects in dye-sensitized photooxidation reactions. Young, R.H.; Wehrly, K.; Martin, R.L., *J. Am. Chem. Soc.* **93**: 5774-9 (1971)
- 71F403 Photo-oxygenation of 4H-pyran-4-thiones and 4H-thiopyran-4-thiones: Reactivity and kinetic characterization. Ishibe, N.; Odani, M.; Sunami, M., *J. Chem. Soc. Pt. B* : 1837-40 (1971)
- 71F512 Collisional deactivation of $O_2(^1\Delta_g)$. Findlay, F.D.; Snelling, D.R., *J. Chem. Phys.* **55**: 545-51 (1971)
- 71F525 Direct measurement of the lifetime of $^1\Delta$ oxygen in solution. Merkel, P.B.; Kearns, D.R., *Chem. Phys. Lett.* **12**: 120-22 (1971)
- 71F577 Chemistry of singlet oxygen. XII. Electronic effects on rate and products of the reaction with olefins. Foote, C.S.; Denny, R.W., *J. Am. Chem. Soc.* **93**: 5162-7 (1971)
- 71F580 Photooxidation of sulfides. Foote, C.S.; Peters, J.W., *23d. Internat. Congr. Pure Appl. Chem., Special Lectures*, London, Butterworths, 1971, Vol. 4, p. 129-54.
- 71F581 Photoinduced reactions-LXIX. Photosensitized oxygenation of cis, cis-1,5-cyclooctadiene and cis-cyclooctene. Matsuura, T.; Horinaka, A.; Yoshida, H.; Butsugan, Y., *Tetrahedron* **27**: 3095-100 (1971)
- 72F027 Deuterium effects on singlet oxygen lifetimes in solutions. A new test of singlet oxygen reactions. Merkel, P.B.; Nilsson, R.; Kearns, D.R., *J. Am. Chem. Soc.* **94**: 1030-1 (1972)
- 72F028 Chemistry of singlet oxygen. XVI. Long lifetime of singlet oxygen in carbon disulfide. Foote, C.; Peterson, E.R.; Lee, K.-W., *J. Am. Chem. Soc.* **94**: 1032-3 (1972)
- 72F116 The dye-sensitized photooxygenation of hexamethyleneammonium hexamethylenedithiocarbamate by xanthene dyes. Yamase, T.; Kokado, H.; Inoue, E., *Bull. Chem. Soc. Jpn.* **45**: 726-31 (1972)
- 72F196 The photoperoxidation of unsaturated organic molecules. Intersystem crossing in 9,10-diphenylanthracene. Stevens, B.; Mills, L.E., *Chem. Phys. Lett.* **15**: 381-2 (1972)
- 72F260 Radiationless decay of singlet molecular oxygen in solution. An experimental and theoretical study of electronic-to-vibrational energy transfer. Merkel, P.B.; Kearns, D.R., *J. Am. Chem. Soc.* **94**: 7244-53 (1972)
- 72F319 Singlet oxygen ($^1\Delta_g$) quenching in the liquid phase by metal(II) chelates. Carlsson, D.I.; Mendenhall, G.D.; Suprunchuk, T.; Wiles, D.M., *J. Am. Chem. Soc.* **94**: 8960-2 (1972)

- 72A019 Dye-sensitized photooxidation of tocopherols. Correlation between singlet oxygen reactivity and vitamin E activity. Grams, G.W.; Eskins, K., *Biochemistry* **11**: 606-8 (1972)
- 72A020 Photoinduced reactions. LIX. Reactivity of singlet oxygen toward methoxybenzenes. Saito, I.; Imuta, M.; Matsuura, T., *Tetrahedron* **28**: 5307-11 (1972)
- 72F510 Substituent effects in dye-sensitized photooxidation reactions of furans. Young, R.H.; Martin, R.L.; Chinh, N.; Mallon, C.; Kayser, R.H., *Can. J. Chem.* **50**: 932-8 (1972)
- 72F512 Kinetic evidence for both quenching and reaction of singlet oxygen with triethylamine in pyridine solution. Smith, W.F., Jr., *J. Am. Chem. Soc.* **94**(1) 186-90 (1972)
- 72F514 On the mechanism of quenching of singlet oxygen by amines. Young, R.H.; Martin, R.L., *J. Am. Chem. Soc.* **94**: 5183-5 (1972)
- 72F515 Role of azide in singlet oxygen reactions: reaction of azide with singlet oxygen. Hasty, N.; Merkel, P.B.; Radlick, P.; Kearns, D.R., *Tetrahedron Lett.* : 49-52 (1972)
- 72F516 Unambiguous evidence for the participation of singlet oxygen ($^1\Delta$) in photodynamic oxidation of amino acids. Nilsson, R.; Merkel, P.B.; Kearns, D.R., *Photochem. Photobiol.* **16**: 117-24 (1972)
- 72F517 The reactivity of isobenzofuran. Young, R.H.; Feriozi, D.T., *J. Chem. Soc. Chem. Commun.* 841-2 (1972)
- 72F518 Inhibition de l'oxygène singulet. Dalle, J.P.; Magous, R.; Mousseron-Cauet, M., *Photochem. Photobiol.* **15**: 411-9 (1972)
- 72F519 An investigation of reaction and quenching mechanisms of singlet molecular oxygen. Martin, R.L., Ph.D., Thesis, Georgetown Univ., Wash. D.C., 1972, 330p.
- 72F520 Photoinduced reactions. LVI. Participation of singlet oxygen in the hydrogen abstraction from a phenol in the photosensitized oxygenation. Matsuura, T.; Yoshimura, N.; Nishinaga, A.; Saito, I., *Tetrahedron* **28**: 4933-8 (1972)
- 72F521 Iodine-sensitized photoformation of singlet oxygen. Olmsted, J., III; Karal, G., *J. Am. Chem. Soc.* **94**: 3305-13 (1972)
- 732066 The possible importance of singlet oxygen quenching reactions in the photostabilization of polyolefins. Carlson, D.J.; Suprunchuk, T.; Wiles, D.M., *J. Polym. Sci., Polym. Lett. Ed.* **11**: 61-5 (1973)
- 733045 Photodynamic inactivation of lysozyme by eosin. Kepka, A.G.; Grossweiner, L.I., *Photochem. Photobiol.* **18**: 49-61 (1973)
- 736061 On the mechanism of quenching of singlet oxygen by amines. III. Evidence for a charge-transfer-like complex. Young, R.H.; Martin, R.L.; Feriozi, D.; Brewer, D.; Kayser, R., *Photochem. Photobiol.* **17**: 233-44 (1973)
- 737014 The determination of rate constants for reaction and lifetimes of singlet oxygen in solution by a flash photolysis technique. Young, R.H.; Brewer, D.; Keller, R.A., *J. Am. Chem. Soc.* **95**: 375-9 (1973)
- 737202 Sensitized photoxygenation: Change from the Type I (radical) to Type II (singlet oxygen) mechanism. Kramer, H.E.A.; Maute, A., *Photochem. Photobiol.* **17**: 413-23 (1973)
- 737333 Energy transfer processes involving ultraviolet stabilizers. Quenching of singlet oxygen. Guillory, J.P.; Cook, C.F., *J. Polym. Sci., Polym. Chem. Ed.* **11**: 1927-37 (1973)
- 737334 Quenching of singlet molecular oxygen by polyolefin additives in carbon disulfide solution. Flood, J.; Russell, K.E.; Wan, J.K.S., *Macromolecules* **6**: 669-71 (1973)
- 737438 On the mechanism of quenching of singlet oxygen in solution. Farnilo, A.; Wilkinson, F., *Photochem. Photobiol.* **18**: 447-50 (1973)
- 737479 Etude cinétique de l'oxydation photosensibilisée d'aldehydes α -éthyleniques en phase liquide à -10°C. Carmier, J.-C.; Deglise, X., C.R. *Hebd. Séances Acad. Sci., Ser. C* **277**: 1187-9 (1973)
- 73F659 Kinetics of photoperoxidation in solution. Stevens, B., *Acc. Chem. Res.* **6**: 90-6 (1973)
- 73F660 Photooxidation of isobenzofurans. A dual mechanism process. Olmsted, J., III; Akashah, T., *J. Am. Chem. Soc.* **95**: 6211-5 (1973)
- 73F662 Mechanisms of singlet oxygen reactions. Intermediates in the reaction of singlet oxygen with dienes. Hasty, N.M.; Kearns, D.R., *J. Am. Chem. Soc.* **95**: 3380-1 (1973)
- 73F664 The reaction of singlet oxygen with α - and β -pinenes. Jefford, C.W.; Boschung, A.F.; Moriarty, R.M.; Rimbault, C.G.; Laffer, M.J.J., *Helv. Chim. Acta* **56**: 2649-59 (1973)
- 743112 On the quenching of singlet oxygen by α -tocopherol. Fahrenholz, S.R.; Doleiden, F.H.; Trozzolo, A.M.; Lamola, A.A., *Photochem. Photobiol.* **20**: 505-9 (1974)
- 743113 Chemistry of singlet oxygen. XVIII. Rates of reaction and quenching of α -tocopherol and singlet oxygen. Foote, C.S.; Ching, T.Y.; Geller, G.G., *Photochem. Photobiol.* **20**: 511-3 (1974)
- 743114 The photoperoxidation of unsaturated organic molecules. XIII. $O_2^{1\Delta_g}$ quenching by α -tocopherol. Stevens, B.; Small, R.D., Jr.; Perez, S.R., *Photochem. Photobiol.* **20**: 515-7 (1974)
- 743115 Reactivity of cholesterol and some fatty acids toward singlet oxygen. Doleiden, F.H.; Fahrenholz, S.R.; Lamola, A.A.; Trozzolo, A.M., *Photochem. Photobiol.* **20**: 519-21 (1974)
- 746110 Fluorescence and quenching of $O_2^{1\Delta_g}$ and $(O_2^{1\Delta_g})_2$ in liquid oxygen. Huestis, D.L.; Black, G.; Edelstein, S.A.; Sharpless, R.L., *J. Chem. Phys.* **60**: 4471-4 (1974)
- 747042 Carotenoid chromophore length and protection against photosensitization. Mathews-Roth, M.M.; Wilson, T.; Fujimori, E.; Krinsky, N.I., *Photochem. Photobiol.* **19**: 217-22 (1974)
- 747044 A standard system to determine the quantum yield of singlet oxygen formation in aqueous solution. Usui, Y.; Kamogawa, K., *Photochem. Photobiol.* **19**: 245-7 (1974)
- 747102 Measurement of the absolute rate constants for singlet molecular oxygen ($^1\Delta_g$) reaction with 1,3-diphenylisobenzofuran and physical quenching by ground state molecular oxygen. Matheson, I.B.C.; Lee, J.; Yamanashi, B.S.; Wolbarsht, M.L., *J. Am. Chem. Soc.* **96**: 3343-8 (1974)
- 747103 Reaction rate of bilirubin with singlet oxygen ($^1\Delta_g$) and its strong enhancement by added base. Matheson, I.B.C.; Curry, N.U.; Lee, J., *J. Am. Chem. Soc.* **96**: 3348-51 (1974)
- 747207 The photoperoxidation of unsaturated organic molecules. X. Solvent effect on $O_2^{1\Delta}$ acceptor reactivity. Stevens, B.; Perez, S.R., *Mol. Photochem.* **6**: 1-7 (1974)
- 747312 Photoperoxidation of unsaturated organic molecules. XIV. $O_2^{1\Delta_g}$ acceptor properties and reactivity. Stevens, B.; Perez, S.R.; Ors, J.A., *J. Am. Chem. Soc.* **96**: 6846-50 (1974)
- 747341 The quenching of singlet oxygen ($^1\Delta_g$) by transition metal chelates. Carlsson, D.J.; Suprunchuk, T.; Wiles, D.M., *Can. J. Chem.* **52**: 3728-37 (1974)
- 747495 Singlet oxygen oxidation of phosphites to phosphates. Bolduc, P.R.; Goe, G.L., *J. Org. Chem.* **39**: 3178-9 (1974)
- 74F640 On the mechanism of quenching by amines: A new method for investigation of interactions with triplet states. Young, R.H.; Brewer, D.; Kayser, R.; Martin, R.; Feriozi, D.; Keller, R.A., *Can. J. Chem.* **52**: 2889-93 (1974)
- 74F641 The photoperoxidation of unsaturated organic molecules. IX. Lipoic acid inhibition of rubrene autoperoxidation. Stevens, B.; Perez, S.R.; Small, R.D., *Photochem. Photobiol.* **19**: 315-6 (1974)
- 74F642 Importance of singlet oxygen in the degradation of rubbers and plastics. Carlsson, D.J.; Wiles, D.M., *Rubber Chem. Technol.* **47**: 991-1004 (1974)
- 74F643 Role of singlet oxygen in some chemiluminescence and enzyme oxidation reactions. Nilsson, R.; Kearns, D.R., *J. Phys. Chem.* **78**: 1681-3 (1974)
- 74F645 Study of the quenching of triplet carbonyl groups and singlet oxygen by light stabilizers. Hrdlovic, P.; Danecek, J.; Karvas, M.; Durmis, J., *Chem. Zvesti* **28**: 792-801 (1974)
- 74F646 A kinetic investigation of the generation and decay of singlet oxygen in dye sensitized systems. Part 1. Brewer, D.R., III, Ph.D., Thesis, Georgetown Univ., Wash. D.C., 1974, 694p.
- 74F647 Reaction of singlet oxygen with 2-methylnorborn-2-ene, 2-methylidenenorbornane, and their 7,7-dimethyl derivatives. The transition state geometry for hydroperoxidation. Jefford, C.W.; Boschung, A.F., *Helv. Chim. Acta* **57**: 2242-57 (1974)

- 74F648 Photosensitized oxidation of di-n-butyl sulphide. Casagrande, M.; Gennari, G.; Cauzzo, G., *Gazz. Chim. Ital.* **104**: 1251-7 (1974)
- 74F649 The photo-peroxidation of unsaturated organic molecules. $O_2^{\cdot 1\Delta_g}$ -acceptor re-encounter probabilities. Stevens, B.; Ors, J.A.; Pinsky, M.L., *Chem. Phys. Lett.* **27**: 157-60 (1974)
- 74R214 Concerning the formation of singlet O_2 during the decomposition of H_2O_2 by catalase. Porter, D.T.; Ingraham, L.L., *Biochim. Biophys. Acta* **334**: 97-102 (1974)
- 752063 Singlet oxygen and polymer photooxidations. I. Sensitizers, quenchers, and reactants. Zweig, A.; Henderson, W.A., Jr., *J. Polym. Sci., Polym. Chem. Ed.* **13**: 717-36 (1975)
- 753071 Chemistry of singlet oxygen. XXI. Kinetics of bilirubin photooxygenation. Foote, C.S.; Ching, T.-Y., *J. Am. Chem. Soc.* **97**: 6209-14 (1975)
- 756223 Untersuchung der selbst-sensibilisierten Photooxidation des Rubrens in verschiedenen Lösungsmitteln. Brauer, H.-D.; Wagener, H., *Ber. Bunsenges. Phys. Chem.* **79**: 597-604 (1975)
- 757088 Radiationless decay of singlet molecular oxygen in solution. II. Temperature dependence and solvent effects. Long, C.A.; Kearns, D.R., *J. Am. Chem. Soc.* **97**: 2018-20 (1975)
- 757147 The quenching of singlet oxygen by amino acids and proteins. Matheson, I.B.C.; Etheridge, R.D.; Kratowich, N.R.; Lee, J., *Photochem. Photobiol.* **21**: 165-71 (1975)
- 757166 Quenching of singlet molecular oxygen in solution by azomethine dyes. Smith, W.F., Jr.; Herkstroeter, W.G.; Eddy, K.L., *J. Am. Chem. Soc.* **97**: 2764-70 (1975)
- 757445 Kinetics and mechanism of nitroxyl radical formation during sensitized amine oxidation. Ivanov, V.B.; Shlyapintokh, V.Ya.; Khvostach, O.M.; Shapiro, A.B.; Rozantsev, E.G., *J. Photochem.* **4**: 313-9 (1975)
- 757485 Photodynamic sensitization by 8-methoxysoralen via the singlet oxygen mechanism. Poppe, W.; Grossweiner, L.I., *Photochem. Photobiol.* **22**: 217-9 (1975)
- 757558 An experimental approach to the estimation of diffusive displacement parameters for bimolecular reactions in solution. Stevens, B.; Williams, R.R., *Chem. Phys. Lett.* **36**: 100-5 (1975)
- 757578 Singlet molecular oxygen and superoxide radical anion. Rosenthal, I., *Isr. J. Chem.* **13**: 86-90 (1975)
- 757623 Reactions of singlet oxygen III. Dye sensitized oxidation of phenothiazine. Nathu Ram; Bansal, W.R.; Uppal, S.S.; Sidhu, K.S., *Proceedings of the Symposium on Singlet Molecular Oxygen*, Bhabha At. Res. Cent., Bombay, 10-13 Feb 1975, p.141-52
- 75F652 Dye-sensitized photooxygenation of tert-butylpyrroles. Lightner, D.A.; Pak, C.-S., *J. Org. Chem.* **40**: 2724-8 (1975)
- 75F653 Chemistry of singlet oxygen. XXII. Photooxidation of nitrones. Ching, T.-Y.; Foote, C.S., *Tetrahedron Lett.* : 3771-4 (1975)
- 75F654 Effect of the aggregate state of the medium on singlet oxygen quenching. Bystritskaya, E.V.; Karpukhin, O.N., *Dokl. Phys. Chem.* **221**: 353-6 (1975) Translated from: *Dokl. Akad. Nauk SSSR* **221**: 1100-3 (1975)
- 75F655 New method for the mass spectrometric study of the kinetics of molecular oxygen consumption in liquid-phase photooxidation processes. Zolotoi, N.B.; Karpov, G.V.; Skurat, V.E., *Kinet. Catal.* **16**: 1229-35 (1975) Translated from: *Kinet. Katal.* **16**: 1415-22 (1975)
- 75F656 Chemistry of singlet oxygen. XX. Mechanism of the sensitized photooxidation of enamines. Foote, C.S.; Dzakpasu, A.A.; Lin, J.W.-P., *Tetrahedron Lett.* : 1247-50 (1975)
- 75F669 Mechanism of photo-oxidation of the porphyrins by oxygen. Byteva, I.M.; Gurinovich, G.P.; Petsol'd, O.M., *Biophysics* **20**: 48-52 (1975) Translated from: *Biofizika* **20**: 51-5 (1975)
- 763011 The photoperoxidation of unsaturated organic molecules. XV. $O_2^{\cdot 1\Delta_g}$ quenching by bilirubin and biliverdin. Stevens, B.; Small, R.D., Jr., *Photochem. Photobiol.* **23**: 33-6 (1976)
- 766072 Chemistry of superoxide ion. III. Quenching of singlet oxygen. Guiraud, H.J.; Foote, C.S., *J. Am. Chem. Soc.* **98**: 1984-6 (1976)
- 767041 Kinetic aspects of dye-sensitized photo-oxygenation. Brkic, D.; Forzatti, P.; Pasquon, I.; Trifiro, F., *J. Photochem.* **5**: 23-32 (1976)
- 767071 Direct and sensitized photooxidation of cyanine dyes. Byers, G.W.; Gross, S.; Henrichs, P.M., *Photochem. Photobiol.* **23**: 37-43 (1976)
- 767105 On the mechanism of the acridine orange sensitized photodynamic inactivation of lysozyme. I. Basic kinetics. Schmidt, H.; Rosenkranz, P., *Z. Naturforsch., C, Biosci.* **31**: 29-39 (1976)
- 767126 The quenching effect of iodide ion on singlet oxygen. Rosenthal, I.; Frimer, A., *Photochem. Photobiol.* **23**: 209-11 (1976)
- 767197 Photooxidation of benzophenone oxime and derivatives. Wamser, C.C.; Herring, J.W., *J. Org. Chem.* **41**: 1476-7 (1976)
- 767247 The photosensitized formation and reaction of singlet oxygen, $O_2^{\cdot 1\Delta_g}$, in aqueous micellar systems. Gorman, A.A.; Lovering, G.; Rodgers, M.A.J., *Photochem. Photobiol.* **23**: 399-403 (1976)
- 767422 Photoperoxidation of unsaturated organic molecules. 16. Excitation energy fission. Stevens, B.; Ors, J.A., *J. Phys. Chem.* **80**: 2164-5 (1976)
- 767570 Photoperoxidation of heterocordianthrone in different solvents. Wagener, H.; Brauer, H.-D., *Mol. Photochem.* **7**: 441-64 (1976)
- 76F900 Nitroso compounds and azo dioxides as quenchers of singlet oxygen ($^1\Delta_g$) and sensitizer triplet states. Singh, P.; Ullman, E.F., *J. Am. Chem. Soc.* **98**: 3018-9 (1976)
- 76F902 Physical properties of singlet oxygen in fluid solvents. Wilkinson, F., *Singlet Oxygen, Reactions with Organic Compounds and Polymers*, B. Ranby and J.F. Rabek (eds.), John Wiley & Sons, New York, N.Y., 1976, p.27-35
- 76F903 The mechanism of quenching of singlet oxygen. Young, R.H.; Brewer, D.R., *Singlet Oxygen, Reactions with Organic Compounds and Polymers*, B. Ranby and J.F. Rabek (eds.), John Wiley & Sons, New York, N.Y., 1976, p.36-47
- 76F909 Singlet oxygen reactions with polyisoprene model molecules: 4-methyl-1-octene and 4,8-dimethyl-4,8-dodecadiene. Chaineaux, J.; Tanielian, C., *Singlet Oxygen, Reactions with Organic Compounds and Polymers*, B. Ranby and J.F. Rabek (eds.), John Wiley & Sons, New York, N.Y., 1976, p.164-73
- 76F921 Antioxidants and stabilizers. LXV. Contribution to the investigation of sensitized photooxidation of phenolic antioxidants. Quenching of singlet oxygen with stilbenequinoid compounds. Taimr, L.; Pospisil, J., *Angew. Makromol. Chem.* **52**: 31-8 (1976)
- 770055 Concerning the use of amines as probes for participation of singlet oxygen in dye-sensitised oxygenation reactions. Davidson, R.S.; Trethewey, K.R., *J. Chem. Soc., Perkin Trans. 2* : 178-82 (1977)
- 777074 Anthracene sulphonate sensitized photooxidation of iodide ion in aqueous solution via singlet oxygen. Rohatgi-Mukherjee, K.K.; Gupta, A.K., *Chem. Phys. Lett.* **46**: 368-71 (1977)
- 777113 Intermediates in the decomposition of aliphatic diazo-compounds. Part 13. Mechanistic studies on the reaction of diaryldiazomethanes with singlet molecular oxygen. Bethell, D.; McKeivor, R., *J. Chem. Soc., Perkin Trans. 2* : 327-33 (1977)
- 777129 The singlet oxygen reactivity of biliverdin. Matheson, I.B.C.; Toledo, M.M., *Photochem. Photobiol.* **25**: 243-8 (1977)
- 777162 Interaction between singlet oxygen ($^1\Delta_g$) and model compounds for polymers. A flash photolytic study. Bortolus, P.; Dellonte, S.; Beggiato, G.; Corio, W., *Eur. Polym. J.* **13**: 185-8 (1977)
- 777176 Die Kinetik der selbst-sensibilisierten Photooxidation des Heterocordianthrone im Bereich niedriger Konzentrationen. Drews, W.; Schmidt, R.; Brauer, H.-D., *J. Photochem.* **6**: 391-403 (1976/77)
- 777221 Energy transfer from luminescent transition metal complexes to oxygen. Demas, J.N.; Harris, E.W.; McBride, R.P., *J. Am. Chem. Soc.* **99**: 3547-51 (1977)
- 777240 Dye sensitized photooxidation of chlorpromazine. Rosenthal, I.; BercoVICI, T.; Frimer, A., *J. Heterocycl. Chem.* **14**: 355-7 (1977)
- 777378 Products formed by photosensitized oxidation of unsaturated fatty acid esters. Terao, J.; Matsushita, S., *J. Am. Oil Chem. Soc.* **54**: 234-8 (1977)

- 777433 Chemistry of singlet oxygen. XXV. Photooxygenation of methionine. Sysak, P.K.; Foote, C.S.; Ching, T.-Y., *Photochem. Photobiol.* **26**: 19-27 (1977)
- 777486 Quenching of singlet oxygen by aliphatic amines. Monroe, B.M., *J. Phys. Chem.* **81**: 1861-4 (1977)
- 777489 Mechanisms of the photosensitized oxidation of tyramine. Seely, G.R., *Photochem. Photobiol.* **26**: 115-23 (1977)
- 777645 Reaction of singlet oxygen with 4-methyl-2,3-dihydro- γ -pyrans. Frimer, A.A.; Bartlett, P.D.; Boschung, A.F.; Jewett, J.G., *J. Am. Chem. Soc.* **99**: 7977-86 (1977)
- 77E617 Photoluminescence of singlet oxygen in solutions of chlorophylls and pheophytins. Krasnovsky, A.A., Jr., *Biophysics* **22**: 960-2 (1977) Translated from: *Biofizika* **22**: 927-8 (1977)
- 77F794 Studies on the sensitized photooxidation of 1,3-dithiol-thiones-(2). Fanghaenel, E.; Lutze, G., *J. Prakt. Chem.* **319**: 875-82 (1977)
- 77F858 Quenching effect of tocopherols on the methyl linoleate photooxidation and their oxidation products. Yamauchi, R.; Matsushita, S., *Agric. Biol. Chem.* **41**: 1425-30 (1977)
- 77F876 I. A study of the kinetics of the 1,2-cycloaddition of singlet oxygen to vinyl ethers. II. An investigation of the reaction of singlet oxygen with adamantlylideneadamantane. Falter, G.R., Ph.D., Thesis, Wayne State Univ., Detroit, MI, 1977, 157p.
- 78A005 Rates of reaction of singlet oxygen with olefins. Monroe, B.M., *J. Phys. Chem.* **82**: 15-8 (1978)
- 78A171 Chemistry of singlet oxygen. XXVI. Photooxygenation of phenols. Thomas, M.J.; Foote, C.S., *Photochem. Photobiol.* **27**: 683-93 (1978)
- 78A174 Production and reaction of singlet oxygen in aqueous micellar solutions using pyrene as photosensitizer. Miyoshi, N.; Tomita, G., *Z Naturforsch., Teil B* **33B**: 622-7 (1978)
- 78A227 Mechanisms of chlorine oxidation of hydrogen peroxide. Held, A.M.; Halko, D.J.; Hurst, J.K., *J. Am. Chem. Soc.* **100**: 5732-40 (1978)
- 78A266 Kinetics and mechanism of the interaction of sterically hindered phenols with singlet oxygen. Snyakin, A.P.; Samsonova, L.V.; Shlyapintokh, V.Ya.; Ershov, V.V., *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **27**: 46-52 (1978) Translated from: *Izv. Akad. Nauk SSSR, Ser. Khim.* : 55-61 (1978)
- 78A275 Photoperoxidation of anthracene sulphonates. Gupta, A.K.; Singh, B.P.; Rohatgi-Mukherjee, K.K., *J. Indian Chem. Soc.* **55**: 382-5 (1978)
- 78A278 On the reactivity of singlet oxygen in aqueous micellar systems. Barboy, N.; Kraljic, I., *J. Photochem.* **9**: 336-7 (1978)
- 78A338 Photo-oxidation of tryptophan sensitized by methylene blue. Smith, G.J., *J. Chem. Soc., Faraday Trans. 2* **74**: 1350-4 (1978)
- 78A344 Singlet oxygen reactions with model compounds of cis and trans polyisoprene containing two units. Tanielian, C.; Chaineaux, J., *J. Photochem.* **9**: 19-32 (1978)
- 78A357 Singlet oxygen reactions with 2,7-dimethyl-2,6-octadiene. Tanielian, C.; Chaineaux, J., *Photochem. Photobiol.* **28**: 487-92 (1978)
- 78A360 Determination of singlet oxygen rate constants in aqueous solutions. Kraljic, I.; Sharpatyi, V.A., *Photochem. Photobiol.* **28**: 583-6 (1978)
- 78E036 The fate of oxygen in the quenching of excited singlet states. Merkel, P.B.; Herkstroeter, W.G., *Chem. Phys. Lett.* **53**: 350-4 (1978)
- 78E143 The lifetime of singlet oxygen ($^1\Delta_g$) in heavy water, a revised value. Matheson, I.B.C.; Lee, J.; King, A.D., *Chem. Phys. Lett.* **55**: 49-51 (1978)
- 78E144 Lifetime and reactivity of singlet oxygen in an aqueous micellar system: A pulsed nitrogen laser study. Gorman, A.A.; Rodgers, M.A.J., *Chem. Phys. Lett.* **55**: 52-4 (1978)
- 78E238 Deactivation of singlet oxygen and triplet pentacene by transition metal complexes. Furue, H.; Russell, K.E., *Can. J. Chem.* **56**: 1595-601 (1978)
- 78E263 A pulse radiolysis study of the triplet sensitized production of singlet oxygen: determination of energy transfer efficiencies. Gorman, A.A.; Lovering, G.; Rodgers, M.A.J., *J. Am. Chem. Soc.* **100**: 4527-32 (1978)
- 78E881 Luminescence of singlet oxygen during energy transfer from photoexcited pigments in solution. Krasnovsky, A.A., Jr., *Izv. Akad. Nauk SSSR, Ser. Fiz.* **42**: 343-8 (1978) Translated from: *Bull. Acad. Sci. USSR, Phys. Ser.* **42**: 96-100 (1978)
- 78E892 Quenching of luminescence of singlet oxygen by porphyrins and metalloporphyrins. Krasnovsky, A.A., Jr.; Venediktov, E.A., *Biophysics* **23**: 294-6 (1978) Translated from: *Biofizika* **23**: 387-9 (1978)
- 78F020 On the reaction of singlet molecular oxygen with N-allylthiourea in aqueous solution. Kraljic, I.; Kramer, H.E.A., *Photochem. Photobiol.* **27**: 9-12 (1978)
- 78F061 Kinetic studies of photosensitized oxygenation by singlet oxygen in aqueous micellar solutions. Usui, Y.; Tsukada, M.; Nakamura, H., *Bull. Chem. Soc. Jpn.* **51**: 379-84 (1978)
- 78F149 Reaction of singlet oxygen with 2-methylnorbornadiene and 2-methylidenenorbornene. Evaluation of electronic and steric effects on the course of hydroperoxidation. Jefford, C.W.; Rimbault, C.G., *J. Org. Chem.* **43**: 1908-12 (1978)
- 78F183 Solvent effect on photosensitized oxidation of iodide ion by anthracene sulphonates. Gupta, A.K.; Rohatgi-Mukherjee, K.K., *Photochem. Photobiol.* **27**: 539-43 (1978)
- 78F201 An investigation of retinal as a source of singlet oxygen. Delmelle, M., *Photochem. Photobiol.* **27**: 731-4 (1978)
- 78F276 Electronic energy transfer from singlet molecular oxygen to carotenoids. Wilkinson, F.; Ho, W.-T., *Spectrosc. Lett.* **11**: 455-63 (1978)
- 78F290 Reaction of singlet oxygen with a norbornadienol ether. Intramolecular interception of a zwitterionic peroxide. Jefford, C.W.; Rimbault, C.G., *J. Am. Chem. Soc.* **100**: 6515-6 (1978)
- 78F404 Protection of chlorophyll a by carotenoid from photodynamic decomposition. Koka, P.; Song, P.-S., *Photochem. Photobiol.* **28**: 509-15 (1978)
- 78F430 Petits cycles. XL. Sur le mechanisme de formation des hydroperoxydes allyliques dans la reaction de l'oxygene singulet avec les vinylcyclopropanes. Rousseau, G.; Le Perche, P.; Conia, J.M., *Tetrahedron* **34**: 3483-94 (1978)
- 78F438 Photosensitized oxidation of pseudoazulenes. Timpe, G.I.; Klokova, E.M.; El'tsov, A.V., *J. Org. Chem. USSR* **14**: 625-30 (1978) Translated from: *Zh. Org. Khim.* **14**: 673-9 (1978)
- 78F464 Preference for syn ene additions of $^1\Delta_g$ to trisubstituted, acyclic olefins. Schulte-Elte, K.H.; Muller, B.L.; Rautenstrauch, V., *Helv. Chim. Acta* **61**: 2777-83 (1978)
- 78F474 Oxydation photochimique d'une amine tertiaire: la dregamine. Herlem, D.; Ouannes, C., *Bull. Soc. Chim. Fr.* : I-451-4 (1978)
- 78F497 The quenching of singlet oxygen by complex salts of O,O-diarylphosphorodithioic acids. Edilashvili, I.L.; Ioseliani, K.B.; Dzhambekov, N.F.; Markova, E.I.; Akhmed-zade, D.A., *Russ. J. Phys. Chem.* **52**: 1197-8 (1978) Translated from: *Zh. Fiz. Khim.* **52**: 2079-80 (1978)
- 78F549 Oxygen luminescence and sensitization of photooxidation reactions in solutions. Byteva, I.M.; Gurinovich, G.P.; Izbavitev, S.P., *J. Appl. Spectrosc.* **29**: 870-2 (1978) Translated from: *Zh. Prikl. Spektroskop.* **29**: 156-8 (1978)
- 78F586 Concerning the inhibition of singlet oxygen attack on compounds containing urazole rings. Kretschmer, G.; Paquette, L.A., *Heterocycles* **11**: 359-62 (1978)
- 78F605 Chemiluminescence and the formation of singlet oxygen in the oxidation of certain polyphenols and quinones. Slawinska, D., *Photochem. Photobiol.* **28**: 453-8 (1978)
- 78F700 Production and quenching of singlet oxidation by retinals. Krasnovsky, A.A., Jr.; Kagan, V.E., *Dokl. Biophys.* **241-243**: 147-50 (1978) Translated from: *Dokl. Akad. Nauk SSSR* **242**: 229-32 (1978)

- 79F1701 Singlet oxygen reactivity of bilirubin and related tetrapyrroles. Lightner, D.A.; Park, Y.-T., *Experientia* **34**: 555-7 (1978)
- 79N003 Photosensitized carcinogen degradation and the possible role of singlet oxygen in carcinogen activation. Greenstock, C.L.; Wiebe, R.H., *Photochem. Photobiol.* **28**: 863-7 (1978)
- 79A010 Photoluminescence of singlet oxygen in pigment solutions. Krasnovsky, A.A., Jr., *Photochem. Photobiol.* **29**: 29-36 (1979)
- 79A050 Quenching of singlet oxygen by nickel complexes. Monroe, B.M.; Mrowca, J.J., *J. Phys. Chem.* **83**: 591-5 (1979)
- 79A085 Rates of reaction of singlet oxygen with sulfides. Monroe, B.M., *Photochem. Photobiol.* **29**: 761-4 (1979)
- 79A086 Chemistry of singlet oxygen. XXVIII. Steric and electronic effects on the reactivity of sulfides with singlet oxygen. Kacher, M.L.; Foote, C.S., *Photochem. Photobiol.* **29**: 765-9 (1979)
- 79A106 The entropy-controlled reactivity of singlet oxygen ($^1\Delta_g$) toward furans and indoles in toluene. A variable-temperature study by pulse radiolysis. Gorman, A.A.; Lovering, G.; Rodgers, M.A.J., *J. Am. Chem. Soc.* **101**: 3050-5 (1979)
- 79A111 The absolute value of the reaction rate constant of bilirubin with singlet oxygen in D_2O . Matheson, I.B.C., *Photochem. Photobiol.* **29**: 875-8 (1979)
- 79A112 Chemical reaction rates of amino acids with singlet oxygen. Matheson, I.B.C.; Lee, J., *Photochem. Photobiol.* **29**: 879-81 (1979)
- 79A113 Oxidopyrromethenes as reactive singlet oxygen acceptors. Measurement of their chemical reaction rates by a laser flash photolysis technique. Matheson, I.B.C.; Lightner, D.A., *Photochem. Photobiol.* **29**: 933-5 (1979)
- 79A241 Photochemistry and photooxidation of tetraphenyl-p-dioxin. George, M.V.; Kumar, C.V.; Scaiano, J.C., *J. Phys. Chem.* **83**: 2452-5 (1979)
- 79A386 The question of the role of singlet oxygen in the dye-sensitised photo-oxygenation of α -oxo-carboxylic acids. Davidson, R.S.; Goodwin, D.; Smith, G., *J. Chem. Soc., Chem. Commun.* : 463-4 (1979)
- 79A434 Mechanism of the antioxidant effect of some salts of O,O-diaryldithiophosphoric acids in photochemical reactions. Isoseliani, K.B.; Edilashvili, I.L.; Dzhanibekov, N.F., *Izv. Akad. Nauk Gruz. SSR, Ser. Khim.* **5**: 334-9 (1979)
- 79A457 Stereoreversed electrophilic additions to 3-norcarenes. Insight into the relative steric demands of singlet oxygen in the ene reaction. Paquette, L.A.; Kretschmer, G., *J. Am. Chem. Soc.* **101**: 4655-63 (1979)
- 79E846 Duration of the luminescence of singlet oxygen in solution following pulsed laser excitation. Salokhiddinov, K.I.; Byteva, I.M.; Dzhagarov, B.M., *Opt. Spectrosc. (USSR)* **47**: 487-90 (1979) Translated from: *Opt. Spektrosk.* **47**: 881-6 (1979)
- 79E970 Investigation of the luminescence of oxygen in solutions by pulsed spectroscopy. Byteva, I.M., *J. Appl. Spectrosc.* **31**: 1041-3 (1979) Translated from: *Zh. Prikl. Spektrosk.* **31**: 333-5 (1979)
- 79F051 Differentiation in singlet oxygenation rates of 2,3-diaryl-2-butenes as a function of cis-trans isomerism. Adam, W.; Carballera, N.; Cheng, C.-C.; Sakamishi, K.; Gleiter, R., *J. Org. Chem.* **44**: 851-3 (1979)
- 79F074 Singlet oxygen reactions with model compounds of cis- and trans-polyisoprene containing one unit. Kinetic aspects. Tanielian, C.; Chaineaux, J., *J. Polym. Sci., Polym. Chem. Ed.* **17**: 715-29 (1979)
- 79F104 On the syntheses and singlet oxygen reactivity of oxodipyrromethene models for bilirubin. Lightner, D.A.; Park, Y.-T., *Tetrahedron* **35**: 463-71 (1979)
- 79F119 Effect of strain on singlet oxygen (1O_2) reactions. 2. Photooxidation of methylenecyclopropanes. Frimer, A.A.; Farkash, T.; Sprecher, M., *J. Org. Chem.* **44**: 989-95 (1979)
- 79F137 Photooxygenation of 3,3-dialkylsubstituted allyl alcohols. Occurrence of syn preference in the ene addition of 1O_2 at E/Z-isomeric allyl alcohols. Schulte-Elte, K.H.; Muller, B.L.; Pamingle, H., *Helv. Chim. Acta* **62**: 816-29 (1979)
- 79F148 The photoperoxidation of unsaturated organic molecules. XVIII. Malachite green and leucomalachite green. Stevens, B.; Kaplan, S.J., *Mol. Photochem.* **9**: 205-11 (1979)
- 79F155 Stereochemical dependence of isotope effects in the singlet oxygen-olefin reaction. Grdina, B.; Orfanopoulos, M.; Stephenson, L.M., *J. Am. Chem. Soc.* **101**: 3111-2 (1979)
- 79F278 Photooxidation of azines. Evidence for a free-radical oxidation initiated by singlet oxygen. Landis, M.E.; Madoux, D.C., *J. Am. Chem. Soc.* **101**: 5106-7 (1979)
- 79F314 Formation of singlet molecular oxygen by 8-methoxypsoralen. de Mol, N.J.; Beijersbergen van Henegouwen, G.M.J., *Photochem. Photobiol.* **30**: 331-5 (1979)
- 79F315 Photochemical decomposition of 1,4-benzodiazepines Nitrazepam. Cornelissen, P.J.G.; Beijersbergen van Henegouwen, G.M.J., *Photochem. Photobiol.* **30**: 337-41 (1979)
- 79F412 The role of geminate reaction in the self-sensitized photooxidation of arylazonaphtholate dyes. Merkel, P.B.; Smith, W.F., Jr., *J. Phys. Chem.* **63**: 2834-40 (1979)
- 79F434 Mechanism of indole-singlet oxygen reactions. Interception of zwitterionic intermediates and "ene" reaction. Saito, I.; Matsugo, S.; Matsuura, T., *J. Am. Chem. Soc.* **101**: 7332-8 (1979)
- 79F463 Photosensitization and quenching of singlet oxygen by pigments and lipids of photoreceptor cells of the retina. Krasnovsky, A.A., Jr.; Kagan, V.E., *FEBS Lett.* **108**: 152-4 (1979)
- 79F473 Photooxygenation product of dibenzo[a,j]perylene. Kajiwara, T.; Fujisawa, S.; Ohno, K.; Harada, Y., *Bull. Chem. Soc. Jpn.* **52**: 2771-7 (1979)
- 79F582 Contribution of singlet oxygen to the photofading of indigo. Kuramoto, N.; Kitao, T., *J. Soc. Dyers Colour.* **95**: 257-61 (1979)
- 79F646 Reaction of 1O_2 with strained olefins. 3. Photooxidation of vinylcyclopropanes. Frimer, A.A.; Roth, D., *J. Org. Chem.* **44**: 3882-7 (1979)
- 79F816 The mechanism of the riboflavin sensitized photodestruction of mitomycin C. Ueda, K.; Mizunoya, T., *J. Biochem. (Tokyo)* **86**: 857-62 (1979)
- 79F824 Mechanism of quenching of singlet molecular oxygen by porphyrins and their metal complexes. Venediktov, E.A.; Krasnovsky, A.A., Jr., *Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol.* **22**: 395-8 (1979)
- 79N020 Quenching of singlet oxygen by sodium azide in reversed micellar systems. Miyoshi, N.; Tomita, G., *Z. Naturforsch., Teil B* **34B**: 339-43 (1979)
- 79N041 Laser photolysis studies of singlet molecular oxygen in aqueous micellar dispersions. Lindig, B.A.; Rodgers, M.A.J., *J. Phys. Chem.* **83**: 1683-8 (1979)
- 80A041 Rate constants for the reaction of singlet oxygen with derivatives of p-aminobenzoic acid. Monroe, B.M., *Biochem. Biophys. Res. Commun.* **93**: 321-5 (1980)
- 80A143 Kinetic and spectroscopic features of some carotenoid triplet states: Sensitization by singlet oxygen. Rodgers, M.A.J.; Bates, A.L., *Photochem. Photobiol.* **31**: 533-7 (1980)
- 80A205 Determination of the lifetime of singlet oxygen in D_2O using 9,10-anthracenedipropionic acid, a water-soluble probe. Lindig, B.A.; Rodgers, M.A.J.; Schaap, A.P., *J. Am. Chem. Soc.* **102**: 5590-3 (1980)
- 80A339 Singlet oxygen and spin trapping with nitrones. Harbour, J.R.; Issler, S.L.; Hair, M.L., *J. Am. Chem. Soc.* **102**: 7778-9 (1980)
- 80A398 Quenching of oxygen luminescence by electron acceptors and donors. Byteva, I.M.; Salokhiddinov, K.I., *Opt. Spectrosc. (USSR)* **49**: 387-90 (1980) Translated from: *Opt. Spektrosk.* **49**: 707-13 (1980)
- 80A430 Sensitized photo-oxidation of polyisoprene: Effect of the presence of hydroxy and hydroperoxy groups. Tanielian, C.; Chaineaux, J., *Eur. Polym. J.* **16**: 619-22 (1980)
- 80C002 Photocalorimetric studies of singlet oxygen reactions. Olmsted, J., III, *J. Am. Chem. Soc.* **102**: 66-71 (1980)

- 80D076 On the production of nitroxide radicals by singlet oxygen reaction: an EPR study. Lion, Y.; Gandin, E.; Van de Vorst, A., *Photochem. Photobiol.* **31**: 305-9 (1980)
- 80E446 The role of anthraquinonoid dyes in the 'catalytic fading' of dye mixtures. Substituent-dependent triplet yield of diaminoanthraquinones. Rembold, M.W.; Kramer, H.E.A., *J. Soc. Dyers Colour.* **96**: 122-6 (1980)
- 80E548 Quenching of the luminescence of singlet molecular oxygen with complexes of porphyrins with highly charged metal ions. Venediktov, E.A.; Krasnovsky, A.A., Jr., *Biophysics* **25**: 347-8 (1980) Translated from: *Biofizika* **25**: 336-7 (1980)
- 80E549 Lifetime of singlet oxygen in different solvents. Byteva, I.M.; Salokhiddinov, K.I., *Biofizika* **25**: 358 (1980)
- 80E558 Photosensitized luminescence of singlet oxygen in solutions at 1588 nm. Salokhiddinov, K.I.; Dzhagarov, B.M.; Byteva, I.M.; Gurinovich, G.P., *Chem. Phys. Lett.* **76**: 85-7 (1980)
- 80E606 Direct observation of singlet oxygen by a thermal lensing technique. Fuke, K.; Ueda, M.; Itoh, M., *Chem. Phys. Lett.* **76**: 372-4 (1980)
- 80E616 Laser induced fluorescence and relaxation of singlet molecular oxygen in the liquid phase. Protz, R.; Maier, M., *J. Chem. Phys.* **73**: 5464-7 (1980)
- 80E761 Yield of singlet oxygen in the quenching of triplet states of aromatic compounds by molecular oxygen. Ivanov, V.B.; Kuprashvili, B.G.; Edilashvili, I.L., *Khim. Vys. Energ.* **14**: 280-2 (1980)
- 80F007 Mechanisms of photooxygenation. 1. Substituent effects on the [2 + 2] cycloaddition of singlet oxygen to vinyl ethers. Zaklika, K.A.; Kaskar, B.; Schaap, A.P., *J. Am. Chem. Soc.* **102**: 386-9 (1980)
- 80F020 Photoinduced reactions. 104. Photooxygenation of magnesium meso-tetraphenylporphyrin. Matsuura, T.; Inoue, K.; Ranade, A.C.; Saito, I., *Photochem. Photobiol.* **31**: 23-6 (1980)
- 80F104 Nature of peroxy intermediates produced by the photosensitized oxygenation of diethyl sulfide and phenylbenzoyl diazomethane: Kinetic studies by co-oxidation with 4-mono substituted diphenyl sulfides. Ando, W.; Kabe, Y.; Miyazaki, H., *Photochem. Photobiol.* **31**: 191-4 (1980)
- 80F111 Reaction of singlet oxygen with α,β -unsaturated ketones and lactones. Ensley, H.E.; Carr, R.V.C.; Martin, R.S.; Pierce, T.E., *J. Am. Chem. Soc.* **102**: 2836-8 (1980)
- 80F151 Ein neues photochromes System von ungewöhnlich hoher thermischer Stabilität. Brauer, H.-D.; Drews, W.; Schmidt, R., *J. Photochem.* **12**: 293-304 (1980)
- 80F169 β -Carotene inhibited photooxidation of polystyrene. 1. Effect of β -carotene on photochemical reactivity of anthracene in hexane solution. Nowakowska, M., *Makromol. Chem.* **181**: 1013-20 (1980)
- 80F204 The photooxidation of mono- and dimethylnaphthalenes by singlet oxygen. van den Heuvel, C.J.M.; Steinberg, H.; de Boer, Th.J., *Recl. Trav. Chim. Pays-Bas* **99**: 109-13 (1980)
- 80F427 Reactions of singlet oxygen. Part V. Dye sensitized photooxidation of m-hydroxydiphenylamine. Ram, N.; Sidhu, K.S., *Can. J. Chem.* **58**: 2073-9 (1980)
- 80F548 Singlet oxygen photo-oxidation of some triazapentalenes. Albini, A.; Bettinetti, G.F.; Minoli, G.; Pietra, S., *J. Chem. Soc., Perkin Trans. I* : 2904-8 (1980)
- 80F634 Mechanism of the photofading of dye. Contribution of singlet oxygen in the catalytic fading of anthraquinone dye mixtures. Kuramoto, N.; Kitao, T., *J. Chem. Technol. Biotechnol.* **30**: 129-35 (1980)
- 80F718 Photochemical processes of 2,6-di-tert-butyl-4-methylphenol. Nowakowska, M.; Kowal, J., *Eur. Polym. J.* **16**: 615-8 (1980)
- 80M378 Controlled generation of singlet oxygen at low temperatures from triphenyl phosphite ozonide. Bartlett, P.D.; Mendenhall, G.D.; Durham, D.L., *J. Org. Chem.* **45**: 4269-71 (1980)
- 80N018 Photophysical and photosensitized generation of singlet molecular oxygen ($^1\Delta_g$) in micellar solutions at elevated pressures. Measurement of singlet molecular oxygen solvent to micelle transfer rates via both molecular diffusion and energy transfer. Matheson, I.B.C.; Massoudi, R., *J. Am. Chem. Soc.* **102**: 1942-8 (1980)
- 80N021 Fluorescein-photosensitized furan oxidation in methanolic and reversed micellar solutions. Part II. Kinetic analysis. Miyoshi, N.; Tomita, G., *Z. Naturforsch., Teil B* **35B**: 107-11 (1980)
- 80R072 On the feasibility of electron transfer to singlet oxygen from mitochondrial components. Peters, G.; Rodgers, M.A.J., *Biochem. Biophys. Res. Commun.* **96**: 770-6 (1980)
- 80R116 Retinal-sensitized photochemiluminescence of trypsin solutions and the effect of 3-hydroxypyridine on it. Shiryaev, V.M.; Fedorovich, I.B.; Sapezhinskii, I.I.; Khristophorov, V.L.; Kuz'min, V.I.; Smirnov, L.D., *Biophysics* **25**: 454-61 (1981) Translated from: *Biofizika* **25**: 439-45 (1980)
- 81A191 Single-electron transfer from NADH analogues to singlet oxygen. Peters, G.; Rodgers, M.A.J., *Biochim. Biophys. Acta* **637**: 43-52 (1981)
- 81A287 Time-resolved determinations of deuterium isotope effects on $O_2(^1\Delta_g)$ lifetimes in solution. Peters, G.; Rodgers, M.A.J., *J. Am. Chem. Soc.* **103**: 6759-61 (1981)
- 81A327 1,4-Dihydropyridine derivatives as deactivators of singlet oxygen. Tirzitis, G.; Byteva, I.M.; Salokhiddinov, K.I.; Gurinovich, G.P.; Dubure, G., *Chem. Heterocycl. Compd.* : 682-4 (1981) Translated from: *Khim. Geterotsikl. Soedin.* : 924-6 (1981)
- 81A363 Rates of reaction of singlet oxygen with some systemic pyrimidine fungicides and related compounds. Harkness, G.J.; Wells, C.H.J., *Pestic. Sci.* **12**: 215-8 (1981)
- 81A403 Photooxidation of arylazonaphthols with singlet oxygen. Becker, H.G.O.; Franze, J., *J. Prakt. Chem.* **323**: 957-64 (1981)
- 81A422 Oxidation of NADH by singlet oxygen generated by triplet flavin. Vekshin, N.L.; Mironov, G.P., *Biophysics* **26**: 974-81 (1981) Translated from: *Biofizika* **26**: 953-9 (1981)
- 81A437 Electronic control of stereoselectivity. 8. The stereochemical course of electrophilic additions to aryl-substituted 9-isopropylidenebenzonorbornenes. Paquette, L.A.; Hertel, L.W.; Gleiter, R.; Boehm, M.C.; Beno, M.A.; Christoph, G.G., *J. Am. Chem. Soc.* **103**: 7106-21 (1981)
- 81E003 Reactivity of singlet oxygen toward strained substrates and a novel non-photochemical method for determination of quenching constants of singlet oxygen. Turro, N.J.; Chow, M.-F.; Kanfer, S.; Jacobs, M., *Tetrahedron Lett.* **22**: 3-6 (1981)
- 81E398 Lifetime of singlet oxygen in different solvents. Salokhiddinov, K.I.; Byteva, I.M.; Gurinovich, G.P., *J. Appl. Spectrosc.* **34**: 561-4 (1981) Translated from: *Zh. Prikl. Spektrosk.* **34**: 892-7 (1981)
- 81E472 Study of the sensitized luminescence of singlet oxygen in solutions in the 1588 nm region. Salokhiddinov, K.I.; Byteva, I.M.; Dzhagarov, B.M.; Gurinovich, G.P., *J. Appl. Spectrosc.* **35**: 888-91 (1981) Translated from: *Zh. Prikl. Spektrosk.* **35**: 280-4 (1981)
- 81E631 Determination of radiative lifetime of the $^1\Delta_g$ state of oxygen in solutions. Krasnovsky, A.A., Jr., *Sov. Phys. Dokl.* **26**: 425-6 (1981) Translated from: *Dokl. Akad. Nauk SSSR* **257**: 1360-2 (1981)
- 81E704 Energy transfer processes accompanying laser excitation of hematoporphyrin in various solvents. Parker, J.G.; Stanbro, W.D., *Johns Hopkins AP Tech. Dig.* **2**: 196-9 (1981)
- 81F018 Syn regioselectivity of the hydroperoxidation of cyclo-alkenes with singlet oxygen. Jefford, C.W.; Rimbault, C.G., *Tetrahedron Lett.* **22**: 91-4 (1981)
- 81F043 Kinetic studies of self-sensitized photooxygenation in H_2O and D_2O of a water-soluble rubrene derivative. Aubry, J.M.; Rigaudy, J.; Chuong, N.K., *Photochem. Photobiol.* **33**: 155-8 (1981)
- 81F311 Nucleophilic oxygen atom transfer reactions by persulfoxide and persulfone. Sawaki, Y.; Ogata, Y., *J. Am. Chem. Soc.* **103**: 5947-8 (1981)
- 81F321 Quenching of singlet oxygen by sensitizer in dye-sensitized photooxygenation. Tanielian, C.; Golder, L., *Photochem. Photobiol.* **34**: 411-4 (1981)

- *1F364 Photoperoxidation of unsaturated organic molecules. 21. Sensitizer yields of $O_2^{1\Delta_g}$. Stevens, B.; Marsh, K.L.; Barltrop, J.A., *J. Phys. Chem.* **85**: 3079-82 (1981)
- *1F445 Rate constants for the reaction of singlet oxygen with conjugated dienes. Monroe, B.M., *J. Am. Chem. Soc.* **103**: 7253-6 (1981)
- *1F534 The mechanism of ene-reactions with singlet oxygen, $^1O_2(^1\Delta_g)$: Dependence of rates on solvent polarity and structure. Gollnick, K.; Hartmann, H.; Paur, H., *Oxygen and Oxy-Radicals in Chemistry and Biology*, M.A.J. Rodgers and E.L. Powers (eds.), Academic Press, New York, NY, 1981, p.379-95
- *1F582 Solvent effect in reaction of singlet oxygen with conjugated dienes. Matsumoto, M.; Kuroda, K., *Synth. Commun.* **11**: 987-92 (1981)
- *1F609 Formation of singlet oxygen by photoexcitation of substituted aromatic azo compounds in oxygenated solution. Gruen, H.; Steffen, H.; Schulte-Frohlinde, D., *J. Soc. Dyers Colour.* **97**: 430-5 (1981)
- *1F622 Intermediates in the reactions of oxazoles with singlet oxygen. Wasserman, H.H.; Pickett, J.E.; Vinnick, F.S., *Heterocycles* **15**: 1069-73 (1981)
- *1N048 Rate parameters for the quenching of singlet oxygen by water-soluble and lipid-soluble substrates in aqueous and micellar systems. Lindig, B.A.; Rodgers, M.A.J., *Photochem. Photobiol.* **33**: 627-34 (1981)
- *1R011 On the acridine and thiazine dye sensitized photodynamic inactivation of lysozyme-singlet oxygen self-quenching by the sensitizers. Schmidt, H.; Al-Ibrahim, A.; Dietzel, U.; Bieker, L., *Photochem. Photobiol.* **33**: 127-30 (1981)
- *1R052 Relation between some photobiological properties of furcoumarins and their extent of singlet oxygen production. de Mol, N.J.; Beijersbergen van Henegouwen, G.M.J., *Photochem. Photobiol.* **33**: 815-9 (1981)
- *1Z251 Rate constants for the decay and reactions of the lowest electronically excited singlet state of molecular oxygen in solution. Wilkinson, F.; Brummer, J.G., *J. Phys. Chem. Ref. Data* **10**: 809-999 (1981)
- *2A059 Steric aspects of the oxidation of thioketones by singlet oxygen. Ramnath, N.; Jayathertha Rao, V.; Ramesh, V.; Ramamurthy, V., *Chem. Lett.* : 89-92 (1982)
- *2A080 Chemistry of singlet oxygen. 37. One-electron oxidation of tetramethylphenylenediamine by singlet oxygen. Manring, L.E.; Foote, C.S., *J. Phys. Chem.* **86**: 1257-9 (1982)
- *2A093 Rates of oxidation of thioketones by singlet oxygen. Ramesh, V.; Ramnath, N.; Jayathertha Rao, V.; Ramamurthy, V., *J. Photochem.* **18**: 109-15 (1982)
- *2A140 Lifetime of singlet oxygen and quenching by NaN_3 in mixed solvents. Miyoshi, N.; Ueda, M.; Fuke, K.; Tanimoto, Y.; Itoh, M.; Tomita, G., *Z. Naturforsch., Teil B* **37B**: 649-52 (1982)
- *2A179 Photoperoxidation of unsaturated organic molecules. 22. Tetraphenylcyclopentadienone. Stevens, B.; Marsh, K.L., *J. Phys. Chem.* **86**: 2553-6 (1982)
- *2A204 Crocetin, a water soluble carotenoid monitor for singlet molecular oxygen. Matheson, I.B.C.; Rodgers, M.A.J., *Photochem. Photobiol.* **36**: 1-4 (1982)
- *2A291 Mechanism of dye-sensitized photooxidation of tryptophan, tryptamine, and their derivatives. Singlet oxygen process in competition with type I process. Inoue, K.; Matsuura, T.; Saito, I., *Bull. Chem. Soc. Jpn.* **55**: 2959-64 (1982)
- *2A322 Photoperoxidation of unsaturated organic molecules. 24. Mixed solvent quenching of $O_2(^1\Delta_g)$ phosphorescence. Stevens, B.; Marsh, K.L., *J. Phys. Chem.* **86**: 4473-6 (1982)
- *2A349 Time-resolved study of the solvent and temperature dependence of singlet oxygen($^1\Delta_g$) reactivity toward enol ethers: Reactivity parameters typical of rapid reversible exciplex formation. Gorman, A.A.; Gould, I.R.; Hamblett, I., *J. Am. Chem. Soc.* **104**: 7098-104 (1982)
- *2A384 Generation and quenching of the luminescence of singlet molecular oxygen by anthraquinone dyes. Venediktov, E.A.; Krasnovsky, A.A., Jr., *Opt. Spectrosc. (USSR)* **52**: 338-9 (1982) Translated from: *Opt. Spektrosk.* **52**: 567-9 (1982)
- 82A412 The quenching of the sensitised luminescence of molecular oxygen by transition metal complexes of different structures. Byteva, I.M.; Salokhiddinov, K.I.; Bondarev, S.L., *Russ. J. Phys. Chem.* **56**: 1056-9 (1982) Translated from: *Zh. Fiz. Khim.* **56**: 1743-7 (1982)
- 82A421 Quenching of singlet oxygen with chlorophylls and porphyrins. Krasnovsky, A.A., Jr.; Venediktov, E.A.; Cherenko, O.M., *Biophysics* **27**: 1009-16 (1982) Translated from: *Biofizika* **27**: 966-72 (1982)
- 82A464 Kinetics of dye-sensitized photodegradation of aqueous phenol. Okamoto, K.; Hondo, F.; Itaya, A.; Kusabayashi, S., *J. Chem. Eng. Jpn.* **15**: 368-75 (1982)
- 82E072 Laser photolysis study of the mechanism of rubrene quenching by molecular oxygen. Darmanyan, A.P., *Chem. Phys. Lett.* **86**: 405-10 (1982)
- 82E104 Lifetime of singlet oxygen in solution directly determined by laser spectroscopy. Hurst, J.R.; McDonald, J.D.; Schuster, G.B., *J. Am. Chem. Soc.* **104**: 2065-7 (1982)
- 82E105 Optical determination of the collisional lifetime of singlet molecular oxygen [$O_2(^1\Delta_g)$] in acetone and deuterated acetone. Parker, J.G.; Stanbro, W.D., *J. Am. Chem. Soc.* **104**: 2067-9 (1982)
- 82E106 Chemistry of singlet oxygen. 36. Singlet molecular oxygen ($^1\Delta_g$) luminescence in solution following pulsed laser excitation. Solvent deuterium isotope effects on the lifetime of singlet oxygen. Ogilby, P.R.; Foote, C.S., *J. Am. Chem. Soc.* **104**: 2069-70 (1982)
- 82E329 Solvent and deuterium isotope effects on the lifetime of singlet oxygen determined by direct emission spectroscopy at 1.27 μm . Jenny, T.A.; Turro, N.J., *Tetrahedron Lett.* **23**: 2923-6 (1982)
- 82E575 Non-exponential decay of the fluorescence from the $^1\Delta_g$ state in liquid oxygen at high excitation intensities. Klingshirn, H.; Faltermeier, B.; Hengl, W.; Maier, M., *Chem. Phys. Lett.* **93**: 485-9 (1982)
- 82F018 Mechanisms for the photooxidation of protoporphyrin IX in solution. Cox, G.S.; Whitten, D.G., *J. Am. Chem. Soc.* **104**: 516-21 (1982)
- 82F068 Wavelength dependent and adiabatic photochemistry of endoperoxides of aromatic hydrocarbons. Schmidt, R.; Drews, W.; Brauer, H.-D., *Z. Naturforsch., Teil A* **37A**: 55-7 (1982)
- 82F102 The effect of solvent on competing hydroperoxide and dioxetane formation on photo-sensitized oxygenation of olefins. Jefford, C.W.; Kohmoto, S., *Helv. Chim. Acta* **65**: 133-6 (1982)
- 82F357 Chemistry of singlet oxygen. 38. Temperature effect on the photooxidation of sulfides. Gu, C.-L.; Foote, C.S., *J. Am. Chem. Soc.* **104**: 6060-3 (1982)
- 82F379 A new water-soluble singlet oxygen probe. Giraud, M.; Valla, A.; Bazin, M.; Santus, R.; Momzikoff, A., *J. Chem. Soc., Chem. Commun.* : 1147-8 (1982)
- 82F446 Wavelength-dependent photostable or photoreversible photochromic system. Schmidt, R.; Drews, W.; Brauer, H.-D., *J. Phys. Chem.* **86**: 4909-13 (1982)
- 82F450 Effects of remote substitution on the photo-oxidation of exocyclic s-cis-butadienes grafted onto bicyclo[2.2.1]heptanes. Thermal and rhodium catalyzed rearrangements of 3,6-dihydro-1,2-dioxines. Hagenbuch, J.-P.; Birbaum, J.-L.; Metral, J.-L.; Vogel, P., *Helv. Chim. Acta* **65**: 887-95 (1982)
- 82F525 Photoreaction of crystal violet in solution. Nakamura, R.; Hida, M., *Sen'i Gakkaishi* **38**: T183-T190 (1982)
- 82F531 Physicochemical study of a series of cyanines. Part V. Photodegradation mechanism. Lougnot, D.J.; Lepaja, S.; Fouassier, J.P., *J. Chim. Phys. Phys.-Chim. Biol.* **79**: 617-28 (1982)
- 82N021 Oxygen transfer rates across water-micelle interfaces derived from measurements of Ni^{2+} quenching of singlet molecular oxygen in aerosol-OT-heptane reverse micelles. Matheson, I.B.C.; Rodgers, M.A.J., *J. Phys. Chem.* **86**: 884-7 (1982)
- 82N027 A laser flash kinetic spectrophotometric examination of the dynamics of singlet oxygen in unilamellar vesicles. Rodgers, M.A.J.; Bates, A.L., *Photochem. Photobiol.* **35**: 473-7 (1982)

- 82N064 Micellar photochemistry. Photooxidations with intramicellar-generated singlet oxygen. Hovey, M.C., *J. Am. Chem. Soc.* **104**: 4196-202 (1982)
- 83A006 Studies on the interactions of transient species formed in liquid phase oxidation of hydrocarbons by laser flash photolysis. I. Interactions of singlet oxygen with ethylbenzene and with products of the oxidation of ethylbenzene. Darmanyan, A.P.; Vidoczy, T.; Irinyi, G.; Gal, D., *Oxid. Commun.* **3**: 15-24 (1983)
- 83A026 Reaction of singlet oxygen with enol esters: A new path implicating a dipolar intermediate. Wilson, S.L.; Schuster, G.B., *J. Am. Chem. Soc.* **105**: 679-81 (1983)
- 83A050 Thermal lensing study of singlet oxygen reactions. Fuke, K.; Ueda, M.; Itoh, M., *J. Am. Chem. Soc.* **105**: 1091-6 (1983)
- 83A063 Quenching of singlet oxygen by 1,3,5-triaryl-2-pyrazolines. Ando, W.; Sato, R.; Yamashita, M.; Akasaka, T.; Miyazaki, H., *J. Org. Chem.* **48**: 542-6 (1983)
- 83A078 Photosensitized oxygenation of oligoisoprenes. Tanielian, C.; Chainiaux, J., *J. Photochem.* **21**: 265-73 (1983)
- 83A094 Formation of superoxide ion via one-electron transfer from electron donors to singlet oxygen. Saito, I.; Matsuura, T.; Inoue, K., *J. Am. Chem. Soc.* **105**: 3200-6 (1983)
- 83A205 L Ascorbic acid quenching of singlet delta molecular oxygen in aqueous media: Generalized antioxidant property of vitamin C. Chou, P.-T.; Khan, A.U., *Biochem. Biophys. Res. Commun.* **115**: 932-7 (1983)
- 83A223 Solvent-induced deactivation of singlet oxygen: Additivity relationships in nonaromatic solvents. Rodgers, M.A.J., *J. Am. Chem. Soc.* **105**: 6201-5 (1983)
- 83A315 Dye-sensitized photooxidation of 2-dimethylamino-5,6-dimethylpyrimidin-4-ol in aqueous solution. Dixon, S.R.; Wells, C.H.J., *Pestic. Sci.* **14**: 444-8 (1983)
- 83A317 A kinetic study of the interaction between bilirubin and thermally produced singlet oxygen. Galliani, G.; Manitto, P.; Monti, D., *Isr. J. Chem.* **23**: 219-22 (1983)
- 83A336 Interaction of singlet oxygen with carotenoids: rate constants of physical and chemical quenching. Krasnovsky, A.A., Jr.; Paramonova, L.I., *Biophysics* **28**: 769-74 (1983) Translated from: *Biofizika* **28**: 725-9 (1983)
- 83A371 Influence of medium on lifetime and reactivity of $^1\text{O}_2$ with 1,3-diphenylisobenzofuran, and the role of $^1\text{O}_2$ in liquid-phase oxidation of ethylbenzene. Darmanyan, A.P.; Vidoczy, T.; Irinii, G.; Gal, D., *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **32**: 2423-9 (1983) Translated from: *Izv. Akad. Nauk SSSR, Ser. Khim.* : 2702-8 (1983)
- 83A394 Sensitized photooxygenation of cis 9-tricosene (housefly sex attractant). Tanielian, C.; Mechlin, R.; Esch, M., *Oxygen Radicals Chem. Biol.*, W. Bors, M. Saran and D. Tait (eds.), de Gruyter, Berlin, Fed. Rep. Germany, Pub. 1984, p.527-30
- 83A395 On the reactivity of singlet oxygen with hemoglobin. Kraljic, I.; Pin, S.; El Mohsni, S.; Alpert, B., *Oxygen Radicals Chem. Biol.*, W. Bors, M. Saran and D. Tait (eds.), de Gruyter, Berlin, Fed. Rep. Germany, Pub. 1984, p.531-4
- 83A397 Dye-sensitized photooxidation of tryptophan. Tanielian, C.; Muller, H.; Golder, L., *Oxygen Radicals Chem. Biol.*, W. Bors, M. Saran and D. Tait (eds.), de Gruyter, Berlin, Fed. Rep. Germany, Pub. 1984, p.551-4
- 83E235 Chemistry of singlet oxygen. 42. Effect of solvent, solvent isotopic substitution, and temperature on the lifetime of singlet molecular oxygen ($^1\Delta_g$). Ogilby, P.R.; Foote, C.S., *J. Am. Chem. Soc.* **105**: 3423-30 (1983)
- 83E398 Nonradiative relaxation of singlet oxygen in solution. Hurst, J.R.; Schuster, G.B., *J. Am. Chem. Soc.* **105**: 5756-60 (1983)
- 83E425 Mechanism of quenching of singlet oxygen luminescence by saturated and unsaturated aliphatic acids. Krasnovsky, A.A., Jr.; Minin, A.A.; Kagan, V.E., *Dokl. Biophys.* **268**: 59-61 (1983) Translated from: *Dokl. Akad. Nauk SSSR* **268**: 1488-91 (1983)
- 83E667 Flash photolysis studies of hemato- and copro-porphyrins in homogeneous and microheterogeneous aqueous dispersions. Reddi, E.; Jori, G.; Rodgers, M.A.J.; Spikes, J.D., *Photochem. Photobiol.* **38**: 639-45 (1983)
- 83E756 Direct measurement of the lifetime of molecular oxygen in the singlet $^1\Delta_g$ state, generated in water by a porphyrin sensitized. Salokhiddinov, K.I.; Dzhagarov, B.M.; Egorova, G.D., *Opt. Spectrosc. (USSR)* **55**: 40-1 (1983) Translated from: *Opt. Spektrosk.* **55**: 71-3 (1983)
- 83E813 Photogeneration of singlet oxygen by psoralens. Krasnovsky, A.A.; Sukhorukov, V.L.; Potapenko, A.Ya., *Bull. Exp. Biol. Med.* **96**: 1259-61 (1983) Translated from: *Byull. Eksp. Biol. Med.* **96**: 59-61 (1983)
- 83E818 Effect of solvent deuteration and of an external heavy atom on the luminescence quenching time of molecular oxygen. Salokhiddinov, K.I.; Bytova, I.M., *Opt. Spectrosc. (USSR)* **55**: 282-4 (1983) Translated from: *Opt. Spektrosk.* **55**: 479-82 (1983)
- 83E844 Pulsed laser excitation of photosensitized oxygen luminescence: Decay kinetics in aqueous solutions. Egorov, S.Yu.; Krasnovsky, A.A., *Biophysics* **28**: 532-4 (1983) Translated from: *Biofizika* **28**: 497-8 (1983)
- 83F028 Photochemical oxidation of thioketones: Steric and electronic aspects. Ramnath, N.; Ramesh, V.; Ramamurthy, V., *J. Org. Chem.* **48**: 214-22 (1983)
- 83F075 Solvent effect on $^1\text{O}_2$ yield and the mechanism of polycyclic hydrocarbon triplet-state quenching by oxygen. Darmanyan, A.P., *Chem. Phys. Lett.* **96**: 383-9 (1983)
- 83F136 Photooxidation of some triaza- and tetraazabenzenopentalenes. Albini, A.; Bettinetti, G.F.; Minoli, G., *J. Org. Chem.* **48**: 1080-3 (1983)
- 83F151 Quenching of singlet oxygen luminescence by fatty acids and lipids. Contribution of physical and chemical mechanisms. Krasnovsky, A.A., Jr.; Kagan, V.E.; Minin, A.A., *FEBS Lett.* **155**: 233-6 (1983)
- 83F166 Singlet oxygen generation for solution kinetics: Clean and simple. Midden, W.R.; Wang, S.Y., *J. Am. Chem. Soc.* **105**: 4129-35 (1983)
- 83F188 Photosensitizing properties of furocoumarins. V. 8-Methoxysoralen: Quantum yields of reaction in aerated and degassed solution: Role of singlet oxygen. Le, Q.T.; Arnaud, R.; Lemaire, J., *C.R. Seances Acad. Sci., Ser. II* **296**: 687-90 (1983)
- 83F195 Chemistry of singlet oxygen. 43. Quenching by conjugated olefins. Manring, L.E.; Kanner, R.C.; Foote, C.S., *J. Am. Chem. Soc.* **105**: 4707-10 (1983)
- 83F196 Chemistry of singlet oxygen. 44. Mechanism of photooxidation of 2,5-dimethylhexa-2,4-diene and 2-methyl-2-pentene. Manring, L.E.; Foote, C.S., *J. Am. Chem. Soc.* **105**: 4710-7 (1983)
- 83F222 The vitamin-B₁₂-derived Co(III)- complex 'pyrocobester' as photosensitizer and as substrate in reactions involving 'singlet oxygen'. Kraeutler, B.; Stepanek, R., *Helv. Chim. Acta* **66**: 1493-512 (1983)
- 83F256 Reaction of retinol and retinal with singlet oxygen. Smith, G.J., *Photochem. Photobiol.* **38**: 119-21 (1983)
- 83F323 Spectroscopic studies of cutaneous photosensitizing agents. IV. The photolysis of benoxaprofen, an anti inflammatory drug with phototoxic properties. Reszka, K.; Chignell, C.F., *Photochem. Photobiol.* **38**: 281-91 (1983)
- 83F335 Generation of $^1\text{O}_2$ from O_2 quenching of the lowest excited singlet and triplet states of 9,10-diphenylanthracene. Drews, W.; Schmidt, R.; Brauer, H.-D., *Chem. Phys. Lett.* **100**: 466-9 (1983)
- 83F406 Mesodiphenylhelianthrene - the most reactive singlet oxygen acceptor. Acs, A.; Schmidt, R.; Brauer, H.-D., *Photochem. Photobiol.* **38**: 527-31 (1983)
- 83F407 Ascorbic acid as a photooxidation inhibitor. Rooney, M.L., *Photochem. Photobiol.* **38**: 619-21 (1983)
- 83N084 Time resolved studies of 1.27 μm luminescence from singlet oxygen generated in homogeneous and microheterogeneous fluids. Rodgers, M.A.J., *Photochem. Photobiol.* **37**: 99-103 (1983)

- 84F171 Singlet molecular oxygen in micellar systems. 1. Distribution equilibria between hydrophobic and hydrophilic compartments. Lee, P.C.; Rodgers, M.A.J., *J. Phys. Chem.* **87**: 4894-8 (1983)
- 84F198 Hematoporphyrin photosensitization of serum albumin and subtilisin BPN'. Richard, P.; Blum, A.; Grossweiner, L.I., *Photochem. Photobiol.* **37**: 287-91 (1983)
- 84F123 Chemistry of singlet oxygen. 46. Cercosporin, a singlet oxygen generator. Dobrowolski, D.C.; Foote, C.S., *Angew. Chem. Int. Ed. Engl.* **22**: 720-1 (1983)
- 84F170 Oxidation of thioketenes by singlet oxygen. Rao, V.J.; Ramamurthy, V.; Schaumann, E.; Nimmagere, H., *J. Org. Chem.* **49**: 615-21 (1984)
- 84F167 Quenching of singlet oxygen by hindered amine light stabilizers. A flash photolytic study. Ballardini, R.; Beggiato, G.; Bortolus, P.; Faucitano, A.; Buttafava, A.; Gratani, F., *Polym. Degradation Stab.* **7**: 41-53 (1984)
- 84F247 Mechanism of singlet oxygen interaction with spin trap α -phenyl-N-tert-butylnitroline. Darmanyan, A.P.; Moger, G., *J. Photochem.* **26**: 269-74 (1984)
- 84F339 Study of carbonyl oxide formation in the reaction of singlet oxygen with diphenyldiazomethane. Casal, H.L.; Sugarnori, S.E.; Scaino, J.C., *J. Am. Chem. Soc.* **106**: 7623-4 (1984)
- 84F366 Strychnine: A fast physical quencher of singlet oxygen ($^1\Delta_g$). Gorman, A.A.; Hamblett, I.; Smith, K.; Standen, M.C., *Tetrahedron Lett.* **25**: 581-4 (1984)
- 84F289 Relaxation of the $^1\Delta_g$ state in pure liquid oxygen and in liquid mixtures of $^{16}\text{O}_2$ and $^{18}\text{O}_2$. Wild, E.; Klingshirn, H.; Maier, M., *J. Photochem.* **25**: 131-43 (1984)
- 84F291 The mechanism of quenching of singlet oxygen $\text{O}_2^*(^1\Delta_g)$ by molecular iodine. Wilkinson, F.; Farmilo, A., *J. Photochem.* **25**: 153-60 (1984)
- 84F296 Optical determination of the rates of formation and decay of $\text{O}_2(^1\Delta_g)$ in H_2O , D_2O and other solvents. Parker, J.G.; Stanbro, W.D., *J. Photochem.* **25**: 545-7 (1984)
- 84F492 Reversible exciplex formation between singlet oxygen, $^1\Delta_g$, and vitamin E. Solvent and temperature effects. Gorman, A.A.; Gould, I.R.; Hamblett, I.; Standen, M.C., *J. Am. Chem. Soc.* **106**: 6956-9 (1984)
- 84F505 Laser flash photolytic determination of triplet yields via singlet oxygen generation. Chattopadhyay, S.K.; Kumar, C.V.; Das, P.K., *J. Photochem.* **24**: 1-9 (1984)
- 84F1065 Solvent dependence of singlet oxygen / substrate interactions in ene-reactions, (4 + 2)- and (2 + 2)-cycloaddition reactions. Gollnick, K.; Griesbeck, A., *Tetrahedron Lett.* **25**: 725-8 (1984)
- 84F075 Photooxidation of 3,4-dihydroxyphenylalanine by hematoporphyrin in aqueous solutions: An electron spin resonance study using 2,2,6,6-tetramethyl-4-piperidone-1-oxyl (tempone). Reszka, K.; Sealy, R.C., *Photochem. Photobiol.* **39**: 293-9 (1984)
- 84F191 Production and quenching of singlet oxygen by the sensitizer in dye-sensitized photo-oxygenations. Tanielian, C.; Golder, L.; Wolff, C., *J. Photochem.* **25**: 117-25 (1984)
- 84F193 Effects of solvent on the rate of bacteriochlorophyll a photo-oxidation. Marsh, K.L.; Connolly, J.S., *J. Photochem.* **25**: 183-95 (1984)
- 84F194 Aspects of photosensitized lignin oxygenation. Forsskahl, I., *J. Photochem.* **25**: 197-209 (1984)
- 84F196 Singlet oxygen oxidation of substituted thiobenzamides. Murray, R.W.; Agarwal, S.K., *J. Photochem.* **25**: 335-43 (1984)
- 84F197 Generation of $^1\text{O}_2$ from oxygen quenching of the lowest excited singlet and triplet states of some aromatic compounds. Brauer, H.-D.; Acs, A.; Drews, W.; Gabriel, R.; Ghaeni, S.; Schmidt, R., *J. Photochem.* **25**: 475-88 (1984)
- 84F199 Singlet oxygen and free-radical oxidation of polydienes and related problems with stabilization: Synergistic and antagonistic effects. Rabek, J.F.; Ranby, B.; Arct, J.; Liu, R., *J. Photochem.* **25**: 519-36 (1984)
- 84F240 Solvent effect on the quantum yield of the self-sensitized photoperoxidation of 1,3-diphenylisobenzofuran. Nowakowska, M., *J. Chem. Soc., Faraday Trans. 1* **80**: 2119-26 (1984)
- 84F244 Photoreactions of 4-(aminomethylene) derivatives of 3-methyl-1-phenyl-2-pyrazoline-5-thione. Rtishchev, N.I.; El'tsov, A.V.; Kvitsko, I.Ya.; Alam, L.V.; Bocharova, E.G., *J. Gen. Chem., USSR* **54**: 134-42 (1984) Translated from: *Zh. Obshch. Khim.* **54**: 152-61 (1984)
- 84F327 The effect of medium polarity on the hematoporphyrin-sensitized photooxidation of L-tryptophan. Reddi, E.; Rodgers, M.A.J.; Spikes, J.D.; Jori, G., *Photochem. Photobiol.* **40**: 415-21 (1984)
- 84F330 Photosensitization of melanin: An electron spin resonance study of sensitized radical production and oxygen consumption. Sealy, R.C.; Sarna, T.; Wanner, E.J.; Reszka, K., *Photochem. Photobiol.* **40**: 453-9 (1984)
- 84F335 Interactions of singlet oxygen with 2,5-dimethyl-2,4-hexadiene in polar and non-polar solvents. Evidence for a vinylog ene-reaction. Gollnick, K.; Griesbeck, A., *Tetrahedron* **40**: 3235-50 (1984)
- 84F350 Is self-sensitized photo-oxidation a significant pathway in the photofading of azo dyes? Albini, A.; Fasani, E.; Pietra, S.; Sulpizio, A., *J. Chem. Soc., Perkin Trans. 2* : 1689-92 (1984)
- 84F381 Singlet oxygen in surface waters. Part I. Furfuryl alcohol as a trapping agent. Haag, W.K.; Hoigne, J.; Gassman, E.; Braun, A.M., *Chemosphere* **13**: 631-40 (1984)
- 84F474 Effect of substituents, solvent, and temperature on the reactivity of the zwitterionic peroxides arising from the photo-oxygenation of 2-(methoxymethylidene)- and 2-(phenoxyethylidene)adamantane. Jefford, C.W.; Grant, H.G.; Jaggi, D.; Boukouvalas, J.; Kohmoto, S., *Helv. Chim. Acta* **67**: 2210-7 (1984)
- 84F477 Mechanism of endoperoxide formation. 3. Utilization of the Young and Carlsson kinetic techniques. Clennan, E.L.; Mehrsheikh-Mohammadi, M.E., *J. Am. Chem. Soc.* **106**: 7112-8 (1984)
- 84F670 Quenching of singlet oxygen by alkaloids and related nitrogen heterocycles. Larson, R.A.; Marley, K.A., *Phytochemistry* **23**: 2351-4 (1984)
- 84F673 Quenching of singlet oxygen by methylene blue. Tanielian, C.; Mechlin, R., *Proc. 10th IUPAC Symp. Photochem.*, Presses Polytechniques Romandes, Lausanne, Switzerland, 1984, p.471-2
- 84N114 Singlet molecular oxygen in micellar systems. 2. Quenching behavior in AOT reverse micelles. Rodgers, M.A.J.; Lee, P.C., *J. Phys. Chem.* **88**: 3480-4 (1984)
- 84N150 Kinetic properties of singlet oxygen in a polymeric microheterogeneous system. Lee, P.C.; Rodgers, M.A.J., *J. Phys. Chem.* **88**: 4385-9 (1984)
- 84P629 Studies on singlet oxygen reaction with ethylene-propylene-ethylideneborborene terpolymer. Part II. Determination of kinetic parameters of photosensitized oxidation of ethylene-propylene-ethylideneborborene terpolymer. Nowakowska, M., *Polym. Photochem.* **4**: 347-59 (1984)
- 85A124 Interaction of singlet oxygen with α - and β -naphthols. Moger, G.; Darmanyan, A.P., *J. Radioanal. Nucl. Chem.* **94**: 9-15 (1985)
- 85A167 The ene reaction of singlet oxygen: Kinetic and product evidence in support of a perepoxide intermediate. Hurst, J.R.; Wilson, S.L.; Schuster, G.B., *Tetrahedron* **41**: 2191-7 (1985)
- 85A176 Studies on singlet oxygen in aqueous solution. Part 1. Formation of singlet oxygen from hydrogen peroxide with two-electron oxidants. Evans, D.F.; Upton, M.W., *J. Chem. Soc., Dalton Trans.* : 1141-5 (1985)
- 85A177 Studies on singlet oxygen in aqueous solution. Part 2. Water-soluble square-planar nickel complexes as quenchers. Botisivali, M.; Evans, D.F.; Missen, P.H.; Upton, M.W., *J. Chem. Soc., Dalton Trans.* : 1147-9 (1985)
- 85A206 Fluorenone oxide: Transient spectroscopy and kinetics of its formation and reactions. Casal, H.L.; Tanner, M.; Werstiuk, N.H.; Scaino, J.C., *J. Am. Chem. Soc.* **107**: 4616-20 (1985)
- 85A261 The unusual reactions of singlet oxygen with isomeric 1,4-di-tert-butoxy-1,3-butadienes. A 2_s + 2_a cycloaddition. Clennan, E.L.; L'Esperance, R.P., *J. Am. Chem. Soc.* **107**: 5178-82 (1985)

- 85A443 Retinal-sensitized photooxidation of rhodopsin. Starostin, A.V.; Fedorovich, I.B.; Ostrovskii, M.A., *Biophysics* **30**: 1082-8 (1985) Translated from: *Biofizika* **30**: 995-9 (1985)
- 85E190 Quantitative aspects of all-trans-retinol singlet and triplet quenching by oxygen. Bhattacharyya, K.; Das, P.K., *Chem. Phys. Lett.* **116**: 326-32 (1985)
- 85E293 Triplet excitation transfer involving β -ionone. A kinetic study by laser flash photolysis. Chattopadhyay, S.K.; Kumar, C.V.; Das, P.K., *Photochem. Photobiol.* **42**: 17-24 (1985)
- 85E591 Thermal-lensing measurements of singlet molecular oxygen (${}^1\Delta_g$): Quantum yields of formation and lifetimes. Rossbroich, G.; Garcia, N.A.; Braslavsky, S.E., *J. Photochem.* **31**: 37-48 (1985)
- 85F145 Singlet oxygen photooxygenation of furans. Isolation and reactions of (4 + 2)-cycloaddition products (unsaturated sec.-ozonides). Gollnick, K.; Griesbeck, A., *Tetrahedron* **41**: 2057-68 (1985)
- 85F148 Reactions of singlet oxygen with polyisoprene and model compounds. Tanielian, C.; Mechlin, R., *Tetrahedron* **41**: 2139-46 (1985)
- 85F150 Dye-sensitized photo-oxygenation of 1,2-cyclohexanediones. Utaka, M.; Nakatani, M.; Takeda, A., *Tetrahedron* **41**: 2163-7 (1985)
- 85F151 Mechanism of oxidation of α,β -unsaturated thiones by singlet oxygen. Pushkara Rao, V.; Ramamurthy, V., *Tetrahedron* **41**: 2169-76 (1985)
- 85F332 Quantum yield measurements of the hematoporphyrin derivative (Hpd) sensitized singlet oxygen production. Murasecco, P.; Oliveros, E.; Braun, A.M.; Monnier, P., *Photobiochem. Photobiophys.* **9**: 193-201 (1985)
- 85F480 Determination of the reaction rate constants of singlet oxygen with dialkyl sulfides. Xu, D.; Xu, B.; Cui, G.; Wu, S., *Huaxue Tongbao* : 17-9 (1985)
- 85F517 Generation and quenching of singlet molecular oxygen by bacteriochlorophyll and bacteriopheophytin a and b. Krasnovsky, A.A., Jr.; Vychezhannina, I.V.; Drozdova, N.N.; Krasnovsky, A.A., *Dokl. Biophys.* **283**: 161-3 (1985) Translated from: *Dokl. Akad. Nauk SSSR* **283**: 474-7 (1985)
- 85F538 Mechanism of interaction of singlet oxygen with spin trap C-phenyl-N-tert-butylnitro. Darmanyan, A.P.; Moger, G., *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **34**: 1163-6 (1985) Translated from: *Izv. Akad. Nauk SSSR, Ser. Khim.* : 1271-5 (1985)
- 85F646 Influence of the nature of the heteroatom in the H-chelate ring on the photochemical properties of aminomethylene derivatives of S or Se substituted pyrazolin-5-ones. Rtishchev, N.I.; El'tsov, A.V.; Kvitko, I.Ya.; Alam, L.V., *J. Gen. Chem., USSR* **55**: 2095-105 (1985) Translated from: *Zh. Obshch. Khim.* **55**: 2358-69 (1985)
- 85F667 Quenching of singlet oxygen generated in tetraphenylporphyrin solutions by retinoid RO 10-9359, ionol and α -tocopherol. Adam, F.; Schwider, P., *Stud. Biophys.* **109**: 95-100 (1985, Pub. 1986))
- 85F675 The influence of specific solvation on the deactivation of singlet molecular oxygen by 1,4-diaminoanthraquinone. Venedikov, E.A.; Perfil'ev, V.A.; Berezin, B.D., *Russ. J. Phys. Chem.* **59**: 1824-6 (1985) Translated from: *Zh. Fiz. Khim.* **59**: 3048-51 (1985)
- 85M098 Importance of single electron-transfer in singlet oxygen reaction in aqueous solution. Oxidation of electron-rich thioanisoles. Inoue, K.; Matsuuwa, T.; Saito, I., *Tetrahedron* **41**: 2177-81 (1985)
- 85M406 Ozonides de phosphite source d'oxygène singulet: rendement, mécanisme. Caminade, A.M.; El Khatib, F.; Koenig, M.; Aubry, J.M., *Can. J. Chem.* **63**: 3203-9 (1985)
- 85N293 O_2 (${}^3\Sigma$) and O_2 (${}^1\Delta$) processes in microheterogeneous systems. Rubio, M.A.; Araya, L.; Abuin, E.B.; Lissi, E.A., *An. Asoc. Quim. Argent.* **73**: 301-9 (1985)
- 85R008 Singlet oxygen generation by hematoporphyrin IX, uroporphyrin I and hematoporphyrin derivative at 546 nm in phosphate buffer and in the presence of egg phosphatidylcholine liposomes. Blum, A.; Grossweiner, L.I., *Photochem. Photobiol.* **41**: 27-32 (1985)
- 86A099 9,10-Dicyanoanthracene-sensitized photooxygenation of α,α' -dimethylstilbenes. Mechanism and kinetics of the competing singlet oxygen and electron transfer photooxygenation reactions. Gollnick, K.; Schnatterer, A., *Photochem. Photobiol.* **43**: 365-78 (1986)
- 86A159 Additions of singlet oxygen to alkoxy-substituted butadienes. An unexpectedly large s-cis/s-trans ratio in an (E,Z)-diene or a kinetic anomeric effect? Clennan, E.L.; L'Esperance, R.P.; Lewis, K.K., *J. Org. Chem.* **51**: 1440-6 (1986)
- 86A198 Determination of the decay rate constant of singlet oxygen (${}^1\Delta_g$) in presence of biomolecules. Rougee, M.; Bensasson, R.V., *C.R. Séances Acad. Sci., Ser. II* **302**: 1223-6 (1986)
- 86A203 Reactivity of sydrones with singlet oxygen. Bhattacharyya, K.; Das, P.K.; George, M.V., *J. Photochem.* **34**: 13-22 (1986)
- 86A216 Is water the best or the worst solvent for [2 + 4] cycloadditions of singlet oxygen to aromatic compounds? Cazin, B.; Aubry, J.-M.; Rigaudy, J., *J. Chem. Soc., Chem. Commun.* : 952-3 (1986)
- 86A264 Time-resolved O_2 (${}^1\Delta_g$) \rightarrow ${}^3\Sigma_g^-$ chemiluminescence upon UV laser photolysis of an aromatic endoperoxide in aqueous solution. Chou, P.-T.; Khan, S.; Frei, H., *Chem. Phys. Lett.* **129**: 463-7 (1986)
- 86A307 Solvent effect and temperature dependence in the interaction of singlet oxygen with α -phenyl-N-tert-butyl nitrone. Darmanyan, A.P.; Moger, G., *J. Radioanal. Nucl. Chem.* **105**: 303-8 (1986)
- 86A406 Flavin mononucleotide-sensitized photooxidation of indoleacetic acid. Miyoshi, N.; Fukuda, M.; Tomita, G., *Photobiochem. Photobiophys.* **11**: 57-65 (1986)
- 86A438 Quenching of singlet oxygen by 4-[2-(N,N-dimethylhydrazono)ethylidene]-2,6-diphenyl-4H-pyran. Akasaka, T.; Nakagawa, M.; Ando, W., *J. Org. Chem.* **51**: 4477-9 (1986)
- 86A540 Singlet oxygen quenching by $Co(SacSac)_2$ complexes. [Sac = dithioacetlyacetone] Darmanyan, A.P.; Karshalykov, K., *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **35**: 2163-5 (1986) Translated from: *Izv. Akad. Nauk SSSR, Ser. Khim.* : 2361-4 (1986)
- 86E108 Singlet oxygen quenching by stable nitroxyl radicals. Darmanyan, A.P.; Tatikolov, A.S., *J. Photochem.* **32**: 157-63 (1986)
- 86E809 Kinetics of singlet oxygen luminescence quenching in polymers. Influence of polymer matrices on the quenching process of the triplet sensitizer by molecular oxygen. Darmanyan, A.P.; Matveev, M.Yu., *Sov. J. Chem. Phys.* **5**: 2768-86 (1990) Translated from: *Khim. Fiz.* **5**: 1488-95 (1986)
- 86E884 Direct determination of the lifetime of singlet oxygen in polymers with saturated and unsaturated bonds. Matveev, M.Yu.; Darmanyan, A.P., *Dokl. Phys. Chem.* **290**: 894-7 (1986) Translated from: *Dokl. Akad. Nauk SSSR* **290**: 897-901 (1986)
- 86F091 Photosensitized oxygenation reactions of phytol and its derivatives. Mihara, S.; Tateba, H., *J. Org. Chem.* **51**: 1142-4 (1986)
- 86F149 Quenching of singlet molecular oxygen by components of media used for isolating chloroplasts and studying photosynthetic activity. Egorov, S.Yu.; Krasnovsky, A.A., *Sov. Plant Physiol.* **33**: 5-8 (1986) Translated from: *Fiziol. Rast. (Moscow)* **33**: 10-4 (1986)
- 86F219 Reactions of singlet oxygen with enol esters. Wilson, S.L.; Schuster, G.B., *J. Org. Chem.* **51**: 2056-60 (1986)
- 86F225 Reaction of singlet oxygen with 2-pyrazoline: Implication for cation radical - superoxide ion pair intermediate. Akasaka, T.; Nakagawa, M.; Nomura, Y.; Sato, R.; Someno, K.; Ando, W., *Tetrahedron* **42**: 3807-12 (1986)
- 86F226 Quenching of singlet oxygen by catecholamines in dye-sensitized photooxidation reactions. Kruck, I., *Z. Phys. Chem. (Leipzig)* **267**: 569-75 (1986)
- 86F316 Photogeneration of singlet oxygen by porphyrins in aqueous solutions. Egorov, S.Yu.; Krasnovsky, A.A., Jr.; Nazarova, O.V.; Yartsev, E.I., *Biophysics* **31**: 376-8 (1986) Translated from: *Biofizika* **31**: 342-3 (1986)
- 86F337 State-selective photochemistry of singlet oxygen precursors: Kinetics and wavelength dependence of the photodissociation of anthracene endoperoxides. Eisenthal, K.B.; Turro, N.J.; Dupuy, C.G.; Hrovat, D.A.; Langan, J.; Jenny, T.A.; Sitzmann, E.V., *J. Phys. Chem.* **90**: 5168-73 (1986)
- 86F510 Photo-oxygenation des dérivés de l'hydroxyméthyl-5-furfural-2. Cottier, L.; Descotes, G.; Nigay, H.; Parron, J.-C.; Gregoire, V., *Bull. Soc. Chim. Fr.* : 844-50 (1986)

- 86P610 Quenching of singlet molecular oxygen by zinc tetraphenylchlorin in solutions. Venedikov, E.A.; Mozhzhukhin, V.V.; Lipatov, N.G.; Berezin, B.D., *Kinet. Catal.* **27**: 847-50 (1986) Translated from: *Kinet. Katal.* **27**: 976-9 (1986)
- 86P622 Investigation of the role of singlet oxygen in the photodegradation of monoazo dyes. Koval'chuk, N.V.; Anisimov, V.M.; Tambieva, O.A.; Titov, V.V., *J. Org. Chem. USSR* **22**: 1746-53 (1986) Translated from: *Zh. Org. Khim.* **22**: 1944-51 (1986)
- 86P670 Quenching of singlet molecular oxygen by pyrrole in solution. Venedikov, E.A.; Balakina, T.A.; Berezin, B.D., *Kinet. Catal.* **27**: 1274-6 (1986) Translated from: *Kinet. Katal.* **27**: 1465-7 (1986)
- 86P678 Singlet oxygen production by UV and visible photoexcited porphyrins under different states of aggregation. Seret, A.; Gandin, E.; Van de Vorst, A., *Photobiochem. Photobiophys.* **12**: 259-66 (1986)
- 86N104 Kinetics of $O_2(^1\Delta_g)$ photo-oxidation reactions in egg-yolk lecithin vesicles. Dearden, S.J., *J. Chem. Soc., Faraday Trans. I* **82**: 1627-35 (1986)
- 86R129 Efficient singlet oxygen inactivation of firefly luciferase. Thompson, A.; Nigro, J.; Seliger, H.H., *Biochem. Biophys. Res. Commun.* **140**: 888-94 (1986)
- 86R210 Determination of the rate constants of reaction between singlet oxygen and DNA bases. Zhang, Y.; Ma, H.; Dai, J., *Kexue Tongbao (Foreign Lang. Ed.)* **31**: 1065-9 (1986)
- 87A035 Reactions of singlet oxygen with alkoxy-substituted dienes. Formation of dioxetanes in the singlet oxygencations of s-cis fixed dienes (*Z,Z*)- and (*E,Z*)-4,5-diethylidene-2,2-dimethyl-1,3-dioxolanes. Clennan, E.L.; Nagraba, K., *J. Org. Chem.* **52**: 294-6 (1987)
- 87A043 Laser flash photokinetic studies of Rose Bengal sensitized photodynamic interactions of nucleotides and DNA. Lee, P.C.C.; Rodgers, M.A.J., *Photochem. Photobiol.* **45**: 79-86 (1987)
- 87A063 Rate constants for interaction of $^1O_2 (^1\Delta_g)$ with azide ion in water. Haag, W.R.; Mill, T., *Photochem. Photobiol.* **45**: 317-21 (1987)
- 87A064 Photokinetic and photophysical measurements of the sensitized photooxidation of the tryptophyl residue in N-acetyl tryptophanamide and in human serum albumin. Reddi, E.; Lambert, C.R.; Jori, G.; Rodgers, M.A.J., *Photochem. Photobiol.* **45**: 345-51 (1987)
- 87A072 Chemistry of singlet oxygen. 49. Photooxidation of thiiranes. Jensen, F.; Foote, C.S., *J. Am. Chem. Soc.* **109**: 1478-85 (1987)
- 87A107 Antiarthritic gold compounds effectively quench electronically excited singlet oxygen. Corey, E.J.; Mehrotra, M.M.; Khan, A.U., *Science* **236**: 68-9 (1987)
- 87A180 Rate constants for reactions for singlet oxygen with phenols and other compounds in water. Scully, F.E., Jr.; Hoigne, J., *Chemosphere* **16**: 681-94 (1987)
- 87A202 Quenching of the luminescence of singlet molecular oxygen in carbon tetrachloride solution by substituted derivatives of pyrrole. Venedikov, E.A.; Perfil'ev, V.A.; Titova, Yu.V.; Karavaev, B.I.; Berezin, B.D., *Biophysics* **32**: 217-20 (1987) Translated from: *Biofizika* **32**: 203-5 (1987)
- 87A318 Singlet oxygen reactions in aqueous solution. Physical and chemical quenching rate constants of crocin and related carotenoids. Manitto, P.; Speranza, G.; Monti, D.; Gramatica, P., *Tetrahedron Lett.* **28**: 4221-4 (1987)
- 87A368 syn- and anti-Dioxasessquinorbornenes. Singlet oxidation of exocyclic S-cis-1,3-butadienes. Clennan, E.L.; Matusuno, K., *J. Org. Chem.* **52**: 3483-5 (1987)
- 87A371 Photophysics of the natural antioxidant quercetin and mechanism of its interaction with singlet oxygen. Darmanyan, A.P.; Kasaikina, O.T.; Khrameeva, N.P., *Khim. Fiz.* **6**: 1083-92 (1987)
- 87E054 Thermal-lensing and phosphorescence studies of the quantum yield and lifetime of singlet molecular oxygen ($^1\Delta_g$) sensitized by hematoporphyrin and related porphyrins in deuterated and non-deuterated ethanol. Redmond, R.W.; Heihoff, K.; Braslavsky, S.E.; Truscott, T.G., *Photochem. Photobiol.* **45**: 209-13 (1987)
- 87E055 Ergosterol (provitamin D2) triplet state: An efficient sensitizer of singlet oxygen, $O_2(^1\Delta_g)$, formation. Gorman, A.A.; Hamblett, I.; Rodgers, M.A.J., *Photochem. Photobiol.* **45**: 215-21 (1987)
- 87E142 Evidence for physical quenching of singlet oxygen ($O_2(^1\Delta_g)$) by dinitrophenols. Gspone, H.E.; Previtali, C.M.; Garcia, N.A., *J. Photochem.* **36**: 247-53 (1987)
- 87E150 Steady-state near-infrared detection of singlet molecular oxygen: A Stern-Volmer quenching experiment with sodium azide. Hall, R.D.; Chignell, C.F., *Photochem. Photobiol.* **45**: 459-64 (1987)
- 87E181 Quenching of singlet oxygen by nickelocene. Rajadurai, S.; Das, P.K., *J. Photochem.* **37**: 33-40 (1987)
- 87E234 Aromatic ketone-naphthalene systems as absolute standards for the triplet-sensitized formation of singlet oxygen, $O_2(^1\Delta_g)$, in organic and aqueous media: A time-resolved luminescence study. Gorman, A.A.; Hamblett, I.; Lambert, C.; Prescott, A.L.; Rodgers, M.A.J.; Spence, H.M., *J. Am. Chem. Soc.* **109**: 3091-7 (1987)
- 87E466 Effect of solvent on the rate constant for the radiative deactivation of singlet molecular oxygen ($^1\Delta_g O_2$). Scurlock, R.D.; Ogilby, P.R., *J. Phys. Chem.* **91**: 4599-602 (1987)
- 87E658 Radiationless deactivation of singlet oxygen ($^1\Delta_g$) by solvent molecules. Schmidt, R.; Brauer, H.-D., *J. Am. Chem. Soc.* **109**: 6976-81 (1987)
- 87E668 Medium effects on the quantum yields for generation of singlet oxygen via anthracene and luminescence lifetime of oxygen in the $^1\Delta_g$ state. Darmanyan, A.P., *Khim. Fiz.* **6**: 1192-8 (1987)
- 87E690 Etude physicochimique d'une serie de cyanines. Partie VII. Rendement en etat triplet et photosensibilisation de l'oxygene singule. Carre, C.; Reichardt, C.; Lougnot, D.J., *J. Chim. Phys. Phys.-Chim. Biol.* **84**: 577-85 (1987)
- 87E941 Effectiveness of the formation of singlet oxygen photosensitised by water-soluble porphyrins. Dzhagarov, B.M.; Salokhiddinov, K.I.; Egorova, G.D.; Gurinovich, G.P., *Russ. J. Phys. Chem.* **61**: 1281-3 (1987) Translated from: *Zh. Fiz. Khim.* **61**: 2450-4 (1987)
- 87E959 Quenching of photosensitized luminescence of singlet molecular oxygen by 1,4-diaminoanthraquinone in solutions. Venedikov, E.A.; Perfil'ev, V.A.; Berezin, B.D., *Opt. Spectrosc. (USSR)* **63**: 400-1 (1987) Translated from: *Opt. Spektrosk.* **63**: 680-1 (1987)
- 87F104 Biological activities of phthalocyanines. VI. Photooxidation of L-tryptophan by selectively sulfonated gallium phthalocyanines: Singlet oxygen yields and effect of aggregation. Wagner, J.R.; Ali, H.; Langlois, R.; Brasseur, N.; van Lier, J.E., *Photochem. Photobiol.* **45**: 587-94 (1987)
- 87F290 Photosensitized generation of singlet molecular oxygen by endogenous photosensitizers of the human lens. Egorov, S.Yu.; Babizhaev, M.A.; Krasnovsky, A.A., Jr.; Shvedova, A.A., *Biophysics* **32**: 184-6 (1987) Translated from: *Biofizika* **32**: 169-71 (1987)
- 87F333 Laser flash-chemiluminescence study of $O_2(^1\Delta)$ photoeliminated from 1,4-dimethylnaphthalene endoperoxide. Chou, P.-T.; Frei, H., *J. Chem. Phys.* **87**: 3843-54 (1987)
- 87F440 An efficient regeneration of singlet oxygen from 2,5-diphenylfuran endoperoxide produced by a dye-sensitized oxygenation. Usui, Y.; Koike, H.; Kurimura, Y., *Bull. Chem. Soc. Jpn.* **60**: 3373-8 (1987)
- 87F479 Generation and deactivation of $^1O_2(^1\Delta)$ by helianthrene and helianthrene-derivatives. Acs, A.; Schmidt, R.; Brauer, H.-D., *Ber. Bunsenges. Phys. Chem.* **91**: 1331-7 (1987)
- 87F480 The role of geminate reaction in the self-sensitized photooxygenation of helianthrene and helianthrene-derivatives encounter- and reencounter probabilities. Brauer, H.-D.; Acs, A.; Schmidt, R., *Ber. Bunsenges. Phys. Chem.* **91**: 1337-46 (1987)
- 87F537 Kinetics of the photosensitized oxidation of polychlorophenols in alkaline aqueous solution. Gspone, H.E.; Previtali, C.M.; Garcia, N.A., *Toxicol. Environ. Chem.* **16**: 23-37 (1987)
- 87F541 Formation d'oxygene singulet $^1\Delta_g$ photosensibilisee par l'hypericine. Caracterisation et etude du mecanisme par spectroscopie laser. Jardon, P.; Lazorthak, N.; Gautron, R., *J. Chim. Phys. Phys.-Chim. Biol.* **84**: 1141-5 (1987)
- 87F542 Azomethine dyes - sensitizers, quenchers and reaction partner for singlet oxygen. Grossmann, N.; Winnig, B.; Weisse, A.; Fanghaenel, E., *J. Prakt. Chem.* **329**: 767-74 (1987)

- 87F554 The photosensitising properties and photostability of stilbene fluorescent whitening agents. Davidson, R.S.; Ismail, G.M.; Lewis, D.M., *J. Soc. Dyers Colour.* **103**: 261-4 (1987)
- 87F569 Reactions involving an amino group: A basic mechanism for interaction between 7-dialkylaminocoumarins and singlet oxygen. Zin'kovskaya, O.V.; Kuznetsova, N.A.; Reznichenko, A.V., *High Energy Chem.* **21**: 296-9 (1987) Translated from: *Khim. Vys. Energ.* **21**: 352-5 (1987)
- 87F639 Interaction of singlet oxygen [$O_2(^1\Delta)$] with aliphatic amines and hydroxylamines. Encinas, M.V.; Lemp, E.; Lissi, E.A., *J. Chem. Soc., Perkin Trans. 2* : 1125-7 (1987)
- 87F655 Quenching of singlet oxygen by cobalt complexes. Vidoczky, T.; Nemeth, S., *Photochemistry and Photophysics of Coordination Compounds*, H. Yersin and A. Vogler (eds.), Springer-Verlag, Berlin, 1987, p. 331-3
- 87F676 Chemistry of singlet oxygen. 48. Isolation and structure of the primary product of photooxygenation of 3,5-di-t-butyl catechol. Jensen, F.; Foote, C.S., *Photochem. Photobiol.* **46**: 325-30 (1987)
- 87M341 Some limitations in the use of biliverdin for the examination of singlet oxygen participation in oxidation reactions. Polewski, K.; Slawinska, D., *Roczn. Akad. Poznaniu* **177**: 35-46 (1987)
- 87N090 Comparative study of the sensitized photooxidation of N-methyl phenothiazine in homogeneous and microheterogeneous media. Oliveros, E.; Pheulpin, P.; Braun, A.M., *Tetrahedron* **43**: 1713-23 (1987)
- 87R032 Photo-properties of a silicon naphthalocyanine: A potential photosensitizer for photodynamic therapy. Firey, P.A.; Rodgers, M.A.J., *Photochem. Photobiol.* **45**: 535-8 (1987)
- 87R074 Photo oxidation effects on β -hydroxybutyrate dehydrogenase: Studies of membrane fragments and intact mitochondria. Robinson, R.S.; Roberts, A.J.; Campbell, I.D., *Photochem. Photobiol.* **45**: 231-4 (1987)
- 87R138 Dye sensitised photo-oxidation of the methyl and phenyl esters of oleic and linoleic acids. Chacon, J.N.; Jamieson, G.R.; Sinclair, R.S., *Chem. Phys. Lipids* **43**: 81-99 (1987)
- 87R227 Quenching of singlet molecular oxygen by the screening pigments melanin and ommochrome. Egorov, S.Yu.; Dontsov, A.E.; Krasnovsky, A.A.; Ostrovskii, M.A., *Biophysics* **32**: 740-2 (1987) Translated from: *Biofizika* **32**: 685-6 (1987)
- 88A026 Interaction of singlet molecular oxygen with disubstituted olefins. Evidence for a physical quenching induced by the hydrocarbon chain. Tanielian, C.; Mechlin, R., *J. Phys. Chem.* **92**: 265-7 (1988)
- 88A105 Singlet molecular oxygen: Not a major product of the reaction between tris(2,2'-bipyridine)ruthenium(3+) and superoxide radical anions. Mulazzani, Q.G.; Ciano, M.; D'Angelantonio, M.; Venturi, M.; Rodgers, M.A.J., *J. Am. Chem. Soc.* **110**: 2451-7 (1988)
- 88A164 Deactivation of singlet molecular oxygen by thiols and related compounds, possible protectors against skin photosensitivity. Rougee, M.; Bensasson, R.V.; Land, E.J.; Pariente, R., *Photochem. Photobiol.* **47**: 485-9 (1988)
- 88A165 Singlet oxygen yields and radical contributions in the dye-sensitised photooxidation in methanol of esters of polyunsaturated fatty acids (oleic, linoleic, linolenic and arachidonic). Chacon, J.N.; McLearie, J.; Sinclair, R.S., *Photochem. Photobiol.* **47**: 647-56 (1988)
- 88A276 Ability of platinum(II) complexes of 2,2'-bipyridine and 1,10-phenanthroline with halides and pseudo-halides to photosensitize the production of 1O_2 . Shukla, S.; Kamath, S.S.; Srivastava, T.S., *J. Photochem. Photobiol. A* **44**: 143-52 (1988)
- 88A284 The production of singlet molecular oxygen by zinc(II) phthalocyanine in ethanol and in unilamellar vesicles. Chemical quenching and phosphorescence studies. Valduga, G.; Nonell, S.; Reddi, E.; Jori, G.; Braslavsky, S.E., *Photochem. Photobiol.* **48**: 1-5 (1988)
- 88A385 Lumiflavin-sensitized photooxygenation of indole. Yoshimura, A.; Ohno, T., *Photochem. Photobiol.* **48**: 561-5 (1988)
- 88A398 Time-resolved reaction of $O_2(^1\Delta)$ with I⁻ in aqueous solution. Braathen, G.; Chou, P.-T.; Frei, H., *J. Phys. Chem.* **92**: 6610-5 (1988)
- 88A427 Identification of both preequilibrium and diffusion limits for reaction of singlet oxygen, $O_2(^1\Delta_g)$, with both physical and chemical quenchers: Variable-temperature, time-resolved infrared luminescence studies. Gorman, A.A.; Hamblett, I.; Lambert, C.; Spencer, B.; Standen, M.C., *J. Am. Chem. Soc.* **110**: 8053-9 (1988)
- 88A507 Kinetics of elementary reactions of singlet oxygen with thiophenes and with their complexes with metals. Darmanyan, A.P.; Sirota, T.V.; Mazaletskii, A.B., *Kinet. Catal.* **29**: 1151-7 (1988) Translated from: *Kinet. Katal.* **29**: 1323-30 (1988)
- 88E018 Singlet oxygen luminescence quantum yields in organic solvents and water. Losev, A.P.; Byteva, I.M.; Gurinovich, G.P., *Chem. Phys. Lett.* **143**: 127-9 (1988)
- 88E452 Triplet energy surfaces of one-way isomerizing olefins as studied by singlet oxygen luminescence technique. Arai, T.; Karatsu, T.; Tsuchiya, M.; Sakuragi, H.; Tokumaru, K., *Chem. Phys. Lett.* **149**: 161-6 (1988)
- 88E479 Singlet-oxygen quenching by carotenoids: Steady-state luminescence experiments. Murasecco-Suardi, P.; Oliveros, E.; Braun, A.M.; Hansen, H.-J., *Helv. Chim. Acta* **71**: 1005-10 (1988)
- 88E657 Silicon naphthalocyanine triplet state and oxygen: A reversible energy-transfer reaction. Firey, P.A.; Ford, W.E.; Sounik, J.R.; Kenney, M.E.; Rodgers, M.A.J., *J. Am. Chem. Soc.* **110**: 7626-30 (1988)
- 88E775 Delayed fluorescence of the Kautsky-Mueller type from liquid solutions of aromatic compounds and oxygen. Nickel, B.; Rodriguez Prieto, M.F., *Ber. Bunsenges. Phys. Chem.* **92**: 1493-503 (1988)
- 88F059 Mechanism of synergistic stabilization by hindered amines. Yang, X.-Z.; Chen, Y.; Dickinson, L.C.; Chien, J.C.W., *Polym. Degrad. Stab.* **20**: 1-35 (1988)
- 88F151 Singlet oxygen production and photoisomerization: Two competitive processes for merocyanine 540 irradiated with visible light. Hoebeke, M.; Seret, A.; Piette, J.; van de Vorst, A., *J. Photochem. Photobiol. B* **1**: 437-46 (1988)
- 88F415 Products and relative reaction rates of the oxidation of tocopherols with singlet molecular oxygen. Neely, W.C.; Martin, J.M.; Barker, S.A., *Photochem. Photobiol.* **48**: 423-8 (1988)
- 88F440 Study on antioxidants and their activity with singlet oxygen by fluorescence spectroscopy. Lang, J.; Shi, J.; Li, G., *Huadong Huagong Xueyuan Xuebao* **14**: 65-70 (1988)
- 88F462 Interaction of singlet oxygen with nitroso- and phenol-type inhibitors. Moger, G.; Darmanyan, A.P., *Magy. Kem. Foly.* **94**: 282-4 (1988)
- 88F485 Oxygen consumption in two aqueous systems: Excited nitrite ions-oxygen and nitrite ions-singlet oxygen. Bilski, P.; Szschlinski, J.; Oleksy, E., *J. Photochem. Photobiol. A* **45**: 269-88 (1988)
- 88F503 Photosensitized formation of singlet molecular oxygen in solutions of water soluble porphyrins. Direct luminescence measurements. Krasnovsky, A.A., Jr.; Egorov, S.Yu.; Nasarova, O.V.; Yartsev, E.I.; Ponomarev, G.V., *Stud. Biophys.* **124**: 123-42 (1988)
- 88F582 Singlet oxygen in the photodegradation of aminoanthraquinones. Rtishchev, N.I.; Studzinskii, O.P.; Ponomareva, R.P., *J. Org. Chem. USSR* **24**: 1150-4 (1988) Translated from: *Zh. Org. Khim.* **24**: 1277-81 (1988)
- 88F650 On the mechanism of quenching of singlet oxygen by chlorinated phenolic pesticides. Palumbo, M.C.; Garcia, N.A., *Toxicol. Environ. Chem.* **17**: 103-16 (1988)
- 88M109 Chemical sources of singlet oxygen. 2. Quantitative generation of singlet oxygen from hydrogen peroxide disproportionation catalyzed by molybdate ions. Aubry, J.M.; Cazin, B., *Inorg. Chem.* **27**: 2013-4 (1988)
- 88N343 Generation of singlet oxygen $^1\Delta_g$ photosensitized by hypericin, kinetic study in non ionic micellar medium. Racinet, H.; Jardon, P.; Gautron, R., *J. Chim. Phys. Phys.-Chim. Biol.* **85**: 971-7 (1988)
- 88P313 Quenching action towards 1O_2 of some nitroxyl radicals and hindered amines and their photostabilizing efficiencies. Cen, J.; Hu, X.; Shen, D.; Zhu, L., *Guangguo Kexue Yu Kuang Huaxue* : 1-8 (1988)

- 88R064 Some prevalent biomolecules as defenses against singlet oxygen damage. Dahl, T.A.; Midden, W.R.; Hartman, P.E., *Photochem. Photobiol.* **47**: 357-62 (1988)
- 88R070 Direct detection of singlet oxygen sensitized by nalidixic acid: The effect of pH and melanin. Dayhaw-Barker, P.; Truscott, T.G., *Photochem. Photobiol.* **47**: 765-7 (1988)
- 88R136 Mechanism of physical quenching of singlet molecular oxygen by chlorophylls and related compounds of biological interest. Tanielian, C.; Wolff, C., *Photochem. Photobiol.* **48**: 277-80 (1988)
- 88R142 Quenching of singlet oxygen by haematoxoporphyrin derivative (and haematoxoporphyrin) and its consequences on the efficiency of photodynamic cancer therapy. Tanielian, C.; Heinrich, G.; Entezami, A., *J. Chem. Soc., Chem. Commun.* : 1197-8 (1988)
- 88R193 Photophysical parameters of pheophorbide a: phosphorescence and generation of singlet molecular oxygen. Krasnovsky, A.A.Jr.; Neverov, K.V.; Egorov, S.Yu.; Reder, B., *Opt. Spectrosc. (USSR)* **64**: 471-4 (1988) Translated from: *Opt. Spektrosk.* **64**: 790-5 (1988)
- 88R194 Photophysical studies of tin(IV)-protoporphyrin: Potential phototoxicity of a chemotherapeutic agent proposed for the prevention of neonatal jaundice. Land, E.J.; McDonagh, A.F.; McGarvey, D.J.; Truscott, T.G., *Proc. Natl. Acad. Sci. U.S.A.* **85**: 5249-53 (1988)
- 88R200 Laser flash photolysis of purpurins: Novel potential photosensitizers of interest in photodynamic therapy. Borland, C.F.; McGarvey, D.J.; Morgan, A.R.; Truscott, T.G., *J. Photochem. Photobiol., B* **2**: 427-34 (1988)
- 88R236 Correlation between the rates of singlet oxygen quenching by phenols and their inhibitory effects on benzo(a)pyrene-induced neoplasia. Bensasson, R.V.; Rougee, M., *C. R. Acad. Sci., Ser. 3* **307**: 807-10 (1988)
- 88Z155 Absolute determination of quantum yields of photosensitization by time resolved thermal lensing. Redmond, R.W.; Braslavsky, S.E., *NATO ASI Ser., Ser. H* **15**: 93-5 (1988)
- 89A099 Trimethyl phosphite traps intermediates in the reaction of $^1\text{O}_2$ and diethyl sulfide. Nahm, K.; Foote, C.S., *J. Am. Chem. Soc.* **111**: 1909-10 (1989)
- 89A105 Generation and transient spectroscopy of substituted diaryl carbonyl oxides. Scaiano, J.C.; McGimpsey, W.G.; Casal, H.L., *J. Org. Chem.* **54**: 1612-6 (1989)
- 89A202 Photochemical reactions. Part 23. Photo-oxygenation of styrenic estrogens: Product characterization and kinetics of the dye-sensitized photo-oxygenation of 9,11-didehydroestrone derivatives. Planas, A.; Lupon, P.; Cascallo, M.; Bonet, J.-J., *Helv. Chim. Acta* **72**: 715-24 (1989)
- 89A235 Concerning the mechanism of reaction of simple conjugated olefin triplet states with oxygen. Gorman, A.A.; Rodgers, M.A.J., *J. Am. Chem. Soc.* **111**: 5557-60 (1989)
- 89A241 Photoenol biradicals as singlet oxygen sensitizers. Redmond, R.W.; Scaiano, J.C., *J. Phys. Chem.* **93**: 5347-9 (1989)
- 89A254 Mechanism and kinetics of the methylene-blue-sensitized photo-oxygenation of polydienes. Tanielian, C.; Mechlin, R., *J. Photochem. Photobiol., A* **48**: 43-51 (1989)
- 89A322 Time-resolved study of the photooxygenation of 3-hydroxyflavone. Studer, S.L.; Brewer, W.E.; Martinez, M.L.; Chou, P.T., *J. Am. Chem. Soc.* **111**: 7643-4 (1989)
- 89A331 Singlet molecular oxygen quenching by saturated and unsaturated fatty-acids and by cholesterol. Vever-Bizet, C.; Dellinger, M.; Brault, D.; Rougee, M.; Bensasson, R.V., *Photochem. Photobiol.* **50**: 321-5 (1989)
- 89A346 Evidence for hole injection from $\text{O}_2(^1\Delta)$ into p-Si. Graetzl, M.; Frei, H., *J. Phys. Chem.* **93**: 7037-41 (1989)
- 89A400 Determination of rate constants of $^1\text{O}_2$ consumption by $^1\text{O}_2$ acceptors in weakly deactivating solvents. Opriel, U.; Seikel, K.; Schmidt, R.; Brauer, H.-D., *J. Photochem. Photobiol., A* **49**: 299-309 (1989)
- 89A506 Lifetime of singlet oxygen in aqueous hydrogen peroxide (BHP). Rodgers, M.A.J.; Shand, M.A., AFWL-TR-89-07, 1989, 24p.; AD-A211545
- 89B024 Triplet state characteristics and singlet oxygen generation properties of anthracycloines. Andreoni, A.; Land, E.J.; Malatesta, V.; McLean, A.J.; Truscott, T.G., *Biochim. Biophys. Acta* **990**: 190-7 (1989)
- 89D112 An ESR study of the visible light photochemistry of givocarcin V. Alegria, A.E.; Krishna, C.M.; Elespuru, R.K.; Riesz, P., *Photochem. Photobiol.* **49**: 257-65 (1989)
- 89E113 A pulse radiolysis based singlet oxygen luminescence facility. Gorman, A.A.; Hamblett, I.; Land, E.J., *J. Am. Chem. Soc.* **111**: 1876-7 (1989)
- 89E158 Anthralin-derived transients. I. The triplet state and the products of its reaction with oxygen in benzene. Bruce, J.M.; Gorman, A.A.; Hamblett, I.; Kerr, C.W.; Lambert, C.; McNeeney, S.P., *Photochem. Photobiol.* **49**: 439-45 (1989)
- 89E232 Thermal lens measurements in liquids on a submicrosecond time scale. Isak, S.J.; Komorowski, S.J.; Merrow, C.N.; Poston, P.E.; Eyring, E.M., *Appl. Spectrosc.* **43**: 419-22 (1989)
- 89E324 Solvent effects on the ability of amines to physically quench singlet oxygen as determined by time-resolved infrared emission studies. Clennan, E.L.; Noe, L.J.; Wen, T.; Szneler, E., *J. Org. Chem.* **54**: 3581-4 (1989)
- 89E365 Dynamics of the triplet state and the reverse proton transfer of 3-hydroxyflavone. Brewer, W.E.; Studer, S.L.; Standiford, M.; Chou, P.T., *J. Phys. Chem.* **93**: 6088-94 (1989)
- 89E388 Influence of heavy atoms on the deactivation of singlet oxygen ($^1\Delta_g$) in solution. Schmidt, R., *J. Am. Chem. Soc.* **111**: 6983-7 (1989)
- 89E505 Chlorin e₆. Spectral-energetic characteristics and generation of singlet molecular oxygen in some homogeneous and heterogeneous systems. Kochubeev, G.A.; Frolov, A.A.; Gurinovich, G.P., *Khim. Fiz.* **8**: 1184-90 (1989)
- 89E585 Rise and decay kinetics of photosensitized singlet oxygen luminescence in water. Measurements with nanosecond time-correlated single photon counting technique. Egorov, S.Yu.; Kamalov, V.F.; Koroteev, N.I.; Krasnovsky, A.A.Jr.; Toleutaev, B.N.; Zinukov, S.V., *Chem. Phys. Lett.* **163**: 421-4 (1989)
- 89E597 Behavior of singlet molecular oxygen ($^1\Delta_g\text{O}_2$) in a polymer matrix: Effects of temperature, matrix rigidity, and molecular composition. Clough, R.L.; Dillon, M.P.; Iu, K.-K.; Ogilby, P.R., *Macromolecules* **22**: 3620-8 (1989)
- 89E700 Direct determination of lifetimes of singlet oxygen in various solvents. Wang, W.; Tang, Y.; Yu, Y., *Yingyong Huaxue* **6**: 54-6 (1989)
- 89E798 Measurement of rate constants for quenching singlet oxygen with a cypridina luciferin analoge (2-methyl-6-[p-methoxyphenyl]-3,7-dihydro-imidazo [1,2-a] pyrazin-3-one) and sodium azide. Mashiko, S.; Ashino, T.; Mizumoto, I.; Suzuki, N.; Nakano, M.; Goto, T.; Inaba, H., *Photomed. Photobiol.* **11**: 191-4 (1989)
- 89F024 Physical quenching and chemical reaction of singlet molecular oxygen with azo dyes. Bortolus, P.; Monti, S.; Albini, A.; Fasani, E.; Pietra, S., *J. Org. Chem.* **54**: 534-40, 54(13): 3246 (1989)
- 89F079 Reaction of singlet oxygen with 2-cyclopenten-1-ones. Kwon, B.-M.; Kanner, R.C.; Foote, C.S., *Tetrahedron Lett.* **30**: 903-6 (1989)
- 89F181 Mixed-ligand complexes of platinum(II) and palladium(II) with 1,10 phenanthroline and aromatic chelating ligands containing OH or SH groups as photosensitizers for the generation of $^1\text{O}_2$. Shukla, S.; Karanath, S.S.; Srivastava, T.S., *J. Photochem. Photobiol., A* **47**: 287-98 (1989)
- 89F238 Singlet oxygen mediated photofragmentation reactions of amino alcohols: A novel oxidative fragmentation involving both superoxide and excited oxygen intermediates. Haugen, C.M.; Whitten, D.G., *J. Am. Chem. Soc.* **111**: 7281-2 (1989)
- 89F260 Quantum yield of singlet molecular oxygen sensitization by copper(II) tetracarboxyphthalocyanine. Lagorio, M.G.; Dicelio, L.E.; San Roman, E.A.; Braslavsky, S.E., *J. Photochem. Photobiol., B* **3**: 615-24 (1989)
- 89F298 Temperature, solvent, and substituent effects on the singlet oxidations of allylic phenyl sulfoxides, sulfones, and sulfides. Clennan, E.L.; Chen, X., *J. Am. Chem. Soc.* **111**: 8212-8 (1989)

- 89F311 Mixed-ligand nickel(II) complexes of 2,2'-bipyridine and 1,10-phenanthroline with 3,4-dimercaptotoluene as quenchers of $^1\text{O}_2$. Shukla, S.; Srivastava, T.S., *J. Photochem. Photobiol. A* **48**: 249-57 (1989)
- 89F349 Singlet oxygen chemistry. IV. Sensitized photooxygenation of 6-substituted-1,4-cyclooctadienes. Chen, D.-L.; Wang, D.-Y.; Xiao, X.-L., *Acta Chim. Sin. (Huaxue Xuebao)* **47**: 147-51 (1989)
- 89F350 Singlet oxygen chemistry. V. Singlet oxygenation of dialkylated phenols. Liu, L.-P.; Wang, D.-Y.; Xiao, X.-L., *Acta Chim. Sin. (Huaxue Xuebao)* **47**: 456-60 (1989)
- 89F358 The comparison of photosensitized oxygenation efficiency for several kinds of sensitizers. Li, Q.-Y.; Xiao, X.-L.; He, H.-Z.; Zhang, L.-H.; Wang, D.-Y., *Ganguang Kexue Yu Kuang Huaxue* : 59-63 (1989)
- 89F368 Octahydropyrrolo[4,3,2-m,n]acridine derivatives. 3. Antiradical, antioxidant, erythrocyte-stabilizing activity and quenching of singlet oxygen. Tirzite, D.Ya.; Kirule, I.E.; Byteva, I.M.; Tirzit, G.D.; Bisenieks, E.A.; Dubur, G.Ya., *Khim.-Farm. Zh.* **23**: 290-2 (1989)
- 89F400 Singlet oxygen and triazolincidone additions to α,β -unsaturated sulfoxides. Akasaka, T.; Misawa, Y.; Goto, M.; Ando, W., *Tetrahedron* **45**: 6657-66 (1989)
- 89F462 Self-sensitized photo-oxidation of platinum(II) complexes of 1,10-phenanthroline with 3,4-dimercaptotoluene and thiosalicylic acid by $^1\text{O}_2$. Shukla, S.; Kamath, S.S.; Srivastava, T.S., *J. Photochem. Photobiol. A* **50**: 199-207 (1989)
- 89F484 Flavin-sensitized photo-oxidation of unsaturated fatty acids. Fukuzumi, S.; Tanii, K.; Tanaka, T., *J. Chem. Soc., Perkin Trans. 2* : 2103-8 (1989)
- 89M015 Chemical sources of singlet oxygen. 3. Peroxidation of water-soluble singlet oxygen carriers with the hydrogen peroxide-molybdate system. Aubry, J.M.; Cazin, B.; Duprat, F., *J. Org. Chem.* **54**: 726-8 (1989)
- 89M038 Evidence for the generation of singlet molecular oxygen during dopa and dopamine peroxidation. Kruk, I.; Lichszeld, K.; Michalska, T.; Bounias, M., *Z. Naturforsch., C Biosci.* **44C**: 39-44 (1989)
- 89M096 The quenching of $^1\text{O}_2(^1\Delta_g)$ by anthracyclines in aqueous solution. Evidence for a charge-transfer mechanism. Manitto, P.; Speranza, G.; Malatesta, V., *Chem. Phys. Lett.* **159**: 310-4 (1989)
- 89M097 Quantification of singlet oxygen generated by thermolysis of 3,3'-(1,4-naphthylidene)dipropionate. Monomol and dimol photoemission and the effects of 1,4-diazabicyclo[2.2.2]octane. Di Mascio, P.; Sies, H., *J. Am. Chem. Soc.* **111**: 2909-14 (1989)
- 89R044 Bacteriochlorophyll a triplet state and its interactions with bacterial carotenoids and oxygen. Borland, C.F.; Cogdell, R.J.; Land, E.J.; Truscott, T.G., *J. Photochem. Photobiol. B* **3**: 237-45 (1989)
- 89R092 Photosensitized oxidation of biomaterials and related model compounds. Davila, J.; Harriman, A., *Photochem. Photobiol.* **50**: 29-35 (1989)
- 89R188 Lycopene as the most efficient biological carotenoid singlet oxygen quencher. Di Mascio, P.; Kaiser, S.; Sies, H., *Arch. Biochem. Biophys.* **274**: 532-8 (1989)
- 90A018 Pressure effect on the lifetime of singlet oxygen in solutions. Okamoto, M.; Tanaka, F.; Teranishi, H., *J. Phys. Chem.* **94**: 669-72 (1990)
- 90A022 Photoreactions of macrocyclic dyes bound to human serum albumin. Davila, J.; Harriman, A., *Photochem. Photobiol.* **51**: 9-19 (1990)
- 90A184 Physical and chemical scavenging of singlet molecular oxygen by tocopherols. Kaiser, S.; Di Mascio, P.; Murphy, M.E.; Sies, H., *Arch. Biochem. Biophys.* **277**: 101-8 (1990)
- 90A300 Kinetic studies on anthralin photooxidation. Mueller, K.; Kanner, R.C.; Foote, C.S., *Photochem. Photobiol.* **52**: 445-50 (1990)
- 90A328 Effect of oxygen-enhanced intersystem crossing on the observed efficiency of formation of singlet oxygen. McLean, A.J.; McGarvey, D.J.; Truscott, T.G.; Lambert, C.R.; Land, E.J., *J. Chem. Soc., Faraday Trans.* **86**: 3075-80 (1990)
- 90A437 Interaction between singlet oxygen and biologically active compounds in aqueous solution. III. Physical and chemical $^1\text{O}_2$ quenching rate constants of 6,6'-diapocarotenoids. Speranza, G.; Manitto, P.; Monti, D., *J. Photochem. Photobiol. B* **8**: 51-61 (1990)
- 90E030 A pulse radiolysis and pulsed laser study of the vitamin D3 triplet state: Lifetime, relaxation and 'nonvertical' excitation. Gorman, A.A.; Hamblett, I.; Lambert, C.; Prescott, A.L., *Photochem. Photobiol.* **51**: 29-35 (1990)
- 90E220 Singlet oxygen formation in a solid organic polymer upon irradiation of the oxygen-polymer charge-transfer band. Ogilby, P.R.; Kristiansen, M.; Clough, R.L., *Macromolecules* **23**: 2698-704 (1990)
- 90E297 Hydrazines: New charge-transfer physical quenchers of singlet oxygen. Clennan, E.L.; Noe, L.J.; Szneler, E.; Wen, T., *J. Am. Chem. Soc.* **112**: 5080-5 (1990)
- 90E312 Quantum yields of triplet and $\text{O}_2(^1\Delta_g)$ formation of 4-thiouridine in water and acetonitrile. Heihoff, K.; Redmond, R.W.; Braslavsky, S.E.; Rougee, M.; Salet, C.; Favre, A.; Bensasson, R.V., *Photochem. Photobiol.* **51**: 634-41 (1990)
- 90E324 Photosensitized formation and quenching of molecular oxygen by monomeric and aggregated pigment molecules of photosynthesizing bacteria. Egorov, S.Yu.; Krasnovsky, A.A., Jr.; Vychevyanina, I.V.; Drozdova, N.N.; Krasnovskii, A.A., *Doklady Biophys.* **310**: 6-10 (1990) Translated from: *Dokl. Akad. Nauk SSSR* **310**: 471-5 (1990)
- 90E374 Laser-induced optoacoustics combined with near-infrared emission. An alternative approach for the determination of intersystem crossing quantum yields applied to porphycenes. Nonell, S.; Aramendia, P.F.; Heihoff, K.; Negri, R.M.; Braslavsky, S.E., *J. Phys. Chem.* **94**: 5879-83 (1990)
- 90E400 Investigations of luminescence quenching of singlet oxygen by organic substances by the mechanism of charge transfer. Venedikov, E.A.; Lipatov, N.G.; Berezin, B.D., *Khim. Fiz.* **9**: 611-5 (1990)
- 90E491 A photophysical study of some purpurins. Macpherson, A.N.; Kessel, D.; Morgan, A.R.; Munro, I.; Truscott, T.G., *J. Chem. Soc., Faraday Trans.* **86**: 3081-5 (1990)
- 90E530 Potential photosensitizers for photodynamic therapy. II. Photophysical properties of [26] porphyrin. Schermann, G.; Schmidt, R.; Voelcker, A.; Brauer, H.-D.; Mertes, H.; Franck, B., *Photochem. Photobiol.* **52**: 741-4 (1990)
- 90E622 Carotenoids, tocopherols and thiols as biological singlet molecular oxygen quenchers. Di Mascio, P.; Devasagayam, T.P.A.; Kaiser, S.; Sies, H., *Biochem. Soc. Trans.* **18**: 1054-6 (1990)
- 90E731 Quenching of the luminescence of singlet molecular oxygen by phthalocyanines and naphthalocyanines. Krasnovsky, A.A., Jr.; Rodgers, M.A.J.; Gal'pern, M.G.; Kenney, M.E.; Luk'yanets, E.A., *Bioorg. Khim.* **16**: 1413-8 (1990)
- 90F065 Kinetic parameters of the reactivity of dihydroxy naphthalenes with singlet oxygen. Croux, S.; Maurette, M.-T.; Hocquaux, M.; Ananides, A.; Braun, A.M.; Oliveros, E., *New J. Chem.* **14**: 161-7 (1990)
- 90F069 Heavy atom quenching of the singlet delta dioxygen molecule ($^1\Delta_g \text{O}_2$). Corey, E.J.; Khan, A.U.; Ha, D.-C., *Tetrahedron Lett.* **31**: 1389-92 (1990)
- 90F111 Intermediates in the ene reactions of singlet oxygen and N-phenyl-1,2,4-triazoline-3,5-dione with olefins. Orfanopoulos, M.; Smouli, I.; Foote, C.S., *J. Am. Chem. Soc.* **112**: 3607-14 (1990)
- 90F157 Chalcogenapyrylium dyes as potential photochemotherapeutic agents. Solution studies of heavy atom effects on triplet yields, quantum efficiencies of singlet oxygen generation, rates of reaction with singlet oxygen, and emission quantum yields. Detty, M.R.; Merkel, P.B., *J. Am. Chem. Soc.* **112**: 3845-55 (1990)
- 90F196 Mixed-ligand palladium(II) and platinum(II) complexes of α -diimine and azide as photosensitizers of $^1\text{O}_2$ production. Kamath, S.S.; Srivastava, T.S., *J. Photochem. Photobiol. A* **52**: 83-9 (1990)
- 90F241 Singlet oxygenation of oxadisiliranes. Syntheses and crystal structure of 1,2,4,3,5-trioxadisilolanes. Ando, W.; Kako, M.; Akasaka, T.; Kabe, Y., *Tetrahedron Lett.* **31**: 4177-80 (1990)

- 90F251 Singlet oxygen photogeneration by ionized and un-ionized derivatives of Rose Bengal and eosin Y in diluted solutions. Amat-Guerri, F.; Lopez-Gonzalez, M.M.C.; Martinez-Utrilla, R.; Sastre, R., *J. Photochem. Photobiol.*, A 53: 199-210 (1990)
- 90F292 Role of exciplex and ion pair in the photosensitized oxygenation of 1,4-diphenyl-1,3-butadiene. Takahashi, Y.; Wakamatsu, K.; Kikuchi, K.; Miyashi, T., *J. Phys. Org. Chem.* 3: 509-18 (1990)
- 90F329 Sensitized photooxygenation. 3. Mechanistic studies on the singlet oxygenation of 5,6-disubstituted 3,4-dihydro-2H-pyrans. Chan, Y.-Y.; Li, X.; Zhu, C.; Liu, X.; Zhang, Y.; Liang, X., *J. Org. Chem.* 55: 5497-504 (1990)
- 90F347 The interaction of singlet molecular oxygen $O_2(^1\Delta_g)$ with indolic derivatives. Distinction between physical and reactive quenching. Palumbo, M.C.; Garcia, N.A.; Arguello, G.A., *J. Photochem. Photobiol.*, B 7: 33-42 (1990)
- 90F360 Mechanisms and rate constants for the quenching of singlet oxygen by nickel complexes. Shiozaki, H.; Nakazumi, H.; Takamura, Y.; Kitao, T., *Bull. Chem. Soc. Jpn.* 63: 2653-8 (1990)
- 90F411 Redox properties of phenols, their relationships to singlet oxygen quenching and to their inhibitory effects on benzo(a)pyrene-induced neoplasia. Scurlock, R.; Rougee, M.; Bensasson, R.V., *Free Radical Res. Commun.* 8: 251-8 (1990)
- 90F473 Quenching of singlet oxygen by alkenes with a hindered double bond. Sharipov, G.L.; Voloshin, A.I.; Kazakov, V.P.; Tolstikov, G.A., *Bull. Acad. Sci. USSR, Div. Chem. Sci.* 38: 233-6 (1990) Translated from: *Izv. Akad. Nauk SSSR, Ser. Khim.* : 286-9 (1990)
- 90F502 Steady-state near-infrared detection of singlet molecular oxygen: A Stern-Volmer quenching experiment with luminol, superoxide dismutase, and Cypridina luciferin analogues. Suzuki, N.; Mizumoto, I.; Toya, Y.; Nomoto, T.; Mashiko, S.; Inaba, H., *Agric. Biol. Chem.* 54: 2783-7 (1990)
- 90F511 The effect of oxygen on the kinetics of the photo-sensitised chemiluminescence of glycylltryptophan. Lozovskaya, E.L.; Sapezhinskii, I.I., *Russ. J. Phys. Chem.* 64: 537-9 (1990) Translated from: *Zh. Fiz. Khim.* 64: 1012-6 (1990)
- 90F518 Singlet molecular oxygen-mediated photooxidation of monochloro and mononitrophenols. A kinetic study. Palumbo, M.C.; Garcia, N.A.; Gutierrez, M.I.; Luiz, M., *Toxicol. Environ. Chem.* 29: 85-94 (1990)
- 90F554 The role of excited singlet molecular oxygen in the photodegradation of functional squarylium dyes. Kuramoto, N., *J. Soc. Dyers Colour.* 106: 181-6 (1990)
- 90M125 Determination of rate constants for quenching singlet oxygen by chemiluminescence technique. Sheremetets, V.V.; Kabalnova, N.N.; Komissarov, V.D.; Mavrodiev, V.K.; Lerman, B.M.; Belogaeva, T.A.; Tolstikov, G.A., *React. Kinet. Catal. Lett.* 41: 251-6 (1990)
- 90N078 Quenching of singlet molecular oxygen ($^1\Delta_g O_2$) in silica gel/cyclohexane heterogeneous systems. A direct time-resolved study. Iu, K.-K.; Thomas, J.K., *J. Am. Chem. Soc.* 112: 3319-25 (1990)
- 90R006 Potential photosensitizers for photodynamic tumor therapy. I. Photophysical properties of two chlorin derivatives. Schermann, G.; Voelcker, A.; Seikel, K.; Schmidt, R.; Brauer, H.-D.; Montfarts, F.-P., *Photochem. Photobiol.* 51: 45-51 (1990)
- 90R128 Effects, quenching mechanisms, and kinetics of carotenoids in chlorophyll-sensitized photooxidation of soybean oil. Lee, S.-H.; Min, D.B., *J. Agric. Food Chem.* 38: 1630-4 (1990)
- 90R134 Quenching of singlet oxygen by human plasma. Kanofsky, J.R., *Photochem. Photobiol.* 51: 299-303 (1990)
- 90R162 EPR studies of trapped singlet oxygen (1O_2) generated during photoirradiation of hypocrellin A. Zang, L.-Y.; Zhang, Z.; Misra, H., *Photochem. Photobiol.* 52: 677-83 (1990)
- 90R164 In vitro photodynamic activity of kryptocyanine. Harriman, A.; Luengo, G.; Gulliya, K.S., *Photochem. Photobiol.* 52: 735-40 (1990)
- 90R212 The photodynamic effect of Rose Bengal on proteins of the mitochondrial inner membrane. Gjulivi, C.; Sarcansky, M.; Rosenfeld, E.; Boveris, A., *Photochem. Photobiol.* 52: 745-51 (1990)
- 90U480 The effects of nickel(II) chelates on photosensitized oxidation of soybean oil. Min, D.B.; Lee, S.H., *Dev. Food Sci.* 24: 939-51 (1990)
- 91A252 Effect of the peptide bond on the singlet-molecular-oxygen-mediated sensitized photo-oxidation of tyrosine and tryptophan dipeptides. A kinetic study. Bertolotti, S.G.; Garcia, N.A.; Arguello, G.A., *J. Photochem. Photobiol.*, B 10: 57-70 (1991)
- 91A271 Pressure effect on the radiationless deactivation of singlet oxygen by 1,4-diazabicyclo[2.2.2]octane in six different solvents. Seikel, K.; Brauer, H.-D., *Ber. Bunsenges. Phys. Chem.* 95: 900-4 (1991)
- 91A297 A laser flash photolysis study of aromatic nitrones. Triplet state properties and reactivity with singlet oxygen. Cyr, D.R.; Mathew, T.; Ashok, K.; Das, P.K.; George, M.V., *J. Photochem. Photobiol.*, A 60: 161-74 (1991)
- 91A311 Singlet oxygen and electron-transfer mechanisms in the dicyanoanthracene-sensitized photooxidation of 2,3-diphenyl-1,4-dioxene. Silverman, S.K.; Foote, C.S., *J. Am. Chem. Soc.* 113: 7672-5 (1991)
- 91A341 Sensitized photo-oxidation of dihydroxybenzenes and chlorinated derivatives. A kinetic study. Martire, D.O.; Braslavsky, S.E.; Garcia, N.A., *J. Photochem. Photobiol.*, A 61: 113-24 (1991)
- 91A394 Oxidation of superoxide radical anion by excited tris(2,2'-bipyridine)ruthenium(II) ion in acetonitrile. A search for singlet molecular oxygen. Mulazzani, Q.G.; D'Angelantonio, M.; Venturi, M.; Rodgers, M.A.J., *J. Phys. Chem.* 95: 9605-8 (1991)
- 91A448 Triplet energies and singlet oxygen reaction rates of pyrazolotriazole azomethine dyes. Douglas, P.; Townsend, S.M.; Ratcliffe, R., *J. Imaging Sci.* 35: 211-8 (1991)
- 91A478 Nanosecond measurements of photosensitized luminescence of singlet molecular oxygen in aqueous solutions saturated by air: Effect of temperature and detergent presence. Zinukov, S.V.; Kamalov, V.F.; Koroteev, N.I.; Krashinsky, A.A., Jr., *Opt. Spectrosc. (USSR)* 70: 460-2 (1991) Translated from *Opt. Spektrosk.* 70: 790-4 (1991)
- 91E003 Photophysical properties of C_{60} . Arbogast, J.W.; Darmanyan, A.P.; Foote, C.S.; Rubin, Y.; Diederich, F.N.; Alvarez, M.M.; Anz, S.J.; Whetten, R.L., *J. Phys. Chem.* 95: 11-2 (1991)
- 91E134 Polymer bound pyrrole compounds. VI. Photophysical properties of monomeric models for polystyrene-bound porphyrins. Nonell, S.; Sese, M.L.; Martire, D.O.; Braslavsky, S.E.; Trull, F.R., *Photochem. Photobiol.* 53: 185-93 (1991)
- 91E297 Charge-transfer state and singlet oxygen ($^1\Delta_g O_2$) production in photoexcited organic molecule-molecular oxygen complexes. Kristiansen, M.; Scurlock, R.D.; Iu, K.-K.; Ogilby, P.R., *J. Phys. Chem.* 95: 5190-7 (1991)
- 91E427 Isotope-dependent quenching of singlet molecular oxygen ($^1\Delta_g$) by ground-state oxygen in several perhalogenated solvents. Afshari, E.; Schmidt, R., *Chem. Phys. Lett.* 184: 128-32 (1991)
- 91E465 The singlet oxygen and carotenoid interaction. Conn, P.F.; Schalch, W.; Truscott, T.G., *J. Photochem. Photobiol.*, B 11: 41-7 (1991)
- 91E570 Measurement of rate constants for quenching singlet oxygen with a Cypridina luciferin analog (2-methyl-6-[p-methoxyphenyl]-3,7-dihydroimidazo[1,2-a]pyrazin-3-one) and sodium azide. Mashiko, S.; Suzuki, N.; Koga, S.; Nakano, M.; Goto, T.; Ashino, T.; Mizumoto, I.; Inaba, H., *J. Biolumin. Chemilumin.* 6: 69-72 (1991)
- 91E697 Structural effects of olefins in the photooxidation with electron-accepting sensitizers. Kinetic approach to reactive intermediates. Konuma, S.; Aihara, S.; Kuriyama, Y.; Misawa, H.; Akaba, R.; Sakuragi, H.; Tokumaru, K., *Chem. Lett.* : 1897-900 (1991)
- 91F098 Self-sensitized photo-oxidation of platinum(II) complexes of 1,10-phenanthroline with 3,4-dimercaptotoluene and thiosalicylic acid by 1O_2 (S. Shukla, S.S. Kamath and T.S. Srivastava, *J. Photochem. Photobiol.*, A: Chem., 50 (1989) 199-207) - a reply. Paul, A.K.; Venkatesan, R.; Srivastava, T.S., *J. Photochem. Photobiol.*, A 56: 399 (1991)
- 91F163 Dye-sensitized photooxygenation of 2,3-dihydrofurans: Competing [2 + 2] cycloadditions and ene reactions of singlet oxygen with a rigid cyclic enol ether system. Gollnick, K.; Knutzen-Mies, K., *J. Org. Chem.* 56: 4017-27 (1991)

- 91F170 Effect of temperature on sulfide photooxidations. Evidence for a reversibly formed exciplex? Clennan, E.L.; Oolman, K.A.; Yang, K.; Wang, D.-X., *J. Org. Chem.* **56**: 4286-9 (1991)
- 91F285 Structure-activity relationship in the quenching reaction of singlet oxygen by tocopherol (vitamin E) derivatives and related phenols. Finding of linear correlation between the rates of quenching of singlet oxygen and scavenging of peroxy and phenoxyl radicals in solution. Mukai, K.; Daifuku, K.; Okabe, K.; Tanigaki, T.; Inoue, K., *J. Org. Chem.* **56**: 4188-92 (1991)
- 91F286 Kinetic implications of remote participation during photooxidation at sulfur. Clennan, E.L.; Yang, K.; Chen, X., *J. Org. Chem.* **56**: 5251-2 (1991)
- 91F300 Oxidation of substituted phenols in the environment: A QSAR analysis of rate constants for reaction with singlet oxygen. Tratnyek, P.G.; Hoigne, J., *Environ. Sci. Technol.* **25**: 1596-604 (1991)
- 91F332 The reactivity of citronellol and α -thujene with singlet oxygen. Rate constants of chemical reaction and physical quenching. El Bouamri, M.; Gorrichon, J.P.; Braun, A.M.; Oliveros, E., *Photochem. Photobiol.* **54**: 619-23 (1991)
- 91F359 Dark and photoinduced interactions between riboflavin and indole auxins. Miskoski, S.; Garcia, N.A., *Collect. Czech. Chem. Commun.* **56**: 1838-49 (1991)
- 91F405 Photo-oxygenation of styrenic estrogens: Structural analysis of 8,9-didehydro-, 6,7-didehydro-, and 9,11-didehydroestrone derivatives and their reactivity towards singlet oxygen. Malet, C.; Planas, A.; Brosa, C.; Piniella, J.F.; Rius, J., *Helv. Chim. Acta* **74**: 1412-24 (1991)
- 91F497 Sensitized photooxygenation. VI. Kinetic studies on the singlet oxygenation of 6-ethyl-3,4-dihydro-2H-pyran-5-carboxylic acid ethyl ester and 6-isopropyl-3,4-dihydro-2H-pyran-5-carboxylic acid ethyl ester. Huang, Z.; Liang, X.; Chan, Y., *Chin. J. Chem.* **9**: 506-11 (1991)
- 91F516 Determination of the chemical reaction rate constants of the quenching agents and singlet oxygen by HPLC. Li, G.; He, S.; Shi, J.; Zhang, J., *Huadong Huagong Xueyuan Xuebao* **17**: 431-6 (1991)
- 91P158 Inhibiting action of polymeric hindered amines on the oxidation of 1,4-polybutadiene by singlet oxygen. Pan, J.-Q., *Polym. Degrad. Stab.* **32**: 219-29 (1991)
- 91R017 Photochemistry of merocyanine 540: The mechanism of chemotherapeutic activity with cyanine dyes. Davila, J.; Harriman, A.; Gulliya, K.S., *Photochem. Photobiol.* **53**: 1-11 (1991)
- 91R020 In vivo and photophysical studies on photooxidative damage to lens proteins and their protection by radioprotectors. Roberts, J.E.; Kinley, J.S.; Young, A.R.; Jenkins, G.; Atherton, S.J.; Dillon, J., *Photochem. Photobiol.* **53**: 33-8 (1991)
- 91R053 Temperature effects on photosensitized processes. Gottfried, V.; Kimel, S., *J. Photochem. Photobiol., B* **8**: 419-30 (1991)
- 91R055 Activity of thiols as singlet molecular oxygen quenchers. Devasagayam, T.P.A.; Sundquist, A.R.; Di Mascio, P.; Kaiser, S.; Sies, H., *J. Photochem. Photobiol., B* **9**: 105-16 (1991)
- 91R127 Formation of 8-hydroxy(deoxy)guanosine and generation of strand breaks at guanine residues in DNA by singlet oxygen. Devasagayam, T.P.A.; Steenken, S.; Obendorf, M.S.W.; Schulz, W.A.; Sies, H., *Biochemistry* **30**: 6283-9 (1991)
- 91R206 Studies on the mechanism of the hematoporphyrin-sensitized photooxidation of 1,3-diphenylisobenzofuran in ethanol and unilamellar liposomes. Reddi, E.; Valduga, G.; Rodgers, M.A.J.; Jori, G., *Photochem. Photobiol.* **54**: 633-7 (1991)
- 91R251 Chemically induced Parkinson's disease. III. A study of a possible role of singlet molecular oxygen in Parkinson's disease. Chacon, J.N.; Truscott, T.G., *J. Photochem. Photobiol., B* **11**: 261-7 (1991)
- 91R262 Deactivation of singlet molecular oxygen by organo-selenium compounds exhibiting glutathione peroxidase activity and by sulfur-containing homologs. Scurlock, R.; Rougee, M.; Bensasson, R.V.; Evers, M.; Dereu, N., *Photochem. Photobiol.* **54**: 733-6 (1991)
- 91U026 Effects, quenching mechanisms, and kinetics of nickel chelates in singlet oxygen oxidation of soybean oil. Lee, S.-H.; Min, D.B., *J. Agric. Food Chem.* **39**: 642-6 (1991)
- 91U128 α , γ - And δ -tocopherol effects on chlorophyll-photosensitized oxidation of soybean oil. Jung, M.Y.; Choe, E.; Min, D.B., *J. Food Sci.* **56**: 807-10, 815 (1991)
- 91U180 Effects of quenching mechanisms of carotenoids on the photosensitized oxidation of soybean oil. Jung, M.Y.; Min, D.B., *J. Am. Oil Chem. Soc.* **68**: 653-8 (1991)
- 91U181 A comparative study on the susceptibilities of soybean, sunflower, and peanut oils to singlet molecular oxygen photooxidation. Neumann, M.M.; Fusero, S.N.; Garcia, N.A., *J. Am. Oil Chem. Soc.* **68**: 662-5 (1991)
- 91U201 Inhibitory effect of amino acids and dipeptides on the sensitized photooxidation of fats. Neumann, M.M.; Fusero, S.N.; Garcia, N.A., *Fette Wiss. Technol.* **93**: 453-6 (1991)
- 92A014 The contribution by diffusion to the cycloaddition reactions of singlet oxygen with furans in solution under high pressure. Okamoto, M., *J. Phys. Chem.* **96**: 245-8 (1992)
- 92A095 Yields of singlet dioxygen produced by the reaction between the excited state of tris(bipyridine)ruthenium(II) and triplet dioxygen in various solvents. Zahir, K.O.; Haim, A., *J. Photochem. Photobiol., A* **63**: 167-72 (1992)
- 92A123 Photobleaching of merocyanine 540: Involvement of singlet molecular oxygen. Singh, R.J.; Feix, J.B.; Kalyanaraman, B., *Photochem. Photobiol.* **55**: 483-9 (1992)
- 92A252 Antioxidant activity of the pyridoindole stobadine. Pulse radiolytic characterization of one-electron-oxidized stobadine and quenching of singlet molecular oxygen. Steenken, S.; Sundquist, A.R.; Jovanovic, S.V.; Crockett, R.; Sies, H., *Chem. Res. Toxicol.* **5**: 355-60 (1992)
- 92A325 Reaction of singlet molecular oxygen with tetramethyl-*o*-xylylene. Redmond, R.W.; Harwig, C.W.; Scaiano, J.C., *J. Photochem. Photobiol., A* **68**: 255-9 (1992)
- 92A386 Singlet oxygen quenching by tertiary alkyl ethers. Boneva, M.; Ivanov, S.K.; Oliveros, E.; Braun, A.M., *J. Photochem. Photobiol., A* **68**: 343-51 (1992)
- 92D008 Photooxidation of organic and inorganic substrates during UV photolysis of nitrite anion in aqueous solution. Bilski, P.; Chignell, C.F.; Szychlinski, J.; Borkowski, A.; Oleksy, E.; Reszka, K., *J. Am. Chem. Soc.* **114**: 549-56 (1992)
- 92D048 Spectroscopic studies of cutaneous photosensitizing agents. XVII. Benzanthrone. Dabestani, R.; Sik, R.H.; Motten, A.G.; Chignell, C.F., *Photochem. Photobiol.* **55**: 533-9 (1992)
- 92D227 Identification by ESR spectroscopy and spectrophotometry of singlet oxygen, formed during autoxidation of catecholamines. Lichszeld, K.; Michalska, T.; Kruk, I., *Z. Phys. Chem. (Munich)* **175**: 117-22 (1992)
- 92E220 Quenching of singlet oxygen in solid organic polymers. Ogilby, P.R.; Dillon, M.P.; Kristiansen, M.; Clough, R.L., *Macromolecules* **25**: 3399-405 (1992)
- 92E225 The quenching of singlet molecular oxygen by carnosine and anserine in aqueous solution. Egorov, S.Yu.; Kurella, E.G.; Boldyrev, A.A.; Krasnovsky, A.A., Jr., *Bioorg. Khim.* **18**: 142-4 (1992)
- 92E256 Collisional deactivation of $O_2^{(1)\Delta_g}$ by solvent molecules. Comparative experiments with $^{16}O_2$ and $^{18}O_2$. Schmidt, R.; Afshari, E., *Ber. Bunsenges. Phys. Chem.* **96**: 788-94 (1992)
- 92E274 Reaction of singlet oxygen with vinyl-substituted porphyrins. A kinetic study by laser flash photolysis. Aveline, B.; Delgado, O.; Brault, D., *J. Chem. Soc., Faraday Trans.* **88**: 1971-6 (1992)
- 92E555 Photophysics and photochemistry of 22π and 26π acetylenecumulene porphyrinoids. Martire, D.O.; Jux, N.; Aramendia, P.F.; Negri, R.M.; Lex, J.; Braslavsky, S.E.; Schaffner, K.; Vogel, E., *J. Am. Chem. Soc.* **114**: 9969-78 (1992)
- 92E654 Assay of lycopene and other carotenoids as singlet oxygen quenchers. Di Mascio, P.; Sundquist, A.R.; Devasagayam, T.P.A.; Sies, H., *Methods Enzymol.* **213**: 429-38 (1992)

- 91F063 Quantum yields of singlet-oxygen production by some natural quinoid fungal metabolites and derivatives. Aminian-Saghafi, T.; Nasini, G.; Caronna, T.; Braun, A.M.; Oliveros, E., *Helv. Chim. Acta* **75**: 531-8 (1992)
- 91F103 Photooxidation of 1,5-dithiacyclooctane. A novel C-S bond cleavage. Sheu, C.; Foote, C.S.; Gu, C.-L., *J. Am. Chem. Soc.* **114**: 3015-21 (1992)
- 91F104 Comparison of the photooxidations of 1,5-dithiacyclooctane, 1,4-dithiane, and pentamethylene sulfide. Another example of remote participation during photooxidation at sulfur? Clennan, E.L.; Wang, D.-X.; Yang, K.; Hodgson, D.J.; Oki, A.R., *J. Am. Chem. Soc.* **114**: 3021-7 (1992)
- 91F181 Influence of the ionic strength on $O_2(^1\Delta_g)$ quenching by azide. Rubio, M.A.; Martire, D.O.; Braslavsky, S.E.; Lissi, E.A., *J. Photochem. Photobiol., A* **66**: 153-7 (1992)
- 92F189 Tellurapyrylium dyes as catalysts for oxidations with hydrogen peroxide and as scavengers of singlet oxygen. Dihydroxytelluranes as mild oxidizing agents. Detty, M.R.; Gibson, S.L., *Organometallics* **11**: 2147-56 (1992)
- 92F192 Photodynamic degradation of vitamin E induced by psoralens. Costantini, C.; d'Ischia, M.; Napolitano, A.; Misuraca, G.; Prota, G., *Biochim. Biophys. Acta* **1116**: 291-6 (1992)
- 92F210 Investigation of 1O_2 generation by water-soluble α -diimine complexes of platinum(II) and palladium(II) with 3,4-dihydroxybenzoic acid as photosensitizers. Anbalagan, V.; Srivastava, T.S., *J. Photochem. Photobiol., A* **66**: 345-53 (1992)
- 92F225 Remote participation during photooxidation at sulfur. Evidence for sulfurane intermediates. Clennan, E.L.; Yang, K., *J. Org. Chem.* **57**: 4477-87 (1992)
- 92F251 Photosensitized oxygenation of olefins with zinc tetraphenylporphine-1,4-diazabicyclo[2.2.2]octane complex. Sakuragi, M.; Ichimura, K.; Sakuragi, H., *Bull. Chem. Soc. Jpn.* **65**: 1944-9 (1992)
- 92F270 Photophysical properties and photo-oxygenation of 2-(4-ethoxycarbonylphenyl)-1,8-dihydropyrrolo[3',2':3,4]cyclopenta[1,2-b]pyridine (EPCP). Dunsbach, R.; Brauer, H.-D.; Schmidt, R.; Wang, J.; Gossauer, A., *J. Photochem. Photobiol., A* **67**: 227-35 (1992)
- 92F402 Physico-chemical modeling of the role of free radicals in photodynamic therapy. I. Utilization of quantum yield data of singlet oxygen formation for the study of the interaction between excited photosensitizer and stable free radicals. Vidoczy, T.; Elzemzam, S.; Gai, D., *Biochem. Biophys. Res. Commun.* **169**: 1548-52 (1992)
- 92F465 Kinetic and thermodynamic aspects of the dye-sensitised photo-oxidation of bioresmethrin and related compounds. Clements, P.; Wells, C.H.J., *Pestic. Sci.* **35**: 305-8 (1992)
- 92F491 Mechanism of the 9,10-dicyanoanthracene-sensitized photooxidation of N,N-dibenzylhydroxylamine in polar and non-polar solvents. Sakurai, T.; Uematsu, Y.; Tanaka, O.; Inoue, H., *J. Chem. Soc., Perkin Trans. 2* : 2163-7 (1992)
- 92F505 Singlet molecular oxygen quantum yield measurements of some porphyrins and metalloporphyrins. Venkatesan, R.; Periasamy, N.; Srivastava, T.S., *Proc. - Indian Acad. Sci., Chem. Sci.* **104**: 713-22 (1992)
- 92M228 The influence of some biological antioxidants on the light emission from the oxytetracycline oxidation reaction. Kruk, I.; Lichszeld, K.; Michalska, T.; Paraskevas, S.M., *Toxicol. Environ. Chem.* **35**: 167-73 (1992)
- 92N179 Singlet oxygen chemistry. VI. The kinetics of photooxidation of 2,5-diphenylbenzofuran (DPBF) induced by hypocrellin A (HA) in micelles. Jiang, X.-F.; Wang, D.-Y., *Acta Chim. Sin. (Huaxue Xuebao)* **50**: 391-6 (1992)
- 92N201 Generation and reactions of singlet oxygen photosensitized by hypericin in dipalmitoylphosphatidylcholine vesicles. Bourig, H.; Eloy, D.; Jardon, P., *J. Chim. Phys. Phys.-Chim. Biol.* **89**: 1391-411 (1992)
- 92N305 Photocatalytic transformation of 2,5-furandimethanol in aqueous ZnO suspensions. Richard, C.; Martre, A.M.; Boule, P., *J. Photochem. Photobiol., A* **66**: 225-34 (1992)
- 92N330 Sensitized photooxidation of mononitrophenols in micellar media. Luiz, M.; Gutierrez, M.I.; Bocco, G.; Bertolotti, S.G.; Garcia, N.A., *Toxicol. Environ. Chem.* **35**: 115-23 (1992)
- 92R054 Synthetic carotenoids, novel polyene polyketones and new capsorubin isomers as efficient quenchers of singlet molecular oxygen. Devasagayam, T.P.A.; Werner, T.; Ippendorf, H.; Martin, H.D.; Sies, H., *Photochem. Photobiol.* **55**: 511-4 (1992)
- 92R076 Quenching of singlet molecular oxygen by phthalocyanines and naphthalocyanines. Krasnovsky, A.A.Jr.; Rodgers, M.A.J.; Galpern, M.G.; Rihter, B.; Kenney, M.E.; Lukjanetz, E.A., *Photochem. Photobiol.* **55**: 691-6 (1992)
- 92R107 A laser flash photolysis study of the triplet states of the red and the blue forms of Blepharisma japonicum pigment. Ghetti, F.; Checucci, G.; Lenci, F.; Heelis, P.F., *J. Photochem. Photobiol., B* **13**: 315-21 (1992)
- 92R313 A proposed function for spermine and spermidine: Protection of replicating DNA against damage by singlet oxygen. Khan, A.U.; Mei, Y.-H.; Wilson, T., *Proc. Natl. Acad. Sci. U.S.A.* **89**: 11426-7 (1992)
- 93A017 Volume of activation for the radiationless deactivation of singlet oxygen in aromatic hydrocarbons. Okamoto, M.; Tanaka, F., *J. Phys. Chem.* **97**: 177-80 (1993)
- 93A050 Reactions of organometallic complexes with singlet oxygen. Photooxidation of Vaska's complex. Selke, M.; Foote, C.S., *J. Am. Chem. Soc.* **115**: 1166-7 (1993)
- 93A103 Hydroxyanthraquinones as sensitizers of electron-transfer-induced photo-oxygenation reactions. Gollnick, K.; Held, S., *J. Photochem. Photobiol., A* **70**: 135-45 (1993)
- 93A244 Production of singlet oxygen ($^1\Delta_g O_2$) by 9,10-dicyanoanthracene and acridine: Quantum yields in acetonitrile. Scurlock, R.D.; Ogilby, P.R., *J. Photochem. Photobiol., A* **72**: 1-7 (1993)
- 93A326 Antioxidant activity of flavonoids: Efficiency of singlet oxygen ($^1\Delta_g$) quenching. Tournaire, C.; Croux, S.; Maurette, M.T.; Beck, I.; Hocquaux, M.; Braun, A.M.; Oliveros, E., *J. Photochem. Photobiol., B* **19**: 205-15 (1993)
- 93A380 Photochromic reactions involving palladium(II) octabutoxynaphthalocyanine and molecular oxygen. Rihter, B.D.; Kenney, M.E.; Ford, W.E.; Rodgers, M.A.J., *J. Am. Chem. Soc.* **115**: 8146-52 (1993)
- 93D154 The photooxidation of diethylhydroxylamine by Rose Bengal in micellar and nonmicellar aqueous solutions. Bilski, P.; Motten, A.G.; Bilksa, M.; Chignell, C.F., *Photochem. Photobiol.* **58**: 11-8 (1993)
- 93E090 Nitroxyl free radical enhancement of the forbidden $O_2(^3\Sigma_g^-) \rightarrow O_2(^1\Delta_g)$ radiative transition in chlorinated hydrocarbon solvents. Belford, R.E.; Seely, G.R.; Gust, D.; Moore, T.A.; Moore, A.; Cherepy, N.J.; Ekbundit, S.; Lewis, J.E.; Lin, S.H., *J. Photochem. Photobiol., A* **70**: 125-33 (1993)
- 93E091 $O_2(^1\Delta_g)$ production and quenching by C_{60} and C_{70} . Black, G.; Dunkle, E.; Dorko, E.A.; Schlie, L.A., *J. Photochem. Photobiol., A* **70**: 147-51 (1993)
- 93E183 Magnetic field effects on excited-state oxygen-organic molecule interactions. Ogilby, P.R.; Sanetra, J., *J. Phys. Chem.* **97**: 4689-94 (1993)
- 93E248 Upconversion of the singlet oxygen 1269 nm emission to the visible. Allan, D.; Gorman, A.A.; Hamblett, I.; Hodgson, B.W.; Lambert, C., *Photochem. Photobiol.* **57**: 893-4 (1993)
- 93E301 Time-resolved measurements of singlet oxygen dimol-sensitized luminescence. Krasnovsky, A.A.Jr.; Foote, C.S., *J. Am. Chem. Soc.* **115**: 6013-6 (1993)
- 93E520 Synthesis and photoproperties of silicon phthalocyanines and silicon naphthalocyanines. Wen, T.-C.; Liu, M.-E.; Young, S.-H.; Lee, Y.-S.; Chen, K.-M.; Kuo, C.-N., *J. Chin. Chem. Soc. (Taipei)* **40**: 141-7 (1993)
- 93E536 Singlet molecular oxygen [$O_2(^1\Delta_g)$] production and quenching by hydroxybiphenyls. Martire, D.O.; Evans, C.; Bertolotti, S.G.; Braslavsky, S.E.; Garcia, N.A., *Chemosphere* **26**: 1691-701 (1993)

- 93E539 Reactions of singlet oxygen with organometallic complexes. 2. Formation of a metastable rhodium-dioxygen complex. Selke, M.; Foote, C.S.; Karney, W.L., *Inorg. Chem.* **32**: 5425-6 (1993)
- 93E684 Mechanism of quenching of singlet molecular oxygen luminescence by pyrroles. Venedikov, E.A.; Titova, Yu.V., *Biofizika* **38**: 1031-4 (1993)
- 93F069 A kinetic study of the photodynamic properties of the xanthene dye merbromin (mercurochrome) and its aggregates with amino acids in aqueous solution. Martinez, G.; Bertolotti, S.G.; Zimerman, O.E.; Martire, D.O.; Braslavsky, S.E.; Garcia, N.A., *J. Photochem. Photobiol.*, **B** **17**: 247-55 (1993)
- 93F128 High reactivity of heterocordianthrones (HCDs) in photoperoxidation and thermal stability of their endoperoxides. Use of HCD as a photosensitizer under sunlight. Motoyoshiya, J.; Masunaga, T.; Harumoto, D.; Ishiguro, S.; Narita, S.; Hayashi, S., *Bull. Chem. Soc. Jpn.* **66**: 1166-71 (1993)
- 93F160 Reaction of singlet oxygen with disiliranes: Dioxygen insertion into Si-Si σ bonds. Ando, W.; Kako, M.; Akasaka, T.; Nagase, S., *Organometallics* **12**: 1514-22 (1993)
- 93F191 The aerobic riboflavin-sensitized photodecomposition of tryptophan, tryptamine and indole acetic acid. Ground-state and photopromoted flavin-indole interactions. Martinez, G.; Zimerman, O.E.; Garcia, N.A., *Collect. Czech. Chem. Commun.* **58**: 1299-308 (1993)
- 93F207 The formation and removal of singlet ($^1\Delta_g$) oxygen in solid organic polymers: Events that may influence polymer photodegradation. Ogilby, P.R.; Kristiansen, M.; Martire, D.O.; Scurlock, R.D.; Taylor, V.L.; Clough, R.L., *Polym. Prepr. (Am. Chem. Soc., Div. Polym. Chem.)* **34**: 133-4 (1993)
- 93F231 Formation of 2,5-dihydroxybenzoic acid during the reaction between $^1\text{O}_2$ and salicylic acid: Analysis by ESR oximetry and HPLC with electrochemical detection. Kalyanaraman, B.; Ramanujam, S.; Singh, R.J.; Joseph, J.; Feix, J.B., *J. Am. Chem. Soc.* **115**: 4007-12 (1993)
- 93F327 Mechanistic study of the photooxidation of squalene sensitized with 2,2'-5',2''-terthiophene and 2,2'-bithiophene. Fujita, H.; Tokiwa, K.; Sayama, K.; Mori, H.; Sasaki, M., *Bull. Chem. Soc. Jpn.* **66**: 2960-4 (1993)
- 93F328 Solvent effect on the reactivity of monosubstituted phenols towards singlet molecular oxygen ($\text{O}_2[^1\Delta_g]$) in alkaline media. Luiz, M.; Gutierrez, M.I.; Bocco, G.; Garcia, N.A., *Can. J. Chem.* **71**: 1247-52 (1993)
- 93F361 Dimethyl 3,3'-(4-methyl-1,3-naphthylene)dipropionate as a singlet oxygen trap in biological systems. Mueller, K.; Ziereis, K., *Arch. Pharm. (Weinheim, Ger.)* **326**: 369-71 (1993)
- 93F367 Sensitized photooxygenations of 3-vinylindole derivatives. Zhang, X.; Khan, S.I.; Foote, C.S., *J. Org. Chem.* **58**: 7839-47 (1993)
- 93F368 Photochemical elimination of singlet oxygen from oxodiperoxomolybdenum(VI)-(hexamethyl phosphoric triamide, water). An adiabatic photoreaction originating from an upper excited electronic state. Seip, M.; Brauer, H.-D., *J. Photochem. Photobiol.*, **A** **76**: 1-5 (1993)
- 93F413 On the $\text{O}_2[^1\Delta_g]$ -mediated photooxidative behaviour of tripeptide glycyl-tyrosyl-alanine in alkaline medium. A kinetic study. Miskosky, S.; Bertolotti, S.G.; Garcia, N.A.; Arguello, G.A., *Amino Acids* **4**: 101-10 (1993)
- 93M161 Kinetic study of the quenching reaction of singlet oxygen by biological hydroquinones and related compounds. Mukai, K.; Itoh, S.; Daifuku, K.; Morimoto, H.; Inoue, K., *Biochim. Biophys. Acta* **1183**: 323-6 (1993)
- 93N051 Quenching of singlet molecular oxygen ($^1\Delta_g\text{O}_2$) in silica gel-solvent heterogeneous system. II. A direct time-resolved study. Iu, K.-K.; Thomas, J.K., *J. Photochem. Photobiol.*, **A** **71**: 55-60 (1993)
- 93R059 Influence of the peptide bond on the singlet molecular oxygen-mediated ($\text{O}_2[^1\Delta_g]$) photooxidation of histidine and methionine dipeptides. A kinetic study. Miskoski, S.; Garcia, N.A., *Photochem. Photobiol.* **57**: 447-52 (1993)
- 93R131 Temporal and spectral separation of singlet oxygen luminescence from near infrared emitting photosensitizers. Naether, D.U.; Gilchrist, J.R.; Gensch, T.; Roeder, B., *Photochem. Photobiol.* **57**: 1056-9 (1993)
- 93R231 A pulsed laser and pulse radiolysis study of amphiphilic chlorophyll derivatives with PDT activity toward malignant melanoma. Fiedor, L.; Gorman, A.A.; Hamblett, I.; Rosenbach-Belkin, V.; Salomon, Y.; Scherz, A.; Tregub, I., *Photochem. Photobiol.* **58**: 506-11 (1993)
- 93R310 Photobiological activities of 1,6-dioxypyrene in pro- and eukaryotic cells. Averbeck, D.; Polasa, K.; Buisson, J.-P.; Bensasson, R.V.; Rougee, M.; Cadet, J.; Ravanat, J.L.; Perin, F.; Vigny, P.; Demerseman, P., *Mutat. Res.* **287**: 165-79 (1993)
- 93U041 Photooxygenation of 3-hydroxyflavone and molecular design of the radiation-hard scintillator based on the excited-state proton transfer. Chou, P.-T.; Martinez, M.L., *Radiat. Phys. Chem.* **41**: 373-8 (1993)
- 93Y025 Interaction of singlet and triplet excited states with the same quencher in micelles. Daraio, M.E.; Aramendia, P.F.; San Roman, I., *Chem. Phys. Lett.* **204**: 415-9 (1993)
- 94A113 Reactivity of singlet oxygen toward amino acids and peptides. Michaeli, A.; Feitelson, J., *Photochem. Photobiol.* **59**: 284-9 (1994)
- 94A126 Photochemical elimination of singlet oxygen from iridium peroxy complexes. Seip, M.; Brauer, H.-D., *J. Photochem. Photobiol.*, **A** **79**: 19-24 (1994)
- 94E153 Curcumin-derived transients: A pulsed laser and pulse radiolysis study. Gorman, A.A.; Hamblett, I.; Srinivasan, V.S.; Wood, P.D., *Photochem. Photobiol.* **59**: 389-98 (1994)
- 94F004 Spectral and photochemical behaviour of newly synthesized 2,2'-dipyridylamine complexes of Pt(II) and Pd(II) with various dioxolenes. Anbalagan, V.; Srivastava, T.S., *J. Photochem. Photobiol.*, **A** **77**: 141-8 (1994)
- 94F024 Photooxidation of sulfenic acid derivatives. 1. Sulfenamides. The surprising behavior of a new class of photooxidation substrates. Clennan, E.L.; Zhang, H., *J. Am. Chem. Soc.* **116**: 809-10 (1994)
- 94F069 Spectral and photochemical properties of curcumin. Chignell, C.F.; Bilski, P.; Reszka, K.J.; Motten, A.G.; Sik, R.H.; Dahl, T.A., *Photochem. Photobiol.* **59**: 295-302 (1994)
- 94F095 Influence of the nuclear substitution on the sensitized photooxidation of model compounds for phenolic-type pesticides. Bocco, G.; Luiz, M.; Gutierrez, M.I.; Garcia, N.A., *J. Prakt. Chem./Chem.-Ztg.* **336**: 243-6 (1994)
- 94R058 Reaction and quenching of singlet molecular oxygen with esters of polyunsaturated fatty acids. Tanielian, C.; Mechlin, R., *Photochem. Photobiol.* **59**: 263-8 (1994)

13. Molecular Formula Index

Br ⁻	Bromide ion 14.4	C ₄ H ₃ BrO	2-Bromofuran 5.21
CCl ₂ D ₂	Dichloromethane- <i>d</i> ₂ 17.17	C ₄ H ₄ O	Furan 5.14
CCl ₃ D	Chloroform- <i>d</i> 17.10	C ₄ H ₄ S	Thiophene 13.167
CHCl ₃	Chloroform 17.9	C ₄ H ₅ N	Pyrrole 6.59
CH ₂ Cl ₂	Dichloromethane 17.16	C ₄ H ₅ N ₃ O	Cytosine 16.20
CH ₄ N ₂ O	Urea 17.77	C ₄ H ₆ CoO ₄	Cobalt(II) acetate 12.126
CH ₄ N ₂ S	Thiourea 13.174	C ₄ H ₆ N ₂	4-Methylimidazole 6.7
CH ₄ O	Methanol 17.40	C ₄ H ₆ NiO ₄	Nickel(II) acetate 12.161
C ₂ D ₃ N	Acetonitrile- <i>d</i> ₃ 17.5	C ₄ H ₆ O	(E)-2-Butenal 2.44 2,3-Dihydrofuran 5.29
C ₂ H ₃ N	Acetonitrile 17.4	C ₄ H ₆ O ₂	(E)-2-Butenoic acid 2.57 3-Butenoic acid 2.59 1,4-Dioxene 2.183 Methacrylic acid 2.252
C ₂ H ₄ O ₂	Acetic acid 17.1	C ₄ H ₈	2-Butene 2.46 (E)-2-Butene 2.47 (Z)-2-Butene 2.48 2-Methylpropene 2.337
C ₂ H ₅ I	Iodoethane 17.28	C ₄ H ₈ N ₂ O	N-Allylurea 17.78
C ₂ H ₅ NO ₂	Glycine 10.15	C ₄ H ₈ N ₂ O ₂	Dimethylglyoxime 15.19
C ₂ H ₅ OS ⁻	2-Hydroxyethylsulfide ion 13.96	C ₄ H ₈ N ₂ O ₃	Glycylglycine 10.16
C ₂ H ₅ O ₃ S ₂ ⁻	2-Mercaptoethanesulfonate ion 13.92	C ₄ H ₈ N ₂ S	Allylthiourea 13.175
C ₂ H ₆ N ₂ S	<i>N</i> -Methylthiourea 13.176	C ₄ H ₈ O	Tetrahydrofuran 17.68 Ethyl vinyl ether 2.224
C ₂ H ₆ O	Ethanol 17.29	C ₄ H ₈ OS	1,4-Thioxane 13.177
C ₂ H ₆ OS	Dimethyl sulfoxide 13.81	C ₄ H ₈ O ₂	Butyric acid 17.6 1,4-Dioxane 17.20
C ₂ H ₆ S	2-Mercaptoethanol 13.95	C ₄ H ₈ O ₆ S ₄ ²⁻	Dimethylsulfide 13.91
C ₂ H ₆ S	Dimethyl sulfide 13.80	C ₄ H ₈ S	Tetrahydrothiophene 13.168
C ₂ H ₇ NS	Cysteamine 13.93	C ₄ H ₈ S ₂	1,4-Dithiane 13.88
C ₂ O ₄ ²⁻	Oxalate ion 17.48	C ₄ H ₉ N	Pyrrolidine 8.96
C ₃ D ₆ O	Acetone- <i>d</i> ₆ 17.3	C ₄ H ₉ NO	2-Methyl-2-nitrosopropane 15.50
C ₃ H ₄ N ₂	Imidazole 6.6	C ₄ H ₁₀ N ₂	Piperazine 8.51
C ₃ H ₄ O	Acrolein 2.1	C ₄ H ₁₀ O	1-Butanol 17.8 2-Methyl-2-propanol 17.62
C ₃ H ₄ O ₂	1,3-Dioxole 2.199	C ₄ H ₁₀ O ₂ S	2,2'-Thiodiethanol 13.164
C ₃ H ₄ O ₄	Malonic acid 17.39	C ₄ H ₁₀ O ₂ S ₂	Dithioerythritol 13.49 Dithiothreitol 13.50
C ₃ H ₆ NO ₂ S ⁻	Cysteine, negative ion 13.68	C ₄ H ₁₀ S	1-Butanethiol 13.51 Diethyl sulfide 13.76
C ₃ H ₆ NS ₂ ⁻	Dimethylthiocarbamate ion 13.79	C ₄ H ₁₀ S ₂	Diethyl disulfide 13.75
C ₃ H ₆ O	Acetone 17.2	C ₄ H ₁₁ N	Butylamine 8.12 <i>tert</i> -Butylamine 8.15 Diethylamine 8.27 Isobutylamine 8.85
C ₃ H ₆ S	Methylthiirane 13.152	C ₄ H ₁₁ NO	<i>O,N</i> -Diethylhydroxylamine 15.21 <i>N,N</i> -Diethylhydroxylamine 15.23
C ₃ H ₇ NO	Acetone oxime 15.2	C ₄ H ₁₁ NO ₃	2-Amino-2-(hydroxymethyl)-1,3-propane-diol 17.59
C ₃ H ₇ NO ₂	<i>N,N</i> -Dimethylformamide 17.31		
C ₃ H ₇ NO ₂	Alanine 10.1		
	β-Alanine 10.2		
C ₃ H ₇ NO ₂ S	Cysteine 13.67		
C ₃ H ₈ O	2-Propanol 17.61		
C ₃ H ₉ N	Isopropylamine 8.28		
	Trimethylamine 8.47		
	Propylamine 8.84		
C ₃ H ₉ NOS	3-Amino-2-hydroxypropanethiol 13.131		
C ₃ H ₉ O ₃ P	Trimethyl phosphite 17.73, 17.74		
C ₄ D ₁₂ N ₂	Tetra(methyl- <i>d</i> ₃)hydrazine 8.42		
C ₄ H ₂ Br ₂ O	2,5-Dibromofuran 5.27		

C ₄ H ₁₂ NO ⁺	N,N-Diethylhydroxylammonium ion	15.24	C ₅ H ₁₃ NO	3-Dimethylamino-1-propanol	8.82
C ₄ H ₁₂ N ₂	Tetramethylhydrazine	8.41	C ₅ H ₁₃ NO ₂	N-Methyldiethanolamine	8.46
C ₅ H ₃ NO	2-Furanitrile	5.63	C ₅ H ₁₄ NO ₆ P	Phosphatidylethanolamine	17.52
C ₅ H ₃ N ₄ O ₃ ⁻	Urate ion	16.64	C ₅ H ₁₄ N ₂ S	2-[(3-Aminopropyl)amino]ethanethiol	
C ₅ H ₄ OS	Pyran-4-thione	13.138			13.94
C ₅ H ₄ O ₂	2-Furaldehyde	5.9	C ₆ Cl ₅ O ⁻	Pentachlorophenoxyde ion	4.179
C ₅ H ₄ O ₃	2-Furancarboxylic acid	5.54	C ₆ D ₆	Benzene-d ₆	3.27
C ₅ H ₄ S ₂	Thiopyran-4-thione	13.172	C ₆ D ₁₂	2,3-Dimethyl-2-butene-d ₁₂	2.54
C ₅ H ₅ N	Pyridine	6.40		Cyclohexane-d ₁₂	17.14
C ₅ H ₅ N ₅	Adenine	16.63	C ₆ HCl ₅ O	Pentachlorophenol	4.140
C ₅ H ₅ N ₅ O	Guanine	16.35	C ₆ H ₂ Cl ₃ O ⁻	2,4,6-Trichlorophenoxyde ion	4.182
C ₅ H ₆	Cyclopentadiene	2.140	C ₆ H ₂ F ₄ O	2,3,5,6-Tetrafluorophenol	4.143
C ₅ H ₆ N ₂ O ₂	Thymine	16.79	C ₆ H ₃ Cl ₂ O ⁻	2,4-Dichlorophenoxyde ion	4.159
C ₅ H ₆ O	2-Methylfuran	5.46		2,6-Dichlorophenoxyde ion	4.160
C ₅ H ₆ O ₂	2-Methoxyfuran	5.42	C ₆ H ₃ Cl ₃ O	2,4,6-Trichlorophenol	4.146
	Furfuryl alcohol	5.78	C ₆ H ₃ N ₂ O ₅ ⁻	2,4-Dinitrophenoxide ion	4.164
C ₅ H ₆ O ₅	2-Ketoglutaric acid	17.51	C ₆ H ₄ BrO ⁻	4-Bromophenoxyde ion	4.153
C ₅ H ₇ N	N-Methylpyrrole	6.64	C ₆ H ₄ ClO ⁻	2-Chlorophenoxyde ion	4.155
C ₅ H ₇ NO	Furylamine	5.79		3-Chlorophenoxyde ion	4.156
C ₅ H ₈	Isoprene	2.42		4-Chlorophenoxyde ion	4.157
	1-Methylcyclobutene	2.98	C ₆ H ₄ Cl ₂	1,3-Dichlorobenzene	3.35
	Cyclopentene	2.156		1,4-Dichlorobenzene	3.36
	(E)-Piperylene	2.328	C ₆ H ₄ Cl ₂ O	2,4-Dichlorophenol	4.113
	(Z)-Piperylene	2.329		2,6-Dichlorophenol	4.114
C ₅ H ₈ NO ₃ S ⁻	N-Acetylcysteine, negative ion	13.70	C ₆ H ₄ FO ⁻	4-Fluorophenoxyde ion	4.165
	2-Mercaptopropionylglycine, negative ion	13.102	C ₆ H ₄ JO ⁻	4-Iodophenoxyde ion	4.168
C ₅ H ₈ O	2,3-Dihydro-5-methylfuran	5.31	C ₆ H ₄ NO ₃ ⁻	2-Nitrophenoxide ion	4.176
	3,4-Dihydropyran	5.90		3-Nitrophenoxide ion	4.177
C ₅ H ₈ O ₂	(E)-2-Methyl-2-butenoic acid	2.58		4-Nitrophenoxide ion	4.178
	4-Pentenoic acid	2.315	C ₆ H ₄ N ₂ O ₅	2,4-Dinitrophenol	4.123
C ₅ H ₉ NO ₃ S	N-Acetyl-L-cysteine	13.69		2,5-Dinitrophenol	4.124
	2-Mercaptopropionylglycine	13.101		2,6-Dinitrophenol	4.125
C ₅ H ₉ N ₃	Histamine	6.5	C ₆ H ₄ O ₂	1,4-Benzoquinone	16.12
C ₅ H ₁₀	2-Methyl-2-butene	2.56	C ₆ H ₄ O ₂ ²⁻	1,4-Benzenediol dianion	4.13
	1-Pentene	2.302	C ₆ H ₄ O ₃	2,5-Furandicarboxaldehyde	5.59
	(E)-2-Pentene	2.304	C ₆ H ₅ Br	Bromobenzene	3.30
	(Z)-2-Pentene	2.305	C ₆ H ₅ Cl	Chlorobenzene	3.31
C ₅ H ₁₀ O	2-Methoxy-2-butene	2.55	C ₆ H ₅ ClO	2-Chlorophenol	4.106
	3-Methyl-2-butene-1-ol	2.60		3-Chlorophenol	4.107
C ₅ H ₁₀ O ₂	Valeric acid	17.79		4-Chlorophenol	4.108
C ₅ H ₁₀ S	Thiane	13.145	C ₆ H ₅ ClO ₂	4-Chlororesorcinol	4.11
C ₅ H ₁₁ N	Piperidine	8.53		Chloro-1,4-benzenediol	4.20
C ₅ H ₁₁ NO ₂ S	Methionine	13.108	C ₆ H ₅ NO	5-Methyl-2-furanitrile	5.64
	Penicillamine	13.116	C ₆ H ₅ NO ₂	Nitrobenzene	3.47
C ₅ H ₁₂	Pentane	17.50	C ₆ H ₅ NO ₃	2-Nitrophenol	4.137
C ₅ H ₁₂ O	1,1-Dimethylethyl methyl ether	17.18		3-Nitrophenol	4.138
C ₅ H ₁₂ S	Butyl methyl sulfide	13.57		4-Nitrophenol	4.139
			C ₆ H ₅ O ⁻	Phenoxyde ion	4.150

C ₆ H ₆	Benzvalene 2.3 Benzene 3.26	C ₆ H ₁₀ O	2,3-Dihydro-4,5-dimethylfuran 5.30 1-Methoxycyclopentene 2.160
C ₆ H ₆ D ₆	(E)-2,3-Dimethyl-2-butene- <i>d</i> ₆ 2.52 (Z)-2,3-Dimethyl-2-butene- <i>d</i> ₆ 2.53	C ₆ H ₁₀ OS	2-Methylthiacyclohex-2-ene-1-oxide 13.143
C ₆ H ₆ O	Phenol 4.92 2-Vinylfuran 5.53	C ₆ H ₁₀ O ₃	3,3-Dimethyl-2-oxobutyric acid 17.7
C ₆ H ₆ O ₂	1,2-Benzenediol 4.4 1,3-Benzenediol 4.10 1,4-Benzenediol 4.12 5-Methylfurfural 5.13 2-Acetyl furan 5.15	C ₆ H ₁₁ NO	5,5-Dimethyl-1-pyrroline-1-oxyl 15.56
C ₆ H ₆ O ₃	5-(Hydroxymethyl)furfural 5.11 Methyl 2-furoate 5.57	C ₆ H ₁₁ N ₃ O ₄	Glycylglycylglycine 10.17
C ₆ H ₇ N	Aniline 9.1	C ₆ H ₁₂	2,3-Dimethyl-1-butene 2.45 2,3-Dimethyl-2-butene 2.51 Cyclohexane 17.13 2-Hexene 2.244 (E)-3-Hexene 2.245 (Z)-3-Hexene 2.246
C ₆ H ₇ NO	4-Aminophenol 4.95		2-Methyl-2-pentene 2.307 3-Methyl-2-pentene 2.308 (E)-3-Methyl-2-pentene 2.309 (Z)-3-Methyl-2-pentene 2.310 (E)-4-Methyl-2-pentene 2.311 (Z)-4-Methyl-2-pentene 2.312
C ₆ H ₇ O ₆ ⁻	Ascorbate ion 16.6	C ₆ H ₁₂ BiN ₂ S ₄	Bis(dimethyldithiocarbamato)bismuth(II) 12.55
C ₆ H ₈	1,3-Cyclohexadiene 2.103	C ₆ H ₁₂ N ₂	Acetone azine 15.3 1,4-Diazabicyclo[2.2.2]octane 8.19 Tetrahydropyrazolo[1,2- <i>a</i>]pyrazole 8.89
C ₆ H ₈ Cl ₂ N ₂ O ₂	<i>trans</i> -1,4-Dichloro-1,4-dinitrosocyclohexane 15.16 <i>cis</i> -1,4-Dichloro-1,4-dinitrosocyclohexane 15.17 1,4-Dichloro-2,3-diazabicyclo[2.2.2]oct-2-ene-2,3-dioxide 15.18	C ₆ H ₁₂ N ₂ NiS ₄	Bis(dimethyldithiocarbamato)nickel(II) 12.56
C ₆ H ₈ N ⁺	1-Methylpyridinium 6.54	C ₆ H ₁₂ N ₄	Hexamethylenetetramine 8.32
C ₆ H ₈ N ₂	<i>o</i> -Phenylenediamine 9.25 Phenylhydrazine 9.32	C ₆ H ₁₂ O	(E)-3-Methyl-2-penten-1-ol 2.319 (Z)-3-Methyl-2-penten-1-ol 2.320 4-Methyl-3-penten-2-ol 2.323 1-Ethoxy-2-methyl-1-propene 2.336
C ₆ H ₈ O	2,4-Dimethylfuran 5.33 2,5-Dimethylfuran 5.34	C ₆ H ₁₂ OS ₂	1,5-Dithiacyclooctane 1-oxide 13.87
C ₆ H ₈ O ₂	Furfuryl methyl ether 5.43 α -Methylfurfuryl alcohol 5.69 2,3-Dioxabicyclo[2.2.2]oct-5-ene 2.182	C ₆ H ₁₂ O ₂	Hexanoic acid 17.38 Ketene diethyl acetal 2.221 (E)-1,2-Diethoxyethene 2.222 (Z)-1,2-Diethoxyethene 2.223
C ₆ H ₈ O ₃	2,5-Bis(hydroxymethyl)furan 5.18	C ₆ H ₁₂ O ₆	Glucose 16.34
C ₆ H ₉ DO	4-Methyl-2,3-dihydropyran-4- <i>d</i> 5.92	C ₆ H ₁₂ S	Thiepane 13.146
C ₆ H ₉ N	2,5-Dimethylpyrrole 6.60	C ₆ H ₁₂ S ₂	1,5-Dithiacyclooctane 13.86
C ₆ H ₉ NO	<i>N</i> -Methylfurfurylamine 5.65	C ₆ H ₁₃ N	Cyclohexylamine 8.17 2-Methylpiperidine 8.59 N-Methylpiperidine 8.60 <i>N,N</i> -Dimethylisobutylamine 8.83
C ₆ H ₉ N ₃ O ₂	Histidine 10.23	C ₆ H ₁₃ NO	3-Hydroxy- <i>N</i> -methylpiperidine 8.70 4-Hydroxy- <i>N</i> -methylpiperidine 8.71
C ₆ H ₉ N ₃ O ₂ S	2-Thiol-L-histidine 13.106	C ₆ H ₁₃ NO ₂ S	Methionine methyl ester 13.111
C ₆ H ₉ OT	4-Methyl-2,3-dihydropyran-4- <i>t</i> 5.93 4-Methyl-2,3-dihydropyran-3- <i>t</i> 5.95 4-Methyl-2,3-dihydropyran-2- <i>t</i> 5.97	C ₆ H ₁₃ NO ₄ S	4-Morpholineethanesulfonic acid 17.45
C ₆ H ₁₀	2,3-Dimethyl-1,3-butadiene 2.39 2-Ethyl-1,3-butadiene 2.41 Cyclohexene 2.116 Methylenecyclopentane 2.153 1-Methylcyclopentene 2.161 Isopropylidene cyclopropane 2.170 1,5-Hexadiene 2.240 (E,E)-2,4-Hexadiene 2.241 (E,Z)-2,4-Hexadiene 2.242 4-Methyl-1,3-pentadiene 2.300	C ₆ H ₁₄ N ₂	Hexahydro-1,2-dimethylpyridazine 8.91 <i>N,N</i> -Dimethyl-1-pyrrolidinamine 8.95
		C ₆ H ₁₄ N ₄ O ₂	Arginine 10.6

C ₆ H ₁₄ O	2,2-Dimethylpropyl methyl ether	17.19	C ₇ H ₈ S	Thioanisole 13.29
C ₆ H ₁₄ S	<i>tert</i> -Butyl ethyl sulfide	13.56		Benzyl mercaptan 13.42
	Diisopropyl sulfide	13.78	C ₇ H ₈ Se	Methylselenobenzene 13.27
	Dipropyl sulfide	13.82	C ₇ H ₈ Te	Methyltellurobenzene 13.28
C ₆ H ₁₅ N	<i>N,N</i> -Dimethyl-1-butanimine	8.14	C ₇ H ₉ N	Benzylamine 8.5
	Triethylamine	8.24		<i>N</i> -Methylaniline 9.14
	Diisopropylamine	8.29	C ₇ H ₁₀	1,3-Cycloheptadiene 2.99
	Dipropylamine	8.86		1-Methyl-1,4-cyclohexadiene 2.107
C ₆ H ₁₅ NO	<i>N,N</i> -Diethylethanamine	8.23	C ₇ H ₁₀ O	2,3-Dimethyl-2-cyclopenten-1-one 2.165
	<i>O,N,N</i> -Triethylhydroxylamine	15.22	C ₇ H ₁₀ O ₂	2-Hydroxy-3-methyl-2-cyclohexen-1-one 2.128
C ₆ H ₁₆ N ₂	1,2-Diethyl-1,2-dimethylhydrazine	8.33		3-Methoxy-2-methyl-2-cyclopenten-1-one 2.167
C ₆ H ₂₆ N ₂	Tetra(2-methylpropyl)hydrazine	8.43		3-Isopropylidene-2-furanone 2.227
C ₇ D ₈	Toluene- <i>d</i> ₈	3.128	C ₇ H ₁₀ O ₃	Methyl 4,5-dihydro-2-methyl-3-furancarboxylate 5.58
C ₇ H ₄ NO ⁻	4-Cyanophenoxy ion	4.158	C ₇ H ₁₀ S	Bicyclo[2.2.1]heptane-2-thione 13.43
C ₇ H ₅ NO	4-Cyanophenol	4.112	C ₇ H ₁₁ N ₃ O	4-Hydroxy-6-methyl-2-(dimethylamino)pyrimidine 16.70
C ₇ H ₅ N ₂ O ₅ ⁻	2-Methyl-4,6-dinitrophenoxide ion	4.173	C ₇ H ₁₁ OT	4,4-Dimethyl-2,3-dihydropyran-3- <i>t</i> 5.94
	4-Methyl-2,6-dinitrophenoxide ion	4.174		4,4-Dimethyl-2,3-dihydropyran-2- <i>t</i> 5.96
C ₇ H ₅ O ₃ ⁻	Salicylate ion	4.35	C ₇ H ₁₂	<i>trans</i> -2-Cyclopropyl-2-butene 2.49
C ₇ H ₆ ClNS	4-Chlorothiobenzamide	13.153		Methylenecyclohexane 2.112
C ₇ H ₆ Cl ₂ O	2,6-Dichloroanisole	3.37		1-Methylcyclohexene 2.122
C ₇ H ₆ NO ₃ ⁻	4-Methyl-2-nitrophenoxide ion	4.175		4-Methylcyclohexene 2.123
C ₇ H ₆ N ₂ O ₂ S	4-Nitrothiobenzamide	13.156		Ethyldenecyclopentane 2.152
C ₇ H ₆ O ₂	Methyl-1,4-benzoquinone	16.13		1,2-Dimethylcyclopentene 2.158
C ₇ H ₇ BrO ₃	Ethyl 5-bromo-2-furoate	5.55		1,5-Dimethylcyclopentene 2.159
C ₇ H ₇ BrS	4-Bromophenyl methyl sulfide	13.7		1-Cyclopropyl-2-methylpropene 2.335
C ₇ H ₇ ClO	2-Chloroanisole	3.34	C ₇ H ₁₂ NS ₂ ⁻	Hexamethylenedithiocarbamate 13.1
C ₇ H ₇ ClO ₂	3-Chloro-5-methoxyphenol	4.111	C ₇ H ₁₂ O	5,6-Dimethyl-3,4-dihydropyran 5.91
C ₇ H ₇ CIS	3-Chlorophenyl methyl sulfide	13.14		2-(Cyclopentylidene)ethanol 2.218
	4-Chlorophenyl methyl sulfide	13.15		3,4-Dimethyl-3-penten-2-one 2.324
C ₇ H ₇ FS	4-Fluorophenyl methyl sulfide	13.21	C ₇ H ₁₃ N	1-(Diethylamino)-2-methylacetylene 8.87
C ₇ H ₇ NO ₃	2-Nitroanisole	3.44		Quinuclidine 8.97
C ₇ H ₇ O ⁻	4-Methylphenoxy ion	4.172	C ₇ H ₁₃ NO	4,5,5-Trimethyl- Δ^1 -pyrroline- <i>N</i> -oxide 15.51
C ₇ H ₇ O ₂ ⁻	2-Methoxyphenoxy ion	4.169		2,4,4-Trimethyl- Δ^1 -pyrroline- <i>N</i> -oxide 15.52
	3-Methoxyphenoxy ion	4.170	C ₇ H ₁₄	1-Heptene 2.228
	4-Methoxyphenoxy ion	4.171		(<i>Z</i>)-2-Heptene 2.229
C ₇ H ₈	Norbornadiene	2.4		2,4-Dimethyl-2-pentene 2.306
	Quadricyclane	17.63	C ₇ H ₁₄ N ₂	2,3-Dimethyl-2,3-diazabicyclo[2.2.1]heptane 8.18
	Toluene	3.127		Diethyl(2-cyanoethyl)amine 8.81
C ₇ H ₈ O	Anisole	3.43	C ₇ H ₁₄ N ₂ O ₃	Glycylmethionine 13.109
	4-Methylphenol	4.135	C ₇ H ₁₄ O	(<i>E</i>)-3,4-Dimethyl-2-penten-1-ol 2.316
C ₇ H ₈ OS	4-(Methylmercapto)phenol	13.120		(<i>Z</i>)-3,4-Dimethyl-2-penten-1-ol 2.317
	2,6-Dimethylpyran-4-thione	13.139		3-Ethyl-2-penten-1-ol 2.318
C ₇ H ₈ O ₂	Methylhydroquinone	4.29	C ₇ H ₁₅ N	1,2-Dimethylpiperidine 8.57
	2-Methoxyphenol	4.129		2,6-Dimethylpiperidine 8.58
	3-Methoxyphenol	4.130	C ₇ H ₁₆	Heptane 17.34
	4-Methoxyphenol	4.131	C ₇ H ₁₆ N ₂	<i>N,N</i> -Dimethyl-1-piperidinamine 8.52
C ₇ H ₈ O ₃	5-(Methoxymethyl)furan-2-carboxaldehyde	5.12		
	2-[2-(1,3-Dioxolanyl)]furan	5.36		
	Ethyl 2-furoate	5.56		

C ₇ H ₁₆ S	<i>tert</i> -Butyl propyl sulfide 13.58 <i>sec</i> -Butyl propyl sulfide 13.59	C ₈ H ₁₀ O	1-Phenylethanol 3.55 2,2-Dimethylcyclohexa-3,5-dien-1-one 2.108 2,3-Dimethylidene-7-oxabicyclo[2.2.1]heptane 2.295 2,4-Dimethylphenol 4.116 2,6-Dimethylphenol 4.117 3,4-Dimethylphenol 4.118
C ₇ H ₁₇ NO	Diethyl(2-methoxyethyl)amine 8.25	C ₈ H ₁₀ OS	4-Methoxyphenyl methyl sulfide 13.22
C ₇ H ₁₈ AuPS	Methylthiolato(triethylphosphino)gold(I) 12.159	C ₈ H ₁₀ O ₂	1,2-Dimethoxybenzene 3.38 1,3-Dimethoxybenzene 3.39 1,4-Dimethoxybenzene 3.40 1,4-Dihydroxy-2,3-dimethylbenzene 4.22 1,4-Dihydroxy-2,5-dimethylbenzene 4.23 1,4-Dihydroxy-2,6-dimethylbenzene 4.24 6,6-Dimethylfulvene endoperoxide 2.150
C ₇ H ₁₈ N ₂	1,1,2-Trimethyl-2-(2-methylpropyl)hydrazine 8.45	C ₈ H ₁₀ O ₃	2-Acetoxy-3-methyl-2-cyclopenten-1-one 2.163 3-Acetoxy-2-methyl-2-cyclopenten-1-one 2.164 2,6-Dimethoxyphenol 4.115
C ₇ H ₁₉ N ₃	Spermidine 8.10	C ₈ H ₁₀ O ₄	5-(2-Hydroxyethoxy)methylfuran-2-carboxaldehyde 5.10
C ₈ F ₁₂ NiS ₄	Bis[1,1,1,4,4,4-hexafluoro-2-butene-2,3-dithiolato]nickel(II) 12.85	C ₈ H ₁₀ S	Methyl 3-methylphenyl sulfide 13.24 Methyl 4-methylphenyl sulfide 13.25 Benzyl methyl sulfide 13.30
C ₈ H ₆ F ₃ NS	4-(Trifluoromethyl)thiobenzamide 13.157	C ₈ H ₁₁ D	<i>endo</i> -2-Methylenenorbornane-3- <i>d</i> 2.13 <i>exo</i> -2-Methylenenorbornane-3- <i>d</i> 2.14
C ₈ H ₆ N ₂ O ₂	3-Phenylsydnone 5.107	C ₈ H ₁₁ N	<i>N,N</i> -Dimethylaniline 9.6 Phenethylamine 8.48
C ₈ H ₆ N ₄	1,2,3-Triazolo[1,2- <i>a</i>]benzotriazole 11.177	C ₈ H ₁₁ NO	Tyramine 4.186
C ₈ H ₆ O	Isobenzofuran 5.80	C ₈ H ₁₁ NO ₂	3,4-Dihydroxyphenethylamine 4.5
C ₈ H ₇ N	Indole 6.8	C ₈ H ₁₁ NO ₃	Noradrenaline 4.6
C ₈ H ₇ NO	5-Hydroxyindole 6.12	C ₈ H ₁₁ N ₃ O ₂	Tetrahydro-2-methyl-5,8-methano-[1,2,4]triazolo[1,2- <i>a</i>]pyridazine-1,3-dione 17.41
C ₈ H ₇ N ₃ O ₂	Luminol 9.33	C ₈ H ₁₂	2-Methylenenorbornane 2.11 2-Methylnorborn-2-ene 2.21 (Cyclopropylmethylene)cyclobutane 2.95 1,3-Cyclooctadiene 2.129 1,5-Cyclooctadiene 2.133 Cyclooctyne 2.139
C ₈ H ₇ O ₂ ⁻	4-Acetylphenoxide ion 4.151	C ₈ H ₁₂ N ₄ O ₃	Histidylglycine 10.20 Glycylhistidine 10.27
C ₈ H ₇ O ₃ ⁻	Methyl salicylate, conjugate base 4.41	C ₈ H ₁₂ NiS ₄	Bis(2-butene-2,3-dithiolato)nickel(II) 12.15
C ₈ H ₈	1,3,5,7-Cyclooctatetraene 2.136 Styrene 3.108	C ₈ H ₁₂ O	2-Methoxybicyclo[2.2.1]cycloheptene 2.20 2,5-Cyclooctadien-1-ol 2.134 2-(Isopropylidene)cyclopentanone 2.155 2-Ethyl-3-methyl-2-cyclopenten-1-one 2.166
C ₈ H ₈ BrO ⁻	4-Bromo-2,6-dimethylphenoxide ion 4.154	C ₈ H ₁₂ OS	2,2,4,4-Tetramethyl-3-thioxocyclobutanone 13.62
C ₈ H ₈ Br ₂	Cyclooctatetraene dibromide 2.32		
C ₈ H ₈ O	Acetophenone 3.2 5,6-Dimethylidene-7-oxa-2-bicyclo[2.2.1]heptene 2.297		
C ₈ H ₈ O ₂	Methyl benzoate 3.57 4'-Hydroxyacetophenone 4.93		
C ₈ H ₈ O ₃	Methyl salicylate 4.40		
C ₈ H ₈ O ₅	Dimethyl 2,5-furandicarboxylate 5.61		
C ₈ H ₉ ClO ₂	2-Chloro-1,4-dimethoxybenzene 3.32 5-Chloro-1,3-dimethoxybenzene 3.33		
C ₈ H ₉ NOS	4-Methoxythiobenzamide 13.154		
C ₈ H ₉ NS	4-Methylthiobenzamide 13.155		
C ₈ H ₉ O ⁻	2,6-Dimethylphenoxide ion 4.162		
C ₈ H ₉ O ₃ ⁻	2,6-Dimethoxyphenoxide ion 4.161		
C ₈ H ₁₀	2-Methylnorborna-2,5-diene 2.6 5-Methylene-2-norbornene 2.22		
C ₈ H ₁₀ BrN	Ethylbenzene 3.41 6,6-Dimethylfulvene 2.149		
C ₈ H ₁₀ ClN	4-Bromo- <i>N,N</i> -dimethylaniline 9.2 3-Chloro- <i>N,N</i> -dimethylaniline 9.3		
C ₈ H ₁₀ N ₂ O	4-Chloro- <i>N,N</i> -dimethylaniline 9.4 <i>N,N</i> -Dimethyl-4-nitrosoaniline 9.7		

C ₈ H ₁₂ O ₂	5-Acetyl-3,4-dihydro-6-methylpyran 5.88 5,5-Dimethyl-1,3-cyclohexanedione 17.15 6-Hydroperoxy-1,4-cyclooctadiene 2.130	C ₈ H ₂₀ N ₂	1,2-Dimethyl-1,2-dipropylhydrazine 8.40 1,1,2-Trimethyl-2-(2,2-dimethylpropyl)hydrazine 8.44
C ₈ H ₁₂ S ₂	2,2,4,4-Tetramethyl-1,3-cyclobutanedithione 13.60	C ₈ H ₂₀ NiO ₄ P ₂ S ₄	Bis(O,O'-diethylphosphorodithiolato)nickel(II) 12.44
C ₈ H ₁₃ N	1- <i>tert</i> -Butylpyrrole 6.61 2- <i>tert</i> -Butylpyrrole 6.62 3- <i>tert</i> -Butylpyrrole 6.63	C ₈ H ₂₁ NO ₆ P ⁺	Phosphatidylcholine 17.53
C ₈ H ₁₃ N ₃ O	4-Hydroxy-5,6-dimethyl-2-(dimethylamino)pyrimidine 16.69	C ₉ H ₆ NO ₂ ⁻	Indole-3-carboxylate ion 6.26
C ₈ H ₁₄	2-Cyclopropyl-3-methyl-2-butene 2.50 1- <i>tert</i> -Butylcyclobutene 2.97 Methylenecycloheptane 2.101 1-Methylcycloheptene 2.102 Ethyldenecyclohexane 2.110 1,2-Dimethylcyclohexene 2.117 1,3-Dimethylcyclohexene 2.118 1,4-Dimethylcyclohexene 2.119 1,6-Dimethylcyclohexene 2.120 Cyclooctene 2.137 2,5-Dimethyl-2,4-hexadiene 2.243	C ₉ H ₇ N	Quinoline 6.71
C ₈ H ₁₄ N ₄ O ₅	Glycylglycylglycylglycine 10.18	C ₉ H ₇ NO	8-Hydroxyquinoline 6.73
C ₈ H ₁₄ NiO ₂ S ₄	Bis[O-(1-methylethyl)carbonodithionato]nickel(II) 12.98	C ₉ H ₇ NO ₂	Indole-2-carboxylic acid 6.25 Indole-3-carboxylic acid 6.27
C ₈ H ₁₄ O	2-(Cyclohexylidene)ethanol 2.216	C ₉ H ₇ N ₃	Pyrazolo[1,2- <i>a</i>]benzotriazole 11.131
C ₈ H ₁₄ OS	2-Methyl-5-isopropylthiacyclopent-2-ene-1-oxide 13.144	C ₉ H ₈	Indene 3.69
C ₈ H ₁₄ O ₂ S ₂	1,2-Dithiolane-3-pentanoic acid 13.90	C ₉ H ₈ N ₂ O ₂	3-(4-Methylphenyl)sydnone 5.104 3-Methyl-4-phenylsydnone 5.105
C ₈ H ₁₄ S	2,2,4,4-Tetramethylcyclobutanethione 13.61	C ₉ H ₈ O ₂	(<i>E</i>)-Cinnamic acid 2.91
C ₈ H ₁₅ O ₂ S ₂ ⁻	Dihydrolipoate ion 13.115	C ₉ H ₈ O ₃	2-Hydroxycinnamic acid 4.69
C ₈ H ₁₆	4-Octene, (<i>E</i>)- 2.287 4-Octene, (<i>Z</i>)- 2.288 2,3,4-Trimethyl-1-pentene 2.303 2,3,4-Trimethyl-2-pentene 2.313 2,4,4-Trimethyl-2-pentene 2.314	C ₉ H ₈ O ₄	Phenylpyruvic acid 17.60
C ₈ H ₁₆ N ₂	1,1'-Bipyrrolidine 8.8 Hexahydropyrazolo[1,2- <i>a</i>][1,2]diazepine 8.88 Octahydropyridazino[1,2- <i>a</i>]pyridazine 8.92	C ₉ H ₉ N	3,4-Dihydroxycinnamic acid 4.68
C ₈ H ₁₆ O	(<i>E</i>)-3,4,4-Trimethyl-2-penten-1-ol 2.321 (<i>Z</i>)-3,4,4-Trimethyl-2-penten-1-ol 2.322	C ₉ H ₁₀	1-Methylindole 6.15 3-Methylindole 6.16
C ₈ H ₁₆ O ₃	1,1,2-Triethoxyethene 2.226	C ₉ H ₉ NO	5-Methoxyindole 6.13 Indole-3-carbinol 6.24
C ₈ H ₁₈	3-Methylheptane 17.35 4-Methylheptane 17.36	C ₉ H ₁₀	5,6-Dimethylidene-2-bicyclo[2.2.1]heptene 2.19 (<i>E</i>)- β -Methylstyrene 3.119 (<i>Z</i>)- β -Methylstyrene 3.120
C ₈ H ₁₈ S	Dibutyl sulfide 13.72 Di- <i>sec</i> -butyl sulfide 13.73 Di- <i>tert</i> -butyl sulfide 13.74 Diisobutyl sulfide 13.77	C ₉ H ₁₀ N ₂	4-(Dimethylamino)benzonitrile 9.22
C ₈ H ₁₉ NO ₂	Di(2-hydroxyethyl)- <i>tert</i> -butylamine 8.26	C ₉ H ₁₀ O	2-Methylacetophenone enol 2.219
		C ₉ H ₁₀ O ₃	3-(<i>p</i> -Hydroxyphenyl)propionic acid 4.34
		C ₉ H ₁₀ O ₅	Furfural diacetate 5.66
		C ₉ H ₁₁ ClOS	3-(3-Chlorophenyl)thio-1-propanol 13.132 3-(4-Chlorophenyl)thio-1-propanol 13.133
		C ₉ H ₁₁ FOS	3-(4-Fluorophenyl)thio-1-propanol 13.134
		C ₉ H ₁₁ NO	<i>p</i> -Dimethylaminobenzaldehyde 9.10
		C ₉ H ₁₁ NO ₂	Ethyl 4-aminobenzoate 9.9 Phenylalanine 10.31
		C ₉ H ₁₁ NO ₃	Tyrosine 4.187
		C ₉ H ₁₁ NO ₄	3-(3,4-Dihydroxyphenyl)-L-alanine 4.183
		C ₉ H ₁₂	2,3-Dimethylenebicyclo[2.2.1]heptane 2.8 (<i>E,E</i>)-1-Phenyl-1,3-pentadiene 2.301
		C ₉ H ₁₂ N ₂ O ₅ S	4-Thiouridine 16.76
		C ₉ H ₁₂ N ₂ O ₆	Uridine 16.80
		C ₉ H ₁₂ O	2,4,6-Trimethylphenol 4.148

C ₉ H ₁₂ OS	2-(4-Methylphenyl)thioethanol 13.97	C ₉ H ₁₅ N ₃ O ₂ S	Ergothioneine 13.104
	3-Phenylthio-1-propanol 13.137		
C ₉ H ₁₂ O ₂	3,4,5,6,7,8-Hexahydro-1-benzopyran-2-one 2.92	C ₉ H ₁₆	Ethyldenedecycloheptane 2.100
	Trimethylhydroquinone 4.31		1-Methylcyclooctene 2.138
	Cyclohexyldenemethyl acetate 2.255		1- <i>tert</i> -Butylcyclopentene 2.157
	4-Methoxy-2,6-dimethylphenol 4.132	C ₉ H ₁₆ NO ₂	2,2,6,6-Tetramethyl-4-piperidone <i>N</i> -oxyl 15.44
C ₉ H ₁₂ O ₃	1,2,3-Trimethoxybenzene 3.52	C ₉ H ₁₇ INO	4-Iodo-2,2,6,6-tetramethylpiperidin-1-oxyl 15.42
	1,2,4-Trimethoxybenzene 3.53		
	1,3,5-Trimethoxybenzene 3.54	C ₉ H ₁₇ N ₄ O	4-Azido-2,2,6,6-tetramethylpiperidin-1-oxyl 15.30
C ₉ H ₁₂ O ₄	Ubiquinol 0 4.21	C ₉ H ₁₈	1-Nonene 2.263
C ₉ H ₁₃ N	3-Phenylpropylamine 8.50		4-Methyl-4-octene 2.289
	<i>N,N</i> ,4-Trimethylaniline 9.19		(E)-4-Methyl-4-octene 2.290
C ₉ H ₁₃ NO	2-Methoxy- <i>N,N</i> -dimethylaniline 9.11		(Z)-4-Methyl-4-octene 2.291
	3-Methoxy- <i>N,N</i> -dimethylaniline 9.12	C ₉ H ₁₈ NO	2,2,6,6-Tetramethylpiperidine- <i>N</i> -oxyl 15.43
	4-Methoxy- <i>N,N</i> -dimethylaniline 9.13	C ₉ H ₁₈ NO ₂	4-Hydroxy-2,2,6,6-tetramethylpiperidine <i>N</i> -oxyl 15.33
C ₉ H ₁₃ NO ₃	Adrenaline 4.9	C ₉ H ₁₈ N ₂	1-(1-Pyrrolidinyl)piperidine 8.62
C ₉ H ₁₃ NS	<i>p</i> -(Dimethylamino)thioanisole 13.6	C ₉ H ₁₈ O ₂	5-Hydroperoxy-4-methyl-3-octene 2.286
C ₉ H ₁₃ N ₃ O ₅	Cytidine 16.18	C ₉ H ₁₈ S	Di- <i>tert</i> -butylthioketone 13.117
C ₉ H ₁₄	(1-Cyclopropylethylidene)cyclobutane 2.94	C ₉ H ₁₉ N	2,2,6,6-Tetramethylpiperidine 8.63
	1,1-Bis(cyclopropyl)-1-propene 2.333	C ₉ H ₁₉ NO	2,2,6,6-Tetramethylpiperidin-4-ol 8.77
C ₉ H ₁₄ N ₃ O ₂ S ⁻	Ergothioneine ion 13.105		2,2,6,6-Tetramethylpiperidin-1-ol 15.28
C ₉ H ₁₄ N ₃ O ₃	2,2,5,5-Tetramethyl-3-cyano-4-nitopyrrolidin-1-oxyl 15.53	C ₉ H ₁₉ N ₂ O	4-Amino-2,2,6,6-tetramethylpiperidine- <i>N</i> -oxyl 15.29
C ₉ H ₁₄ N ₃ O ₇ P	2'-Deoxycytidine-5'-monophosphate 16.22	C ₉ H ₂₀ N ₂	4-Amino-2,2,6,6-tetramethylpiperidine 8.54
C ₉ H ₁₄ N ₃ O ₈ P	Cytidine 5'-monophosphate 16.19	C ₁₀ H ₆ O ₃	5-Hydroxy-1,4-naphthoquinone 4.90
C ₉ H ₁₄ N ₄ O ₃	Histidinyl-β-alanine 10.3		6-Hydroxy-1,4-naphthoquinone 4.91
	Carnosine 10.24	C ₁₀ H ₇ BrO	2-(4-Bromophenyl)furan 5.22
C ₉ H ₁₄ O	2-Ethyl-3-methyl-2-cyclohexen-1-one 2.127		3-(4-Bromophenyl)furan 5.23
	6-Methoxy-1,4-cyclooctadiene 2.131	C ₁₀ H ₇ ClO	2-(4-Chlorophenyl)furan 5.25
C ₉ H ₁₄ O ₂	2-[(1,1-Dimethylethoxy)methyl]furan 5.35	C ₁₀ H ₇ FO	3-(4-Fluorophenyl)furan 5.41
	Acetoxyxymethylidenecyclohexane 2.111	C ₁₀ H ₇ NO ₃	2-(4-Nitrophenyl)furan 5.49
	(E,Z)-4,5-Diethylidene-2,2-dimethyl-1,3-dioxolane 2.197	C ₁₀ H ₈	Naphthalene 3.74
	(Z,Z)-4,5-Diethylidene-2,2-dimethyl-1,3-dioxolane 2.198	C ₁₀ H ₈ NO ₂ ⁻	Indole-3-acetate ion 6.20
C ₉ H ₁₄ O ₃	5-Ethoxycarbonyl-3,4-dihydro-6-methylpyran 5.98	C ₁₀ H ₈ O	1-Naphthol 4.88
C ₉ H ₁₄ O ₄	5-Methoxymethyl-2-(dimethoxymethyl)furan 5.32		2-Naphthol 4.89
C ₉ H ₁₄ S	3,3-Dimethylbicyclo[2.2.1]heptane-2-thione 13.44		2-Phenylfuran 5.50
	3,5,5-Trimethyl-2-cyclohexene-1-thione 13.65	C ₁₀ H ₉ N	3-Phenylfuran 5.51
C ₉ H ₁₅ NO ₃ S	Captopril 13.130	C ₁₀ H ₈ O ₂	1,5-Dihydroxynaphthalene 4.82
C ₉ H ₁₅ N ₃ O	2-(Diethylamino)-4-hydroxy-6-methylpyrimidine 16.68		1,6-Dihydroxynaphthalene 4.83
			1,7-Dihydroxynaphthalene 4.84
			1,8-Dihydroxynaphthalene 4.85
			2,6-Dihydroxynaphthalene 4.86
			2,7-Dihydroxynaphthalene 4.87
		C ₁₀ H ₉ N	2-Naphthylamine 9.24
		C ₁₀ H ₉ NO	6-Methoxyquinoline 6.74
		C ₁₀ H ₉ NO ₂	Indole-3-acetic acid 6.21

$C_{10}H_{10}$	2-Methyl-5-(1-methylethyl)-1,3-cyclohexadiene	2.106	$C_{10}H_{14}N_5O_7P$	Adenosine 5'-monophosphate	16.2
$C_{10}H_{10}Fe$	Ferrocene	12.144		2'-Deoxyguanosine 5'-monophosphate	16.23
$C_{10}H_{10}N_2O_2$	4-Methyl-3-(4-methylphenyl)sydnone	5.103	$C_{10}H_{14}N_5O_8P$	Guanosine 5'-monophosphate	16.38
$C_{10}H_{10}Ni$	Nickelocene	12.165	$C_{10}H_{14}NiO_4$	Bis(acetylacetone)nickel(II)	12.10
$C_{10}H_{10}O$	2,3,5,6-Tetramethylidene-7-oxabicyclo[2.2.1]heptane	2.296	$C_{10}H_{14}NiO_4 \cdot 2H_2O$	Bis(acetylacetone)nickel(II) dihydrate	12.11
$C_{10}H_{10}O_3$	2,3,5,6-Tetramethylidene-7-oxa-2-bicyclo[2.2.1]heptane monoendoperoxide	2.357	$C_{10}H_{14}NiS_4$	Bis(dithioacetylacetone)nickel(II)	12.74
$C_{10}H_{10}O_4$	4-Hydroxy-3-methoxycinnamic acid	4.70	$C_{10}H_{14}O$	2-Cyclopentylidenecyclopentanone	2.154
$C_{10}H_{11}N$	1,3-Dimethylindole	6.10		2- <i>tert</i> -Butylphenol	4.120
	2,3-Dimethylindole	6.11		4- <i>tert</i> -Butylphenol	4.121
$C_{10}H_{11}NO_2$	5,6-Dimethoxyindole	6.9	$C_{10}H_{14}OS$	3-(4-Methylphenyl)thio-1-propanol	13.13 <i>b</i>
$C_{10}H_{12}$	4,5,6,7-Tetrahydro-4,7-methanoindene	2.254	$C_{10}H_{14}O_2$	<i>p</i> - <i>tert</i> -Butylcatechol	4.8
	β,β -Dimethylstyrene	3.114		4-Methoxy-2,3,6-trimethylphenol	4.134
$C_{10}H_{12}N_2$	Tryptamine	10.34	$C_{10}H_{14}O_2S$	3-(4-Methoxyphenyl)thio-1-propanol	13.135
$C_{10}H_{12}O$	Anethole	3.116	$C_{10}H_{14}O_4$	1,2,3,4-Tetramethoxybenzene	3.49
$C_{10}H_{12}OS$	(E)-2-Phenylsulfinyl-2-butene	13.54		1,2,3,5-Tetramethoxybenzene	3.50
	(Z)-2-Phenylsulfinyl-2-butene	13.55		1,2,4,5-Tetramethoxybenzene	3.51
	1-Phenylsulfinyl-2-methylpropene	2.338	$C_{10}H_{14}O_4Zn$	Bis(acetylacetone)zinc(II)	12.13
$C_{10}H_{12}O_5$	2,5-Bis(ethoxycarbonyl)furan	5.60	$C_{10}H_{14}S$	(Butylthio)benzene	13.8
	Diethyl 3,4-furandicarboxylate	5.62		4,6,6-Trimethylbicyclo[3.1.1]hept-3-ene-2-thione	13.48
$C_{10}H_{13}ClS$	1-(Butylthio)-3-chlorobenzene	13.9	$C_{10}H_{14}S_4$	1,5,7-Trimethyl-3-methylidene-2,6,9,10-tetrathiaadamantane	2.253
	1-(Butylthio)-4-chlorobenzene	13.10	$C_{10}H_{15}N$	4-Phenylbutylamine	8.49
$C_{10}H_{13}FS$	1-(Butylthio)-4-fluorobenzene	13.11		<i>N,N</i> ,2,4-Tetramethylaniline	9.17
$C_{10}H_{13}NO_3$	Tyrosine, methyl ester	4.190	$C_{10}H_{15}NO_2$	3,4-Dimethoxy- <i>N,N</i> -dimethylaniline	9.5
$C_{10}H_{13}N_5O_4$	Adenosine	16.1	$C_{10}H_{15}N_2O_7P$	3'-Deoxythymidine 5'-monophosphate	
	2'-Deoxyguanosine	16.37		16.25	
$C_{10}H_{13}N_5O_5$	Guanosine	16.36	$C_{10}H_{15}N_2O_8P$	Thymidine 5'-monophosphate	16.78
$C_{10}H_{13}O^-$	4-(<i>tert</i> -Butyl)phenoxide ion	4.163	$C_{10}H_{15}N_5O_4$	Glycyl-L-histidylglycine	10.19
$C_{10}H_{14}$	2,3-Dimethylenebicyclo[2.2.2]octane	2.33		Glycylglycyl-L-histidine	10.28
	(Dicyclopropylmethylidene)cyclopropane		$C_{10}H_{16}$	7,7-Dimethyl-2-methylenenorbornane	2.9
	2.168			2,7,7-Trimethylnorborn-2-ene	2.23
$C_{10}H_{14}CoO_4$	Bis(acetylacetone)cobalt(II)	12.7		2-Carene	2.26
$C_{10}H_{14}CoS_4$	Bis(dithioacetylacetone)cobalt(II)	12.73		4-Carene	2.27
$C_{10}H_{14}CuO_4$	Bis(acetylacetone)copper(II)	12.8		3-Carene	2.28
$C_{10}H_{14}N_2$	Nicotine	8.93		α -Thujene	2.30
$C_{10}H_{14}N_2O$	α -(4-Pyridyl)- <i>tert</i> -butyl nitrone	15.48		α -Terpinene	2.105
$C_{10}H_{14}N_2O_2$	α -(4-Pyridyl 1-oxide)- <i>N</i> - <i>tert</i> -butylnitronone	15.47		Terpinolene	2.121
				Limonene	2.124
$C_{10}H_{14}N_2O_5$	Thymidine	16.77		Octalin	2.261
$C_{10}H_{14}N_5O_6P$	2'-Deoxyadenosine 5'-monophosphate			2,6-Dimethyl-2,4,6-octatriene	2.285
	16.21			α -Pinene	2.326
				β -Pinene	2.327
				1,1-Bis(cyclopropyl)-2-methyl-1-propene	
				2.334	

$C_{10}H_{16}N_2$	N,N,N',N' -Tetramethyl- <i>o</i> -phenylenediamine 9.26	$C_{10}H_{20}N_2$	1,1'-Bipiperidine 8.7 1,2-Bis(diethylamino)acetylene 8.30
	N,N,N',N' -Tetramethyl- <i>m</i> -phenylenediamine 9.27	$C_{10}H_{20}N_2NiS_4$	Bis(diethyldithiocarbamato)nickel(II) 12.40
	N,N,N',N' -Tetramethyl- <i>p</i> -phenylenediamine 9.31	$C_{10}H_{20}O$	(<i>E</i>)-3,7-Dimethyl-2-octen-1-ol 2.292 (<i>Z</i>)-3,7-Dimethyl-2-octen-1-ol 2.293 Citronellol 2.294
$C_{10}H_{16}N_2NiO_2$	Bis(4-imino-2-pentanonato)nickel(II) 12.97	$C_{10}H_{20}O_4$	Tetraethoxyethene 2.225
$C_{10}H_{16}N_3O_6S^-$	Glutathione, negative ion 13.100	$C_{10}H_{21}N$	1,2,2,6,6-Pentamethylpiperidine 8.61
$C_{10}H_{16}N_4O_2$	Homocarnosine 10.26	$C_{10}H_{21}NO$	1,2,2,6,6-Pentamethyl-4-piperidinol 8.72
$C_{10}H_{16}N_4O_3$	Anserine 10.25	$C_{10}H_{24}N_2$	1,2-Dimethyl-1,2-dibutylhydrazine 8.34 1,1-Dimethyl-2,2-di(2-methylpropyl)hydrazine 8.37 1,2-Dimethyl-1,2-di(2-methylpropyl)hydrazine 8.38
$C_{10}H_{16}N_5O_{13}P_3$	Adenosine triphosphate 16.3	$C_{10}H_{26}N_4$	Spermine 8.11
$C_{10}H_{16}O$	Pulegone 2.115	$C_{10}H_{26}ON_2$	Tetrahydro-2,2,6,6-tetramethylpyrazolo[1,2- <i>a</i>]pyrazole 8.90
$C_{10}H_{16}OS$	3,3,6,6-Tetramethylthiacyclohept-4-yne 1-oxide 13.149	$C_{11}H_6O_3$	Angelicin 5.1 Psoralen 5.86
$C_{10}H_{16}OSi$	2-(Trimethylsiloxy)norborna-2,5-diene 2.7	$C_{11}H_7NO$	2-(4-Cyanophenyl)furan 5.26
$C_{10}H_{16}O_2$	6,6-Dimethylcyclohex-1-enyl acetate 2.126	$C_{11}H_8O_2$	2-Methyl-1,4-naphthoquinone 16.44
$C_{10}H_{16}O_2S$	3,3,6,6-Tetramethylthiacyclohept-4-yne 1,1-dioxide 13.148	$C_{11}H_{10}$	1-Methylnaphthalene 3.86
$C_{10}H_{16}O_3$	6-Ethyl-3,4-dihydropyran-5-carboxylic acid ethyl ester 5.101	$C_{11}H_{10}Cl_2N_2O_2$	Azo dioxide 6 15.20
$C_{10}H_{16}S$	Thiofenchone 13.46 Thiocamphor 13.47 3,3,6,6-Tetramethylthiacyclohept-4-yne 13.147	$C_{11}H_{10}NO_2^-$	Indole-3-propionate ion 6.29
$C_{10}H_{17}N_3O_6S$	Glutathione 13.99	$C_{11}H_{10}O$	2-(4-Methylphenyl)furan 5.47 3-(4-Methylphenyl)furan 5.48
$C_{10}H_{18}$	<i>p</i> -Menth-1-ene 2.125 Ethylidenecyclooctane 2.135 (<i>E</i>)-2,6-Dimethyl-2,6-octadiene 2.275 (<i>Z</i>)-2,6-Dimethyl-2,6-octadiene 2.276 2,7-Dimethyl-2,6-octadiene 2.277	$C_{11}H_{10}O_2$	2-(4-Methoxyphenyl)furan 5.44 3-(4-Methoxyphenyl)furan 5.45 α -Phenylfurfuryl alcohol 5.70
$C_{10}H_{18}MnO_6^+$	Bis(acetylacetonato)manganese(III) 12.9	$C_{11}H_{11}NO_2$	Indole-3-propionic acid 6.30
$C_{10}H_{18}O$	2-(Cyclooctylidene)ethanol 2.217 Linalool 2.280 (<i>E</i>)-3,7-Dimethyl-2,6-octadien-1-ol 2.281 (<i>E</i>)-3,7-Dimethyl-2,7-octadien-1-ol 2.282 (<i>Z</i>)-3,7-Dimethyl-2,7-octadien-1-ol 2.283	$C_{11}H_{11}N_3$	1,3-Dimethylpyrazolo[1,2- <i>a</i>]benzotriazole 11.132
$C_{10}H_{18}OSi$	2-(Trimethylsiloxy)norborn-2-ene 2.24	$C_{11}H_{11}N_3O_2$	1,2,5,6,6 <i>a</i> ,6 <i>b</i> ,6 <i>c</i> ,6 <i>d</i> -Octahydro-1,6[1',2']-4-methyl-3,5-dioxo-1,2,4-triazolo[1,3]cyclopropa[1,2,3- <i>cd</i>]cyclopropa[<i>gh</i>]pentalene 17.70
$C_{10}H_{18}O_2$	7-Hydroperoxy-2,7-dimethyl-2,5-octadiene 2.274 Hydroperoxy-2,6-dimethyl-2,6-octadiene 2.278 6-Hydroperoxy-2,7-dimethyl-2,7-octadiene 2.279	$C_{11}H_{11}N_3S$	4-(Aminomethylene)-3-methyl-1-phenyl-2-pyrazoline-5-thione 13.142
$C_{10}H_{18}O_3$	Monohydroperoxy-3,7-dimethyl-2,6-octadien-1-ol 2.284	$C_{11}H_{11}N_3Se$	4-(Aminomethylene)-3-methyl-1-phenyl-2-pyrazoline-5-selone 13.141
$C_{10}H_{18}S$	3,3,6,6-Tetramethylthiacyclohept-4-ene 13.150 Di- <i>tert</i> -butylthioketene 13.165	$C_{11}H_{12}$	2-Benzyl-1,3-butadiene 2.43 1,2,3,4-Tetrahydro-1,4-methanonaphthalene 3.72
		$C_{11}H_{12}N_2O$	Indole-3-propionamide 6.28
		$C_{11}H_{12}N_2O_2$	Indole, 1,2,3-trimethyl-5-nitro-Tryptophan 10.36
		$C_{11}H_{13}Cl$	<i>m</i> -Chloro- α,β,β -trimethylstyrene 3.109 <i>p</i> -Chloro- α,β,β -trimethylstyrene 3.110
		$C_{11}H_{13}ClS$	4'-Chloropivalothiophenone 13.127

C ₁₁ H ₁₃ FS	4'-Fluoropivalothiophenone	13.128	C ₁₁ H ₂₂ O ₂	Undecanoic acid	17.76
C ₁₁ H ₁₃ N	1,2,3-Trimethylindole	6.18	C ₁₁ H ₂₃ NO	<i>N</i> -(2-Hydroxyethyl)-2,2,6,6-tetramethylpiperidine	8.68
C ₁₁ H ₁₃ N ₃ O	Tryptophanamide	10.43	C ₁₁ H ₂₆ N ₂	<i>N,N</i> -Diethyl-1,7-heptanediamine	8.31
C ₁₁ H ₁₄	3-Methyl-1-phenyl-2-butene	3.45	C ₁₂ H ₂ Cl ₆ NiS ₄ .C ₁₆ H ₃₆ N	Bis[3,4,6-trichloro-1,2-benzenedithiolato]nickelate(I), tetrabutylammonium salt	12.117
C ₁₁ H ₁₄ N ₂ O ₂	<i>N</i> -Methoxycarbonyltryptamine	10.35	C ₁₂ H ₈ O ₄	8-Methoxypсорален	5.87
C ₁₁ H ₁₄ N ₂ O ₄	L-Tyrosylglycine	4.76	C ₁₂ H ₉ ClS	1-Chloro-4-(phenylthio)benzene	13.16
	Glycyltyrosine	4.189	C ₁₂ H ₉ NO	Phenoxyazine	6.38
C ₁₁ H ₁₄ OS	2-Methyl-3-phenylsulfinyl-2-butene	13.53	C ₁₂ H ₉ NO ₂ S	1-Nitro-4-(phenylthio)benzene	13.31
C ₁₁ H ₁₄ S	Pivalothiophenone	13.126	C ₁₂ H ₉ NS	Phenothiazine	13.121
C ₁₁ H ₁₅ NO	Phenyl- <i>N-tert</i> -butylnitrone	15.46	C ₁₂ H ₉ N ₃ O ₂	4-Nitroazobenzene	11.61
C ₁₁ H ₁₅ NOS	4-[(Phenylmethyl)thio]morpholine	13.112	C ₁₂ H ₉ O ⁻	2-Phenylphenoxyde	4.180
C ₁₁ H ₁₅ NO ₂	3,5-Diacetyl-1,4-dihydropyridine	6.41		4-Phenylphenoxyde ion	4.181
	Ethyl 4-(dimethylamino)benzoate	9.21	C ₁₂ H ₉ O ₂ ⁻	2,2'-Dihydroxybiphenyl, conjugate base	
C ₁₁ H ₁₅ NO ₄ S	2-Sulfonatophenyl- <i>N-tert</i> -butylnitrone			4.166	
	15.49			2,5-Dihydroxybiphenyl, conjugate base	
C ₁₁ H ₁₆	Nopadiene	2.25	C ₁₂ H ₁₀ N ₂	Azobenzene	11.57
	2,3-Dimethylenebicyclo[3.2.2]nonane	2.31	C ₁₂ H ₁₀ N ₃ S ⁺	Thionine cation	11.114
	(Dicyclopropylmethyldiene)cyclobutane		C ₁₂ H ₁₀ O	2'-Acetonaphthone	3.1
	2.96			2-Hydroxybiphenyl	4.141
	1,1,2-Tricyclopropylethene	2.169	C ₁₂ H ₁₀ O ₂	2,5-Dihydroxybiphenyl	4.30
C ₁₁ H ₁₆ O	2- <i>tert</i> -Butyl-4-hydroxyanisole	4.122		2,2'-Dihydroxybiphenyl	4.128
C ₁₁ H ₁₆ OS	1-(Butylthio)-4-methoxybenzene	13.12	C ₁₂ H ₁₀ S	Diphenyl sulfide	13.34
	4-(4-Methylphenyl)thio-1-butanol	13.52	C ₁₂ H ₁₀ S ₂	Diphenyl disulfide	13.84
C ₁₁ H ₁₆ O ₂	4-Methoxy-2,3,5,6-tetramethylphenol	4.133	C ₁₂ H ₁₀ Se ₂	Diphenyldiselenide	13.83
C ₁₁ H ₁₆ O ₅	Pentamethoxybenzene	3.48	C ₁₂ H ₁₀ Te ₂	Diphenylditelluride	13.85
C ₁₁ H ₁₆ S	1-(Butylthio)-4-methylbenzene	13.13	C ₁₂ H ₁₁ N	Diphenylamine	9.16
	Methyl 4- <i>tert</i> -butylphenyl sulfide	13.20	C ₁₂ H ₁₁ NO	3-Hydroxydiphenylamine	4.142
C ₁₁ H ₁₇ N	<i>N,N</i> ,2,4,6-Pentamethylaniline	9.15	C ₁₂ H ₁₁ N ₂ O ₃ ⁻	Nalidixic acid, anion	16.46
C ₁₁ H ₁₇ NS	<i>N,N</i> -Diethylbenzenemethanesulfenamide		C ₁₂ H ₁₂	1,2-Dimethylnaphthalene	3.75
	13.35			1,3-Dimethylnaphthalene	3.76
C ₁₁ H ₁₇ N ₂ O ₄ ⁺	α -(1-Methyl-4-pyridyl)- <i>tert</i> -butyl nitrone			1,4-Dimethylnaphthalene	3.77
	15.45			1,6-Dimethylnaphthalene	3.78
C ₁₁ H ₁₇ N ₂ O ₈ P	Nicotinamide mononucleotide, reduced			1,7-Dimethylnaphthalene	3.79
	16.51			1,8-Dimethylnaphthalene	3.80
C ₁₁ H ₁₈ O	3-Methyl-2-pentyl-2-cyclopenten-1-one			2,3-Dimethylnaphthalene	3.81
	2.162			2,6-Dimethylnaphthalene	3.82
C ₁₁ H ₁₈ O ₃	6-Isopropyl-3,4-dihydropyran-5-carboxylic acid ethyl ester	5.99		2,7-Dimethylnaphthalene	3.83
C ₁₁ H ₁₈ S	2,2,6,6-Tetramethylcyclohexylthioketene		C ₁₂ H ₁₂ N ⁺	<i>N</i> -Methyl-4-phenylpyridinium	6.55
	13.166		C ₁₂ H ₁₂ NO ₂ ⁻	Indole-3-butyrate ion	6.22
C ₁₁ H ₁₉ N ₃ O	Dimethirimol	16.66	C ₁₂ H ₁₂ N ₂	Benzidine	9.23
	Ethirimol	16.67	C ₁₂ H ₁₂ N ₂ O ₃	Nalidixic acid	16.45
C ₁₁ H ₂₀ OSi	6-(Trimethylsiloxy)-1,4-cyclooctadiene		C ₁₂ H ₁₂ OS	3-(4-Methoxyphenyl)-2-cyclopentene-1-thione	13.66
	2.132				
C ₁₁ H ₂₁ N	<i>N</i> -Cyclohexylpiperidine	8.56			
C ₁₁ H ₂₂ N ₂ O ₃ S	L-Methionyl-L-leucine	13.107			

C ₁₂ H ₁₂ O ₂	α -Benzylfurfuryl alcohol 5.72 4-Hydroxymethyl-2-(phenylmethyl)furan 5.74 [2,2](2,5)Furanophane 5.77	C ₁₂ H ₁₈ OS	5-(4-Methylphenyl)thio-1-pentanol 13.118
C ₁₂ H ₁₂ S	1,1-Dimethyl-2-naphthalenethione 13.113	C ₁₂ H ₁₈ O ₂	Durohydroquinone monoethyl ether 4.126
C ₁₂ H ₁₃ N	<i>m</i> -Cyano- α,β,β -trimethylstyrene 3.111 <i>p</i> -Cyano- α,β,β -trimethylstyrene 3.112	C ₁₂ H ₁₈ O ₃	4-(1-Hydroxy-2,2-dimethyl)propyl-2-methoxyphenol 4.127
C ₁₂ H ₁₃ NO	(E)-1-Methyl-3-(β -methoxylvinyl)indole 2.247 (Z)-1-Methyl-3-(β -methoxylvinyl)indole 2.248	C ₁₂ H ₁₈ O ₆	Hexamethoxybenzene 3.42 (Cyclohexylidene)cyclohexane 2.109
C ₁₂ H ₁₃ NO ₂	3-Indolebutyric acid 6.23	C ₁₂ H ₂₀	1,1-Bis(cyclobutyl)-2-methyl-1-propene 2.332
C ₁₂ H ₁₄	4,5,6,7-Tetrahydrospirocyclopropane[4,7]methanoindene 2.345	C ₁₂ H ₂₀ O	2- <i>tert</i> -Butylfuran 5.24 2,5-Di- <i>tert</i> -butylfuran 5.28 (Cyclohexylidene)cyclohexanol 2.113
C ₁₂ H ₁₄ Cl ₃ MnO ₆	Bis(acetylacetonato)trichloroacetato-manganese(III) 12.12	C ₁₂ H ₂₀ OSi	7,7-Dimethyl-2-(trimethylsiloxy)-norborna-2,5-diene 2.5
C ₁₂ H ₁₄ N ⁺	<i>N</i> -Methyl-4-phenyl-2,3-dihydropyridinium 6.53	C ₁₂ H ₂₀ S	1,3,3,7,7-Pentamethylbicyclo[2.2.1]heptane-2-thione 13.45
C ₁₂ H ₁₄ N ₂ O ₂	Tryptophan, methyl ester 10.42	C ₁₂ H ₂₂ O	(E,E)-8,10-Dodecadienol 2.204
(C ₁₂ H ₁₅ N) _n	Peranax 45 6.72	C ₁₂ H ₂₂ O ₂	(E,E)-1,4-Di(<i>tert</i> -butoxy)-1,3-butadiene 2.36 (E,Z)-1,4-Di(<i>tert</i> -butoxy)-1,3-butadiene 2.37 (Z,Z)-1,4-Di(<i>tert</i> -butoxy)-1,3-butadiene 2.38
C ₁₂ H ₁₅ N	<i>N</i> -Methyl-4-phenyl-1,2,3,6-tetrahydropyridine 8.94	C ₁₂ H ₂₂ O ₁₁	Sucrose 16.75
C ₁₂ H ₁₅ NO	5-Methoxy-1,2,3-trimethylindole 6.14	C ₁₂ H ₂₄ O ₂	Dodecanoic acid 17.24
C ₁₂ H ₁₆	Tetramethyl- <i>o</i> -xylylene 2.104 3, α,β,β -Tetramethylstyrene 3.121 4, α,β,β -Tetramethylstyrene 3.122	C ₁₂ H ₂₆	Dodecane 17.23
C ₁₂ H ₁₆ ClMnO ₆	Bis(acetylacetonato)chloroacetatomanganese(III) 12.6	C ₁₂ H ₂₆ O ₄ S	Dodecyl sulfate, sodium salt 17.25
C ₁₂ H ₁₆ N ₂ O ₅ S	5-S-Cysteinyldopa 4.71	C ₁₂ H ₂₇ N	Tributylamine 8.13
C ₁₂ H ₁₆ O	<i>m</i> -Methoxy- α,β,β -trimethylstyrene 3.117 <i>p</i> -Methoxy- α,β,β -trimethylstyrene 3.118	C ₁₂ H ₂₇ O ₃ P	Tributyl phosphite 17.71
C ₁₂ H ₁₆ OS	4'-Methoxypivalothiophenone 13.129	C ₁₂ H ₂₈ N ₂	1,1-Dimethyl-2,2-di(2,2-dimethylpropyl)hydrazine 8.35 1,2-Dimethyl-1,2-di(2,2-dimethylpropyl)hydrazine 8.36 1,2-Dimethyl-1,2-dipentylhydrazine 8.39
C ₁₂ H ₁₆ O ₂	2,3-Dihydro-5-hydroxy-2,4,6,7-tetramethylbenzofuran 4.37	C ₁₂ H ₂₈ NiO ₄ P ₂ S ₄	Bis(O,O'-diisopropylphosphorodithiolato)nickel(II) 12.52
C ₁₂ H ₁₆ S	2,3-Dimethyl-2-butenyl phenyl sulfide 13.17	C ₁₃ H ₈ CINOSe	7-Chloro-2-phenyl-1,2-benzisoselenazol-3-one 13.36
C ₁₂ H ₁₇ MnO ₆	Acetatobis(acetylacetonato)manganese(III) 12.1	C ₁₃ H ₈ Cl ₂ N ₂	Diazodi(4-chlorophenyl)methane 11.17
C ₁₂ H ₁₇ NO ₄	Diethyl 2,4-dimethylpyrrole-3,5-dicarboxylate 6.66	C ₁₃ H ₈ Cl ₂ S	4,4'-Dichlorothiobenzophenone 13.159
C ₁₂ H ₁₇ NS	1-[(Phenylmethyl)thio]piperidine 13.125	C ₁₃ H ₈ FNOSe	7-Fluoro-2-phenyl-1,2-benzisoselenazol-3-one 13.37
C ₁₂ H ₁₈	Hexamethylbicyclo[2.2.0]hexa-2,5-diene 2.29 (Dicyclopropylmethylidene)cyclopentane 2.151	C ₁₃ H ₈ N ₂	9-Diazofluorene 11.63
C ₁₂ H ₁₈ O	2-(Cyclohexylidene)cyclohexanone 2.114	C ₁₃ H ₈ N ₂ O ₃ Se	7-Nitro-2-phenyl-1,2-benzisoselenazol-3-one 13.39
		C ₁₃ H ₈ O	Phenalen-1-one 3.97
		C ₁₃ H ₉ BrN ₂	Diazo(4-bromophenyl)phenylmethane 11.15

C ₁₃ H ₉ ClS	4-Chlorothiobenzophenone	13.158	C ₁₃ H ₁₈ OS	2,3-Dimethyl-2-but enyl 4-methoxyphenyl sulfide	13.18
C ₁₃ H ₉ NOS	2-Phenyl-1,2-benzisothiazol-3-one	13.41	C ₁₃ H ₁₈ O ₂	Acetoxymethylenadamantane	2.2
C ₁₃ H ₉ NOSe	Ebselen	13.40		2,3-Dihydro-5-hydroxy-2,2,4,6,7-pentamethylbenzofuran	4.36
C ₁₃ H ₁₀ NO ⁻	Benzophenone oximate anion	15.13		Adamantylidenemethyl acetate	2.256
C ₁₃ H ₁₀ N ₂	Diazodiphenylmethane	11.16	C ₁₃ H ₁₈ S	2,3-Dimethyl-2-but enyl 4-methylphenyl sulfide	13.19
C ₁₃ H ₁₀ O	Benzophenone	3.58	C ₁₃ H ₁₉ N	p-(N,N-Dimethylamino)-α,β,β-trimethylstyrene	3.115
C ₁₃ H ₁₁ NO	Diphenyl nitrone	15.10	C ₁₃ H ₁₉ NO ₂	3,5-Diacetyl-1,4-dihydro-2,6-dimethylpyridine	6.42
	Benzophenone oxime	15.14	C ₁₃ H ₁₉ NO ₄	3,5-Diethoxycarbonyl-1,4-dihydro-2,6-dimethylpyridine	6.47
C ₁₃ H ₁₁ NO ₂	2-Hydroxyphenyl phenyl nitrone	15.7	C ₁₃ H ₁₉ NS ₂	N-[(5-Methyl-2-(methylthio)-3-thienyl)methylene]cyclohexanamine	13.64
C ₁₃ H ₁₁ NS	10-Methylphenothiazine	13.123		3-[(Cyclohexylamino)methylene]-5-ethyl-2-thiophenethione	13.170
C ₁₃ H ₁₁ N ₃	3,6-Diaminoacridine	11.1	C ₁₃ H ₂₀	2-Isopropylideneadamantane	2.353
C ₁₃ H ₁₁ N ₃ O	2 Methyl 6 phenylimidazo[1,2- <i>a</i>]pyrazin-3-one	16.41	C ₁₃ H ₂₀ CoN ₅ O ₅	Hydroxybis(dimethylglyoximato)pyridinecobalt(II)	12.146
C ₁₃ H ₁₁ O ⁻	2-Benzylphenoxide ion	4.152	C ₁₃ H ₂₀ O	β-Ionone	2.61
C ₁₃ H ₁₂ O	2-Methoxybiphenyl	3.59	C ₁₃ H ₂₀ OS	6-(4-Methylphenyl)thio-1-hexanol	13.103
	2-Benzylphenol	4.98	C ₁₃ H ₂₂ NO ₃	2,2,6,6-Tetramethyl-4-piperidinol-1-oxyl crotonate	15.36
C ₁₃ H ₁₂ OS	1-Methoxy-4-(phenylthio)benzene	13.23		2,2,6,6-Tetramethyl-4-piperidinol-1-oxyl methacrylate	15.40
C ₁₃ H ₁₂ O ₃	2-[(4-Ethoxycarbonyl)phenyl]furan	5.40	C ₁₃ H ₂₄	(E)-2,6,9-Trimethyl-1,6-decadiene	2.171
C ₁₃ H ₁₂ S	1-Methyl-4-(phenylthio)benzene	13.26		(Z)-2,6,9-Trimethyl-1,6-decadiene	2.172
	Benzyl phenyl sulfide	13.33		(E)-2,6-Dimethyl-1,6-undecadiene	2.359
C ₁₃ H ₁₂ Se	Benzyl phenyl selenide	13.32		(Z)-2,6-Dimethyl-1,6-undecadiene	2.360
C ₁₃ H ₁₄ N ₂ O ₂	9-Methyl-6-nitro-2,3,4,9-tetrahydrocarbazole	6.3	C ₁₃ H ₂₄ NO ₃	2,2,6,6-Tetramethyl-4-piperidinol-1-oxyl butyrate	15.35
C ₁₃ H ₁₄ N ₂ O ₃	<i>N</i> -Acetyltryptophan	10.37	C ₁₃ H ₂₅ NO ₂	<i>N</i> -(2-Acetoxyethyl)-2,2,6,6-tetramethylpiperidine	8.69
C ₁₃ H ₁₄ O ₂	α-Phenethylfurfuryl alcohol	5.71	C ₁₃ H ₂₇ NO	1-Butoxy-2,2,6,6-tetramethylpiperidine	8.55
C ₁₃ H ₁₄ S	1,1,3-Trimethyl-2-naphthalenethione	13.114	C ₁₄ H ₈ Br ₂ NiO ₄ .2H ₂ O	Bis(5-bromo-2-hydroxybenzaldehydato)nickel(II) dihydrate	12.14
C ₁₃ H ₁₅ N	9-Methyl-2,3,4,9-tetrahydrocarbazole	6.2	C ₁₄ H ₈ Cl ₂	9,10-Dichloroanthracene	3.7
C ₁₃ H ₁₅ NO ₂	7-(Diethylamino)coumarin	5.6	C ₁₄ H ₈ O ₂	Naphtho[1,8- <i>bc</i> :5,4- <i>b'</i> , <i>c'</i>]dipyran	3.94
C ₁₃ H ₁₅ N ₃ O ₂	<i>N</i> -Acetyl-L-tryptophanamide	10.44	C ₁₄ H ₈ O ₆ S ₂ ²⁻	1,5-Anthracenedisulfonate ion	3.17
C ₁₃ H ₁₅ N ₃ O ₃	Tryptophylglycine	10.21	C ₁₄ H ₉ BrN ₂ O ₄	1,5-Diaminobromo-4,8-dihydroxyanthraquinone	11.12
	Glycyltryptophan	10.41	C ₁₄ H ₉ Cl	1-Chloroanthracene	3.4
C ₁₃ H ₁₆	α-Cyclopropyl-β,β-trimethylstyrene	3.113		9-Chloroanthracene	3.5
C ₁₃ H ₁₆ FN ₂ O ₃	2-(4-Fluoro-3-nitrophenyl)-2,5,5-trimethylpyrrolidin-1-oxyl	15.55			
C ₁₃ H ₁₆ N ₂	Dehydrostobadine	6.58			
C ₁₃ H ₁₆ N ₂ O	5'-Oxo-3'-ethyl-5,4'-dimethyl-1',5'-dihydro-(2,2')-dipyrromethene	7.107			
	5'-Oxo-4-ethyl-3,5-dimethyl-1',5'-dihydro-(2,2')-dipyrromethene	7.108			
C ₁₃ H ₁₆ N ₂ O ₂	L-Tryptophan, ethyl ester	10.40			
C ₁₃ H ₁₆ O ₂	1,2,3,4-Tetrahydro-5,8-dimethoxy-1,4-methanonaphthalene	3.73			
C ₁₃ H ₁₇ NO ₄	3,5-Diethoxycarbonyl-2,6-dimethylpyridine	6.52			
	<i>N</i> -Acetyl-L-tyrosine, ethyl ester	4.188			
C ₁₃ H ₁₈ N ₂	Stobadine	6.56			

$C_{14}H_9NO_2$	1-Aminoanthraquinone	11.3	$C_{14}H_{13}NO$	4-Methylphenyl phenyl nitrone	15.8
$C_{14}H_9NO_3$	2-Aminoanthraquinone	11.4		Phenyl 4-methylphenyl nitrone	15.9
$C_{14}H_9O_3^-$	1-Amino-4-hydroxyanthraquinone	11.5		α -Phenyl-N-benzylnitrone	15.12
	1,8-Dihydroxy-9-anthrone, conjugate base	4.3		Benzophenone oxime O-methyl ether	15.15
$C_{14}H_9O_3S^-$	1-Anthracenesulfonate ion	3.18	$C_{14}H_{13}NOS$	2-(Methylthio)-N-phenylbenzamide	13.3
	2-Anthracenesulfonate ion	3.19	$C_{14}H_{13}NOSe$	2-(Methylseleno)-N-phenylbenzamide	13.2
$C_{14}H_{10}$	Anthracene	3.3	$C_{14}H_{13}N_3O_2$	2-Methyl-6-(4-methoxyphenyl)imidazo[1,2-a]pyrazin-3-one	16.40
$C_{14}H_{10}N_2O_2$	1,4-Diaminoanthraquinone	11.8	$C_{14}H_{14}$	1-(β,β -Dimethylvinyl)naphthalene	3.87
	1,5-Diaminoanthraquinone	11.9	$C_{14}H_{14}N_3^+$	3,6-Diamino-10-methylacridinium	11.2
	1,8-Diaminoanthraquinone	11.10	$C_{14}H_{14}O_3$	4-Acetoxymethyl-2-(phenylmethyl)furan	
	2,6-Diaminoanthraquinone	11.11		5.75	
	3,4-Diphenylsydnone	5.102	$C_{14}H_{14}S$	Dibenzyl sulfide	13.71
$C_{14}H_{10}NiO_4 \cdot 2H_2O$	Bis(2-hydroxybenzaldehydato)nickel(II) dihydrate	12.88	$C_{14}H_{15}NO$	<i>N,N</i> -Dibenzylhydroxylamine	15.11
$C_{14}H_{10}O_3$	1,8-Dihydroxy-9-anthrone	4.2	$C_{14}H_{15}O_2^-$	4-Methyl-1-naphthalenepropionate ion	3.91
$C_{14}H_{11}NO_2Se$	7-Methoxy-2-phenyl-1,2-benzisoselenazol-3-one	13.38	$C_{14}H_{15}O_4^-$	4-Methyl-1-naphthalenepropionate ion endoperoxide	3.92
$C_{14}H_{12}$	(Z)-Stilbene	3.99	$C_{14}H_{16}$	1,2,3,4-Tetrahydro-9-isopropylidene-1,4-methanonaphthalene	2.260
$C_{14}H_{12}Cl_4$	5,6,7,8-Tetrachloro-1,2,3,4-tetrahydro-9-isopropylidene-1,4-methanonaphthalene	2.258	$C_{14}H_{16}N_2O_3$	<i>N</i> -Acetyltryptophan methyl ester	10.38
$C_{14}H_{12}CoS_4$	Bis[4-methyl-1,2-benzenedithiolato]cobalt(II) tetrabutylammonium salt	12.99	$C_{14}H_{16}O_2$	α -(3-Phenylpropyl)furfuryl alcohol	5.73
$C_{14}H_{12}F_4$	5,6,7,8-Tetrafluoro-1,2,3,4-tetrahydro-9-isopropylidene-1,4-methanonaphthalene	2.259	$C_{14}H_{16}O_3$	5-Ethoxycarbonyl-3,4-dihydro-6-phenylpyran	5.100
$C_{14}H_{12}N_2NiO_2$	Bis[2-(iminomethyl)phenolato]nickel(II)		$C_{14}H_{16}O_4$	4-Methyl-1-naphthalenepropionic acid endoperoxide	3.93
	12.96		$C_{14}H_{17}NO_2$	7-(Diethylamino)-4-methylcoumarin	5.7
$C_{14}H_{12}N_2NiO_4$	Bis(2-hydroxybenzaldchyde oximato)nickel(II)	12.89	$C_{14}H_{17}N_3O_3$	L-Alanyl-L-tryptophan	10.39
$C_{14}H_{12}N_2NiS_4$	Bis(phenyldithiocarbamato)nickel(II)	12.107	$C_{14}H_{17}O_4^-$	6-Hydroxy-2,5,7,8-tetramethylchroman-2-carboxylate ion	4.44
$C_{14}H_{12}N_2O$	Diazo(4-methoxyphenyl)phenylmethane	11.20	$C_{14}H_{18}N_2O$	5'-Oxo-3',4'-diethyl-5-methyl-1',5'-dihydro-(2.2')-dipyrrromethene	7.101
$C_{14}H_{12}N_2O_6S_2^{2-}$	4,4'-Diaminostilbene-2,2'-disulfonate ion	3.107		5'-Oxo-3'-ethyl-4',3,5-trimethyl-1',5'-dihydro-(2.2')-dipyrrromethene	7.103
$C_{14}H_{12}N_4O_2$	1,4,5,8-Tetraaminoanthraquinone	11.14		5'-Oxo-4',4,5-trimethyl-3'-ethyl-1',5'-dihydro-(2.2')-dipyrrromethene	7.104
$C_{14}H_{12}NiS_4$	Bis[4-methyl-1,2-benzenedithiolato]nickel(II) tetrabutylammonium salt	12.100	$C_{14}H_{19}NO_3S$	CBZ-L-Methionine methyl ester	13.110
$C_{14}H_{12}O$	(E)-2,3-Diphenyloxirane	17.49	$C_{14}H_{19}N_3O_5$	Glycyl-L-tyrosyl-L-alanine	4.1
$C_{14}H_{12}OS$	4-Methoxythiobenzophenone	13.162	$C_{14}H_{20}N_2$	Caged hydrazine	8.22
$C_{14}H_{12}S$	(E)-2,3-Diphenylthiirane	13.151	$C_{14}H_{20}O_2$	6-Hydroxy-2,2,5,7,8-pentamethylchroman	
				4.49	
				1-Adamantylideneethyl acetate	2.220
			$C_{14}H_{21}BrO$	4-Bromo-2,6-di- <i>tert</i> -butylphenol	4.105
			$C_{14}H_{21}ClO$	2,6-Di- <i>tert</i> -butyl-4-chlorophenol	4.110
			$C_{14}H_{21}NO_4$	3,5-Diethoxycarbonyl-1,4-dihydro-2,4,6-trimethylpyridine	6.51
			$C_{14}H_{22}$	2-Isobutylideneadamantane	2.355
			$C_{14}H_{22}O$	2,6-Di- <i>tert</i> -butylphenol	4.99

$C_{14}H_{22}OS$	3-Hydroxy-3,4-dimethylpentyl <i>p</i> -methylphenyl sulfide	13.119	$C_{15}H_{12}O_6$	Dihydroluteolin	4.59
$C_{14}H_{22}O_2$	3,5-Di- <i>tert</i> -butylcatechol	4.7	$C_{15}H_{12}O_7$	Dihydroquercetin	4.63
	2,5-Di- <i>tert</i> -butylhydroquinone	4.14	$C_{15}H_{12}S_3$	4,5-Diphenyl-1,3-dithiane-2-thione	13.89
$C_{14}H_{26}$	4,8-Dimethyl-4,8-dodecadiene	2.201	$C_{15}H_{13}N$	3-Methyl-2-phenylindole	6.17
	(<i>E,E</i>)-4,8-Dimethyl-4,8-dodecadiene	2.202	$C_{15}H_{14}$	(<i>E</i>)- α -Methylstilbene	3.104
$C_{14}H_{26}NO_3$	2,2,6,6-Tetramethyl-4-piperidinol-1-oxyl pentanoate	15.41		(<i>Z</i>)- α -Methylstilbene	3.105
$C_{14}H_{26}O$	2-(Cyclododecylidene)ethanol	2.215	$C_{15}H_{14}N_2$	Diazodi(4-methylphenyl)methane	11.18
$C_{14}H_{26}O_2$	Monohydroperoxides of 4,8-dimethyl-4,8-dodecadiene	2.203	$C_{15}H_{14}O_2S$	4,4'-Dimethoxythiobenzophenone	13.160
$C_{14}H_{28}CoN_2S_4$	Bis(diisopropylthiocarbamato)cobalt(II)		$C_{15}H_{14}O_6$	Catechin	4.67
	12.46		$C_{15}H_{14}S$	4,4'-Dimethylthiobenzophenone	13.161
$C_{14}H_{28}CuN_2S_4$	Bis(diisopropylthiocarbamato)copper(II)		$C_{15}H_{15}NO_2$	Coumarin LD-490	8.3
	12.47		$C_{15}H_{15}N_3O_2$	1,2,5,6,6a,6b,6c,6d-Octahydro-1,6[1',2']-4-methyl-3,5-dioxo-1,2,4-triazolo[1,3]cyclopropa[1,2,3- <i>cd</i>]cyclopropa[<i>gh</i>]pentalene	17.69
$C_{14}H_{28}FeN_2S_4^+$	Bis(diisopropylthiocarbamato)iron(III)		$C_{15}H_{18}N_2$	1-Isopropylamino-4-phenylaminobenzene	
	12.48			9.30	
$C_{14}H_{28}MnN_2S_4$	Bis(diisopropylthiocarbamato)manganese(II)		$C_{15}H_{18}N_4O_4$	L-Tryptophylglycylglycine	10.22
	12.49		$C_{15}H_{18}N_4O_5$	Mitomycin C	16.7
$C_{14}H_{28}N_2NiS_4$	Bis(diisopropylthiocarbamato)nickel(II)		$C_{15}H_{19}N_3O$	4-Benzylxy-5,6-dimethyl-2-(dimethylamino)pyrimidine	16.65
	12.50		$C_{15}H_{20}N_2O$	<i>N</i> -Acetylstobadine	6.57
$C_{14}H_{28}N_2S_4Zn$	Bis(diisopropylthiocarbamato)zinc(II)		$C_{15}H_{21}CoO_6$	Tris(acetylacetone)cobalt(III)	12.173
	12.51		$C_{15}H_{21}CrO_6$	Tris(acetylacetone)chromate(III)	12.172
$C_{14}H_{28}O_2$	Tetradecanoic acid	17.67	$C_{15}H_{21}FeO_6$	Tris(acetylacetone)iron(III)	12.174
$C_{14}H_{31}CoN_5O_5$	Hydroxybis(dimethylglyoximato)triethylamine-cobalt(II)	12.147	$C_{15}H_{21}MnO_6$	Tris(acetylacetone)manganese(III)	12.175
$C_{15}H_9O_2^-$	9-Anthroate ion	3.21	$C_{15}H_{22}O_2$	7-(1,1-Dimethylethyl)-3,4-dihydro-2,2-dimethyl-1-benzopyran-6-ol	4.54
$C_{15}H_{10}O_3$	3-Hydroxyflavone	4.74	$C_{15}H_{23}NO_4$	1,4-Dihydro-2,6-dimethyl-3,5-di(1-methylethoxycarbonyl)pyridine	6.49
$C_{15}H_{10}O_4$	5,7-Dihydroxyflavone	4.73	$C_{15}H_{24}$	Caryophyllene	2.34
$C_{15}H_{10}O_5$	Galangin	4.65	$C_{15}H_{24}O$	2,6-Di- <i>tert</i> -butyl-4-methylphenol	4.102
$C_{15}H_{10}O_6$	Luteolin	4.60	$C_{15}H_{24}O_2$	3,5-Di- <i>tert</i> -butyl-4-hydroxybenzyl alcohol	
	Kaempferol	4.64		4.32	
	Fisetin	4.75		2,4-Di- <i>tert</i> -butyl-5-methoxyphenol	4.100
$C_{15}H_{10}O_7$	Quercetin	4.61		2,6-Di- <i>tert</i> -butyl-4-methoxyphenol	4.101
$C_{15}H_{11}NO$	2,5-Diphenyloxazole	6.31	$(C_{15}H_{25}NO_4)_n$	Tinuvin 622	8.67
$C_{15}H_{11}NO_2$	1-Amino-2-methylanthraquinone	11.6	$C_{15}H_{26}$	2,6,10-Trimethyl-2,6,10-dodecatriene	2.210
$C_{15}H_{12}$	9-Methylanthracene	3.13	$C_{15}H_{26}O_2$	Hydroperoxy-2,6,10-trimethyl-2,6,10-dodecatriene	2.209
$C_{15}H_{12}N_2O_2$	3-(4-Methylphenyl)-4-phenylsydnone		$C_{15}H_{26}O_4$	Dihydroperoxy-2,6,10-trimethyl-2,6,10-dodecatriene	2.208
	5.106		$C_{15}H_{34}ClN$	Dodecyltrimethylammonium chloride	17.26
$C_{15}H_{12}N_2O_3$	1,4-Diamino-2-methoxyanthraquinone		$C_{16}H_{10}Br_2O$	2,5-Bis(4-bromophenyl)furan	5.16
	11.13		$C_{16}H_{10}Cl_2O$	2,5-Bis(4-chlorophenyl)furan	5.17
$C_{15}H_{12}O$	9-Methoxyanthracene	3.12	$C_{16}H_{10}N_2O_2$	Indigo	11.69
$C_{15}H_{12}O_5$	4',5,7-Trihydroxyflavanone	4.72			

C ₁₆ H ₁₁ BrN ₂ O	1-(4-Bromophenylazo)-2-naphthol	11.94	C ₁₆ H ₁₄ O ₄	2,3-Di(4-hydroxyphenyl)-1,4-dioxene 2.189
C ₁₆ H ₁₁ ClN ₂ O	4-(4-Chlorophenylazo)-1-naphthol	11.87	C ₁₆ H ₁₄ O ₄ ²⁻	1,4-Naphthalenedipropionate ion 3.88
	1-(4-Chlorophenylazo)-2-naphthol	11.95	C ₁₆ H ₁₅ N ₃ O ₂	1-(Phenylazo)-1-(phenylaminocarbonyl)-2-propanone 11.62
C ₁₆ H ₁₁ FN ₂ O	1-(4-Fluorophenylazo)-2-naphthol	11.97	C ₁₆ H ₁₆	1,1-Diphenyl-2-methylpropene 3.46 <i>trans</i> - α,α' -Dimethylstilbene 3.102 <i>cis</i> - α,α' -Dimethylstilbene 3.103
C ₁₆ H ₁₁ IN ₂ O	1-(4-Iodophenylazo)-2-naphthol	11.99	C ₁₆ H ₁₆ ClN ₃ O	2-(2'-Hydroxy-3'-chloro-5'- <i>tert</i> -butylphenyl)benzotriazole 4.97
C ₁₆ H ₁₁ N ₃ O ₃	4-(4-Nitrophenylazo)-1-naphthol	11.91	C ₁₆ H ₁₆ N ₂ NiO ₄	Bis(2'-hydroxyacetophenone oximato)nickel(II) 12.87
	1-(4-Nitrophenylazo)-2-naphthol	11.103	C ₁₆ H ₁₆ N ₂ NiS ₄	Bis(4-methylphenyldithiocarbamato)nickel(II) 12.103
C ₁₆ H ₁₂ CINO	4-Methyl-2-(3-chlorophenyl)-5-phenyloxazole	6.33	C ₁₆ H ₁₆ O ₂	4,4'-Dimethoxystilbene 3.100 α,β -Dimethoxystilbene 3.101
	4-Methyl-2-(4-chlorophenyl)-5-phenyloxazole	6.34	C ₁₆ H ₁₇ NO ₂	2,3,6,7-Tetrahydro-9-methyl[1]benzopyran[6,7,8- <i>ij</i>]quinolizine 8.4
C ₁₆ H ₁₂ Cl ₂ O ₂	2,3-Di(3-chlorophenyl)-1,4-dioxene	2.186	C ₁₆ H ₁₇ N ₅ O ₄	4-(Diethylamino)-2',4'-dinitroazobenzene 11.52
	2,3-Di(4-chlorophenyl)-1,4-dioxene	2.187	C ₁₆ H ₁₈ CuN ₂ S ₄	[[3,3]-[1,2-Ethanediylbis(nitrilomethylidyne)]bis[5-ethyl-2-thiophenethionato]]copper(II) 12.138
C ₁₆ H ₁₂ N ₂ Na ₂ NiO ₈ S ₂	[<i>N,N'</i> -Ethylenebis(5-sulfosalicylideneiminato)]nickelate(II) disodium salt	12.143	C ₁₆ H ₁₈ N ₃ S ⁺	Methylene Blue cation 11.113
C ₁₆ H ₁₂ N ₂ O	4-Phenylazo-1-naphthol	11.92	C ₁₆ H ₁₈ N ₄ O ₂	4-(Diethylamino)-3'-nitroazobenzene 11.54 4-(Diethylamino)-4'-nitroazobenzene 11.55
	1-Phenylazo-2-naphthol	11.104	C ₁₆ H ₁₈ NiO ₃ S	Aqua[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II) 12.4
C ₁₆ H ₁₂ N ₂ O ₂	1-(4-Hydroxyphenylazo)-2-naphthol	11.98	C ₁₆ H ₁₉ N ₃	4-(Diethylamino)azobenzene 11.56
C ₁₆ H ₁₂ N ₂ O ₃	4-Methyl-2-(4-nitrophenyl)-5-phenyloxazole	6.37	C ₁₆ H ₂₀ N ₂ O	5'-Oxo-4'-vinyl-4-ethyl-3',5-trimethyl-1',5'-dihydro-(2,2')-dipyromethene 7.105 5'-Oxo-4-ethyl-3'-ethenyl-3,5,4'-trimethyl-1',5'-dihydro-(2,2')-dipyromethene 7.106
C ₁₆ H ₁₂ O	2,3-Diphenylfuran	5.37	C ₁₆ H ₂₀ N ₂ S ₄	(Z,Z)-3,3'-[1,2-Ethanediylbis(iminomethylidyne)]bis[5-ethyl-2-thiophenethione] 13.171
	2,5-Diphenylfuran	5.38	C ₁₆ H ₂₀ O ₂	5,8-Dimethoxy-1,2,3,4-tetrahydro-9-isopropylidene-1,4-methanonaphthalene 2.257
	3,4-Diphenylfuran	5.39	C ₁₆ H ₂₂ NO ₃	2,2,6,6-Tetramethyl-4-piperidinol-1-oxyl benzoate 15.34
C ₁₆ H ₁₃ NO	2,5-Diphenyl-4-methyloxazole	6.32	C ₁₆ H ₂₂ O ₄	3-(6-Hydroxy-2,5,7,8-tetramethylchroman-2-yl)propionic acid 4.66
C ₁₆ H ₁₃ N ₃ O	1-(<i>p</i> -Aminophenylazo)-2-naphthol	11.93		
C ₁₆ H ₁₄	1,4-Diphenyl-1,3-butadiene	2.40		
	9,10-Dimethylnanthracene	3.10		
	1-Methyl-2-phenylindene	3.71		
C ₁₆ H ₁₄ CoN ₂ O ₂	2,2'-[1,2-Ethanediylbis(nitrilomethylidyne)]-bis[phenolato]cobalt(II)	12.139		
C ₁₆ H ₁₄ N ₂ Na ₂ NiO ₈ S ₂	[<i>N,N'</i> -Propylenebis(5-sulfosalicylideneiminato)]nickelate(II) disodium salt	12.169		
C ₁₆ H ₁₄ N ₂ Na ₂ NiO ₂	2,2'-[1,2-Ethanediylbis(nitrilomethylidyne)]-bis[phenolato]nickel(II)	12.140		
C ₁₆ H ₁₄ N ₂ O ₂	1,4-Bis(methylamino)anthraquinone	11.7		
C ₁₆ H ₁₄ NiO ₆ .2H ₂ O	Bis(2-hydroxy-5-methoxybenzaldehydato)nickel(II) dihydrate	12.92		
C ₁₆ H ₁₄ O ₂	9,10-Dimethoxyanthracene	3.8		
	2,3-Diphenyl-1,4-dioxene	2.194		

C ₁₆ H ₂₄ O ₂	4-Acetyl-2,6-di- <i>tert</i> -butylphenol	4.94	C ₁₇ H ₁₇ N ₃ O ₂	1-(4-Methylphenylazo)-1-(phenylaminocarbonyl)-2-propanone
C ₁₆ H ₂₄ O ₃	Methyl 3,5-di- <i>tert</i> -butyl-4-hydroxybenzoate	4.39		11.60
C ₁₆ H ₂₆ O	2,6-Di- <i>tert</i> -butyl-4-methylanisole	3.29	C ₁₇ H ₁₈ N ₄	4-[N-(2-Cyanoethyl)-N-ethylamino]azobenzene
C ₁₆ H ₂₆ O ₂	1,3-Dimethoxy-4,6-di- <i>tert</i> -butylbenzene	3.28	C ₁₇ H ₁₈ O ₄	4-Methyl-1,3-naphthalenedipropionic acid
	2,5-Di- <i>tert</i> -amylhydroquinone	4.16		3.89
C ₁₆ H ₃₂ O ₂	Hexadecanoic acid	17.37	C ₁₇ H ₁₉ ClN ₂ S	Chlorpromazine
C ₁₇ H ₁₀ O	Benzanthrone	3.25	C ₁₇ H ₁₉ ClO	2-(3-Chlorophenoxy)methylidene)adamantane
C ₁₇ H ₁₂ N ₂ PtS ₂	[4-Methyl-1,2-benzenedithiolato](1,10-phenanthroline)platinum(II)	12.155		2.350
C ₁₇ H ₁₂ OS	2,6-Diphenylpyran-4-thione	13.140		2-(4-Chlorophenoxy)methylidene)adamantane
C ₁₇ H ₁₂ S ₂	2,6-Diphenylthiopyran-4-thione	13.173		2.351
C ₁₇ H ₁₃ Cl ₂	1,4-Bis(4-chlorophenyl)-1,3-cyclopentadiene	2.141	C ₁₇ H ₂₀ N ₄ O ₆	Riboflavin
C ₁₇ H ₁₃ F ₂	1,4-Bis(4-fluorophenyl)-1,3-cyclopentadiene	2.142	C ₁₇ H ₂₁ N ₃ O	4-(Diethylamino)-4'-methoxyazobenzene
C ₁₇ H ₁₃ N ₃ O ₃	1-(4-Methyl-2-nitrophenylazo)-2-naphthol	11.101		11.53
C ₁₇ H ₁₄ Cl	1-(4-Chlorophenyl)-4-phenyl-1,3-cyclopentadiene	2.146	C ₁₇ H ₂₁ N ₃ O ₂	4-[N-Ethyl-N-(2-hydroxyethyl)amino]-4'-methoxyazobenzene
C ₁₇ H ₁₄ N ₂ NiS ₂	[4-Methyl-1,2-benzenedithiolato](2,2'-bipyridine)nickel(II)	12.153	C ₁₇ H ₂₂ NO ₂	4-Hydroxy-4-(2-phenylethynyl)-2,2,6,6-tetramethylpiperidine N-oxyl
C ₁₇ H ₁₄ N ₂ O	1-Methoxy-4-(phenylazo)naphthalene	11.59	C ₁₇ H ₂₃ NO ₂	3,4,6,7,9,10-Hexahydro-3,3,6,6-tetramethyl-1,8-acridinedione
	1,4-Naphthoquinone N-methyl-N-phenylhydrazone	11.83	C ₁₇ H ₂₃ NO ₃	Atropine
	4-(4-Methylphenylazo)-1-naphthol	11.90	C ₁₇ H ₂₄ N ₂ O	5'-Oxo-3',4',4-triethyl-3,5-dimethyl-1',5'-dihydro-(2,2')-dipyrromethene
	1-(4-Methylphenylazo)-2-naphthol	11.102	C ₁₇ H ₂₄ N ₄ NiO ₂	1,1'-(7,13-Dimethyl-1,4,8,12-tetraazacyclopentadeca-4,7,12,15-tetraene-6,14-diyl)bis[ethanoato]nickel(II)
C ₁₇ H ₁₄ N ₂ O ₂	4-(4-Methoxyphenylazo)-1-naphthol	11.89	C ₁₇ H ₂₅ NO ₃	Eucatropine
	1-(4-Methoxyphenylazo)-2-naphthol	11.100	C ₁₇ H ₂₆ O ₃	3,5-Di- <i>tert</i> -butyl-4-hydroxyphenyl propionate
C ₁₇ H ₁₄ O ₂	α,α -Diphenylfurfuryl alcohol	5.67	C ₁₈ H ₁₁ NO ₂	Quinophthalone
C ₁₇ H ₁₅	1,4-Diphenyl-1,3-cyclopentadiene	2.147	C ₁₈ H ₁₂	Benz[a]anthracene
C ₁₇ H ₁₅ NO	4-Methyl-2-(4-methylphenyl)-5-phenyloxazole	6.36	C ₁₈ H ₁₂ O	Tetracene
C ₁₇ H ₁₅ NO ₂	4-Methyl-2-(4-methoxyphenyl)-5-phenyloxazole	6.35	C ₁₈ H ₁₄ N ₃ O ₆ S ⁻	2-Phenylcyclopenta[b][1]benzopyran
C ₁₇ H ₁₅ NO ₃ S	(Z)-1-(Phenylsulfonyl)-3-(β -methoxylvinyl)indole	2.249		5-Methoxy-4-[2-(methylsulfonyl)-4-nitrophenyl]azo-1-naphthoxide ion
C ₁₇ H ₁₅ N ₃ O ₂	4-(4'-Methoxyphenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one	11.148	C ₁₈ H ₁₅ ClO	1-(4-Chlorophenyl)-4-(4-methoxyphenyl)-1,3-cyclopentadiene
C ₁₇ H ₁₅ N ₅	1,3-Dimethylpyrazolo[1',2':2,3][1,2,3]-triazolo[4,5- <i>a</i>]phenazin-4-iium	6.39	C ₁₈ H ₁₅ N	2.145
C ₁₇ H ₁₆ N ₄ O	4-(4'-Aminophenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one	11.133	C ₁₈ H ₁₅ P	Triphenylamine
C ₁₇ H ₁₆ O ₃	2-(4-Methoxyphenyl)-3-phenyl-1,4-dioxene	2.195	C ₁₈ H ₁₆	Triphenylphosphine
				1,2,5,6,6 <i>a</i> ,6 <i>b</i> ,6 <i>c</i> ,6 <i>d</i> -Octahydro-1,6- <i>o</i> -benzenobenzo[1,3]cyclopropa[1,2,3- <i>cd</i>]cyclopropa[<i>gh</i>]pentalene
			C ₁₈ H ₁₆ N ₂	2.181
				N,N'-Diphenyl- <i>p</i> -phenylenediamine
				9.29

$C_{18}H_{16}N_2O_8S_2^{2-}$	4,4'-Diacetamidostilbene-2,2'-disulfonate ion	3.106	$C_{18}H_{22}O$	2-(4-Methylphenoxy(methylidene)adamantane
$C_{18}H_{16}O$	2,5-Bis(4-methylphenyl)furan	5.20	$C_{18}H_{22}O_2$	2-(4-Methoxyphenoxy(methylidene)adamantane
	1-(4-Methoxyphenyl)-4-phenyl-1,3-cyclopentadiene	2.148		2.352
$C_{18}H_{16}O_2$	α -Benzhydrylfuryl alcohol	5.68	$C_{18}H_{23}NNiO_2S$	Ethylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II)
	5-Benzoyl-3,4-dihydro-6-phenylpyran	5.89		12.141
$C_{18}H_{16}O_3$	2,5-Bis(4-methoxyphenyl)furan	5.19	$C_{18}H_{24}O_2$	<i>exo,exo</i> -2,3-Dioxymethyl-7-adamantylidenenorbornane
$C_{18}H_{16}O_6S_2^{2-}$	9,10-Anthracenediethanesulfonate ion	3.15	$C_{18}H_{26}O_2$	2,10
$C_{18}H_{17}NO$	β -Phenyl- α -pyrrolidinylbenzeneethanol	8.2		1,2,3,8,9,10-Hexahydro-3,3,5,6,8,8-hexamethylbenzo[1,2- <i>b</i> :4,3- <i>b'</i>]dipyran
$C_{18}H_{17}N_3O$	1-[4-(Dimethylamino)phenylazo]-2-naphthol	11.96		5.2
				2,3,4,7,8,9-Hexahydro-2,2,5,7,7,10-hexamethylbenzo[1,2- <i>b</i> :4,5- <i>b'</i>]dipyran
$C_{18}H_{18}N_2NiO_2$	2,2'-(1,2-Ethanediylbis(nitriloethylidyne))bis[phenolato]nickel(II)	12.137	$C_{18}H_{28}N_2S_3$	5.3
$C_{18}H_{18}N_3O_7$	2,5-Di-(4-hydroxy-3-nitrophenyl)-2,5-dimethylpyrrolidin-1-oxyl	15.54		3,4-Bis[(cyclohexylamino)methylene]dihydro-2,5-thiophenedithione
$C_{18}H_{18}N_4O$	4-(4'-Dimethylaminophenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one	11.150	$C_{18}H_{30}O$	13.169
$C_{18}H_{18}O_2$	1,4-Di(4,4'-dimethoxyphenyl)-1,3-butadiene	2.35	$C_{18}H_{30}O_2$	2,4,6-Tri- <i>tert</i> -butylphenol
	2,3-Di(4-methylphenyl)-1,4-dioxene	2.193		4- <i>tert</i> -Butoxy-2,6-di- <i>tert</i> -butylphenol
$C_{18}H_{18}O_4$	2,3-Di(3-methoxyphenyl)-1,4-dioxene	2.190	$C_{18}H_{32}O_2$	4.147
	2,3-Di(4-methoxyphenyl)-1,4-dioxene	2.191	$C_{18}H_{33}O_3P$	Linolenic acid
$C_{18}H_{19}NO_2$	2-(4-Dimethylaminophenyl)-3-phenyl-1,4-dioxene	2.192	$C_{18}H_{34}O_2$	2.267
$C_{18}H_{20}B_2N_{12}Ni$	Bis[hydrotris(1-pyrazolyl)borato]nickel(II)	12.86	$C_{18}H_{36}CoN_2S_4$	Oleic acid
				12.26
$C_{18}H_{20}N_2NiO_4$	Bis[2'-hydroxy-4'-methylacetophenone oximato]nickel(II)	12.93	$C_{18}H_{36}CuN_2S_4$	Bis(dibutylthiocarbamato)cobalt(II)
$C_{18}H_{20}N_4O$	4-[<i>N</i> -(2-Cyanoethyl)- <i>N</i> -ethylamino]-4'-methoxyazobenzene	11.50		12.27
$C_{18}H_{20}O_2$	3-Hydroxy-1,3,5(10),9(11)-estratetraen-17-one	16.31	$C_{18}H_{36}N_2NiS_4$	Bis(dibutylthiocarbamato)copper(II)
$C_{18}H_{20}O_4$	6,11-Dimethyl4,6,8,10,12-hexadecapentaene-2,3,14,15-tetraone (<i>all-E</i>)	2.234		12.28
$C_{18}H_{22}N_2$	1-Cyclohexylamino-4-phenylaminobenzene	9.28	$C_{18}H_{36}N_2S_4Zn$	Bis(dibutylthiocarbamato)nickel(II)
$C_{18}H_{22}N_4Ni$	Bis[<i>N</i> -methyl-7-(methylimino)-1,3,5-cycloheptatrien-1-aminato]nickel(II)	12.102		12.29
$C_{18}H_{22}NiO_6S_2$	Bis(2,4,6-trimethylbenzenesulfonato)nickel(II)	12.118	$C_{18}H_{36}O_2$	Stearic acid
			$C_{19}H_{13}NO$	17.66
			$C_{19}H_{13}O_2S^-$	Phenyl fluorenyl nitrone
			$C_{19}H_{14}N_2NiS_2$	15.6
				9-(Phenylsulfonyl)fluoren-9-yl anion
				13.98
			$C_{19}H_{14}S$	[4-Methyl-1,2-benzenedithiolato](1,10-phenanthroline)nickel(II)
				12.154
			$C_{19}H_{15}NO$	4-Phenylthiobenzophenone
			$C_{19}H_{16}N_2O_2$	13.163
				Triphenyl nitrone
				15.5
			$C_{19}H_{18}N_2O_2$	2-(4-Ethoxycarbonylphenyl)-1,8-dihydropyrrolo[3',2':3,4]cyclopenta[1,2- <i>b</i>]pyridine
				6.70
			$C_{19}H_{19}$	1,4-Dioxene, 2-(4-cyanophenyl)-3-(4-dimethylaminophenyl)-
				2.184
				1,4-Bis(4-methylphenyl)-1,3-cyclopentadiene
				2.144

C ₁₉ H ₁₉ O ₂	1,4-Bis(4-methoxyphenyl)-1,3-cyclopentadiene	2.143	C ₂₀ H ₂₀ O ₂	1,2,5,6,6a,6b,6c,6d-Octahydro-9,12-dimethoxy-1,6-o- benzenobenzo[1,3]cyclopropa[1,2,3- cd]cyclopropa[gh]pentalene	2.180
C ₁₉ H ₂₁ NO ₄	Boldine	8.20	C ₂₀ H ₂₀ O ₇	Tangeretin	4.62
C ₁₉ H ₂₂ N ₂	Bis(2,4,6-trimethylphenyl)diazomethane	11.19	C ₂₀ H ₂₂ N ₄ O	4-(4'-Amino-2',3',5',6'- tetramethylphenyl)imino-3-methyl-1- phenyl-2-pyrazolin-5-one	11.134
C ₁₉ H ₂₂ N ₂ O ₃	Dregamine	8.101		4-(4'-Diethylaminophenyl)imino-3-methyl-1- 1-phenyl-2-pyrazolin-5-one	11.147
C ₁₉ H ₂₂ O ₂	3-Methoxyestra-1,3,5(10),8-tetraen-17-one	16.28		4-(4'-Dimethylamino-3',5'- dimethylphenyl)imino-3-methyl-1- phenyl-2-pyrazolin-5-one	11.149
	3-Methoxy-1,3,5(10),9(11)-estratetraen-17-one	16.32	C ₂₀ H ₂₂ N ₄ O ₂	2,4-Dihydro-4-[4-[(2-hydroxyethyl)ethylamino]phenyl]imino- 5-methyl-2-phenylpyrazol-3-one	11.130
C ₁₉ H ₂₂ O ₃	<i>endo</i> -7-Adamantylidenenorbornane-2,3-dicarboxylic acid anhydride	2.15	C ₂₀ H ₂₂ O ₃	3-Acetoxy-1,3,5(10),9(11)-estratetraen-17-one	16.29
	<i>exo</i> -7-Adamantylidenenorbornane-2,3-dicarboxylic acid anhydride	2.16	C ₂₀ H ₂₄ ClN ₃ O	2-(3',5'-Di- <i>tert</i> -butyl-2'-hydroxyphenyl)-5-chlorobenzotriazole	4.109
C ₁₉ H ₂₂ O ₄	Dimethyl 4-methyl-1,3-naphthalenedipropionate	3.90	C ₂₀ H ₂₄ N ₂ NiO ₂	Bis[2-[(1-methylethylimino)methyl]-phenolato]nickel(II)	12.101
C ₁₉ H ₂₄ NO ₂	4-Hydroxy-4-(2-naphthyl)-2,2,6,6-tetramethylpiperidine-1-oxyl	15.31	C ₂₀ H ₂₄ N ₂ O ₂	1,4-Dioxene, 2,3-di(4-dimethylaminophenyl)-	2.188
C ₁₉ H ₃₀ O ₂	Methyl arachidonate	2.213	C ₂₀ H ₂₄ O ₄	Crocin	2.93
C ₁₉ H ₃₂ O ₂	Methyl linolenate	2.269	C ₂₀ H ₂₆ NO	Bis(4- <i>tert</i> -butylphenyl)nitroxide	15.26
C ₁₉ H ₃₄ O ₂	Methyl linoleate	2.265	C ₂₀ H ₂₆ N ₂ O	N,N-Diethyl-3,5-dimethyl-4-(4'-oxo-2',6'-dimethylcyclohexadienyliden)aminoaniline	11.49
C ₁₉ H ₃₆ O ₂	Methyl oleate	2.272	C ₂₀ H ₂₆ O	4-Phenyl-2,6-di- <i>tert</i> -butylphenol	4.103
C ₁₉ H ₃₈ O ₂	Methyl stearate	16.43	C ₂₀ H ₂₇ NNiO ₂ S	Butylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II)	12.120
C ₂₀ H ₂ Cl ₄ I ₄ O ₅ ²⁻	Rose Bengal dianion	11.66	C ₂₀ H ₂₈	Adamantylideneadamantane	2.356
C ₂₀ H ₆ Br ₄ O ₅ ²⁻	Eosin dianion	11.65	C ₂₀ H ₂₈ O	<i>exo,exo</i> -2,3-(2 ¹ -Oxatrimethylene)-7-adamantylidenenorbornane	2.12
C ₂₀ H ₈ Br ₂ HgNa ₂ O ₆	Mercurochrome	11.64		11-(Z)-Retinal	2.339
C ₂₀ H ₁₂ N ₂ Na ₂ NiO ₈ S ₂	[N,N'- <i>o</i> -Phenylenebis(5-sulfosalicylideneiminato)]nickelate(II) disodium salt	12.167		13-(Z)-Retinal	2.340
C ₂₀ H ₁₂ N ₄ Zn	Porphinatozinc(II)	7.71		(all-E)-Retinal	2.341
C ₂₀ H ₁₄	9-Phenylanthracene	3.14	C ₂₀ H ₂₈ O ₂	(all-E)-Retinoic acid	2.342
C ₂₀ H ₁₄ N ₂ NiO ₂	2,2'-(1,2-Phenylenebis(nitrilomethylidyne))-bis[phenolato]nickel(II)	12.166	C ₂₀ H ₃₀ O	(all-E)-Retinol	2.343
C ₂₀ H ₁₄ O	1,3-Diphenylisobenzofuran	5.83	C ₂₀ H ₃₀ O ₂	17-Hydroxy-4-methyl-4-androsten-3-one	16.4
C ₂₀ H ₁₆	9,10-Dimethylbenz[a]anthracene	3.24		Abiotic acid	2.325
C ₂₀ H ₁₆ O ₄ ²⁻	9,10-Anthracenedipropionate ion	3.16	C ₂₀ H ₃₂ O ₂	Arachidonic acid	2.212
C ₂₀ H ₁₇ NOS	N-Phenyl-2-(phenylmethyl)thiobenzamide	13.5	C ₂₀ H ₃₄	2,6,10,14-Tetramethyl-2,6,10,14-hexadecatetraene	2.235
C ₂₀ H ₁₇ NOSe	N-Phenyl-2-(phenylmethyl)selenobenzamide	13.4	C ₂₀ H ₃₄ AuO ₉ PS	Tetra-O-acetylglucose-1-thiolato(triethylphosphino)gold(I)	12.170
C ₂₀ H ₂₀ N ₂ O ₄	2,3-Di(4-acetaminophenyl)-1,4-dioxene	2.185			

C ₂₀ H ₃₄ O ₂	Ethyl linolenate	2.268	C ₂₁ H ₂₉ N ₇ O ₁₄ P ₂	Nicotinamide adenine dinucleotide, reduced
C ₂₀ H ₄₀ O	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	2.238		16.48
C ₂₀ H ₄₀ O ₂	Eicosanoic acid	17.27	C ₂₁ H ₃₀ N ₇ O ₁₇ P ₃	Nicotinamide-adenine dinucleotide phosphate, reduced
C ₂₁ H ₁₄	1,2-Diphenylindene	3.70		16.50
C ₂₁ H ₁₇ ClN ₂	3-(4-Chlorophenyl)-1,5-diphenyl-2-pyrazoline	11.121	C ₂₁ H ₃₀ O ₃	7-Dehydroandrosterone-3-acetate
	5-(4-Chlorophenyl)-1,3-diphenyl-2-pyrazoline	11.122	C ₂₁ H ₃₁ NO ₄	3,5-Dicyclohexyloxycarbonyl-1,4-dihydro-2,6-dimethylpyridine
C ₂₁ H ₁₇ DN ₂	5-Deutero-1,3,5-triphenyl-2-pyrazoline	11.123	C ₂₁ H ₃₂ O ₂	Pregnenolone
C ₂₁ H ₁₈ BrNO	N-Benzhydryl-4-bromomethylbenzamide	3.22	C ₂₁ H ₃₅ NO ₄	3,5-Dihexyloxycarbonyl-1,4-dihydro-2,6-dimethylpyridine
C ₂₁ H ₁₉ BrCl ₂ N ₂ S ₂	5,5'-Dichloro-3,3-diethyl-2,2'-thiacarbocyanine bromide	11.163	C ₂₁ H ₃₆ N ₇ O ₁₆ P ₃ S	Coenzyme A
C ₂₁ H ₁₉ N ₃ O ₂	1,4-Dihydro-2,6-dimethyl-3,5-diphenylaminocarbonylpyridine	6.44	C ₂₁ H ₄₂ O	Methyl phytol ether
C ₂₁ H ₂₀ N ₂ O	4-[2-(N,N-Dimethylhydrazono)ethylidene]-2,6-diphenylpyran	15.1	C ₂₂ H ₁₄	Dibenz[a,h]anthracene
C ₂₁ H ₂₀ O ₆	Curcumin	4.77		Pentacene
C ₂₁ H ₂₁ N	Tribenzylamine	8.6	C ₂₂ H ₁₆ N ₂ O	4-(2',3'-Benzo-4'-oxocyclohexadienyliden)amino-N-phenylaniline
C ₂₁ H ₂₂ N ₂ O ₂	Strychnine	8.99		11.84
C ₂₁ H ₂₂ N ₆	Pyrazole dye 4	11.24	C ₂₂ H ₁₈ O	1,3-Diphenyl-5,6-dimethylisobenzofuran
C ₂₁ H ₂₃ N ₇	Pyrazole dye 6	11.21		5.82
C ₂₁ H ₂₄ N ₄ O	4-(4'-Diethylamino-2'-methylphenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one	11.144	C ₂₂ H ₂₀ N ₂ O	3-(4-Methylphenyl)-1,5-diphenyl-2-pyrazoline
				11.126
C ₂₁ H ₂₄ N ₄ O ₂	2,4-Dihydro-4-[4-[ethyl(2-hydroxyethyl)amino]-2-methylphenyl]imino-5-methyl-2-phenylpyrazol-3-one	11.129		5-(4-Methylphenyl)-1,3-diphenyl-2-pyrazoline
				11.127
C ₂₁ H ₂₅ NO ₄	Glaucine	8.21	C ₂₂ H ₂₂ Cl ₃ N ₅ O ₂	3-(4-Methoxyphenyl)-1,5-diphenyl-2-pyrazoline
C ₂₁ H ₂₆ N ₂ O ₃	Vincamine	8.100		11.124
C ₂₁ H ₂₆ N ₇ O ₁₃ P ₂ ⁺	Nicotinamide adenine dinucleotide	16.47	C ₂₂ H ₂₃ N ₃ O ₂	3,4-Dihydro-2,4,6-trimethyl-3,5-diphenylaminocarbonylpyridine
C ₂₁ H ₂₆ O ₃	2-Hydroxy-4-octyloxybenzophenone	4.43		6.45
	4-(1,1,3,3-Tetramethylbutyl)phenyl salicylate	4.144	C ₂₂ H ₂₄ BrN ₅ O ₂	2,4-Dihydro-2-(4-bromophenyl)-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamylpyrazol-3-one
C ₂₁ H ₂₈ N ₆ O ₁₅ P ₂	Nicotinamide hypoxanthine dinucleotide, reduced	16.42		11.135
C ₂₁ H ₂₈ N ₇ O ₁₇ P ₃	Nicotinamide adenine dinucleotide phosphate	16.49	C ₂₂ H ₂₄ ClN ₅ O ₂	2,4-Dihydro-2-(3-chlorophenyl)-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamylpyrazol-3-one
				11.136
C ₂₁ H ₂₈ O	4-Benzyl-2,6-di- <i>tert</i> -butylphenol	4.104		2,4-Dihydro-2-(4-chlorophenyl)-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamylpyrazol-3-one
C ₂₁ H ₂₈ O ₄	<i>endo</i> -2,3-Di(methoxycarbonyl)-7-adamantylidenenorbornane	2.17		11.137
	<i>exo</i> -2,3-Di(methoxycarbonyl)-7-adamantylidenenorbornane	2.18	C ₂₂ H ₂₄ N ₂ O ₂	2,4-Di[2-methyl-4-(dimethylamino)phenyl]squarylium
				11.47
			C ₂₂ H ₂₄ N ₆	Pyrazole dye 3
				11.22

$C_{22}H_{24}N_6O_4$	2,4-Dihydro-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2-(4-nitrophenyl)pyrazol-3-one	11.145	$C_{23}H_{20}N_2$	4,5-Dihydro-1,5-diphenyl-3-(2-phenylethenyl)pyrazole	11.119
$C_{22}H_{25}N_5O_2$	2,4-Dihydro-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2-phenylpyrazol-3-one	11.128	$C_{23}H_{21}IN_2S_2$	3,8,3',10'-Di(1,3-propanediyl)-2,2'-thiacarbocyanine iodide	11.156
$C_{22}H_{26}Br_2N_2NiO_2$	Bis[2-[(butylimino)methyl]-4-bromophenolato]nickel(II)	12.17	$C_{23}H_{23}BrCl_2N_2S_2$	5,5'-Dichloro-3,9,3'-triethyl-2,2'-thiacarbocyanine bromide	11.164
$C_{22}H_{26}N_4O$	4-(4'-Diethylamino-2',6'-dimethylphenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one	11.140	$C_{23}H_{23}IN_2$	1,1'-Diethyl-2,2'-cyanine iodide	11.45
	4-(4'-Dimethylamino-2',3',5',6'-tetramethylphenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one	11.151	$C_{23}H_{23}IN_2S_2$	3,3'-Diethyl-2,2'-thiadicarbocyanine iodide	11.173
$C_{22}H_{26}O_3$	Bioresmethrin	5.76	$C_{23}H_{23}NO_2$	3,5-Dibenzoyl-1,4-dihydro-2,6-dimethylpyridine	6.43
$C_{22}H_{27}N_5O_3S$	2,4-Dihydro-4-[4-[(2-methylsulfamylethyl)ethylamino]-2-methylphenyl]imino-5-methyl-2-phenylpyrazol-3-one	11.78	$C_{23}H_{23}NO_4$	3,5-Diethoxycarbonyl-1,4-dihydro-2,6-diphenylpyridine	6.50
$C_{22}H_{28}N_2NiO_2$	Bis[2-[(butylimino)methyl]phenolato]nickel(II)	12.19	$C_{23}H_{25}BrN_2S_2$	3,9,3'-Triethyl-2,2'-thiacarbocyanine bromide	11.171
	Bis[2-[(1,1-dimethylethylimino)methyl]phenolato]nickel(II)	12.57	$C_{23}H_{25}N_3$	4-Diethylaminobenzaldehyde diphenylhydrazone	15.4
	Bis[2-[(1-methylpropylimino)methyl]phenolato]nickel(II)	12.105	$C_{23}H_{26}N_2$	Leucomalachite Green	11.77
$C_{22}H_{29}NNiO_2S$	Cyclohexylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II)	12.130	$C_{23}H_{26}N_2O_4$	Brucine	8.9
$C_{22}H_{29}N_3O$	2-(2'-Hydroxy-3',5'-di- <i>tert</i> -pentylphenyl)benzotriazole	4.96	$C_{23}H_{27}N_5O_2$	2,4-Dihydro-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2-(3-methylphenyl)pyrazol-3-one	11.143
$C_{22}H_{32}O_2$	Docosahexaenoic acid	2.200	$C_{23}H_{27}N_5O_3$	2,4-Dihydro-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2-(3-methoxyphenyl)pyrazol-3-one	11.142
	Retinyl acetate	2.344	$C_{23}H_{28}O_4$	2,2'-Spirobis[3,4-dihydro-6-hydroxy-4,4,7-trimethyl-1-benzopyran]	4.185
$C_{22}H_{32}O_6$	6,11-Dimethyl-2,2,15,15-tetramethoxy-4,6,8,10,12-hexadecapentaene-3,14-dione (<i>all-E</i>)	2.233	$C_{23}H_{29}N_3O_2$	<i>N,N</i> -Diethyl-4-[<i>N</i> -(phenylaminocarbonyl)-pivaloylmethylene]aminoaniline	11.112
$C_{22}H_{38}O_2$	2,5-Di- <i>sec</i> -octylhydroquinone	4.19	$C_{23}H_{30}N_2O_8$	4,4'-Dimethyl-3,3',5,5'-tetracarbethoxy-2,2'-dipyrrymethane	6.65
$C_{22}H_{42}O_2$	Phytol acetate	2.239	$C_{23}H_{30}O_3$	RO 10-9359	2.262
$C_{22}H_{44}O$	Ethyl phytol ether	2.236	$C_{23}H_{32}O_2$	2,2'-Methylenebis(6- <i>tert</i> -butyl-4-methylphenol)	4.136
$C_{22}H_{44}O_2$	Docosanoic acid	17.22	$C_{23}H_{32}O_3$	16-Dehydropregnolone-3-acetate	16.61
$C_{23}H_{17}H_5O_9S_3^{2-}$	<i>N</i> -[5-Hydroxy-8-[(2-(methylsulfonyl)-4-nitrophenyl)azo]-1-naphthalenyl-1,3-benzenedisulfonamide, conjugate dibase	11.86	$C_{23}H_{36}N_2O_2$	4-(<i>N</i> -Dipivaloylmethylene)amino-3, <i>N,N</i> -triethylaniline	11.67
$C_{23}H_{18}H_5O_9S_3^-$	<i>N</i> -[5-Hydroxy-8-[(2-(methylsulfonyl)-4-nitrophenyl)azo]-1-naphthalenyl-1,3-benzenedisulfonamide, conjugate base	11.85	$C_{23}H_{46}$	(<i>Z</i>)-9-Tricosene	2.349
			$C_{24}H_{16}N_2NiO_2$	2,2'-[1,8-Naphthylenebis(nitrilomethylidyne)]-bis[phenolato]nickel(II)	12.160
			$C_{24}H_{18}O$	2,4,6-Triphenylphenol	4.149
			$C_{24}H_{19}N_3O_2$	1-(4-Methylphenylazo)-3-(phenylaminocarbonyl)-2-naphthol	11.80

$C_{24}H_{20}CoO_4P_2S_4$	$C_{24}H_{32}N_2NiO_4$
Bis(O,O' - diphenylphosphorodithiolato)cobalt(II) 12.68	Bis[2-[(butylimino)methyl]-4-methoxyphenolato]nickel(II) 12.18
$C_{24}H_{20}CrO_4P_2S_4$	$C_{24}H_{36}O_2$
Bis(O,O' - diphenylphosphorodithiolato)chromium(II) 12.67	Phenyl linoleate 2.266
$C_{24}H_{20}CuO_4P_2S_4$	$C_{24}H_{38}O_2$
Bis(O,O' - diphenylphosphorodithiolato)copper(II) 12.69	Phenyl oleate 2.273
$C_{24}H_{20}NiO_4P_2S_4$	$C_{24}H_{42}N_2O_6$
Bis(O,O' - diphenylphosphorodithiolato)nickel(II) 12.71	Bis(2,2,6,6-tetramethyl-4-piperidinol-1-oxyl)-1,6-hexanedioate 15.38
$C_{24}H_{20}O_4P_2PbS_4$	$C_{24}H_{44}CoO_4P_2S_4$
Bis(O,O' - diphenylphosphorodithiolato)lead(II) 12.70	Bis(O,O' - dicyclohexylphosphorodithiolato)cobalt(II) 12.36
$C_{24}H_{20}O_4P_2S_4Zn$	$C_{24}H_{44}NiP_2S_4$
Bis(O,O' - diphenylphosphorodithiolato)zinc(II) 12.72	Bis(dicyclohexylphosphinodithiolato)nickel(II) 12.35
$C_{24}H_{22}CoN_4S_4$	$C_{24}H_{50}N_4$
Bis[5-ethyl-3-[(3-pyridinylmino)methyl]-2-thiophenethionato]cobalt(II) 12.84	N,N' -Bis[4-(2,2,6,6-tetramethylpiperidinyl)]-1,6-hexanediamine 8.64
$C_{24}H_{25}BrN_2O$	$(C_{24}H_{50}N_4)_n$
1-(4-Bromophenyl)-4,4,8,8-tetramethyl-2,3,4,5,7,8,9,10-octahydropyrrolo[4,3,2- <i>m,n</i>]acridine-10-one 6.67	Poly(N,N' -Bis[4-(2,2,6,6-tetramethylpiperidinyl)]-1,6-hexanediamine) 8.65
$C_{24}H_{25}IN_2S_2$	$C_{25}H_{19}N$
3,3'-Diethyl-8,9-(1,3-propanediyl)-2,2'-thiacarbocyanine iodide 11.170	1,2-Diphenyl-4-methylcyclopenta[b]quinoline 6.4
$C_{24}H_{26}N_2O$	$C_{25}H_{20}N_4O_4$
1-Phenyl-4,4,8,8-tetramethyl-2,3,4,5,7,8,9,10-octahydropyrrolo[4,3,2- <i>m,n</i>]acridine-10-one 6.69	1-(4-Aminocarbonylphenylazo)-3-(2-methoxyphenylaminocarbonyl)-2-naphthol 11.79
$C_{24}H_{26}N_6O_2$	$C_{25}H_{23}BF_4N_4S_2$
Pyrazole dye 2 11.155	5,5'-Dicyano-3,9,3'-triethyl-2,2'-thiacarbocyanine tetrafluoroborate 11.165
$C_{24}H_{28}N_2O_4$	$C_{25}H_{24}N_2O_2$
2,4-Di[2-hydroxy-4-(diethylamino)phenyl]squarylium 11.46	4,5-Dihydro-5-(4-methoxyphenyl)-3-[2-(4-methoxyphenyl)ethenyl]-1-phenylpyrazole 11.120
$C_{24}H_{28}O_4$	$C_{25}H_{24}N_6S$
6,6'-Diapo- ψ,ψ -carotenedioic acid 2.174 9-cis-6,6'-Diapo- ψ,ψ -carotenedioic acid 2.179	Pyrazole dye 5 11.25
$C_{24}H_{30}N_4O$	$C_{25}H_{24}O_3$
4-(4'-Diethylamino-2'-methylphenyl)imino-3-(2-methyl-2-propyl)-1-phenyl-2-pyrazolin-5-one 11.141	3-Benzoyloxy-1,3,5(10),9(11)-estratetraen-17-one 16.30
$C_{24}H_{31}N_3O_2$	$C_{25}H_{25}ClN_2$
<i>N,N</i> -Diethyl-3-methyl-4-[<i>N</i> -(phenylaminocarbonyl)-pivaloylmethylene]aminoaniline 11.111	1,1'-Diethyl-2,2'-carbocyanine chloride 11.41
$C_{24}H_{32}NO_3S_2^-$	$C_{25}H_{25}IN_2O_2$
10-(12-Sulfonatododecyl)phenothiazine 13.124	3,3'-Diethyl-2,2'-oxatricarbocyanine iodide 11.109
	$C_{25}H_{25}IN_2S_2$
	3,3'-Diethyl-2,2'-thiatricarbocyanine iodide 11.174
	$C_{25}H_{25}N_2^+$
	1,1'-Diethyl-4,4'-carbocyanine 11.42
	$C_{25}H_{25}N_3O_2$
	4-[<i>N</i> -Benzoyl(phenylaminocarbonyl)-methylene]amino- <i>N,N</i> -diethylaniline 11.28
	$C_{25}H_{26}O_5S$
	3-Tosyloxy-1,3,5(10),9(11)-estratetraen-17-one 16.33
	$C_{25}H_{28}N_2O_2$
	1-(4-Methoxyphenyl)-4,4,8,8-tetramethyl-2,3,4,5,7,8,9,10-octahydropyrrolo[4,3,2- <i>m,n</i>]acridine-10-one 6.68
	$C_{25}H_{30}ClN_3$
	Crystal Violet 11.44

C ₂₅ H ₃₀ O ₄	Bixin	2.177	C ₂₆ H ₃₂ N ₂ NiO ₂	Bis[2-
C ₂₅ H ₃₂ N ₄ O	4-(4'-Diethylamino-2',6'-dimethylphenyl)imino-3-(2-methyl-2-propyl)-1-phenyl-2-pyrazolin-5-one	11.139		[(cyclohexylimino)methyl]phenolato]nickel(II) 12.24
C ₂₅ H ₃₄ O ₃	4-Dodecyloxy-2-hydroxybenzophenone	4.42	C ₂₆ H ₃₂ NiO ₆	Bis[3,5-di(1-methylethyl)salicylato]nickel(II) 12.60
C ₂₆ H ₁₄ O ₂	Anthra[1,9- <i>bc</i> :4,10- <i>b'c'</i>]dichromene	3.20	C ₂₆ H ₃₂ O ₄	<i>trans</i> -Methylbixin 2.175
	Benzo[1,2,3- <i>kl</i> :4,5,6- <i>k'l'</i>]dixanthene	3.56		Methylbixin 2.176
C ₂₆ H ₁₆	9,9'-Bifluorenylidene	3.66	C ₂₆ H ₃₃ N ₃ NaO ₆ S ₂	Merocyanine 540 11.39
C ₂₆ H ₁₇ Cl	1-Chloro-9,10-diphenylanthracene	3.6	C ₂₆ H ₃₄ O ₂	Phenyl linolenate 2.270
C ₂₆ H ₁₈	9,10-Diphenylanthracene	3.11	C ₂₆ H ₃₆ CoN ₂ S ₄	Bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenethionato)cobalt(II) 12.20
C ₂₆ H ₁₉ NO	1,1,3-Triphenylisoindole <i>N</i> -oxyl	15.25	C ₂₆ H ₃₆ CuN ₂ O ₂ S ₂	Bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenonato)copper(II) 12.22
C ₂₆ H ₂₀ CoN ₂ O ₂	Bis[2-[(phenylimino)methyl]phenolato]cobalt(II)	12.108	C ₂₆ H ₃₆ N ₂ NiS ₄	Bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenethionato]nickel(II) 12.21
C ₂₆ H ₂₀ CuN ₂ O ₂	Bis[2-[(phenylimino)methyl]phenolato]copper(II)	12.109	C ₂₆ H ₃₆ N ₂ S ₄	<i>N,N'</i> -[Dithiobis[(5-ethyl-2,3-thiophenediyi)-methylidyne]]biscyclohexanamine 13.63
C ₂₆ H ₂₀ N ₂ NiO ₂	Bis[2-[(phenylimino)methyl]phenolato]nickel(II)	12.110	C ₂₆ H ₃₆ N ₂ S ₄ Zn	Bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenethionato]zinc(II) 12.23
C ₂₆ H ₂₀ N ₂ NiS ₄	Bis(diphenyldithiocarbamato)nickel(II)	12.64	C ₂₆ H ₃₆ O ₂	Phenyl arachidonate 2.214
C ₂₆ H ₂₂ N ₄ NiO ₂	Bis[2-hydroxybenzaldehyde phenylhydrazone]nickel(II)	12.90	C ₂₆ H ₃₆ O ₆	8,13-Dimethyl-2,2,19,19-tetramethoxy-4,6,8,10,12,14,16-eicosahexa-3,18-dione (<i>all-E</i>) 2.211
C ₂₆ H ₂₆ ClIN ₂ S	3-Methyl-2-[7-(5-chloro-1,3,3-trimethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)-benzothiazolium iodide	11.34	C ₂₆ H ₃₈ N ₂ O ₆	Bis(2,2,6,6-tetramethyl-4-piperidinol-1-oxyl)terephthalate 15.39
C ₂₆ H ₂₆ FIN ₂ S	3-Methyl-2-[7-(5-fluoro-1,3,3-trimethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)-benzothiazolium iodide	11.35	C ₂₆ H ₄₄ O ₂	Tocol 4.48
C ₂₆ H ₂₇ IN ₂ S	3-Methyl-2-[7-(1,3,3-trimethyl-2-indolylidene)-1,3,5-heptatrienyl]benzothiazolium, iodide	11.37	C ₂₇ H ₂₁ N	1-Methylamino-9,10-diphenylanthracene 9.20
C ₂₆ H ₂₇ N ₃ O ₂	4-[<i>N</i> -Benzoyl(phenylaminocarbonyl)methylene]-amino- <i>N,N</i> -diethyl-3-methylaniline	11.27	C ₂₇ H ₂₃ ClN ₂ S ₂	3,3'-Diethyl-4,5,4',5'-dibenzo-2,2'-thiacyanine chloride 11.105
C ₂₆ H ₃₀ CoN ₄ O ₅ P	Hydroxybis(dimethylglyoximato)triphenylphosphinecobalt(II)	12.148	C ₂₇ H ₂₇ N ₇	Pyrazole dye 7 11.23
C ₂₆ H ₃₂ CoO ₆	Bis[3,5-di(1-methylethyl)salicylato]cobalt(II)	12.59	C ₂₇ H ₂₈ N ₂ O ₂	4-(Dibenzoylmethylene)amino-3, <i>N,N</i> -triethylaniline 11.116
			C ₂₇ H ₂₈ N ₂ O ₉	5-Iminodaunomycin 4.81
			C ₂₇ H ₂₉ IN ₂ OS	3-Methyl-2-[7-(5-methoxy-1,3,3-trimethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)-benzothiazolium iodide 11.36
			C ₂₇ H ₂₉ IN ₂ S	3-Methyl-2-[7-(1,3,3,5-tetramethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)benzothiazolium, iodide 11.38

$C_{27}H_{29}NO_{10}$	Daunomycin	4.79	$C_{28}H_{28}CuO_4P_2S_4$	
$C_{27}H_{29}NO_{11}$	Adriamycin	4.80		Bis[O,O' -di(4-methylphenyl)phosphorodithiolato]copper(II)
$C_{27}H_{29}N_3O_2$	4-[<i>N</i> -Benzoyl(phenylaminocarbonyl)-methylene]amino- <i>N,N</i> -diethyl-3,5-dimethylaniline	11.26		12.62
$C_{27}H_{30}N_4$	5-[<i>p</i> -(Dimethylamino)phenyl]-3-[<i>p</i> -(dimethylamino)styryl]-1-phenyl-2-pyrazoline	11.118	$C_{28}H_{28}N_2O_3S_3$	3,3'-Diethyl-2,2'-thiacarbocyanine toluenesulfonate 11.166
$C_{27}H_{30}N_4O_2$	4-[<i>N</i> -Di(phenylaminocarbonyl)methylene]-amino-3, <i>N,N</i> -triethylaniline	11.115	$C_{28}H_{28}NiO_4P_2S_4$	Bis[O,O' -di(4-methylphenyl)phosphorodithiolato]nickel(II) 12.63
$C_{27}H_{30}O_{16}$	Rutin	4.58	$C_{28}H_{29}N_2O_2S_2^-$	Squarylium Dye 3a 11.32
$C_{27}H_{44}$	3,5-Cholestadiene	2.89	$C_{28}H_{29}N_5O_2$	4-(4'-Diethylamino-2',6'-dimethylphenyl)imino-3-(benzoylamino)-1-phenyl-2-pyrazolin-5-one 11.138
$C_{27}H_{44}O$	Vitamin D ₃	2.361	$C_{28}H_{32}O_4$	6,10,15,19-Tetramethyl-4,6,8,10,12,14,16,18,20-tetracosanonaene-2,3,22,23-tetraone (<i>all-E</i>) 2.348
$C_{27}H_{46}O$	Cholesterol	16.14	$C_{28}H_{42}NiO_3S$	Aqua[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) 12.5
$C_{27}H_{46}O_2$	δ -Tocopherol	4.46	$C_{28}H_{42}O_2S$	2,2'-Thiobis[4-(1,1,3,3-tetramethylbutyl)phenol] 4.145
$C_{28}H_{12}Cl_8NiS_4$			$C_{28}H_{43}N$	4,4'-Di- <i>tert</i> -octyl diphenylamine 9.19
	Bis[1,2-di(2,4-dichlorophenyl)-1,2-ethenedithiolato]nickel(II)	12.37	$C_{28}H_{43}NNiO_2S$	Ammine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) 12.2
	Bis[1,2-di(3,4-dichlorophenyl)-1,2-ethenedithiolato]nickel(II)	12.38		Dodecylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II) 12.134
$C_{28}H_{14}O_2$	Heterocoerdianthrone	3.68	$C_{28}H_{44}O$	Ergosterol 16.26
$C_{28}H_{16}$	Dibenzo[<i>a,j</i>]perylene	3.63	$C_{28}H_{46}O$	4-Methyl-4-cholest-3-one 2.90
	Helianthrene	3.67	$C_{28}H_{48}O_2$	5,7-Dimethyltocol 4.51
$C_{28}H_{16}Br_4NiS_4$				β -Tocopherol 4.52
	Bis[1,2-di(2-bromophenyl)-1,2-ethenedithiolato]nickel(II)	12.25		γ -Tocopherol 4.53
$C_{28}H_{16}Cl_4NiS_4$			$C_{28}H_{50}O_3$	β -Tocopherol hydroquinone 4.26
	Bis[1,2-di(2-chlorophenyl)-1,2-ethenedithiolato]nickel(II)	12.33		γ -Tocopherol hydroquinone 4.27
	Bis[1,2-di(4-chlorophenyl)-1,2-ethenedithiolato]nickel(II)	12.34	$C_{28}H_{52}N_2O_4$	Tinuvin 770 8.78
$C_{28}H_{20}N_2O$	4-(2',3'-Benzo-4'-oxocyclohexadienyliden)amino- <i>N,N</i> -diphenylaniline	11.82	$C_{28}H_{52}N_2O_6$	Bis(2,2,6,6-tetramethyl-4-piperidinol-1-oxyl)1,10-decanedioate 15.37
$C_{28}H_{20}NiS_4$	Bis[1,2-diphenyl-1,2-ethenedithiolato]nickel(II)	12.66	$C_{29}H_{20}O$	Tetraphenylcyclopentadienone 3.60
$C_{28}H_{20}O$	Tetraphenylfuran	5.52	$C_{29}H_{26}O_{10}$	Cercosporin 16.55
$C_{28}H_{20}O_2$	Tetraphenyl- <i>p</i> -dioxin	2.196	$C_{29}H_{27}NO_9$	Aminocercosporin 16.54
$C_{28}H_{22}N_4O$	4-(4'-Diphenylaminophenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one	11.152	$C_{29}H_{30}N_2O_8$	Diaminocercosporin 16.56
$C_{28}H_{22}NiO_4$	Bis[2-hydroxy-5-methylbenzophenonato]nickel(II)	12.94	$C_{29}H_{31}Cl_2IN_2$	5,5'-Dichloro-1,1',3,3,3',3'-hexamethyltricarbocyanine, iodide 11.178
$C_{28}H_{22}O_2$	1,4-Dimethoxy-9,10-diphenylanthracene	3.9		
$C_{28}H_{28}CoO_4P_2S_4$				
	Bis[O,O' -di(4-methylphenyl)phosphorodithiolato]cobalt(II)	12.61		

$C_{29}H_{31}F_2IN_2$	5,5'-Difluoro-1,1',3,3,3',3'-hexamethyltricarbocyanine, iodide 11.179	$C_{29}H_{43}Se_2^+$	2,6-Bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)selenopyran-4-ylidene]-3-propenyl]selenopyrylium 11.160
$C_{29}H_{32}BBtF_4N_2$	2-[7-(4-Bromo-1,3,3-trimethyl-2-indol-2-ylidene)-1,3,5-heptatrienyl]-1,3,3-trimethylindolium, tetrafluoroborate 11.70	$C_{29}H_{43}Te_2^+$	2,6-Bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)telluropyran-4-ylidene]-3-propenyl]telluropyrylium 11.161
$C_{29}H_{32}BClF_4N_2$	2-[7-(4-Chloro-1,3,3-trimethyl-2-indol-2-ylidene)-1,3,5-heptatrienyl]-1,3,3-trimethylindolium, tetrafluoroborate 11.71	$C_{29}H_{50}O_2$	α -Tocopherol 4.50
$C_{29}H_{32}BF_4IN_2$	2-[7-(4-Iodo-1,3,3-trimethyl-2-indol-2-ylidene)-1,3,5-heptatrienyl]-1,3,3-trimethylindolium, tetrafluoroborate 11.73	$C_{29}H_{52}O_3$	α -Tocopherol hydroquinone 4.28
$C_{29}H_{32}I_2N_2$	2-[7-(4-Iodo-1,3,3-trimethyl-2-indol-2-ylidene)-1,3,5-heptatrienyl]-1,3,3-trimethylindolium, iodide 11.72	$C_{30}H_{16}O_8$	Hypericin 16.60
$C_{29}H_{32}O_8$	1,1'-Spirobis[2,3-dihydro-5,6-di(methoxycarbonyl)-3,3-dimethylindene] 3.98	$C_{30}H_{18}O_2$	Dimethylhomocoerdianthrone 3.95
$C_{29}H_{33}ClN_2O_4$	1,1',3,3,3',3'-Hexamethyltricarbocyanine, perchlorate 11.183	$C_{30}H_{22}O_9$	Rubellin A 16.8
$C_{29}H_{33}FN_2$	1,1',3,3,3',3'-Hexamethyltricarbocyanine, fluoride 11.181	$C_{30}H_{24}Cl_2N_6Ru$	Tris(2,2'-bipyridine)ruthenium(II) dichloride 12.177
$C_{29}H_{33}IN_2$	1,1',3,3,3',3'-Hexamethyltricarbocyanine, iodide 11.182	$C_{30}H_{24}CoN_6^{2+}$	Tris(2,2'-bipyridine)cobalt(II) ion 12.176
$C_{29}H_{40}O_4$	2,2'-Spirobis[7- <i>tert</i> -butyl-3,4-dihydro-6-hydroxy-4,4-dimethylbenzopyran] 4.184	$C_{30}H_{24}O_{10}$	Elsinochrome A 16.10
$C_{29}H_{42}O_3$	2',4'-Di- <i>tert</i> -butylphenyl 3,5-di- <i>tert</i> -butyl-4-hydroxybenzoate 4.38	$C_{30}H_{26}O_{10}$	Elsinochrome B 16.9
$C_{29}H_{43}OSe^+$	2,6-Bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)selenopyran-4-ylidene]-3-propenyl]pyrylium 11.157	$C_{30}H_{28}N_4NiO_4$	Nickel(II) deuteroporphyrin 7.95
$C_{29}H_{43}OTe^+$	2,6-Bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)telluropyran-4-ylidene]-3-propenyl]pyrylium 11.158	$C_{30}H_{28}N_6O_2$	4,4'-[2,3,5,6-Tetramethylphenylenebis]-3-methyl-1-phenyl-2-pyrazolin-5-one 11.153
$C_{29}H_{43}O_2^+$	4,4'-(1,3-Propenyl)bis[2,6-di(1,1-dimethylethyl)pyrylium 11.159	$C_{30}H_{28}O_{10}$	Methylcercosporin 16.58
$C_{29}H_{43}SSe^+$	2,6-Bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)selenopyran-4-ylidene]-3-propenyl]thiopyrylium 11.175	$C_{30}H_{30}FN_3O_5S$	4-[2',3'-Benzo-4'-oxo-5'-(2-methoxy-5-fluorosulfonyl)phenyl]amino- <i>N,N</i> -diethylaniline 11.81
$C_{29}H_{43}S_2^+$	2,6-Bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)thiopyran-4-ylidene]-3-propenyl]thiopyrylium 11.176	$C_{30}H_{30}NO$	Bis[4-(1-methyl-1-phenylethyl)phenyl]nitroxide 15.27
		$C_{30}H_{30}N_2O_3S_3$	3,3'-Diethyl-8,10-ethanediyl-2,2'-thiacarbocyanine toluenesulfonate 11.169
		$C_{30}H_{30}O_{10}$	Phleichrome 16.53
		$C_{30}H_{32}N_2O_3S_3$	3,3'-Diethyl-8,10-dimethyl-2,2'-thiacarbocyanine toluenesulfonate 11.168
		$C_{30}H_{32}N_2O_5S_3$	3,3'-Diethyl-5,5'-dimethoxy-2,2'-thiacarbocyanine toluenesulfonate 11.167
		$C_{30}H_{40}O$	β - <i>apo</i> -8'-Carotenal 2.62
		$C_{30}H_{42}O$	β - <i>apo</i> -8'-Carotenol 2.86
		$C_{30}H_{42}O_2$	1,2-Ethandiylidenebis[3,5-di(1,1-dimethylethyl)-2,5-cyclohexadien-4-one] 17.12
		$C_{30}H_{44}$	C_{30} -Carotene analog 2.77

$C_{30}H_{44}O_4$	C ₃₀ -Capsorubin 2.205 C ₃₀ -Epiisocapsorubin 2.206 C ₃₀ -Isocapsorubin 2.207	C ₃₂ H ₃₂ N ₂ O ₃ S ₃ 1,1'-Diethyl-4,4'-carbocyanine toluenesulfonate 11.43
$C_{30}H_{46}O_2$	Ergosterol acetate 16.27	C ₃₂ H ₃₂ N ₄ O ₄ 7-Ethenyl-3,8,13,17-tetramethylporphine-2,18-dipropanoic acid 7.93
$C_{30}H_{47}NNiO_2S$	Ethylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) 12.142	C ₃₂ H ₃₄ N ₄ O ₄ Deuteroporphyrin, dimethyl ester 7.94
$C_{30}H_{50}$	Squalene 2.346	C ₃₂ H ₃₆ N ₂ O ₃ S ₃ 3,9,3'-Triethyl-5,5'-dimethoxy-2,2'-thiacarbocyanine toluenesulfonate 11.172
$C_{30}H_{52}O_2$	α -Tocopherol methyl ether 5.4 5,7-Diethyltocol 4.45	C ₃₂ H ₄₄ O ₂ Ethyl β -apo-8'-carotenoate 2.85
$C_{30}H_{54}O_2$	2,5-Di-sec-dodecylhydroquinone 4.17	C ₃₂ H ₄₄ O ₆ 6,10,15,19-Tetramethyl-2,2,23,23-tetramethoxytetracosa-4,6,8,10,12,14,16,18,20-nonaene-3,22-dione (<i>all-E</i>) 2.347
$C_{30}H_{56}N_2O_4$	Di(1,2,2,6,6-pentaamethyl-4-piperidinol)-1,10-decanedioate 8.74	C ₃₂ H ₄₄ O ₁₄ emi-Crocin 2.178
$C_{31}H_{29}BrN_2S_2$	3,9,3'-Triethyl-4,5,4',5'-dibenzo-2,2'-thiacarbocyanine bromide 11.106	C ₃₂ H ₅₁ CoNO ₂ S Butylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]cobalt(II) 12.121
$C_{31}H_{30}O_{10}$	Dimethylcercosporin 16.57	C ₃₂ H ₅₁ NNiO ₂ S Butylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) 12.122
$C_{31}H_{32}IN_3$	1,3,3,1',3',3'-Hexamethyl-8,10-indolo-2,2'-carbocyanine iodide 11.40	C ₃₂ H ₅₆ O ₂ 5,7-Di(1-methylethyl)tocol 4.47
$C_{31}H_{37}IN_2$	1,1',3,3,3',3',5,5'-Octamethyltricarbocyanine, iodide 11.184	C ₃₃ H ₃₀ CoN ₅ O ₅ Hydroxybis(diphenylglyoximato)pyridine-cobalt(II) 12.149
$C_{31}H_{37}IN_2O_2$	5,5'-Dimethoxy-1,1',3,3,3',3'-hexamethyltricarbocyanine, iodide 11.180	C ₃₃ H ₃₄ N ₄ O ₆ Biliverdin 7.7
$C_{31}H_{38}N_4$	Pyrazole, 4,5-dihydro-5-[4-(diethylamino)phenyl]-3-[2-[4-(diethylamino)phenyl]ethenyl]-1-phenyl 11.117	C ₃₃ H ₃₆ N ₄ O ₆ Bilirubin 7.5
$C_{31}H_{38}N_4O_2$	Etiobiliverdin-IV γ 7.3	C ₃₃ H ₄₀ N ₂ O ₉ Reserpine 8.98
$C_{31}H_{40}N_4O_2$	Etiobilirubin-IV γ 7.4	C ₃₃ H ₄₀ N ₄ O ₆ Mesobilirubin IX 7.9
$C_{31}H_{49}NNiO_2S$	Propylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) 12.168	C ₃₃ H ₄₁ BF ₄ N ₂ O 2-[7-(1,3,3-Trimethyl-2-indol-2-ylidene)-1-[4-(2,2-dimethoxyethyl)-1,3,5-heptatrienyl]-1,1,3-trimethylindolium, tetrafluoroborate 11.74
$C_{31}H_{50}O_2$	Stigmasteryl acetate 16.74	C ₃₃ H ₄₂ O ₁₉ Troxerutin 4.57
$C_{31}H_{52}O_2$	Sitosteryl acetate 16.73	C ₃₃ H ₆₀ N ₄ O ₆ Tri(2,2,6,6-tetramethyl-4-piperidinol) nitrilotriacetate 8.80
$C_{31}H_{52}O_3$	α -Tocopheryl acetate 16.11	C ₃₄ H ₂₂ O ₂ 1,2,3,10,11,12-Hexahydrodindeno[5,6-a:5',6'-j]perylene-8,17-dione 3.65
$C_{31}H_{54}O_2$	α -Tocopherol ethyl ether 5.5 7- <i>tert</i> -Butyl-5-methyltocol 4.55 8- <i>tert</i> -Butyl-5-methyltocol 4.56	C ₃₄ H ₃₂ CdN ₄ O ₄ Cadmium(II) protoporphyrin 7.75
$C_{32}H_{12}AlClN_8O_9S_3$	Chloroaluminum(III) sulfophthalocyanine 7.43	C ₃₄ H ₃₂ CoN ₄ O ₄ Cobalt(II) protoporphyrin 7.76
$C_{32}H_{16}F_{12}NiS_4$	Bis[1,2-di(4-trifluoromethylphenyl)-1,2-ethenedithiolato]nickel(II) 12.76	C ₃₄ H ₃₂ CuN ₄ O ₄ Copper(II) protoporphyrin 7.77
$C_{32}H_{22}O$	1,3,4,7-Tetraphenylisobenzofuran 5.85	C ₃₄ H ₃₂ N ₄ NiO ₄ Nickel(II) protoporphyrin 7.80
$C_{32}H_{28}NiO_4S_4$	Bis[1,2-di(4-methoxyphenyl)-1,2-ethenedithiolato]nickel(II) 12.54	C ₃₄ H ₃₂ N ₄ O ₄ Zn Zinc(II) protoporphyrin IX 7.81

$C_{34}H_{34}N_4O_4$	Protoporphyrin IX	7.73	$C_{36}H_{24}CoN_6^{2+}$	
$C_{34}H_{34}NiO_6$	Bis[4-(1,1-dimethylethyl)phenylsalicylato]nickel(II)	12.58	Tris(1,10-phenanthroline)cobalt(II) ion	
$C_{34}H_{36}CoN_4O_4$	Cobalt(II) mesoporphyrin	7.82	12.178	
$C_{34}H_{36}N_4NiO_4$	Nickel(II) mesoporphyrin	7.87	$C_{36}H_{24}N_{10}Ni$	
$C_{34}H_{36}N_4O_4$	8-Ethenyl-13-ethyl-3,7,12,17-tetramethylporphine-2,18-dipropanoic acid	7.90	Bis[2,3-dihydro-N-2-pyridinyl-3-(2-pyridinylimino)isoindol-1-amine]nickel(II)	12.45
$C_{34}H_{36}N_4O_5$	Monohydroxyethyl vinyl deuteroporphyrin	7.92	$C_{36}H_{26}N_5O_7S^-$	
$C_{34}H_{38}N_4O_6$	Hematoporphyrin IX	7.72	2-Methyl-6-[4-[2-[3-carboxy-4-(6-hydroxy-3-xanthenon-9-yl)-phenylthiocarbamylethoxy]phenyl]imidazo[1,2-a]pyrazin-3-one	16.39
$C_{34}H_{42}B_2F_8N_3$	2-[7-(1,3,3-Trimethyl-2-indol-2-ylidene)-1-[4-(1-piperidinio)-1,3,5-heptatrienyl]-1,3,3-trimethylindolium, bis(tetrafluoroborate)	11.75	$C_{36}H_{26}O_2$	1,2,3,4,11,12,13,14-Octahydrodinaphtho[2,3-a:2',3'-j]perylene-9,19-dione
$C_{34}H_{47}NNiO_2S$	Aniline[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II)	12.3		3.62
$C_{34}H_{48}O_4$	C_{34} -Capsorubin	2.231	$C_{36}H_{32}N_2O_3S_3$	3,3'-Diethyl-4,5,4',5'-dibenzo-2,2'-thiacarbocyanine toluenesulfonate
	C_{34} -Epiisocapsorubin	2.232	11.107	
$C_{34}H_{50}O_2$	Cholestryl benzoate	16.15	$C_{36}H_{36}CoS_4$	Bis[1,2-di(4-ethylphenyl)-1,2-ethenedithiolato]cobalt(II)
$C_{34}H_{53}NNiO_2S$	Cyclohexylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II)	12.131	12.41	
$C_{34}H_{55}NNiO_5S$	Triethanolamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II)	12.171	$C_{36}H_{36}N_4NiO_4$	Nickel(II) protoporphyrin, dimethyl ester
$C_{34}H_{56}NiO_8P_2$	Bis[O-ethyl-3,5-di-(1,1-dimethylethyl)-4-hydroxybenzylphosphonato]nickel(II)	12.83	7.79	
$C_{35}H_{25}Te_2^+$	2,6-Diphenyl-4-(2,6-diphenyltelluropyran-4-ylidene)methyltelluropyrylium	11.162	$C_{36}H_{36}NiS_4$	Bis[1,2-di(4-ethylphenyl)-1,2-ethenedithiolato]nickel(II)
$C_{35}H_{32}MgN_4O_5$	Protochlorophyllide	7.37	12.42	
$C_{35}H_{38}N_4O_6$	Biliverdin, dimethyl ester	7.8	$C_{36}H_{36}PtS_4$	Bis[1,2-di(4-ethylphenyl)-1,2-ethenedithiolato]platinum(II)
$C_{35}H_{40}N_4O_6$	Bilirubin IX dimethyl ester	7.6	12.43	
$C_{35}H_{50}$	C_{35} -Carotene analog	2.78	$C_{36}H_{38}N_4O_4$	Protoporphyrin IX, dimethyl ester
$C_{35}H_{62}O_3$	Octadecyl 3-(3',5'-di- <i>tert</i> -butyl-4'-hydroxyphenyl)propionate	4.33	7.78	
$(C_{35}H_{66}N_8)_n$	Chimasorb 944	8.16	$C_{36}H_{40}CoN_4O_4$	Cobalt(II) mesoporphyrin IX, dimethyl ester
$C_{36}H_{16}CuN_8O_8$	Copper(II) tetracarboxyphthalocyanine	7.45	7.86	
			$C_{36}H_{40}N_4NiS_4$	Bis[1,2-di(dimethylaminophenyl)-1,2-ethenedithiolato]nickel(II)
			12.39	
			$C_{36}H_{40}N_4O_4$	8-Ethenyl-13-ethyl-3,7,12,17-tetramethylporphine-2,18-dipropanoic acid, dimethyl ester
			7.91	
			$C_{36}H_{42}N_4O_4$	Mesoporphyrin IX, dimethyl ester
			7.85	
			$C_{36}H_{54}N_2NiO_2$	2,2'-[1,2-Ethanediylbis(nitrilodecyldyne)]bis[4-methylphenolato]nickel(II)
			12.136	
			$C_{36}H_{66}N_4O_6$	Tri(1,2,2,6,6-pentamethyl-4-piperidinol) nitrolotriacetate
			8.75	
			$C_{37}H_{30}BrIrOP_2$	Bromocarbonylbis(triphenylphosphine)iridium
			12.119	
			$C_{37}H_{30}ClIrOP_2$	Carbonylchlorobis(triphenylphosphine)iridium
			12.123	

C ₃₇ H ₃₀ ClOP ₂ Rh		C ₃₉ H ₆₀ O ₄	Ubiquinol 10	4.25
Carbonylchlorobis(triphenylphosphine)-rhodium	12.124	C ₄₀ H ₂₄	Mesodiphenylhelianthrene	3.64
C ₃₇ H ₃₀ IIrOP ₂	Carbonyliodobis(triphenylphosphine)iridium	C ₄₀ H ₄₄ NiO ₁₂ S ₄	Bis[1,2-di(3,4,5-trimethoxyphenyl)-1,2-ethenedithiolato]nickel(II)	12.79
12.125		C ₄₀ H ₄₈ CdO ₄ P ₂ S ₆	Bis[2,2'-thiobis[O,O'-di(4- <i>tert</i> -butylphenyl)phosphorodithiolato]cadmium(II)	12.111
C ₃₇ H ₄₂ N ₄ O ₂	Zinc(II) etiopurpurin ethyl ester	C ₄₀ H ₄₈ CoO ₄ P ₂ S ₆	Bis[2,2'-thiobis[O,O'-di(4- <i>tert</i> -butylphenyl)phosphorodithiolato]cobalt(II)	12.112
C ₃₇ H ₄₆ Si ₂	1,1,2,2-Tetrakis(2,4,6-trimethylphenyl)disilirane	C ₄₀ H ₄₈ NiO ₄ P ₂ S ₆	Bis[2,2'-thiobis[O,O'-di(4- <i>tert</i> -butylphenyl)phosphorodithiolato]nickel(II)	12.114
C ₃₇ H ₄₈ N ₆ O ₁₂ S ₂	Bilirubin ditaurate	C ₄₀ H ₄₈ O ₄ P ₂ PbS ₆	Bis[2,2'-thiobis[O,O'-di(4- <i>tert</i> -butylphenyl)phosphorodithiolato]lead(II)	12.113
C ₃₈ H ₃₀ N ₄ NiO ₂	Bis[2-[(4-(phenylamino)phenyl)imino]methyl]-phenolato]nickel(II)	C ₄₀ H ₄₈ O ₄ P ₂ S ₆ Zn	Bis[2,2'-thiobis[O,O'-di(4- <i>tert</i> -butylphenyl)phosphorodithiolato]zinc(II)	12.115
C ₃₈ H ₄₃ FeN ₄ O ₆	Iron(III) (acetato)mesoporphyrin IX, dimethyl ester	C ₄₀ H ₅₀ N ₄	Octaethyl[22]porphyrin-(2.2.2.2) (di- <i>trans</i>)	7.98
C ₃₈ H ₄₄ MnN ₄ O ₄ ⁺	Manganese(III) mesoporphyrin diethyl ester	C ₄₀ H ₅₂ CoO ₄ P ₂ S ₄	Bis[O,O'-di-(4- <i>tert</i> -butylphenyl)phosphorodithiolato]cobalt(II)	12.30
7.84		C ₄₀ H ₅₂ CuO ₄ P ₂ S ₄	Bis[O,O'-di-(4- <i>tert</i> -butylphenyl)phosphorodithiolato]copper(II)	12.31
C ₃₈ H ₄₅ N ₅ O ₅	(Z)-Dimethyl 7-[2-(dimethylamino)-2-oxoethyl]-8-ethylidene-7,8-dihydro-3,7,12,17-tetramethylporphine-2,18-dipropionate	C ₄₀ H ₅₂ NiO ₄ P ₂ S ₄	Bis[O,O'-di-(4- <i>tert</i> -butylphenyl)phosphorodithiolato]nickel(II)	12.32
7.89	7.89	C ₄₀ H ₅₂ OSi ₂	2,2,3,3-Tetrakis(2,6-diethylphenyl)oxadisilirane	17.46
C ₃₈ H ₄₇ N ₅ O ₅	(Z)-Dimethyl 7-[2-(dimethylamino)-2-oxoethyl]-8-ethyl-7,8-dihydro-3,7,12,17-tetramethylporphine-2,18-dipropionate	C ₄₀ H ₅₂ O ₂	β,β -Carotene-4,4'-dione	2.83
7.88		C ₄₀ H ₅₂ O ₄	Astaxanthin	2.82
C ₃₈ H ₆₀ CoN ₂ O ₂	Bis[2-[(dodecylimino)methyl]phenolato]cobalt(II)	C ₄₀ H ₅₄ O	β,β -Caroten-4-one	2.88
12.80		C ₄₀ H ₅₆	α -Carotene	2.63
C ₃₈ H ₆₀ CuN ₂ O ₂	Bis[2-[(dodecylimino)methyl]phenolato]copper(II)		9-cis- β -Carotene	2.64
12.81			15,15'-(Z)- β -Carotene	2.65
C ₃₈ H ₆₀ N ₂ NiO ₂	Bis[2-[(dodecylimino)methyl]phenolato]nickel(II)		β -Carotene	2.66
12.82			γ -Carotene	2.76
C ₃₈ H ₆₀ N ₂ NiO ₄	Bis[2'-hydroxy-4'-methyldodecanophenone oximate]nickel(II)		Lycopene, (<i>all-E</i>)	2.250
12.95		C ₄₀ H ₅₆ O	β,β -Caroten-3-ol	2.87
C ₃₈ H ₆₄ NiO ₈ P ₂	Bis[O-butyl-3,5-di-(1,1-dimethylethyl)-4-hydroxybenzylphosphonato]nickel(II)			
12.16				
C ₃₈ H ₇₀ O ₂	2,5-Di-sec-hexadecylhydroquinone			
4.18				
C ₃₉ H ₃₄ Cl ₃ N ₃ O ₄ S ₂	IR 140			
11.33				
C ₃₉ H ₃₉ MgN ₅ O ₇	Protochlorophyllide 2-amino-2-(methoxycarbonyl)ethyl ester			
	7.36			

$C_{40}H_{56}O_2$	Xanthophyll 2.75 β,β -Carotene-3,3'-diol, (3 <i>R</i> ,3' <i>R</i>)- 2.79 β,β -Carotene-4,4'-diol, <i>all-E</i> - 2.81 Dihydroxylycopene 2.251	$C_{43}H_{58}N_4O_4$ Methyl acetal of oxidized octaethylporphyrin ethyl ester 7.16
$C_{40}H_{56}O_4$	Violaxanthin 2.80 C_{40} -Epiisocapsorubin 2.84	$C_{44}H_{24}Cl_2N_4O_{12}S_4Sn^{4-}$ Dichloro[5,10,15,20-tetrakis(4-sulfonatophenyl)porphinatostannate(IV) ion 7.58
$C_{40}H_{58}$	Neurosporene 2.70	$C_{44}H_{24}CuI_4N_4$ 5,10,15,20-Tetrakis(4-iodophenyl)porphine, copper(II) 7.55
$C_{40}H_{60}$	ζ -Carotene 2.72	$C_{44}H_{26}$ Mesodiphenylbenzheianthrene 3.129
$C_{40}H_{62}$	15- <i>cis</i> -Phytofluene 2.73	$C_{44}H_{26}N_4O_{12}S_4^{4-}$ 5,10,15,20-Tetrakis(4-sulfonatophenyl)porphine 7.57
$C_{40}H_{64}$	15-(<i>Z</i>)-Phytoene 2.74	$C_{44}H_{28}AlClN_4$ 5,10,15,20-Tetraphenylporphinato(chloro)aluminum(III) ion 7.61
$C_{40}H_{67}NNiO_2S$	Didodecylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II) 12.132 Dodecylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) 12.135	$C_{44}H_{28}CdN_4$ 5,10,15,20-Tetraphenylporphinatocadmium(II) 7.60
$C_{40}H_{68}N_2O_5$	A12 8.66	$C_{44}H_{28}ClFcN_4$ Chloro(5,10,15,20-tetraphenylporphinato)iron(III) 7.49
$C_{41}H_{50}N_4O_2$	2,3,7,8,12,13,17,18-Octaethyl-5-porphinepropenoic acid ethyl ester 7.97	$C_{44}H_{28}Cl_2N_4Sn$ Dichloro(5,10,15,20-tetraphenylporphinato)tin(IV) 7.64
$C_{41}H_{52}Cl_2N_4O_2Sn$	Tin(IV) dichloro{octaethylidihydropurpurin ethyl ester} 7.29	$C_{44}H_{28}CoN_4$ 5,10,15,20-Tetraphenylporphinatocobalt(II) 7.62
$C_{41}H_{52}N_4O_2$	Octaethylporpurin ethyl ester 7.30	$C_{44}H_{28}CuN_4$ 5,10,15,20-Tetraphenylporphinatocupper(II) 7.63
$C_{41}H_{54}N_4O_2$	Octaethylidihydropurpurin ethyl ester 7.28	$C_{44}H_{28}MgN_4$ 5,10,15,20-Tetraphenylporphinatomagnesium(II) 7.65
$C_{41}H_{58}O_2$	Spheroidenone 2.69	$C_{44}H_{28}MnN_4^+$ Manganese(III) 5,10,15,20-tetraphenylporphyrin 7.66
$C_{41}H_{60}O$	Spheroidene 2.68	$C_{44}H_{28}N_4Zn$ 5,10,15,20-Tetraphenylporphinatozinc(II) 7.68
$C_{42}H_{28}$	Rubrene 3.125	$C_{44}H_{28}NiN_4$ Nickel(II) 5,10,15,20-tetraphenylporphyrin 7.67
$C_{42}H_{38}N_4Ni$	Bis[N-(4-methylphenyl)-7-[(4-methylphenyl)imino]-1,3,5-cycloheptatrien-1-aminato]nickel(II) 12.104	$C_{44}H_{30}CdN_4$ Cadmium(II) tetraphenylchlorin 7.51
$C_{42}H_{52}CoO_4P_2S_4$	2,2'-Methylenebis[<i>O,O'</i> -di(4- <i>tert</i> -butylphenyl)dithiophosphate]cobalt(II) 12.156	$C_{44}H_{30}CuN_4$ Copper(II) tetraphenylchlorin 7.52
$C_{42}H_{52}NiO_4P_2S_4$	2,2'-Methylenebis[<i>O,O'</i> -di(4- <i>tert</i> -butylphenyl)dithiophosphate]nickel(II) 12.158	$C_{44}H_{30}N_4$ 5,10,15,20-Tetraphenylporphine 7.59
$C_{42}H_{52}O_4P_2PbS_4$	2,2'-Methylenebis[<i>O,O'</i> -di(4- <i>tert</i> -butylphenyl)dithiophosphate]lead(II) 12.157	$C_{44}H_{30}N_4Zn$ Zinc(II) tetraphenylchlorin 7.53
$C_{42}H_{54}O_2^{2-}$	4,4'-(4,7-Dihydro-5,6-dimethylisobenzofuran-1,3-diyl)bis(benzene- <i>p</i> -decanoate ion) 5.81	$C_{44}H_{30}O$ Hexaphenylisobenzofuran 5.84
$C_{42}H_{58}O_6$	Fucoxanthin 2.67	$C_{44}H_{32}N_4$ Tetraphenylchlorin 7.50
$C_{42}H_{60}O_2$	Spirilloxanthin 2.71	$C_{44}H_{34}N_4$ <i>trans</i> -Tetraphenylbacteriochlorin 7.54
$C_{42}H_{72}N_2O_5$	Tinuvin 144 8.73	$C_{44}H_{46}N_4$ 2,3,10,11,16,17,24,25-Octaethyl-5,6,7,8,19,20,21,22-octadehydro[26]porphyrin-(2.4.2.4) 7.100
$C_{43}H_{47}N_2NaO_6S_2$		
	IR 125 11.29	

C ₄₄ H ₄₈ BF ₄ N ₃	2-[7-(1,3,3-Trimethyl-2-indol-2-ylidene)-1-[4-[3-(1,3,3-trimethyl-2-indolylidene)-2-propenyl]-1,3,5-heptatrienyl]-1,1,3-trimethylindolium, tetrafluoroborate	11.76	C ₅₀ H ₅₀ N ₆ NiO ₄ Nickel(II) deuteroporphyrin bis(piperidine)	7.96
C ₄₄ H ₅₆ NNiS ₄	Bis(1,2-diphenyl-1,2-ethanedithionato)nickelate(I), tetrabutylammonium salt	12.65	C ₅₀ H ₆₆ N ₈ O ₂ Si ₃ Bis(tripropylsiloxy)silicon phthalocyanine	7.41
C ₄₄ H ₆₄ O ₂₄	Crocin	2.173	C ₅₀ H ₆₈ Decapreno-β-carotene	2.230
C ₄₅ H ₂₈ FeN ₅ S	5,10,15,20-Tetraphenylporphinato(thiocyanato-S)iron(III)	7.70	C ₅₀ H ₇₂ N ₈ O ₅ S Magenta dye 9	11.154
C ₄₅ H ₅₉ N ₄ .ClO ₄	4-[1,5,5-Tri(4-diethylaminophenyl)-pentadienylidene]anilinium perchlorate	11.110	C ₅₀ H ₇₂ O ₂ P-438 (9 conjugated bonds)	2.299
C ₄₆ H ₂₄ O ₈ ⁴⁻	Rubrene-2,3,8,9-tetracarboxylate ion	3.126	C ₅₀ H ₇₄ O ₂ P-422 (8 conjugated bonds)	2.298
C ₄₆ H ₃₁ FeN ₄ O ₂	Iron(III) (acetato)5,10,15,20-tetraphenylporphyrin	7.47	C ₅₁ H ₆₇ CoN ₆ O ₁₂ Hexamethyl Coα Coβ-dicyano-7-de(carboxymethyl)-7,8-didehydrocobyrinate	16.16
C ₄₆ H ₃₁ MnN ₄ O ₂	Acetato(5,10,15,20-tetraphenylporphinato)manganese(III)	7.48	C ₅₃ H ₄₈ ClN ₃ O ₈ S ₂ IR 132	11.108
C ₄₆ H ₃₈ CoN ₄ O ₅ P	Hydroxybis(diphenylglyoximato)-triphenylphosphinecobalt(II)	12.150	C ₅₄ H ₈₈ N ₂ O ₄ 2,4-Di[2-methyl-4-(N-methyl-N-octadecylamino)phenyl]squarylium	11.48
C ₄₆ H ₅₈ O ₄	1,2-Ethanediylidenebis[3-[2-hydroxy-5-methyl-3-(1,1-dimethylethyl)benzyl]-5-(1,1-dimethylethyl)]-2,5-cyclohexadien-4-one	17.11	C ₅₅ H ₇₀ MgN ₄ O ₅ Protochlorophyll	7.39
C ₄₇ H ₃₉ ClO ₆	2,3,2',3'-Bis(1,4,10,13-trideca-4,6,8,10-tetraen)tetrayl[4,6-diphenylpyrylium] perchlorate	11.31	C ₅₅ H ₇₀ MgN ₄ O ₆ Chlorophyll b	7.12
C ₄₈ H ₃₆ CuN ₄ O ₄	5,10,15,20-Tetrakis(4-methoxyphenyl)porphinatocuppper(II)	7.56	C ₅₅ H ₇₂ MgN ₄ O ₅ Chlorophyll a	7.11
C ₄₈ H ₄₄ NNiO ₄ S ₄	Bis[1,2-di(4-methoxyphenyl)-1,2-ethanedithionato]nickelate(I), tetrabutylammonium salt	12.53	C ₅₅ H ₇₂ MgN ₄ O ₆ Bacteriochlorophyll b	7.2
C ₄₈ H ₅₀ N ₈	Tetra- <i>tert</i> -butylphthalocyanine	7.46	C ₅₅ H ₇₂ N ₄ O ₅ Protopheophytin	7.38
C ₄₈ H ₅₂ F ₁₂ NNiS ₄	Bis[1,2-di(4-trifluoromethylphenyl)-1,2-ethanedithionato]nickelate(I), tetrabutylammonium salt	12.75	C ₅₅ H ₇₂ N ₄ O ₆ Pheophytin b	7.27
C ₄₈ H ₆₆ N ₄ ²⁺	[26] Porphyrin	7.99	C ₅₅ H ₇₄ MgN ₄ O ₆ Bacteriochlorophyll a	7.1
C ₄₈ H ₆₈ OSi ₂	2,2,3,3-Tetrakis(2,6-diisopropylphenyl)oxadisilirane	17.47	C ₅₅ H ₇₄ N ₄ O ₅ Pheophytin a	7.26
			C ₅₅ H ₇₄ N ₄ O ₆ Bacteriopheophytin b	7.33
			C ₅₅ H ₇₆ N ₄ O ₆ Bacteriopheophytin a	7.32
			C ₅₆ H ₇₃ N ₅ O ₈ S ₂ IR 144	11.30
			C ₅₆ H ₈₀ NNiO ₁₂ S ₄ Bis[1,2-di(3,4,5-trimethoxyphenyl)-1,2-ethanedithionato]nickelate(I), tetrabutylammonium salt	12.77
			C ₅₆ H ₈₄ NiO ₄ S ₂ Bis(2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)phenolato]nickel(II)	12.116
			C ₅₆ H ₉₆ NiO ₄ Pd Bis[2'-hydroxy-4'- <i>tert</i> -butyloctadecanophenone oximato]palladium(II)	12.91
			C ₅₇ H ₇₂ MgN ₄ O ₅ 4-Vinylprotochlorophyll	7.34

$C_{57}H_{74}N_4O_5$	4-Vinylprotopheophytin	7.35	Cl_2Ni	Nickel(II) dichloride	12.162
C_{60}	Fullerene-C ₆₀	17.32	$Cl_2Ni.6H_2O$	Nickel(II) dichloride hexahydrate	12.163
$C_{60}H_{80}$	Dodecapreno- β -carotene	2.358	Co^{2+}	Cobalt(II) ion	12.128
$C_{64}H_{58}N_8$	2,11,20,29-Tetra- <i>tert</i> -butylnaphthalocyanine 7.22		$CoH_{18}N_6^{3+}$	Hexaamminecobalt(III) ion	12.145
$C_{64}H_{80}N_8O_8$	1,4,8,11,15,18,22,25- Octabutoxyphthalocyanine	7.42	Cu^{2+}	Copper(II) ion	12.129
$C_{66}H_{62}AlN_8OSi$	Bis(<i>tri-n</i> -hexyloxysiloxy)-2,3- naphthalocyaninatoaluminum	7.24	DO^-	Hydroxide- <i>d</i> ion	14.8
$C_{66}H_{62}GaN_8OSi$	Bis(<i>tri-n</i> -hexyloxysiloxy)-2,3- naphthalocyaninatogallium	7.25	D_2O_2	Deuterium peroxide	14.6
C_{70}	Fullerene-C ₇₀	17.33	HN_3	Hydrogen azide	14.3
$C_{72}H_{41}AlN_8O$	[1,10,19,28-Tetraphenylnaphthalocyanine]- hydroxyaluminum(III)	7.23	H_4N_2	Hydrazine hydrate	14.7
$C_{72}H_{86}N_8O_2Si_3$	Bis(triisobutylsiloxy)silicon 2,3- naphthalocyanine	7.20	Γ	Iodide ion	14.9
$C_{74}H_{66}N_8O_2Si_3$	Bis(tribenzylsiloxy)silicon phthalocyanine 7.40		I_2	Iodine	14.10
$C_{76}H_{70}N_8O_{10}$	Mesoporphyrin di[4- (diphenylmethylaminocarbonyl-2- nitrophenylmethyl] ester	7.83	NH_3	Ammonia	14.1
$C_{80}H_{89}N_8O_8Pd$	1,6,10,15,19,24,28,33- Octabutoxynaphthalocyaninatopalladium(II) 7.21		NO_2^-	Nitrite ion	14.11
$C_{84}H_{102}N_8O_2Si_2Sn$	Bis(<i>tri-n</i> -hexyloxysiloxy)-2,3- naphthalocyaninatotin	7.19	N_3^-	Azide ion	14.2
$C_{84}H_{102}N_8O_2Si_3$	Bis(<i>tri-n</i> -hexyloxysiloxy)-2,3- naphthalocyaninatosilicon	7.18	Ni^+	Nickel(I) ion	12.164
$C_{85}H_{103}N_4NiO_{12}S_4$	Bis{1,2-di(3,4,5-trimethoxyphenyl)-1,2- ethanedithionato}nickelate(I), 4-[1,5,5- tri(4-diethylaminophenyl)- pentadienylidene]anilinium salt	12.78	O_2	Oxygen	14.12
$C_{88}H_{56}Fe_2N_8O$	μ -Oxobis[5,10,15,20- tetraphenylporphinatoiron(II)]	7.69	$^{18}O_2$	Oxygen-18	14.13
$C_{90}H_{74}N_8O_2Si_3$	Bis(tribenzylsiloxy)silicon 2,3- naphthalocyanine	7.17	O_2^-	Superoxide radical anion	14.14
Cl^-	Chloride ion	14.5			
$Cl_2Co.6H_2O$	Cobalt(II) chloride hexahydrate	12.127			
$Cl_2Mn.6H_2O$	Manganese(II) chloride hexahydrate	12.151			
$Cl_2Mn.4H_2O$	Manganese(II) chloride tetrahydrate	12.152			

14. Chemical Name Index

Entry numbers for Tables 2-17 are printed in bold face following the names. Entry numbers are printed in smaller type, not bold, for entries in which the chemical species have been used as reference substrates and relative rates have been determined. For example, Acetone azine appears in Table 15 (entry 15.3) but rate constants for another substrate relative to k_r for acetone azine are found in entry 2.40. There is no entry in the tables for 9-Acetyl-2,3,4,9-tetrahydrocarbazole but relative data are found in entries 6.2, 6.3, 6.10, 6.14, 6.18 and 6.19.

- A12 **8.66**
- A 31 **8.64**
- Abietic acid **2.325**
- Acetaldehyde, (2,6-diphenylpyran-4-ylidene)-, dimethyl hydrazone **15.1**
- Acetamide, *N*-[4-[[4-(diethylamino)-2-methylphenyl]imino]-4,5-dihydro-5-oxo-1-(4-bromophenyl)pyrazol-3-yl]- **11.135**
- Acetamide, *N*-[4-[[4-(diethylamino)-2-methylphenyl]imino]-4,5-dihydro-5-oxo-1-(3-chlorophenyl)pyrazol-3-yl]- **11.136**
- Acetamide, *N*-[4-[[4-(diethylamino)-2-methylphenyl]imino]-4,5-dihydro-5-oxo-1-(4-chlorophenyl)pyrazol-3-yl]- **11.137**
- Acetamide, *N*-[4-[[4-(diethylamino)-2-methylphenyl]imino]-4,5-dihydro-5-oxo-1-(3-methoxyphenyl)pyrazol-3-yl]- **11.142**
- Acetamide, *N*-[4-[[4-(diethylamino)-2-methylphenyl]imino]-4,5-dihydro-5-oxo-1-(3-methylphenyl)pyrazol-3-yl]- **11.143**
- Acetamide, *N*-[4-[[4-(diethylamino)-2-methylphenyl]imino]-4,5-dihydro-5-oxo-1-(4-nitrophenyl)pyrazol-3-yl]- **11.145**
- Acetamide, *N*-[4-[[4-(diethylamino)-2-methylphenyl]imino]-4,5-dihydro-5-oxo-1-phenylpyrazol-3-yl]- **11.128**
- Acetamide, *N*-[4-[[4-(diethylamino)-2-methylphenyl]imino]-4,5-dihydro-5-oxo-1-(2,4,6-trichlorophenyl)pyrazol-3-yl]- **11.146**
- Acetatobis(acetylacetato)manganese(III) **12.1**
- Acetato(7,12-diethyl-3,8,13,17-tetramethylporphine-2,18-dipropanoato)ferrate(III), dimethyl ester **7.74**
- Acetatoiron(III) tetraphenylporphyrin **7.47**
- Acetato(5,10,15,20-tetraphenylporphinato)iron(III) **7.47**
- Acetato(5,10,15,20-tetraphenylporphinato)manganese(III) **7.48**
- Acetic acid **17.1**
- 2'-Acetonaphthone **3.1**
- Acetone **17.2**
- Acetone d_6 **17.3**
- Acetone azine **15.3, 2.40**
- Acetone oxime **15.2**

- Acetonitrile **17.4**
- Acetonitrile- d_3 **17.5**
- Acetophenone **3.2**
- Acetophenone, 4'-hydroxy- **4.93**
- N*-(2-Acetoxyethyl)-2,2,6,6-tetramethylpiperidine **8.69**
- 2-Acetoxy-3-methyl-2-cyclopenten-1-one **2.163**
- 3-Acetoxy-2-methyl-2-cyclopenten-1-one **2.164**
- Acetoxymethyleneadamantane **2.2**
- Acetoxymethyleneadamantane **2.111**
- 4-Acetoxymethyl-2-(phenylmethyl)furan **5.75**
- N*-Acetylcysteine, negative ion **13.70**
- N*-Acetyl-L-cysteine **13.69**
- 4-Acetyl-2,6-di-*tert*-butylphenol **4.94**
- 5-Acetyl-3,4-dihydro-6-methylpyran **5.88**
- Acetylene, 1,2-bis(diethylamino) **8.30**
- Acetylene, 1-(diethylamino)-2-methyl- **8.87**
- 2-Acetyl furan **5.15**
- N*-Acetyl-2,3,4,4*a*,5,9*b*-hexahydro- 2,8-dimethylpyrido[4,3-*b*]indole **6.57**
- 2-Acetyl naphthalene **3.1**
- 3-Acetoxy-1,3,5(10),9(11)-estratetraen-17-one **16.29**
- 4-Acetylphenol **4.93**
- 4-Acetylphenoxide ion **4.151**
- N*-Acetylstobadine **6.57**
- 9-Acetyl-2,3,4,9-tetrahydrocarbazole **6.2, 6.3, 6.10, 6.14, 6.18, 6.19**
- N*-Acetyltryptophan **10.37**
- N*-Acetyl-L-tryptophanamide **10.44**
- N*-Acetyltryptophan methyl ester **10.38**
- N*-Acetyl-L-tyrosine, ethyl ester **4.188**
- Acridine, 3,6-diamino- **11.1**
- 1,8-Acidinedione, 3,4,6,7,9,10-hexahydro-3,3,6,6-tetramethyl- **6.1**
- Acridinium, 3,6-diamino-10-methyl- **11.2**
- Acriflavine cation **11.2**
- Acrolein **2.1**
- Adamantane, 2-(hydroxymethylene)-, acetate **2.2**
- Adamantylideneadamantane **2.356, 13.112**
- 1-Adamantylideneethyl acetate **2.220**
- Adamantylidenemethyl acetate **2.256, 2.126**
- endo*-7-Adamantylidene norbornane-2,3-dicarboxylic acid anhydride **2.15**
- exo*-7-Adamantylidene norbornane-2,3-dicarboxylic acid anhydride **2.16**
- Adenine **16.63**
- Adenosine **16.1**
- Adenosine monophosphate **16.2**
- Adenosine 5'-monophosphate **16.2**
- Adenosine triphosphate **16.3**
- Adenosine triphosphate, ester with 1- β -D-ribofuranosyl-3-pyridinecarboxamide **16.50**
- 5'-Adenylic acid **16.2**

- Adrenaline **4.9**
 Adriamycin **4.80**
 Alanine **10.1**
 β -Alanine **10.2**
 β -Alanine, L-histidyl- **10.3**
 L-Alanine, glycyl-L-tyrosyl- **4.1**
 β -Alanyl-L-histidine **10.24**
 β -Alanyl-1-methylhistidine **10.25**
 L-Alanyl-L-tryptophan **10.39**
 Albumin **10.4**
 Alloocimine **2.285**
 Allylthiourea **13.175**
 N-Allylurea **17.78**
 Aluminum, tetranaphtho[2,3-*b*:2',3'-*g*:2'',3''-*l*:2'',3''-*q*]porphyrinatobis(trihexylsilanolato)- **7.24**
 Aluminum(III) chloro(5,10,15,20-tetraphenylporphyrin) **7.61**
 Aluminum(III) hydroxy[1,10,19,28-tetraphenylnaphthalocyanine] **7.23**
 1-Aminoanthraquinone **11.3**
 2-Aminoanthraquinone **11.4**
 1-(4-Aminocarbonylphenylazo)-3-(2-methoxyphenylaminocarbonyl)-2-naphthol **11.79**
 Aminocercosporin **16.54**
 5-Amino-2,3-dihydro-1,4-phthalazinedione **9.33**
 2-Aminoethanethiol **13.93**
 3-(2-Aminoethyl)indole **10.34**
 p -(2-Aminoethyl)phenol **4.186**
 1-Amino-4-hydroxyanthraquinone **11.5**
 2-Amino-2-(hydroxymethyl)-1,3-propanediol **17.59**
 3-Amino-2-hydroxypropanethiol **13.131**
 2-Amino-6-hydroxypurine **16.35**
 1-Amino-2-methylantraquinone **11.6**
 α -(Aminomethyl)-3,4-dihydroxybenzyl alcohol **4.6**
 4-(Aminomethylene)-3-methyl-1-phenyl-2-pyrazoline-5-selone **13.141**
 4-(Aminomethylene)-3-methyl-1-phenyl-2-pyrazoline-5-thione **13.142**
 2-Amino-4-(methylthio)butanoic acid **13.108, 4.30, 4.76, 4.92, 4.152, 4.153, 4.157, 4.159, 4.160, 4.165, 4.166, 4.168, 4.180, 4.181, 4.182, 4.187, 4.189, 10.21, 10.36, 10.41**
 2-Aminonaphthalene **9.24**
 4-Aminophenol **4.95**
 1-(*p*-Aminophenylazo)-2-naphthol **11.93**
 4-(4'-Aminophenyl)imino-3-methyl-1-phenyl-2-pyrazolinone **11.133**
 3-Aminophthalhydrazide **9.33**
 1-Aminopropane **8.84**
 2-[*(3*-Aminopropyl)amino]ethanethiol **13.94**
N-(3-Aminopropyl)-1,4-butanediamine **8.10**
 6-Aminopurine **16.63**
 4-Amino-2-pyrimidinone **16.20**
N-[3-(Aminosulfonyl)phenyl]-3-[3-cyano-5-hydroxy-phenylpyrazol-4-yl]azo]-4-methoxybenzenesulfonamide **11.21, 11.22, 11.23, 11.25, 11.155**
 4-(4'-Amino-2',3',5',6'-tetramethylphenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one **11.134**
 4-Amino-2,2,6,6-tetramethylpiperidine **8.54**
 4-Amino-2,2,6,6-tetramethylpiperidine-*N*-oxyl **15.29**
 Ammine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) **12.2**
 Ammonia **14.1**
 Amylene **2.56, 2.122, 2.161, 2.307, 2.309, 2.310, 3.110, 15.13, 15.14, 15.15**
 4-Androsten-3-one, 17-hydroxy-4-methyl- **16.4**
 Androst-5-en-17-one, 3-(acetyloxy)-(3 β) **16.5**
 Anethole **3.116**
 Angelicin **5.1**
 Aniline **9.1**
 Aniline, 4-bromo-*N,N*-dimethyl- **9.2**
 Aniline, 3-chloro-*N,N*-dimethyl- **9.3**
 Aniline, 4-chloro-*N,N*-dimethyl- **9.4**
 Aniline, 3,4-dimethoxy-*N,N*-dimethyl- **9.5**
 Aniline, *N,N*-dimethyl- **9.6**
 Aniline, *N,N*-dimethyl-4-nitroso- **9.7**
 Aniline, 4-(1,2-dimethyl-1-propenyl)-*N,N*-dimethyl- **3.111**
 Aniline, *N,N*-diphenyl- **9.8**
 Aniline, 4-(ethoxycarbonyl)- **9.9**
 Aniline, 4-formyl-*N,N*-dimethyl- **9.10**
 Aniline, 2-methoxy-*N,N*-dimethyl- **9.11**
 Aniline, 3-methoxy-*N,N*-dimethyl- **9.12**
 Aniline, 4-methoxy-*N,N*-dimethyl- **9.13**
 Aniline, *N*-methyl- **9.14**
 Aniline, *N,N*,2,4,6-pentamethyl- **9.15**
 Aniline, *N*-phenyl- **9.16**
 Aniline, *N,N*,2,4-tetramethyl- **9.17**
 Aniline, 4-(1,1,3,3-tetramethylbutyl)-*N*-[4-(1,1,3,3-tetramethylbutyl)phenyl]- **9.18**
 Aniline, *N,N*,4-trimethyl- **9.19**
 Aniline[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) **12.3**
m-Anilinophenol **4.142**
 Anisole **3.43**
 Anisole, 2,6-dichloro- **3.37**
 Anisole, 4-(methylthio)- **13.22**
 2-Anisyl-3-methyl-2-butene **3.118, 3.115, 3.123**
 1-Anisyl-1-propene **3.116**
 Anserine **10.25**
 Anthracenamine, *N*-methyl-9,10-diphenyl- **9.20**
 Anthracene **3.3, 3.23, 3.61, 3.77, 4.61, 4.102, 6.8, 6.15, 6.16, 6.24, 6.25, 6.27**
 Anthracene, 1-chloro- **3.4**
 Anthracene, 9-chloro- **3.5**

- Anthracene, 1-chloro-9,10-diphenyl- 3.6
 Anthracene, 9,10-dichloro- 3.7
 Anthracene, 9,10-dimethoxy- 3.8
 Anthracene, 1,4-dimethoxy-9,10-diphenyl- 3.9
 Anthracene, 9,10-dimethyl- 3.10, 3.124, 3.125, 4.1, 4.10, 4.11,
 4.12, 4.20, 4.153, 4.154, 4.157, 4.159, 4.160, 4.162, 4.165,
 4.166, 4.168, 4.175, 4.176, 4.180, 4.181, 4.182, 11.16, 11.17,
 11.60, 11.62, 11.63, 11.77, 11.79, 11.80, 11.90, 11.92,
 11.101, 11.103, 11.104, 12.28, 12.123, 13.175
 Anthracene, 9,10-diphenyl- 3.11, 4.50, 11.78, 11.128, 11.129,
 11.130, 11.135, 11.136, 11.137, 11.142, 11.143, 11.144,
 11.145, 11.146, 11.147, 13.76
 Anthracene, 9-methoxy- 3.12
 Anthracene, 9-methyl- 3.13
 Anthracene, N-Methylamino-9,10-diphenyl- 9.20
 Anthracene, 9-phenyl- 3.14
 Anthracene-9-carboxylate ion 3.21
 9,10-Anthracenediethanesulfonate ion 3.15, 12.143, 12.167,
 12.169
 9,10-Anthracenedipropionate ion 3.16
 1,5-Anthracenedisulfonate ion 3.17
 Anthracene-9,10-diyl diethyl disulfonate ion 3.15, 12.143,
 12.167, 12.169
 1-Anthracenesulfonate ion 3.18
 2-Anthracenesulfonate ion 3.19
 Anthra[1,9-*bc*:4,10-*b'c'*]dichromene 3.20
 Anthralin 4.2
 Anthralin conjugate base 4.3
 9,10-Anthraquinone, 1-amino- 11.3
 9,10-Anthraquinone, 2-amino- 11.4
 9,10-Anthraquinone, 1-amino-4-hydroxy- 11.5
 9,10-Anthraquinone, 1-amino-2-methyl- 11.6
 9,10-Anthraquinone, 1,4-bis(methylamino)- 11.7
 9,10-Anthraquinone, 1,2-diamino- 11.3, 11.4, 11.5, 11.8, 11.9,
 11.10, 11.11, 11.13, 11.14
 9,10-Anthraquinone, 1,4-diamino- 11.8
 9,10-Anthraquinone, 1,5-diamino- 11.9
 9,10-Anthraquinone, 1,8-diamino- 11.10
 9,10-Anthraquinone, 2,6-diamino- 11.11
 9,10-Anthraquinone, 1,5-diaminobromo-4,8-dihydroxy-
 11.12
 9,10-Anthraquinone, 1,4-diamino-2-methoxy- 11.13
 9,10-Anthraquinone, 1,4,5,8-tetraamino- 11.14
 9-Anthroate ion 3.21
 Anthrone, 1,8-dihydroxy- 4.2
 Anthrone, 1,8-dihydroxy-, conjugate base 4.3
 8'-Apo- β,Ψ -carotenol 2.86
 Aposuperoxide dismutase 10.5
 Aqua[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II) 12.4
 Aqua[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) 12.5
 Arachidic acid 17.27
 Arachidonic acid 2.212
 Arginine 10.6
 Ascorbate ion 16.6
 Astaxanthin 2.82
 ATCC 9760 10.30
 ATEE 4.188
 5,6,6a,7-Tetrahydro-1,10-dimethoxy-6-methyldibenzo[*de,g*]quinoline-2,9-diol 8.20
 5,6,6a,7-Tetrahydro-1,2,9,10-tetramethoxy-6-methyldibenzo[*de,g*]quinoline-2,9-diol 8.21
 Atropine 8.1
 Auranofin 12.170
 1-Azabicyclo[2.2.2]octane 8.97
 8-Azabicyclo[3.2.1]octan-3-ol, 8-methyl-, α -(hydroxymethyl)benceneacetate 8.1
 Azepine-1-carbodithioic acid, hexahydro-, ion(1-) 13.1
 Azide ion 14.2, 8.77
 4-Azido-2,2,6,6-tetramethylpiperidin-1-oxyl 15.30
 Azirino[2',3':3,4]pyrrolo[1,2-*a*]indole-4,7-dione, 6-amino-8-[(aminocarbonyl)oxy]methyl]-1,1a,2,8,8a,8b-hexahydro-8a-methoxy-5-methyl- 16.7
 Azobenzene 11.57
 Azobenzene, 4-[*N*-(2-cyanoethyl)-*N*-ethylamino]- 11.51
 Azobenzene, 4-[*N*-(2-cyanoethyl)-*N*-ethylamino]-4'-methoxy- 11.50
 Azobenzene, 4-(diethylamino)- 11.56, 11.53, 11.90, 11.102
 Azobenzene, 4-(diethylamino)-2',4'-dinitro- 11.52
 Azobenzene, 4-(diethylamino)-4'-methoxy- 11.53
 Azobenzene, 4-(diethylamino)-3'-nitro- 11.54
 Azobenzene, 4-(diethylamino)-4'-nitro- 11.55
 Azobenzene, 4-[*N*-ethyl-*N*-(2-hydroxyethyl)amino]-4'-methoxy- 11.58
 Azobenzene, 4-nitro- 11.61
 Azo dioxide 6 15.20
 Azo dye 10 11.21, 11.22, 11.23, 11.24, 11.25, 11.155
 Bacteriochlorin, tetraphenyl- (*trans*) 7.54
 Bacteriochlorophyll *a* 7.1
 Bacteriochlorophyll *b* 7.2
 Bacteriopheophytin *a* 7.32
 Bacteriopheophytin *b* 7.33
 Basic Violet 3 11.44
 Behenic acid 17.22
 Benzaldehyde, 4-diethylamino-, diphenylhydrazone 15.4
 Benzaldehyde, 4-(dimethylamino)- 9.10
 Benzamide, 4-(bromomethyl)-*N*-(diphenylmethyl)- 3.22
 Benzamide, *N*-[4-[[4-(diethylamino)-2,6-dimethylphenyl]imino]-4,5-dihydro-5-oxo-1-phenyl-1-pyrazol-3-yl]- 11.138
 Benzamide, 4-[3-(2-methoxyphenylaminocarbonyl)-2-hydroxy-1-naphthyl]- 11.79
 Benzamide, 2-(methylseleno)-*N*-phenyl- 13.2
 Benzamide, 2-(methylthio)-*N*-phenyl- 13.3

- Benzamide, *N*-phenyl-2-(phenylmethyl)seleno- 13.4
 Benzamide, *N*-phenyl-2-(phenylmethyl)thio- 13.5
 1,2-Benzanthracene 3.23
 2,3-Benzanthracene 3.124, 3.12, 7.53, 8.72, 8.77, 9.18, 15.46
 Benz[*a*]anthracene 3.23
 Benz[*a*]anthracene, 9,10-dimethyl- 3.24, 3.124, 3.125
 Benz[*b*]anthracene 3.124, 3.12, 7.53, 8.72, 8.77, 9.18, 15.46
 Benz[*de*]anthracen-7-one 3.25
 Benzanthrone 3.25
 Benzenamine, *N,N*-dimethyl-4-(methylthio)- 13.6
 Benzenamine, 4-(1,2-dimethyl-1-propenyl)-*N,N*-dimethyl- 3.115
 Benzenamine, *N*-(diphenylmethylen)-, *N*-oxide 15.5
 Benzenamine, *N*-fluoren-9-ylidene, *N*-oxide 15.6
 Benzenamine, *N*-[(2-hydroxyphenyl)methylene]-, *N*-oxide 15.7
 Benzenamine, 4-methyl-*N*-(phenylmethylen)-, *N*-oxide 15.8
 Benzenamine, *N*-[(4-methylphenyl)methylene]-, *N*-oxide 15.9
 Benzenamine, *N*-(phenylmethylen)-, *N*-oxide 15.10
 Benzenamine, 4-(1,1,3,3-tetramethylbutyl)-*N*-[4-(1,1,3,3-tetramethylbutyl)phenyl]- 9.18
 Benzene 3.26
 Benzene-*d*₆ 3.27
 Benzene, 1-benzyl-2-hydroxy- 4.98
 Benzene, 1,5-bis(1,1-dimethylethyl)-2,4-dimethoxy- 3.28
 Benzene, 1,3-bis(1,1-dimethylthyl)-2-methoxy-5-methyl- 3.29
 Benzene, bromo- 3.30
 Benzene, 1-bromo-4-(diazophenylmethyl)- 11.15
 Benzene, 1-bromo-4-(dimethylamino)- 9.2
 Benzene, 1-bromo-4-(methylthio)- 13.7
 Benzene, 1,1'-(1,3-butadiene-1,4-diyl)bis[4-methoxy- 2.35
 Benzene, (2-butenylsulfinyl)- (*E*) 13.54
 Benzene, (2-butenylsulfinyl) (*Z*) 13.55
 Benzene, (butylthio)- 13.8
 Benzene, 1-(butylthio)-3-chloro- 13.9
 Benzene, 1-(butylthio)-4-chloro- 13.10
 Bcnznc, 1-(butylthio)-4-fluoro- 13.11
 Benzene, 1-(butylthio)-4-methoxy- 13.12
 Benzene, 1-(butylthio)-4-methyl- 13.13
 Benzene, chloro- 3.31
 Benzene, 1-chloro-4-(butylthio)- 13.10
 Benzene, 2-chloro-1,4-dimethoxy- 3.32
 Benzene, 5-chloro-1,3-dimethoxy- 3.33
 Benzene, 1-chloro-3-(1,2-dimethyl-1-propenyl)- 3.109
 Benzene, 1-chloro-4-(1,2-dimethyl-1-propenyl)- 3.110,
 3.121, 3.122
 Benzene, 1-chloro-2-methoxy- 3.34
 Benzene, 1-chloro-3-(methylthio)- 13.14
 Benzene, 1-chloro-4-(methylthio)- 13.15
 Benzene, 1-chloro-4-(phenylthio)- 13.16
 Benzene, (1-cyclopropyl-2-methyl)propenyl- 3.113
 Benzene, 1,1'-(diazomethylene)bis- 11.16, 11.15, 11.17,
 11.18, 11.20
 Benzene, 1,1'-(diazomethylene)bis[4-chloro- 11.17
 Benzene, 1,1'-(diazomethylene)bis[4-methyl- 11.18
 Benzene, 1,1'-(diazomethylene)bis[2,4,6-trimethyl- 11.19
 Benzene, 1,3-dichloro- 3.35
 Benzene, 1,4-dichloro- 3.36
 Benzene, 2,6-dichloro-1-methoxy- 3.37
 Benzene, 1,4-dihydroxy-2-methyl- 4.29
 Benzene, 1,2-dimethoxy- 3.38
 Benzene, 1,3-dimethoxy- 3.39
 Benzene, 1,4-dimethoxy- 3.40
 Benzene, 1,1'-(1,2-dimethoxy-1,2-ethenediyl)bis- 3.101
 Benzene, [(2,3-dimethyl-2-butenyl)thio]- 13.17
 Benzene, [(2,3-dimethyl-2-butenyl)thio]-4-methoxy- 13.19
 Benzene, [(2,3-dimethyl-2-butenyl)thio]-4-methyl- 13.19
 Benzene, 1,1'-(1,2-dimethyl-1,2-ethenediyl)bis-, (*E*)-
 3.102, 3.103
 Benzene, 1,1'-(1,2-dimethyl-1,2-ethenediyl)bis-, (*Z*)- 3.101
 Benzene, 1-(1,1-dimethylethyl)-4-(methylthio)- 13.20
 Benzene, (1,2-dimethyl-1-propenyl)- 3.123, 2.96
 Benzene, 1-(1,2-dimethyl-1-propenyl)-3-methoxy- 3.117
 Benzene, 1-(1,2-dimethyl-1-propenyl)-4-methoxy- 3.118,
 3.115, 3.123
 Benzene, 1-(1,2-dimethyl-1-propenyl)-3-methyl- 3.121
 Benzene, 1-(1,2-dimethyl-1-propenyl)-4-methyl- 3.122,
 3.109, 3.111, 3.112, 3.117, 3.118, 3.123
 Benzene, 1,1'-ethenediylbis[4-methoxy- 3.100
 Benzene, 1,1'-ethenylidenebis[4-methoxy- 3.100
 Benzene, ethyl- 3.41
 Benzene, 1-fluoro-4-(butylthio)- 13.11
 Benzene, 1-fluoro-4-(methylthio)- 13.21
 Benzene, hexamethoxy- 3.42
 Benzene, methoxy- 3.43
 Benzene, 1-methoxy-4-(butylthio)- 13.12
 Benzene, 1-methoxy-4-(diazophenylmethyl)- 11.20
 Benzene, 1-methoxy-4-(methylthio)- 13.22
 Benzene, 1-methoxy-2-nitro- 3.44
 Benzene, 1-methoxy-4-(phenylthio)- 13.23
 Benzene, 1-methoxy-4-(1-propenyl)- 3.116
 Benzene, (3-methyl-2-butenyl)- 3.45
 Benzene, [(3-methyl-2-butenyl)sulfinyl] 13.53
 Benzene, 1-methyl-4-(butylthio)- 13.13
 Benzene, 1,1'-(1-methyl-1,2-ethenediyl)bis-, (*E*)- 3.104
 Benzene, 1,1'-(1-methyl-1,2-ethenediyl)bis-, (*Z*)- 3.105
 Benzene, 1-methyl-3-(methylthio)- 13.24
 Benzene, 1-methyl-4-(methylthio)- 13.25
 Benzene, 1-methyl-4-(phenylthio)- 13.26
 Benzene, (2-methyl-1-propenyl)- 3.114

- Benzene, 1,1'-(2-methyl-1-propenylidene)bis- **3.46**
 Benzene, [(2-methyl-2-propenyl)sulfinyl] **2.338**
 Benzene, methylseleno- **13.27**
 Benzene, methyltelluro- **13.28**
 Benzene, (methylthio)- **13.29**
 Benzene, [(methylthio)methyl]- **13.30**
 Benzene, nitro- **3.47**
 Benzene, 1-nitro-4-(phenylthio)- **13.31**
 Benzene, 1,3-pentadienyl-, (*E,E*)- **2.301**
 Benzene, 2,4-pentadienyl- **2.43**
 Benzene, pentamethoxy- **3.48**
 Benzene, [(phenylmethyl)seleno]- **13.32**
 Benzene, [(phenylmethyl)thio]- **13.33**
 Benzene, 1-propenyl-, (*E*)- **3.119**
 Benzene, 1-propenyl-, (*Z*)- **3.120**
 Benzene, 1,2,3,4-tetramethoxy- **3.49**
 Benzene, 1,2,3,5-tetramethoxy- **3.50**
 Benzene, 1,2,4,5-tetramethoxy- **3.51**, 3.42, 3.48, 3.50
 Benzene, 1,1'-thiobis- **13.34**, 13.16, 13.23, 13.26, 13.31
 Benzene, 1,2,3-trimethoxy- **3.52**
 Benzene, 1,2,4-trimethoxy- **3.53**, 3.51
 Benzene, 1,3,5-trimethoxy- **3.54**
 Benzeneacetic acid, α -hydroxy-, 1,2,2,6-tetramethyl-4-piperidinyl ester **8.76**
 Benzenecarboxylic acid, α -(hydroxymethyl)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl ester **8.1**
 Benzenecarbothioamide, 4-methoxy- **13.154**
 Benzenedecanoic acid, 4,4'-(4,7-dihydro-5,6-dimethyl-1,3-isobenzofurandiyi)bis-, dianion **5.81**
 1,2-Benzenediamine **9.25**
 1,4-Benzenediamine, N^4 -(6-amino-3-phenylpyrazolo[5,1-*c*]-1,2,4-triazol-7-ylidene)- N^1,N^1 -diethyl-3-methyl- **11.21**
 1,4-Benzenediamine, *N*-cyclohexyl-*N'*-phenyl- **9.28**
 1,4-Benzenediamine, N^1,N^1 -diethyl-3-methyl- N^4 -(6-methyl-3-phenylpyrazolo[5,1-*c*]-1,2,4-triazol-7-ylidene)- **11.22**
 1,4-Benzenediamine, N^1,N^1 -diethyl-3-methyl- N^4 -(6-phenylamino)-3-phenylpyrazolo[5,1-*c*]-1,2,4-triazol-7-ylidene)- **11.23**
 1,4-Benzenediamine, N^1,N^1 -diethyl-3-methyl- N^4 -(3-phenylpyrazolo[5,1-*c*]-1,2,4-triazol-7-ylidene)- **11.24**
 1,4-Benzenediamine, N^1,N^1 -diethyl-3-methyl- N^4 -[3-phenyl-6-(2-thienyl)pyrazolo[5,1-*c*]-1,2,4-triazol-7-ylidene]- **11.25**
 1,4-Benzenediamine, *N*-(1-methylethyl)-*N'*-phenyl- **9.30**
 1,4-Benzenediamine, *N,N,N',N'*-tetramethyl- **9.31**
 1,2-Benzenediol **4.4**
 1,2-Benzenediol, 4-(2-aminoethyl)- **4.5**
 1,2-Benzenediol, 4-(2-amino-1-hydroxyethyl)- **4.6**
 1,2-Benzenediol, 3,5-di-(1,1-dimethylethyl)- **4.7**
 1,2-Benzenediol, 4-(1,1-dimethylethyl)- **4.8**
 1,2-Benzenediol, 4-(2-methylamino-1-hydroxyethyl)- **4.9**
 1,3-Benzenediol **4.10**
 1,3-Benzenediol, 4-chloro- **4.11**
 1,4-Benzenediol **4.12**
 1,4-Benzenediol, 2,6-bis(1,1-dimethylethyl)-, 4-propanoate **4.15**
 1,4-Benzenediol, chloro- **4.20**
 1,4-Benzenediol, 2,5-di-*tert*-butyl- **4.14**
 1,4-Benzenediol, 2,5-di(2,2-dimethylpropyl)- **4.16**
 1,4-Benzenediol, 2,5-di-*sec*-dodecyl- **4.17**
 1,4-Benzenediol, 2,5-di-*sec*-hexadecyl- **4.18**
 1,4-Benzenediol, 2,3-dimethoxy-5-methyl- **4.21**
 1,4-Benzenediol, 2,3-dimethyl- **4.22**
 1,4-Benzenediol, 2,5-dimethyl- **4.23**
 1,4-Benzenediol, 2,6-dimethyl- **4.24**
 1,4-Benzenediol, 2,5-di-*sec*-octyl- **4.19**
 1,4-Benzenediol, 2-(3,7,11,15,19,23-hexamethyl-2,6,10,14,18,22-tetracosahexaenyl)-5,6-dimethoxy-3-methyl- (*all-E*) **4.25**
 1,4-Benzenediol, 2-(3-hydroxy-3,7,11,15-tetramethylhexadecyl)-3,6-dimethyl- **4.26**
 1,4-Benzenediol, 2-(3-hydroxy-3,7,11,15-tetramethylhexadecyl)-5,6-dimethyl- **4.27**
 1,4-Benzenediol, 2-(3-hydroxy-3,7,11,15-tetramethylhexadecyl)-3,5,6-trimethyl- **4.28**
 1,4-Benzenediol, phenyl- **4.30**
 1,4-Benzenediol, trimethyl- **4.31**
 1,4-Benzenediol dianion **4.13**
 1,3-Benzenedisulfonamide, *N*-[5-hydroxy-8-[[2-(methylsulfonyl)-4-nitrophenyl]azo]-1-naphthalenyl]-conjugate base **11.85**
 1,3-Benzenedisulfonamide, *N*-[5-hydroxy-8-[[2-(methylsulfonyl)-4-nitrophenyl]azo]-1-naphthalenyl]-conjugate dibase **11.86**
 Benzeneethanol, β -phenyl- α -pyrrolidinyl- **8.2**
 Benzenemethanamine **8.5**
 Benzenemethanamine, *N*-hydroxy-*N*-(phenylmethyl)- **15.11**
 Benzenemethanamine, *N*-(phenylmethylen)-, *N*-oxide **15.12**
 Benzenemethanesulfonamide, *N,N*-diethyl- **13.35**
 Benzenemethanol, 3,5-bis(1,1-dimethylethyl)-4-hydroxy- **4.32**
 Benzenemethanol, α -(1,1-dimethylethyl)-4-hydroxy-3-methoxy- **4.127**
 Benzenemethanol, α -methyl- **3.55**
 Benzenepropanamide, α -[(4-diethylamino)-2,6-dimethylphenyl]imino- β -oxo-*N*-phenyl- **11.26**
 Benzenepropanamide, α -[(4-diethylamino)-2-methylphenyl]imino- β -oxo-*N*-phenyl- **11.27**
 Benzenepropanamide, α -[(4-diethylamino)phenyl]imino- β -oxo-*N*-phenyl- **11.28**
 Benzenepropanamine **8.50**

- 3-Benzene propanoic acid, 3,5-bis(1,1-dimethylethyl)-4-hydroxy-, octadecyl ester **4.33**
- 3-Benzene propanoic acid, 4-hydroxy- **4.34**
- Benzene sulfonamide, *N*-[3-(aminosulfonyl)phenyl]-3-[3-cyano-5-hydroxy-1-phenylpyrazol-4-yl)azo]-4-methoxy- **11.21, 11.22, 11.23, 11.24, 11.25, 11.155**
- Benzene sulfonate ion, 2,2'-(1,2-ethenediyil)bis[5-(acetylamino)-] **3.106**
- Benzene sulfonate ion, 2,2'-(1,2-ethenediyil)bis[5-amino-] **3.107**
- N*-Benzhydryl-4-bromomethylbenzamide **3.22**
- α -Benzhydrylfuryl alcohol **5.68**
- Benzidine **9.23**
- Benz[e]indolium, 2-[7-(1,3-dihydro-1,1-dimethyl-3-(sulfobutyl)benz[e]indol-2-ylidene]-1,3,5-heptatrienyl]-1,1-dimethyl-3-(sulfobutyl)-, hydroxide, inner salt, Na salt **11.29**
- Benz[e]indolium, 2-[2-[3-[(1,3-dihydro-1,1-dimethyl-3-(3-sulfopropyl)benz[e]indol-2-ylidene]ethylidene]-2-[4-(ethoxycarbonyl)-1-piperidinyl]-1-cyclopenten-1-yl]ethenyl]-1,1-dimethyl-3-(3-sulfopropyl)-, hydroxide, inner salt, compound with triethylamin **11.30**
- 1,2-Benzisoselenazol-3-one, 7-chloro-2-phenyl- **13.36**
- 1,2-Benzisoselenazol-3-one, 7-fluoro-2-phenyl- **13.37**
- 1,2-Benzisoselenazol-3-one, 7-methoxy-2-phenyl- **13.38**
- 1,2-Benzisoselenazol-3-one, 7-nitro-2-phenyl- **13.39**
- 1,2-Benzisoselenazol-3-one, 2-phenyl **13.40**
- 1,2-Benzisothiazol-3-one, 2-phenyl- **13.41**
- Benzoate ion, 2-hydroxy- **4.35**
- Benzo[1,3]cyclopropa[1,2,3-*cd*]cyclopropa[*gh*]pentalene, 1,2,5,6,6a,6b,6c,6d-octahydro-1,6-*o*-benzeno- **2.181**
- Benzo[1,3]cyclopropa[1,2,3-*cd*]cyclopropa[*gh*]pentalene, 1,2,5,6,6a,6b,6c,6d-octahydro-9,12-dimethoxy-1,6-*o*-benzeno- **2.180**
- Benzo[1,2-*b*:4,3-*b'*]dipyran, 1,2,3,8,9,10-hexahydro-3,3,5,6,8,8-hexamethyl- **5.2**
- Benzo[1,2-*b*:4,5-*b'*]dipyran, 2,3,4,7,8,9-hexahydro-2,2,5,7,7,10-hexamethyl- **5.3**
- Benzo[1,2,3-*k*l:4,5,6-*k'l'*]dixanthene **3.56**
- 3,4-Benzofuran **5.80**
- Benzo[c]furan **5.80**
- Benzofuran-5-ol, 2,3-dihydro-2,2,4,6,7-pentamethyl- **4.36**
- Benzofuran-5-ol, 2,3-dihydro-2,4,6,7-tetramethyl- **4.37**
- Benzoic acid, 4-amino-, ethyl ester **9.9**
- Benzoic acid, 3,5-bis(1,1-dimethylethyl)-4-hydroxy-, 2,4-bis(1,1-dimethylethyl)phenyl ester **4.38**
- Benzoic acid, 3,5-bis(1,1-dimethylethyl)-4-hydroxy-, methyl ester **4.39**
- Benzoic acid, 4-(dimethylamino)-, ethyl ester **9.21**
- Benzoic acid, methyl ester **3.57**
- Benzo[e]naphtho[2',3':5,6]fluoreno[1,9ab]oxepin-5,10,19-trione, 5c,8,8a,16-tetrahydro-1,8,11,15,18-pentahydroxy-13-methyl- **16.8**
- Benzonitrile, 4-(dimethylamino)- **9.22**
- Benzonitrile, 3-(1,2-dimethyl-1-propenyl)- **3.111**
- Benzonitrile, 4-(1,2-dimethyl-1-propenyl)- **3.112**
- Benzonitrile, 4-hydroxy- **4.112**
- 4-(2',3'-Benzo-4'-oxocyclohexadienyliden)amino-*N,N*-diphenylaniline **11.82**
- 4-(2',3'-Benzo-4'-oxocyclohexadienyliden)amino-*N*-phenylaniline **11.84**
- 4-[2',3'-Benzo-4'-oxo-5'-(2-methoxy-5-fluorosulfonyl)phenyl]amino-*N,N*-diethylaniline **11.81**
- Benzo[*ghi*]perylene-4,11-dione, 1-acetyl-1,2-dihydro-5,10-dihydroxy-2-(1-hydroxyethyl)-3,7,8,12-tetramethoxy- **16.9**
- Benzo[*ghi*]perylene-4,11-dione, 1,2-diacetyl-1,2-dihydro-5,10-dihydroxy-3,7,8,12-tetramethoxy-, *trans*- **16.10**
- Benzophenone **3.58**
- Benzophenone, 4-dodecyloxy-2-hydroxy- **4.42**
- Benzophenone, 2-hydroxy-4-octyloxy- **4.43**
- Benzophenone oximate anion **15.13**
- Benzophenone oxime **15.14**
- Benzophenone oxime *O*-methyl ether **15.15**
- 1-Benzopyran, 3,4-dihydro-6-methoxy-2,5,7,8-tetramethyl-2-(4,8,12-trimethyltridecyl)- **5.4**
- 1-Benzopyran, 6-ethoxy-3,4-dihydro-2,5,7,8-tetramethyl-(4,8,12-trimethyltridecyl)- **5.5**
- 1-Benzopyran-2-carboxylic acid, 3,4-dihydro-6-hydroxy-2,5,7,8-tetramethyl-, ion(1-) **4.44**
- 1-Benzopyran-6-ol, 5,7-diethyl-3,4-dihydro-2-methyl-2-(4,8,12-trimethyltridecyl)- **4.45**
- 1-Benzopyran-6-ol, 3,4-dihydro-2,8-dimethyl-2-(4,8,12-trimethyltridecyl)- **4.46**
- 1-Benzopyran-6-ol, 3,4-dihydro-2-methyl-5,7-di(1-methylethyl)-2-(4,8,12-trimethyltridecyl)- **4.47**
- 1-Benzopyran-6-ol, 3,4-dihydro-1-methyl-1-(4,8,12-trimethyltridecyl)- **4.48**
- 1-Benzopyran-6-ol, 3,4-dihydro-2,2,5,7,8-pentamethyl- **4.49**
- 1-Benzopyran-6-ol, 3,4-dihydro-2,5,7,8-tetramethyl-2-(4,8,12-trimethyltridecyl)- **4.50**
- 1-Benzopyran-6-ol, 3,4-dihydro-2,5,7,8-tetramethyl-2-(4,8,12-trimethyltridecyl)-, acetate **16.11**
- 1-Benzopyran-6-ol, 3,4-dihydro-2,5,7-trimethyl-2-(4,8,12-trimethyltridecyl)- **4.51**
- 1-Benzopyran-6-ol, 3,4-dihydro-2,5,8-trimethyl-2-(4,8,12-trimethyltridecyl)- **4.52**
- 1-Benzopyran-6-ol, 3,4-dihydro-2,7,8-trimethyl-2-(4,8,12-trimethyltridecyl)- **4.53**
- 1-Benzopyran-6-ol, 7-(1,1-dimethylethyl)-3,4-dihydro-2,2-dimethyl- **4.54**
- 1-Benzopyran-6-ol, 7-(1,1-dimethylethyl)-3,4-dihydro-2-methyl-2-(4,8,12-trimethyltridecyl)- **4.55**
- 1-Benzopyran-6-ol, 8-(1,1-dimethylethyl)-3,4-dihydro-2-methyl-2-(4,8,12-trimethyltridecyl)- **4.56**

- 1-Benzopyran-6-ol, 2,2'-spirobis[7-*tert*-butyl-3,4-dihydro-4,4-dimethyl- 4.184
- 1-Benzopyran-6-ol, 2,2'-spirobis[3,4-dihydro-4,4,7-trimethyl- 4.185
- 1-Benzopyran-4-one, 2-[3,4-bis(2-hydroxyethoxy)phenyl]-3-[6-*O*-(6-deoxy- α -L-mannopyranosyl- β -D-glycopyranosyl)oxy]-5-hydroxy-7-(2-hydroxyethoxy)- 4.57
- 1-Benzopyran-4-one, 3-[[6-*O*-(6-deoxy- α -mannopyranosyl)- β -glucopyranosyl]oxy]-2-(3,4-dihydroxyphenyl)-5,7-dihydroxy- 4.58
- 1-Benzopyran-4-one, 3,7-dihydroxy-2-(3,4-dihydroxyphenyl)- 4.75
- 1-Benzopyran-4-one, 2-(2,3-dihydroxyphenyl)-2,3-dihydro-5,7-dihydroxy-, (*S*) 4.59
- 1-Benzopyran-4-one, 2-(3,4-dihydroxyphenyl)-5,7-dihydroxy- 4.60
- 1-Benzopyran-4-one, 2-(3,4-dihydroxyphenyl)-3,5,7-trihydroxy- 4.61
- 1-Benzopyran-4-one, 3-hydroxy-2-phenyl- 4.74
- 1-Benzopyran-4-one, 5,6,7,8-tetramethoxy-2-(4-methoxyphenyl)- 4.62
- 1-Benzopyran-4-one, 3,5,7-trihydroxy-2,3-dihydro-2-(3,4-dihydroxyphenyl)- 4.63
- 1-Benzopyran-4-one, 3,5,7-trihydroxy-2-(4-hydroxyphenyl)- 4.64
- 1-Benzopyran-4-one, 3,5,7-trihydroxy-2-phenyl- 4.65
- [1]Benzopyrano[6,7,8-*ij*]quinolizin-11-one, 2,3,6,7-tetrahydro- 8.3
- [1]Benzopyrano[6,7,8-*ij*]quinolizin-11-one, 2,3,6,7-tetrahydro-9-methyl- 8.4
- 1-Benzopyran-2-propanoic acid, 3,4-dihydro-6-hydroxy-2,5,7,8-tetramethyl- 4.66
- 1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro- (2*R-trans*)- 4.67
- 1-Benzopyrylium, 8-[5-(6,7-dihydro-2,4-diphenyl-1-benzopyran-8-yl)-2,4-pentadienylidene]-5,6,7,8-tetrahydro-2,4-diphenyl-, perchlorate 11.31
- 1,4-Benzoquinone 16.12
- 1,4-Benzoquinone, methyl- 16.13
- Benzothiazolium, 2-[(2-butoxy-3-[(3-ethyl-2-benzothiazolylidene)methyl]-4-oxo-2-cyclobuten-1-ylidene)methyl]-3-ethyl- 11.32
- Benzothiazolium, 5-chloro-2-[2-3-[(5-chloro-3-ethyl-2-benzothiazolylidene)ethylidene]-2-(diphenylamino)-1-cyclopenten-1-yl]ethenyl]-3-ethyl-, perchlorate 11.33
- Benzothiazolium, 5-chloro-2-[(5-chloro-3-ethyl-2-benzothiazolylidene)methyl]-1-butenyl]-3-ethyl-, bromide 11.164
- Benzothiazolium, 5-chloro-2-[3-(5-chloro-3-ethylbenzothiazolylidene)-1-propenyl]-3-ethyl-, bromide 11.163
- Benzothiazolium, 2-[7-(5-chloro-1,3,3-trimethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)-3-methyl-, iodide 11.34
- Benzothiazolium, 5-cyano-2-[(5-cyano-3-ethyl-2-benzothiazolylidene)methyl]-1-butenyl]-3-ethyl-, tetrafluoroborate 11.165
- Benzothiazolium, 3-ethyl-2-[(3-ethyl-2-benzothiazolylidene)ethyl]-1-methylethenyl-, toluenesulfonate 11.168
- Benzothiazolium, 3-ethyl-2-[7-(3-ethyl-2-benzothiazolylidene)-1,3,5-heptatrienyl]-, iodide 11.174
- Benzothiazolium, 3-ethyl-2-[(3-ethyl-2-benzothiazolylidene)methyl]-1-butenyl-, bromide 11.171
- Benzothiazolium, 3-ethyl-2-[(3-ethyl-2-benzothiazolylidene)methyl]-1-cyclopentenyl-, iodide 11.170
- Benzothiazolium, 3-ethyl-2-[(3-ethyl-2-benzothiazolylidene)methyl]-1-cyclopenten-1-yl-, toluenesulfonate 11.169
- Benzothiazolium, 3-ethyl-2-[(3-ethyl-2-benzothiazolylidene)-1,3-pentadienyl]-, iodide 11.173
- Benzothiazolium, 3-ethyl-2-[3-(3-ethyl-2-benzothiazolylidene)-1-propenyl], toluenesulfonate 11.166
- Benzothiazolium, 3-ethyl-2-[(3-ethyl-5-methoxy-2-benzothiazolylidene)methyl]-1-butenyl]-5-methoxy-, toluenesulfonate 11.172
- Benzothiazolium, 3-ethyl-2-[3-(3-ethyl-5-methoxybenzothiazolylidene)-1-propenyl]-5-methoxy-, toluenesulfonate 11.167
- Benzothiazolium, 2-[7-(5-fluoro-1,3,3-trimethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)-3-methyl-, iodide 11.35
- Benzothiazolium, 2-[7-(5-methoxy-1,3,3-trimethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)-3-methyl-, iodide 11.36
- Benzothiazolium, 3-methyl-2-[7-(1,3,3-trimethyl-2-indolylidene)-1,3,5-heptatrienyl]-, iodide 11.37
- Benzothiazolium, 2-[7-(1,3,3,5-tetramethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)-3-methyl-, iodide 11.38
- Benzotriazole, 2-(2'-hydroxy-3'-chloro-5'-*tert*-butylphenyl)- 4.97
- Benzotriazole, 2-(2'-hydroxy-3',5'-di-*tert*-pentylphenyl)- 4.96
- 3-Benzoxazolepropanesulfonic acid, 2-[4-(1,3-dibutyltetrahydro-4,6-dioxo-2-thioxo-5-pyrimidinylidene)-2-but enylidene]-, sodium salt 11.39
- Benzoxazolium, 3-ethyl-2-[7-(3-ethyl-2-benzoxazolylidene)-1,3,5-heptatrienyl]-, iodide 11.109
- 5-Benzoyl-3,4-dihydro-6-phenylpyran 5.89

- 3-Benzoyloxy-1,3,5(10),9(11)-estratetraen-17-one **16.30**
- 4-[*N*-Benzoyl(phenylaminocarbonyl)methylene]amino-
N,N-diethylaniline **11.28**
- 4-[*N*-Benzoyl(phenylaminocarbonyl)methylene]amino-
N,N-diethyl-3,5-dimethylaniline **11.26**
- 4-[*N*-Benzoyl(phenylaminocarbonyl)methylene]amino-
N,N-diethyl-3-methylaniline **11.27**
- Benzvalene **2.3**
- Benzyl alcohol, α -(aminomethyl)-3,4-dihydroxy- **4.6**
- Benzylamine **8.5**
- Benzylamine, *N*-benzyl-*N*-hydroxy- **15.11**
- 2-Benzyl-1,3-butadiene **2.43**
- 4-Benzyl-2,6-di-*tert*-butylphenol **4.104**
- α -Benzylfurfuryl alcohol **5.72**
- p,p'*-Benzylidenebis(*N,N*-dimethylaniline) **11.77**
- Benzyl mercaptan **13.42**
- Benzyl methyl sulfide **13.30**
- 4-Benzylxy-5,6-dimethyl-2-(dimethylamino)pyrimidine
16.65
- 2-Benzylphenol **4.98**
- 2-Benzylphenoxyde ion **4.152**
- Benzyl phenyl selenide **13.32**
- Benzyl phenyl sulfide **13.33**
- 2-(Benzylseleno)-*N*-phenylbenzamide **13.4**
- Benzyl sulfide **13.71**
- Biadamantylidene **2.356, 13.112**
- Bicyclo[2.2.1]hepta-2,5-diene **2.4**
- Bicyclo[2.2.1]hepta-2,5-diene, 7,7-dimethyl-2-
(trimethylsiloxy)- **2.5**
- Bicyclo[2.2.1]hepta-2,5-diene, 2-methyl- **2.6**
- Bicyclo[2.2.1]hepta-2,5-diene, 2-(trimethylsiloxy)- **2.7**
- Bicyclo[2.2.1]heptane, 2,3-bis(methylene)- **2.8, 2.19, 2.140,**
2.295, 2.296, 2.297, 2.357
- Bicyclo[2.2.1]heptane, 7,7-dimethyl-2-methylene- **2.9**
- Bicyclo[2.2.1]heptane, 2,3-dioxymethyl-7-
tricyclo[3.3.1.1^{3,7}]decylidene- (*exo,exo*) **2.10**
- Bicyclo[2.2.1]heptane, 2-methylene- **2.11, 2.9, 2.13, 2.14,**
2.21, 2.23
- Bicyclo[2.2.1]heptane, 2,3-(2¹-oxatrimethylene)-7-
tricyclo[3.3.1.1^{3,7}]decylidene- (*exo,exo*) **2.12**
- Bicyclo[2.2.1]heptane-2-*d*, 3-methylene-, *endo*- **2.13**
- Bicyclo[2.2.1]heptane-2-*d*, 3-methylene-, *exo*- **2.14**
- Bicyclo[2.2.1]heptane-2,3-dicarboxylic acid, 7-
tricyclo[3.3.1.1^{3,7}]decylidene-, anhydride
(*endo,endo*) **2.15**
- Bicyclo[2.2.1]heptane-2,3-dicarboxylic acid, 7-
tricyclo[3.3.1.1^{3,7}]decylidene-, anhydride (*exo,exo*)
2.16
- Bicyclo[2.2.1]heptane-2,3-dicarboxylic acid, 7-
tricyclo[3.3.1.1^{3,7}]decylidene-, dimethyl ester
(*endo,endo*) **2.17**
- Bicyclo[2.2.1]heptane-2,3-dicarboxylic acid, 7-
tricyclo[3.3.1.1^{3,7}]decylidene-, dimethyl ester
(*exo,exo*) **2.18**
- Bicyclo[2.2.1]heptane-2-thione **13.43**
- Bicyclo[2.2.1]heptane-2-thione, 3,3-dimethyl- **13.44**
- Bicyclo[2.2.1]heptane-2-thione, 1,3,3,7,7-pentamethyl-
13.45
- Bicyclo[2.2.1]heptane-2-thione, 1,3,3-trimethyl- **13.46**
- Bicyclo[2.2.1]heptane-2-thione, 1,7,7-trimethyl- **13.47**
- Bicyclo[2.2.1]hept-2-ene, 5,6-bis(methylene)- **2.19**
- Bicyclo[2.2.1]hept-2-ene, 7,7-dimethyl-2-(trimethylsiloxy)-
2.5, 2.7, 2.24
- Bicyclo[2.2.1]hept-2-ene, 2-methoxy- **2.20**
- Bicyclo[2.2.1]hept-2-ene, 2-methyl- **2.21, 2.23**
- Bicyclo[2.2.1]hept-2-ene, 5-methylene- **2.22, 2.6, 2.11, 2.21,**
2.153
- Bicyclo[2.2.1]hept-2-ene, 2,7,7-trimethyl- **2.23, 2.9, 2.11**
- Bicyclo[2.2.1]hept-2-ene, 2-(trimethylsiloxy)- **2.24**
- Bicyclo[3.1.1]hept-2-ene, 2-ethenyl-6,6-dimethyl- **2.25**
- Bicyclo[4.1.0]hept-2-ene, 3,7,7-trimethyl- **2.26**
- Bicyclo[4.1.0]hept-2-ene, 4,7,7-trimethyl- **2.27**
- Bicyclo[4.1.0]hept-3-ene, 3,7,7-trimethyl- **2.28**
- Bicyclo[3.1.1]hept-3-ene-2-thione, 4,6,6-trimethyl- **13.48**
- Bicyclo[2.2.0]hexa-2,5-diene, hexamethyl- **2.29**
- Bicyclo[3.1.0]hex-2-ene, 4-methyl-1-(1-methylethyl)-,
(*1R*)- **2.30**
- Bicyclo[3.2.2]nonane, 6,7-bis(methylene)- **2.31**
- Bicyclo[4.2.0]octa-2,4-diene, 7,8-dibromo- **2.32**
- Bicyclo[2.2.2]octane, 2,3-bis(methylene)- **2.33**
- Bicyclo[7.2.0]undec-4-ene, 4,11,11-trimethyl-8-methylene-
[*IR*-(*IR*^{*},*4E,9S*^{*})]- **2.34**
- Bicyclo[7.2.0]undec-4-ene, 4,11,11-trimethyl-8-methylene-
[*IR*-(*IR*^{*},*4Z,9S*^{*})]- **2.34**
- Bifluorenylidene **3.66**
- 9,9'-Bifluorenylidene **3.66**
- 2,3'-Biindolium, 1,1,3-trimethyl-2'-(2-(1,3,3-
trimethylindolylidene)methyl)-, iodide **11.40**
- Biline-1,19-dione, 3,8,12,17-tetraethyl-22,24-dihydro-
2,7,13,18-tetramethyl- **7.3**
- Biline-1,19-dione, 3,8,12,17-tetraethyl-10,22,23,24-
tetrahydro-2,7,13,18-tetramethyl- **7.4**
- Biline-8,12-dipropanoic acid, 2,17-diethenyl-
1,10,19,22,23,24-hexahydro-3,7,13,18-tetramethyl-
1,19-dioxo- **7.5, 3.60**
- Biline-8,12-dipropanoic acid, 2,17-diethenyl-
1,10,19,22,23,24-hexahydro-3,7,13,18-tetramethyl-
1,19-dioxo-, dimethyl ester **7.6**
- Biline-8,12-dipropanoic acid, 2,17-diethenyl-1,19,22,24-
tetrahydro-3,7,13,18-tetramethyl-1,19-dioxo- **7.7**
- Biline-8,12-dipropanoic acid, 2,17-diethenyl-1,19,22,24-
tetrahydro-3,7,13,18-tetramethyl-1,19-dioxo- **7.7**

- (15*E*)-Biline-8,12-dipropanoic acid, 2,17-diethenyl-1,19,22,24-tetrahydro-3,7,13,18-tetramethyl-1,19-dioxo-, dimethyl ester **7.8**
- Bilirubin **7.5**, **3.60**
- Bilirubin ditaurate **7.10**
- Bilirubin IX **7.5**, **3.60**
- Bilirubin IX dimethyl ester **7.6**
- Biliverdin **7.7**
- Biliverdin, dimethyl ester **7.8**
- Biliverdin IX **7.7**
- Bioresmethrin **5.76**
- o,o'*-Biphenol **4.128**
- Biphenyl, 4,4'-diamino- **9.23**
- Biphenyl, 2,2'-dihydroxy- **4.128**
- Biphenyl, 2,2'-dihydroxy-, conjugate base **4.166**
- Biphenyl, 2,5-dihydroxy-, conjugate base **4.167**
- Biphenyl, 2-hydroxy-, conjugate base **4.180**
- Biphenyl, 4-hydroxy-, conjugate base **4.181**
- Biphenyl, 2-methoxy- **3.59**
- 1,1'-Biphenyl-2,5-diol **4.30**
- 1,1'-Biphenyl-2-ol **4.141**
- [1,1'-Biphenyl]-4-ol, 2,6-bis(1,1-dimethylethyl)- **4.103**
- 1,1'-Bipiperidine **8.7**
- 1,1'-Bipyrrolidine **8.8**
- Bis(acetylacetato)acetatomanganese(III) **12.1**
- Bis(acetylacetato)chloroacetatomanganese(III) **12.6**
- Bis(acetylacetato)cobalt(II) **12.7**
- Bis(acetylacetato)copper(II) **12.8**
- Bis(acetylacetato)manganese(III) **12.9**
- Bis(acetylacetato)nickel(II) **12.10**
- Bis(acetylacetato)nickel(II) dihydrate **12.11**
- Bis(acetylacetato)trichloroacetatomanganese(III) **12.12**
- Bis(acetylacetato)zinc(II) **12.13**
- N,N*-Bis(3-aminopropyl)-1,4-butanediamine **8.11**
- Bis(5-bromo-2-hydroxybenzaldehydato)nickel(II) dihydrate **12.14**
- 2,5-Bis(4-bromophenyl)furan **5.16**
- Bis(2-butene-2,3-dithiolato)nickel(II) **12.15**
- Bis[*O*-butyl-3,5-di-(1,1-dimethylethyl)-4-hydroxybenzylphosphonato]nickel(II) **12.16**
- Bis[2-[(butylimino)methyl]-4-bromophenolato]nickel(II) **12.17**
- Bis[2-[(butylimino)methyl]-4-methoxyphenolato]nickel(II) **12.18**
- Bis[2-[(butylimino)methyl]phenolato]nickel(II) **12.19**
- Bis(4-*tert*-butylphenyl)nitroxide **15.26**
- 1,4-Bis(4-chlorophenyl)-1,3-cyclopentadiene **2.141**
- 2,5-Bis(4-chlorophenyl)furan **5.17**
- 1,1-Bis(cyclobutyl)-2-methyl-1-propene **2.332**
- 3,4-Bis[(cyclohexylamino)methylene]dihydro-2,5-thiophenedithione **13.169**
- Bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenethionato]cobalt(II) **12.20**
- Bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenethionato]nickel(II) **12.21**
- Bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenethionato]zinc(II) **12.23**
- Bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenonato]copper(II) **12.22**
- Bis[2-[(cyclohexylimino)methyl]phenolato]nickel(II) **12.24**
- 1,1-Bis(cyclopropyl)-2-methyl-1-propene **2.334**
- 1,1-Bis(cyclopropyl)-1-propene **2.333**
- Bis[1,2-di(2-bromophenyl)-1,2-ethenedithiolato]nickel(II) **12.25**
- Bis(dibutylthiocarbamato)cobalt(II) **12.26**
- Bis(dibutylthiocarbamato)copper(II) **12.27**
- Bis(dibutylthiocarbamato)nickel(II) **12.28**
- Bis(dibutylthiocarbamato)zinc(II) **12.29**
- Bis[*O,O'*-di-(4-*tert*-butylphenyl)phosphorodithiolato]cobalt(II) **12.30**
- Bis[*O,O'*-di-(4-*tert*-butylphenyl)phosphorodithiolato]copper(II) **12.31**
- Bis[*O,O'*-di-(4-*tert*-butylphenyl)phosphorodithiolato]nickel(II) **12.32**
- Bis[1,2-di(2-chlorophenyl)-1,2-ethenedithiolato]nickel(II) **12.33**
- Bis[1,2-di(4-chlorophenyl)-1,2-ethenedithiolato]nickel(II) **12.34**
- Bis(dicyclohexylphosphinodithiolato)nickel(II) **12.35**
- Bis(*O,O'*-dicyclohexylphosphorodithiolato)cobalt(II) **12.36**
- Bis[1,2-di(2,4-dichlorophenyl)-1,2-ethenedithiolato]nickel(II) **12.37**
- Bis[1,2-di(3,4-dichlorophenyl)-1,2-ethenedithiolato]nickel(II) **12.38**
- Bis[1,2-di(dimethylaminophenyl)-1,2-ethenedithiolato]nickel(II) **12.39**
- 1,2-Bis(diethylamino)acetylene **8.30**
- Bis(diethylthiocarbamato)nickel(II) **12.40**
- Bis[1,2-di(4-ethylphenyl)-1,2-ethenedithiolato]cobalt(II) **12.41**
- Bis[1,2-di(4-ethylphenyl)-1,2-ethenedithiolato]nickel(II) **12.42**
- Bis[1,2-di(4-ethylphenyl)-1,2-ethenedithiolato]platinum(II) **12.43**
- Bis(*O,O'*-diethylphosphorodithiolato)nickel(II) **12.44**
- Bis[2,3-dihydro-*N*-2-pyridinyl-3-(2-pyridinylimino)isoindol-1-amine]nickel(II) **12.45**
- Bis(diisopropylthiocarbamato)cobalt(II) **12.46**
- Bis(diisopropylthiocarbamato)copper(II) **12.47**
- Bis(diisopropylthiocarbamato)iron(III) **12.48**
- Bis(diisopropylthiocarbamato)manganese(II) **12.49**
- Bis(diisopropylthiocarbamato)nickel(II) **12.50**
- Bis(diisopropylthiocarbamato)zinc(II) **12.51**

- Bis[diisopropylidithiophosphato]nickel(II) **12.52**
 Bis(*O,O'*-diisopropylphosphorodithiolato)nickel(II) **12.52**
 Bis[1,2-di(4-methoxyphenyl)-1,2-ethanedithionato]nickelate(I), tetrabutylammonium salt **12.53**
 Bis[1,2-di(4-methoxyphenyl)-1,2-ethanedithiolato]nickel(II) **12.54**
 1,4-Bis(*N,N*-dimethylamino)benzene **9.31**
 3,7-Bis(dimethylamino)phenothiazinium **11.113**
N-[4-Bis[4-(dimethylamino)phenyl]methylene]2,5-cyclohexadien-1-ylidene]-*N*-methylmethanaminium chloride **11.44**
 Bis(dimethylidithiocarbamato)bismuth(II) **12.55**
 Bis(dimethylidithiocarbamato)nickel(II) **12.56**
 Bis(1,2-dimethyl-1,2-ethanedithiolato)nickel(II) **12.15**
 2,6-Bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)selenopyran-4-ylidene]-3-propenyl]pyrylium **11.157**
 2,6-Bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)selenopyran-4-ylidene]-3-propenyl]selenopyrylium **11.160**
 2,6-Bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)selenopyran-4-ylidene]-3-propenyl]thiopyrylium **11.175**
 2,6-Bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)telluropyran-4-ylidene]-3-propenyl]pyrylium **11.158**
 2,6-Bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)telluropyran-4-ylidene]-3-propenyl]telluropyrylium **11.161**
 2,6-Bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)thiopyran-4-ylidene]-3-propenyl]thiopyrylium **11.176**
 Bis[2-[(1,1-dimethylethylimino)methyl]phenolato]nickel(II) **12.57**
 Bis[4-(1,1-dimethylethyl)phenylsalicylato]nickel(II) **12.58**
 Bis[3,5-di(1-methylethyl)salicylato]cobalt(II) **12.59**
 Bis[3,5-di(1-methylethyl)salicylato]nickel(II) **12.60**
 Bis[*O,O'*-di(4-methylphenyl)phosphorodithiolato]cobalt(II) **12.61**
 Bis[*O,O'*-di(4-methylphenyl)phosphorodithiolato]copper(II) **12.62**
 Bis[*O,O'*-di(4-methylphenyl)phosphorodithiolato]nickel(II) **12.63**
 Bis(diphenyldithiocarbamato)nickel(II) **12.64**
 Bis(1,2-diphenyl-1,2-ethanedithionato)nickelate(I), tetrabutylammonium salt **12.65**
 Bis(1,2-diphenyl-1,2-ethanedithiolato)nickel(II) **12.66**
 Bis(*O,O'*-diphenylphosphorodithiolato)chromium(II) **12.67**
 Bis(*O,O'*-diphenylphosphorodithiolato)cobalt(II) **12.68**
 Bis(*O,O'*-diphenylphosphorodithiolato)copper(II) **12.69**
 Bis(*O,O'*-diphenylphosphorodithiolato)lead(II) **12.70**
 Bis(*O,O'*-diphenylphosphorodithiolato)nickel(II) **12.71**
 Bis(*O,O'*-diphenylphosphorodithiolato)zinc(II) **12.72**
 Bis(dithioacetylacetone)cobalt(II) **12.73**
 Bis(dithioacetylacetone)nickel(II) **12.74**
 Bis[1,2-di(trifluoromethyl)-1,2-ethenedithiolato]nickel(II) **12.85**
 Bis[1,2-di(4-trifluoromethylphenyl)-1,2-ethanedithionato]nickelate(I), tetrabutylammonium salt **12.75**
 Bis[1,2-di(4-trifluoromethylphenyl)-1,2-ethanedithiolato]nickel(II) **12.76**
 Bis[1,2-di(3,4,5-trimethoxyphenyl)-1,2-ethanedithionato]nickelate(I), tetrabutylammonium salt **12.77**
 Bis[1,2-di(3,4,5-trimethoxyphenyl)-1,2-ethanedithionato]nickelate(I), 4-[1,5,5-tri(4-diethylaminophenyl)pentadienyliidene]anilinium salt **12.78**
 Bis[1,2-di(3,4,5-trimethoxyphenyl)-1,2-ethanedithiolato]nickel(II) **12.79**
 Bis[2-[(dodecylimino)methyl]phenolato]cobalt(II) **12.80**
 Bis[2-[(dodecylimino)methyl]phenolato]copper(II) **12.81**
 Bis[2-[(dodecylimino)methyl]phenolato]nickel(II) **12.82**
 9,10-Bis(ethanesulfonate)anthracene **3.15, 12.143, 12.167, 12.169**
 2,5-Bis(ethoxycarbonyl)furan **5.60**
 Bis[*O*-ethyl-3,5-di-(1,1-dimethylethyl)-4-hydroxybenzylphosphonato]nickel(II) **12.83**
 Bis[5-ethyl-3-[(3-pyridinylimino)methyl]-2-thiophenethionato]cobalt(II) **12.84**
 1,4-Bis(4-fluorophenyl)-1,3-cyclopentadiene **2.142**
 Bis[1,1,1,4,4,4-hexafluoro-2-butene-2,3-dithiolato]nickel(II) **12.85**
 Bis[hydrotrois(1-pyrazolyl)borato]nickel(II) **12.86**
 Bis(2'-hydroxyacetophenone oximato)nickel(II) **12.87**
 Bis(2-hydroxybenzaldehydato)nickel(II) dihydrate **12.88**
 Bis(2-hydroxybenzaldehyde oximato)nickel(II) **12.89**
 Bis[2-hydroxybenzaldehyde phenylhydrazone]nickel(II) **12.90**
 Bis[2'-hydroxy-4'-*tert*-butyloctadecanophenone oximato]palladium(II) **12.91**
 7,12-Bis(1-hydroxyethyl)-3,8,13,17-tetramethylporphine-2,18-dipropanoic acid **7.72**
 Bis(2-hydroxy-5-methoxybenzaldehydato)nickel(II) dihydrate **12.92**
 Bis[2'-hydroxy-4'-methylacetophenone oximato]nickel(II) **12.93**
 Bis[2-hydroxy-5-methylbenzophenonato]nickel(II) **12.94**
 Bis[2'-hydroxy-4'-methyldodecanophenone oximato]nickel(II) **12.95**
 2,5-Bis(hydroxymethyl)furan **5.18, 10.2, 10.3, 10.23, 10.24, 10.26, 13.104, 13.106, 16.37, 16.64, 17.77**
 Bis[(2-hydroxy-5-methylphenyl)phenylmethanonato]nickel(II) **12.94**
 Bis[2-(iminomethyl)phenolato]nickel(II) **12.96**
 Bis(4-imino-2-pantanone)nickel(II) **12.97**

- 1,2-Bis(isopropylidene)-3,5-cyclohexadiene 2.104
 2,5-Bis(methoxycarbonyl)furan 5.61
 1,4-Bis(4-methoxyphenyl)-1,3-cyclopentadiene 2.143
 1,1-Bis(4-methoxyphenyl)ethylene 3.100
 2,5-Bis(4-methoxyphenyl)furan 5.19
 1,4-Bis(methylamino)anthraquinone 11.7
 Bis[4-methyl-1,2-benzenedithiolato]cobalt(II) tetrabutylammonium salt 12.99
 Bis[4-methyl-1,2-benzenedithiolato]nickel(II) tetrabutylammonium salt 12.100
 Bis[O-(1-methylethyl)carbonodithionato]nickel(II) 12.98
 Bis[2-[(1-methylethylimino)methyl]phenolato]nickel(II) 12.101
 Bis[N-methyl-7-(methylelimino)-1,3,5-cycloheptatrien-1-aminato]nickel(II) 12.102
 1,4-Bis(4-methylphenyl)-1,3-cyclopentadiene 2.144
 Bis(4-methylphenyldithiocarbamato)nickel(II) 12.103
 Bis[4-(1-methyl-1-phenylethyl)phenyl]nitroxide 15.27
 2,5-Bis(4-methylphenyl)furan 5.20
 Bis(4-methylphenyl)methanethione 13.161
 Bis[N-(4-methylphenyl)-7-[(4-methylphenyl)imino]-1,3,5-cycloheptatrien-1-aminato]nickel(II) 12.104
 Bis[2-[(1-methylpropylimino)methyl]phenolato]nickel(II) 12.105
 Bismuth(II) bis(dimethyldithiocarbamate) 12.55
 Bis(2,4-pentanedionato)cobalt(II) 12.7
 Bis(2,4-pentanedionato)copper(II) 12.8
 Bis(2,4-pentanedionato)zinc(II) 12.13
 Bis(2,4-pentanedithionato)cobalt 12.73
 Bis[2-pentene-2,4-dithiolato]nickel(II) 12.74
 1,4-Bis(N-phenylamino)benzene 9.29
 Bis[2-[(4-phenylamino)phenyl]imino]methyl]phenolato]nickel(II) 12.106
 Bis(phenyldithiocarbamato)nickel(II) 12.107
 Bis[2-[(phenylimino)methyl]phenolato]cobalt(II) 12.108
 Bis[2-[(phenylimino)methyl]phenolato]copper(II) 12.109
 Bis[2-[(phenylimino)methyl]phenolato]nickel(II) 12.110
 9,10-Bis(2-sulfonatoethyl)anthracene 3.15, 12.143, 12.167, 12.169
 Bis(2,2,6,6-tetramethyl-4-piperidinol-1-oxyl) 1,10-decanedioate 15.37
 Bis(2,2,6,6-tetramethyl-4-piperidinol-1-oxyl) 1,6-hexanedioate 15.38
 Bis(2,2,6,6-tetramethyl-4-piperidinol-1-oxyl) terephthalate 15.39
N,N'-Bis[4-(2,2,6,6-tetramethylpiperidinyl)]-1,6-hexanediamine 8.64
 Bis[2,2'-thiobis[O,O'-di(4-*tert*-butylphenyl)phosphorodithiolato]cadmium(II) 12.111
 Bis[2,2'-thiobis[O,O'-di(4-*tert*-butylphenyl)phosphorodithiolato]cobalt(II) 12.112
 Bis[2,2'-thiobis[O,O'-di(4-*tert*-butylphenyl)phosphorodithiolato]lead(II) 12.113
 Bis[2,2'-thiobis[O,O'-di(4-*tert*-butylphenyl)phosphorodithiolato]nickel(II) 12.114
 Bis[2,2'-thiobis[O,O'-di(4-*tert*-butylphenyl)phosphorodithiolato]zinc(II) 12.115
 Bis(2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)phenolato]nickel(II) 12.116
 Bis(tribenzylsiloxy)silicon 2,3-naphthalocyanine 7.17
 Bis(tribenzylsiloxy)silicon phthalocyanine 7.40
 Bis[3,4,6-trichloro-1,2-benzenedithiolato]nickelate(I), tetrabutylammonium salt 12.117
 2,3,2',3'-Bis(1,4,10,13-trideca-4,6,8,10-tetraen)tetrayl[4,6-diphenylpyrylium] perchlorate 11.31
 Bis(tri-*n*-hexyloxysiloxy)-2,3-naphthalocyaninatoaluminum 7.24
 Bis(tri-*n*-hexyloxysiloxy)-2,3-naphthalocyaninatogallium 7.25
 Bis(tri-*n*-hexyloxysiloxy)-2,3-naphthalocyaninatosilicon 7.18
 Bis(tri-*n*-hexyloxysiloxy)-2,3-naphthalocyaninatotin 7.19
 Bis(trihexylsiloxy)silicon 2,3-naphthalocyanine 7.18
 Bis(trihexylsiloxy)tin 2,3-naphthalocyanine 7.19
 Bis(triisobutylsiloxy)silicon 2,3-naphthalocyanine 7.20
 Bis(2,4,6-trimethylbenzenesulfonato)nickel(II) 12.118
 Bis(2,4,6-trimethylphenyl)diazomethane 11.19
 Bis(propylsiloxy)silicon phthalocyanine 7.41
 Bixin 2.177
 Blepharismin 16.59
 Boldine 8.20
 Bromide ion 14.4
 Bromobenzene 3.30
 Bromocarbonylbis(triphenylphosphine)iridium 12.119
 4-Bromo-2,6-di-*tert*-butylphenol 4.105
 1-Bromo-4-(dimethylamino)benzene 9.2
 4-Bromo-*N,N*-dimethylaniline 9.2
 4-Bromo-2,6-dimethylphenoxide ion 4.154
 2-Bromofuran 5.21
 4-Bromophenoxy ion 4.153
 1-(4-Bromophenylazo)-2-naphthol 11.94
 2-(4-Bromophenyl)furan 5.22
 3-(4-Bromophenyl)furan 5.23
 4-Bromophenyl methyl sulfide 13.7
p-Bromophenyl methyl sulfide 13.7
 1-(4-Bromophenyl)-4,4,8,8-tetramethyl-2,3,4,5,7,8,9,10-octahydronaphthalene[4,3,2-*m,n*]acridine-10-one 6.67
p-Bromothioanisole 13.7
 2-[7-(4-Bromo-1,3,3-trimethyl-2-indol-2-ylidene)-1,3,5-heptatrienyl]-1,3,3-trimethylindolinium, tetrafluoroborate 11.70
 Brucine 8.9
 Buckminsterfullerene 17.32
 1,3-Butadiene, 1,4-di(4,4'-dimethoxyphenyl)- 2.35

- 1,3-Butadiene, 1,4-di(1,1-dimethylethoxy)-, (*E,E*)- 2.36
 1,3-Butadiene, 1,4-di(1,1-dimethylethoxy)-, (*E,Z*)- 2.37
 1,3-Butadiene, 1,4-di(1,1-dimethylethoxy)-, (*Z,Z*)- 2.38
 1,3-Butadiene, 2,3-dimethyl- 2.39
 1,3-Butadiene, 1,4-diphenyl- 2.40, 2.35
 1,3-Butadiene, 2-ethyl- 2.41
 1,3-Butadiene, 2-methyl- 2.42
 1,3-Butadiene, 2-(phenylmethyl)- 2.43
 Butanamide, 2-(4-methylphenylazo)-3-oxo-*N*-phenyl- 11.60
 Butanamide, 3-oxo-*N*-phenyl-2-(phenylazo)- 11.62
 1-Butaneamine 8.12
 1,4-Butanediamine, *N*-(3-aminopropyl)- 8.10
 1,4-Butanediamine, *N,N'*-bis(3-aminopropyl)- 8.11
 Butanedioic acid, polymer with 4-hydroxy-2,2,6,6-tetramethylpiperidineethanol 8.67
 2,3-Butanediol, 1,4-dimercapto-, *erythro*- 13.49
 2,3-Butanediol, 1,4-dimercapto-, *threo*- 13.50
 1-Butanethiol 13.51
 Butanoic acid 17.6
 Butanoic acid, 3,3-dimethyl-2-oxo- 17.7
 Butanoic acid, 4-(methylthio)-2-[[[(phenylmethyl)carbonyl]amino]-, methyl ester 13.110
 1-Butanol 17.8
 1-Butanol, 4-(4-methylphenyl)thio- 13.52
tert-Butanol 17.62
 (*E*)-2-Butenal 2.44
trans-2-Butenal 2.44
 1-Butene, 2,3-dimethyl- 2.45
 2-Butene 2.46
 2-Butene, (*E*) 2.47, 2.48
 2-Butene, (*Z*) 2.48
 2-Butene, 2-anisyl-3-methyl- 3.118, 3.115, 3.123
 2-Butene, 2-cyclopropyl-, (*E*) 2.49
 2-Butene, 2-cyclopropyl-3-methyl- 2.50
 2-Butene, 2,3-dimethyl- 2.51, 2.38, 2.50, 2.52, 2.53, 2.56, 2.103, 2.117, 2.140, 2.168, 2.170, 2.184, 2.185, 2.186, 2.187, 2.188, 2.189, 2.190, 2.191, 2.192, 2.193, 2.194, 2.195, 2.196, 2.223, 2.225, 2.303, 2.310, 2.332, 2.334, 3.10, 3.11, 3.28, 3.66, 3.125, 4.3, 4.100, 5.34, 7.5, 7.65, 8.83, 9.1, 13.86, 13.145
 2-Butene, 2,3-dimethyl-, (*d*₁₂)- 2.54
 2-Butene, 2,3-dimethyl-*d*₆, (*E*)- 2.52
 2-Butene, 2,3-dimethyl-*d*₆, (*Z*)- 2.53
 2-Butene, 2,3-di(α -naphthyl)-, (*E*)- 3.84
 2-Butene, 2,3-di(α -naphthyl)-, (*Z*)- 3.84
 2-Butene, 2,3-di(β -naphthyl)-, (*E*)- 3.85
 2-Butene, 2,3-di(β -naphthyl)-, (*Z*)- 3.85
 2-Butene, 2,3-diphenyl-, (*E*)- 3.102, 3.103
 2-Butene, 2,3-diphenyl-, (*Z*)- 3.103
 2-Butene, 2-methoxy- 2.55
 2-Butene, 2-methyl- 2.56, 2.122, 2.161, 2.307, 2.309, 2.310, 3.110, 15.2, 15.13, 15.14, 15.15
 2-Butene, [(2-methyl-3-(phenylsulfinyl)]- 13.53
 2-Butene, 2-(phenylsulfinyl)- (*E*) 13.54
 2-Butene, 2-(phenylsulfinyl)- (*Z*) 13.55
 (*E*)-2-Butene 2.47, 2.48
 (*Z*)-2-Butene 2.48
 2-Butenoic acid, 2-methyl-, (*E*) 2.58
 3-Butenoic acid 2.59
 (*E*)-2-Butenoic acid 2.57
 2-Buten-1-ol, 3-methyl- 2.60
 1-Buten-3-one, 1-(2,6,6-trimethyl-1-cyclohexen-1-yl)- 2.61
 4-*tert*-Butoxy-2,6,6-tetramethylpiperidine 8.55
 Butyl alcohol 17.8
tert-Butyl alcohol 17.62
 Butylamine 8.12
 Butylamine, *N,N*-dibutyl- 8.13
 Butylamine, *N,N*-dimethyl- 8.14
tert-Butylamine 8.15
 Butylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II) 12.120
 Butylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]cobalt(II) 12.121
 Butylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) 12.122
 Butylated hydroxytoluene 4.102, 4.99, 4.116, 4.117, 4.118, 4.147
p-tert-Butylcatechol 4.8
 1-*tert*-Butylcyclobutene 2.97
 1-*tert*-Butylcyclohexene 2.97, 2.98, 2.102, 2.122, 2.138, 2.141
 1-*tert*-Butylcyclopentene 2.157
n-Butyldimethylamine 8.14
tert-Butyl ethyl sulfide 13.56
 2-*tert*-Butylfuran 5.24
 2-*tert*-Butyl-4-hydroxyanisole 4.122
n-Butyl mercaptan 13.51
tert-Butyl methyl ether 17.18
 Butyl methyl sulfide 13.57
 7-*tert*-Butyl-5-methyltocol 4.55
 8-*tert*-Butyl-5-methyltocol 4.56
 2-*tert*-Butylphenol 4.120
 4-*tert*-Butylphenol 4.121
 4-(*tert*-Butyl)phenoxide ion 4.163
N-tert-Butyl- α -phenylnitronite 15.46
 2-Butyl propyl sulfide 13.59
sec-Butyl propyl sulfide 13.59
tert-Butyl propyl sulfide 13.58
 1-*tert*-Butylpyrrole 6.61
 2-*tert*-Butylpyrrole 6.62
 3-*tert*-Butylpyrrole 6.63

<i>sec</i> -Butyl sulfide	13.73	9- <i>cis</i> - β -Carotene	2.64
<i>p-tert</i> -Butylthioanisole	13.20	15,15'- <i>cis</i> - β -Carotene	2.65
(Butylthio)benzene	13.8	15,15'-(<i>Z</i>)- β -Carotene	2.65
1-(Butylthio)-3-chlorobenzene	13.9	15- <i>cis</i> - β -Carotene	2.65
1-(Butylthio)-4-chlorobenzene	13.10	15(<i>Z</i>)- ψ -Carotene, 7,7',8,8',11,11',12,12'-octahydro-	2.74
1-(Butylthio)-4-fluorobenzene	13.11	α -Carotene	2.63
1-(Butylthio)-4-methoxybenzene	13.12	(<i>all-E</i>)- ψ -Carotene	2.250
1-(Butylthio)-4-methylbenzene	13.13	<i>all-trans</i> - ψ -Carotene	2.250
Butyric acid	17.6	β -Carotene	2.66, 2.183
C ₆₀	17.32	β,β -Carotene, 3'-(acetoxy)-6',7'-didehydro-5,6-epoxy-	
C ₇₀	17.33	5,5',6,6',7,8-hexahydro-3,3',5'-trihydroxy-8-oxo-, (3S,3'S,5R,5'R,6S,6'R)-	2.67
Cadmium(II) bis[2,2'-thiobis[O,O'-di(4- <i>tert</i> -butylphenyl)dithiophosphate]]	12.111	β,ψ -Carotene	2.76
Cadmium(II) protoporphyrin	7.75	ψ -Carotene, 3,4-didehydro-1,2,7',8'-tetrahydro-1-methoxy-	
Cadmium(II) tetraphenylchlorin	7.51	(<i>all-E</i>)-	2.68
Cadmium(II) tetraphenylporphyrin	7.60	ψ -Carotene, 3,4-didehydro-1,2,7',8'-tetrahydro-1-methoxy-	
Caffeic acid	4.68	2-oxo- (<i>all-E</i>)-	2.69
Caged hydrazine	8.22	ψ -Carotene, 7,8-dihydro- (<i>all-E</i>)-	2.70
Canthaxanthin	2.83	ψ -Carotene, dihydroxy-, (<i>all-E</i>)-	2.251
Caproic acid	17.38	ψ -Carotene, 3,3',4,4'-tetradehydro-1,1',2,2'-tetrahydro-1,1'-	
Captopril	13.130	dimethoxy- (<i>all-E</i>)-	2.71
Carbamic acid, [2-(indol-3-yl)ethyl]-, methyl ester	10.35	ψ -Carotene, 7,8,11,12-tetrahydro-	2.72
Carbazole, 9-acetyl-2,3,4,9-tetrahydro-	6.2, 6.3, 6.10, 6.14, 6.18, 6.19	ψ,ψ -Carotene, 7,7',8,8',11,12-hexahydro-, 15- <i>cis</i> -	2.73
Carbazole, 2,3,4,9-tetrahydro-9-methyl-	6.2	(<i>E</i>)- β,ε -Carotene, 3,3'-dihydroxy- (3R,3'R,6'R)-	2.75
Carbazole, 2,3,4,9-tetrahydro-9-methyl-6-nitro-	6.3	γ -Carotene	2.76
4-Carbethoxyaniline	9.9	ζ -Carotene	2.72
2,2'-Carbocyanine, 1,1'-diethyl-, chloride	11.41	Carotene analog, C-30	2.77
4,4'-Carbocyanine, 1,1'-diethyl-	11.42	Carotene analog, C-35	2.78
4,4'-Carbocyanine, 1,1'-diethyl-, toluenesulfonate	11.43	(3R,3'R,6'R)- β,ε -Carotene-3,3'-diol	2.75
3-Carbomethoxy-4,5-dihydro-2-methylfuran	5.58	<i>all-trans</i> - β,β -Carotene-4,4'-diol	2.81
3-Carbomethoxy-2-methyl-4,5-dihydrofuran	5.58	β,β -Carotene-3,3'-diol, (3R,3'R)-	2.79
Carbonate hydrolyase	10.7	β,β -Carotene-3,3'-diol, 5,6:5',6'-diepoxy-5,5',6,6'-	
Carbonic anhydrase	10.7	tetrahydro- (<i>all-E</i>)-	2.80
Carbon seventy-atom molecule	17.33	β,β -Carotene-4,4'-diol, <i>all-E</i> -	2.81
Carbon sixty-atom molecule	17.32	β -Carotene-4,4'-dione, 3,3'-dihydroxy-	2.82
Carbonylbis(triphenylphosphine)iridium(I) chloride	12.123	β,β -Carotene-4,4'-dione	2.83
Carbonylbis(triphenylphosphine)rhodium(I) chloride	12.124	κ,κ -Carotene-6,6'-dione, 2,2'-dihydroxy- (2S,2'S,5R,5'R)	
Carbonylchlorobis(triphenylphosphine)iridium	12.123	2.84	
Carbonylchlorobis(triphenylphosphine)rhodium	12.124	β -apo-8-Carotenoic acid, ethyl ester	2.85
Carbonyliodobis(triphenylphosphine)iridium	12.125	β -apo-8'-Carotenol	2.86
[1-Carboxy-2-[2-mercaptopimidazol- 4(5)yl]ethyl]trimethylammonium ion	13.104	β,β -Caroten-3-ol	2.87
[1-Carboxy-2-[2-mercaptopimidazol- 4(5)yl]ethyl]trimethylammonium ion(1-)	13.105	β,β -Caroten-4-one	2.88
2-Carene	2.26	Carvomenthene	2.125
3-Carene	2.28	Caryophyllene	2.34
4-Carene	2.27	(-)Caryophyllene	2.34
Carnosine	10.24	β -Caryophyllene	2.34
L-Carnosine	10.24	γ -Caryophyllene	2.34
β -apo-8'-Carotenal	2.62	Catechin	4.67
		(+)-(2R,3S)-Catechin	4.67
		Catechol	4.4
		Catechol, 3,5-di- <i>tert</i> -butyl-	4.7
		CBZ-L-Methionine methyl ester	13.110

- C₃₀-Capsorubin **2.205**
C₃₄-Capsorubin **2.231**
C-30 Carotene **2.77**
C-35 Carotene **2.78**
C₃₀-Carotene analog **2.77**
C₃₅-Carotene analog **2.78**
C₃₀-Epiisocapsorubin **2.206**
C₃₄-Epiisocapsorubin **2.232**
C₄₀-Epiisocapsorubin **2.84**
Cercosporin **16.55**
Chalcogenopyrylium dye 1a **11.159**
Chimasorb 944 **8.16**
Chloride ion **14.5**
Chloroaluminum(III) sulfophthalocyanine **7.43**
2-Chloroanisole **3.34**
1-Chloroanthracene **3.4**
9-Chloroanthracene **3.5**
Chlorobenzene **3.31**
Chloro-1,4-benzenediol **4.20**
1-Chloro-3-(butylthio)benzene **13.9**
1-Chloro-4-(butylthio)benzene **13.10**
5-Chloro-2-[2-[3-[(5-chloro-3-ethyl- 2-
benzothiazolylidene)ethylidene]-2-(diphenylamino)-
1-cyclopenten-1-yl]ethenyl]-3-ethylbenzothiazolium,
perchlorate **11.33**
4-Chloro-1,3-dihydroxybenzene **4.11**
2-Chloro-1,4-dimethoxybenzene **3.32**
5-Chloro-1,3-dimethoxybenzene **3.33**
1-Chloro-3-(dimethylamino)benzene **9.3**
2-Chloro-10-dimethylaminopropylphenothiazine **13.122**
3-Chloro-*N,N*-dimethylaniline **9.3**
4-Chloro-*N,N*-dimethylaniline **9.4**
2-Chloro-*N,N*-dimethylphenothiazine-10-propanamine
13.122
1-Chloro-9,10-diphenylanthracene **3.6**
Chloroform **17.9**
Chloroform-*d* **17.10**
Chlorohydroquinone **4.20**
3-Chloro-5-methoxyphenol **4.111**
2-Chlorophenol **4.106**
3-Chlorophenol **4.107**
4-Chlorophenol **4.108**
2-Chlorophenoxyde ion **4.155**
3-Chlorophenoxyde ion **4.156**
4-Chlorophenoxyde ion **4.157**
2-(3-Chlorophenoxy)methylidene)adamantane **2.350**
2-(4-Chlorophenoxy)methylidene)adamantane **2.351**
1-(4-Chlorophenylazo)-2-naphthol **11.95**
4-(4-Chlorophenylazo)-1-naphthol **11.87**
7-Chloro-2-phenyl-1,2-benzisosenazol-3-one **13.36**
3-(4-Chlorophenyl)-1,5-diphenyl-2-pyrazoline **11.121**
5-(4-Chlorophenyl)-1,3-diphenyl-2-pyrazoline **11.121**
2-(4-Chlorophenyl)furan **5.25**
1-(4-Chlorophenyl)-4-(4-methoxyphenyl)-1,3-
cyclopentadiene **2.145**
3-Chlorophenyl methyl sulfide **13.14**
4-Chlorophenyl methyl sulfide **13.15**
m-Chlorophenyl methyl sulfide **13.14**
1-(4-Chlorophenyl)-4-phenyl-1,3-cyclopentadiene **2.146**
4-Chlorophenyl phenyl sulfide **13.16**
1-Chloro-4-(phenylthio)benzene **13.16**
3-(3-Chlorophenyl)thio-1-propanol **13.132**
3-(4-Chlorophenyl)thio-1-propanol **13.133**
Chlorophyll *a* **7.11**
Chlorophyll *b* **7.12**
4'-Chloropivalothiophenone **13.127**
4-Chlororesorcinol **4.11**
Chloro(5,10,15,20-tetraphenylporphinato)iron(III) **7.49**
m-Chlorothioanisole **13.14**
4-Chlorothiobenzamide **13.153**
4-Chlorothiobenzophenone **13.158**
2-[7-(4-Chloro-1,3,3-trimethyl-2-indol-2-
yldene)-1,3,5-heptatrienyl]-1,3,3-trimethylindolinium
tetrafluoroborate **11.71**
m-Chloro- α,β,β -trimethylstyrene **3.109**
p-Chloro- α,β,β -trimethylstyrene **3.110, 3.121, 3.122**
Chlorpromazine **13.122**
3,5-Cholestadiene **2.89**
 $\Delta^{3,5}$ -Cholestadiene **2.89**
4-Cholesten-3-one, 4-methyl- **2.90**
Cholesterol **16.14, 2.213, 2.265, 2.269, 2.272**
Cholestryl benzoate **16.15**
6-Chromanol, 2,8-dimethyl-2-(4,8,12-trimethyltridecyl)
4.46
6-Chromanol, 2,5,8-trimethyl-2-(4,8,12-trimethyltridecyl)
4.52
6-Chromanol, 2,7,8-trimethyl-2-(4,8,12-trimethyltridecyl)
4.53
Chromium(II) bis(*O,O'*-diphenyldithiophosphate) **12.67**
Chromium(III) tris(acetylacetone) **12.172**
Chrysin **4.73**
C.I. 12055 **11.104, 11.94, 11.95, 11.96, 11.97, 11.98, 11.99,**
11.100, 11.102, 11.103
C.I. 12120 **11.101**
C.I. 42555 **11.44**
C.I. 60700 **11.6**
C.I. 60710 **11.5**
C.I. 61500 **11.7**
C.I. 62015 **11.13**
C.I. 64500 **11.14**
Cinnamic acid, 3,4-dihydroxy- **4.68**
(*E*)-Cinnamic acid **2.91**
Cinnamic acid, 2-hydroxy- **4.69**

- Cinnamic acid, 4-hydroxy-3-methoxy- 4.70
trans-Cinnamic acid 2.91
 C_{30} -Isocapsorubin 2.207
Citronellol 2.294
Cleland's Reagent 13.50
Cobalt(II), bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenethionato]- 12.20
Cobalt(II), bis[5-ethyl-3-[(3-pyridinylimino)methyl]-2-thiophenethionato]- 12.84
Cobalt, bis(cyano)[tetramethyl-12,13-didehydro-2,18-bis(2-methoxy-2-oxoethyl)-3,5,8,8,13,15,18,19-octamethyl-3,7,12,17-corrin tetrapropanoato]- 16.16
Cobalt, [[2,2'-(1,2-ethanediyl)bis(nitrilomethylidyne)]bis(phenolato)](2-*N,N',O,O'*-) 12.139
Cobalt(II) acetate 12.126
Cobalt(II) bis(acetylacetone) 12.7
Cobalt(II) bis(dibutylthiocarbamate) 12.26
Cobalt(II) bis[*O,O'*-di(4-*tert*-butylphenyl)dithiophosphate] 12.30
Cobalt(II) bis[*O,O'*-dicyclohexylthiophosphate] 12.36
Cobalt(II) bis(*O,O'*-dicyclohexylphosphorodithiolato)- 12.36
Cobalt(II) bis[1,2-di(4-ethylphenyl)-1,2-ethenedithiolato]- 12.41
Cobalt(II) bis(diisopropylthiocarbamate) 12.46
Cobalt(II) bis[3,5-diisopropylsalicylate] 12.59
Cobalt(II) bis[*O,O'*-di(4-methylphenyl)dithiophosphate] 12.61
Cobalt(II) bis(*O,O'*-diphenylthiophosphate) 12.68
Cobalt(II) bis(dithioacetylacetone) 12.73
Cobalt(II) bis[2-(*N*-dodecylformimidoyl)phenol] 12.80
Cobalt(II) bis[2-(*N*-phenylformimidoyl)phenol] 12.108
Cobalt(II) bis[2,2'-thiobis[*O,O'*-di(4-*tert*-butylphenyl)dithiophosphate]] 12.112
Cobalt(II) bis[μ -toluene-3,4-dithiolato]TBA₂ 12.99
Cobalt(II) butylamine[2,2'-thiobis(4-*tert*-octyl)phenolate] 12.121
Cobalt(II) chloride hexahydrate 12.127
Cobalt(II) 2,2'-[ethylenebis(nitrilomethylidyne)]diphenol 12.139
Cobalt(II) hydroxybis(2,3-butanedione dioximato)triethylamine 12.147
Cobalt(II) ion 12.128
Cobalt(II) mesoporphyrin 7.82
Cobalt(II) mesoporphyrin IX, dimethyl ester 7.86
Cobalt(II) 2,2'-methylenebis[*O,O'*-di(4-*tert*-butylphenyl)dithiophosphate] 12.156
Cobaltous acetate 12.126
Cobalt(II) protoporphyrin 7.76
Cobalt(II) 5,10,15,20-tetraphenylporphyrin 7.62
Cobalt(III) tris(acetylacetone) 12.173
Cobalt(II) tris(2,2'-bipyridine) 12.176
Cobalt(II) tris(1,10-phenanthroline) 12.178
Cobrynic acid, bis(cyano)-7-de(carboxymethyl)-7,8-didehydro-, hexamethyl ester 16.16
Coenzyme A 16.17
Copper(II), bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenonato]- 12.22
Copper(II), [[3,3]-[1,2-ethanediyl]bis(nitrilomethylidyne)]bis[5-ethyl-2-thiophenethionato]]- 12.138
Copper(II) bis(acetylacetone) 12.8
Copper(II) bis(dibutylthiocarbamate) 12.27
Copper(II) bis[*O,O'*-di(4-*tert*-butylphenyl)dithiophosphate] 12.31
Copper(II) bis(diisopropylthiocarbamate) 12.47
Copper(II) bis[*O,O'*-di(4-methylphenyl)dithiophosphate] 12.62
Copper(II) bis(*O,O'*-diphenylthiophosphate) 12.69
Copper(II) bis[2-(*N*-dodecylformimidoyl)phenol] 12.81
Copper(II) bis[2-(*N*-phenylformimidoyl)phenol] 12.109
Copper(II) ion 12.129
Copper(II) protoporphyrin 7.77
Copper(II) tetracarboxyphthalocyanine 7.45
Copper(II) tetra(4-iodophenyl)porphyrin 7.55
Copper(II) tetra(4-methoxyphenyl)porphyrin 7.56
Copper(II) tetraphenylchlorin 7.52
Copper(II) tetraphenylporphyrin 7.63
Copper(II) 5,10,15,20-tetraphenylporphyrin 7.63
Coumarin 1 5.7
Coumarin 102 8.4
Coumarin 460 5.7
Coumarin 47 5.7
Coumarin, 7-(diethylamino)- 5.6
Coumarin, 7-(diethylamino)-4-methyl- 5.7
Coumarin, 3,4,5,6,7,8-hexahydro- 2.92
Coumarin LD-490 8.3
C₁₈-Polyene-tetrone 2.234
C₂₈-Polyene-tetrone 2.348
C₁₈-Polyene-tetrone-diacetal 2.233
C₂₂-Polyene-tetrone-diacetal 2.211
C₂₈-Polyene-tetrone-diacetal 2.347
p-Cresol 4.135
Crocetin 2.93
 α -Crocetin 2.93
Crocin 2.173
Crocin 3 2.178
(E)-Crotonaldehyde 2.44
(E)-Crotonic acid 2.57
Cryptoxanthin 2.87
Crystal Violet 11.44
Curcumin 4.77
Cu,Zn-Erythrocuprein 10.14
2,2'-Cyanine, 1,1'-diethyl-, iodide 11.45

- 4-Cyano-*N,N*-dimethylaniline **9.22**
 4-[*N*-(2-Cyanoethyl)-*N*-ethylamino]azobenzene **11.51**
 4-[*N*-(2-Cyanoethyl)-*N*-ethylamino]-4'-methoxyazobenzene **11.50**
 2-Cyanofuran **5.63**
 Cyanohemoglobin **7.13**
 4-Cyanophenol **4.112**
 4-Cyanophenoxy ion **4.158**
 2-(4-Cyanophenyl)-3-(4-dimethylaminophenyl)-1,4-dioxene **2.184**
 2-(4-Cyanophenyl)furan **5.26**
m-Cyano- α,β,β -trimethylstyrene **3.111**
p-Cyano- α,β,β -trimethylstyrene **3.112**
 Cyclobutane, (1-cyclopropylethylidene)- **2.94**
 Cyclobutane, (cyclopropylmethylene)- **2.95**
 Cyclobutane, (dicyclopropylmethylidene)- **2.96**, **2.49**, **2.50**, **2.94**, **2.95**, **2.151**, **2.168**, **2.169**, **2.333**, **2.334**, **2.335**, **3.113**
 1,3-Cyclobutanedithione, 2,2,4,4-tetramethyl- **13.60**
 Cyclobutanethione, 2,2,4,4-tetramethyl- **13.61**
 Cyclobutanone, 2,2,4,4-tetramethyl-3-thioxo- **13.62**
 Cyclobutendiylium, 1,3-bis[4-(diethylamino)-2-hydroxyphenyl]-2,4-dihydroxy- **11.46**, **12.66**
 Cyclobutendiylium, 1,3-bis[4-(dimethylamino)-2-methylphenyl]-2,4-dihydroxy- **11.47**
 Cyclobutendiylium, 1,3-bis[4-(*N*-methyl-*N*-octadecylamino)-2-methylphenyl]-2,4-dihydroxy- **11.48**
 Cyclobutene, 1-*tert*-butyl- **2.97**
 Cyclobutene, 1-methyl- **2.98**
 2-(Cyclododecylidene)ethanol **2.215**
 1,3-Cycloheptadiene **2.99**
 Cycloheptane, ethylidene- **2.100**
 Cycloheptane, methylene- **2.101**
 Cycloheptene, 1-methyl- **2.102**
 1,3-Cyclohexadiene **2.103**
 1,3-Cyclohexadiene, 5,6-bis(1-methylethylidene)- **2.104**
 1,3-Cyclohexadiene, 1-methyl-4-(1-methylethyl)- **2.105**
 1,3-Cyclohexadiene, 2-methyl-5-(1-methylethyl)- **2.106**
 1,4-Cyclohexadiene, 1-methyl- **2.107**
 2,4-Cyclohexadien-1-one, 6,6-dimethyl- **2.108**
 2,5-Cyclohexadien-1-one, 4-[[4-(diethylamino)-2,5-dimethylphenyl]imino]-3,5-dimethyl- **11.49**
 2,5-Cyclohexadien-4-one, 4,4'-(1,2-ethanediylidene)bis[2,6-bis(1,1-dimethylethyl)- **17.12**
 2,5-Cyclohexadien-4-one, 1,2-ethanediylidenebis[3-[2-hydroxy-5-methyl-3-(1,1-dimethylethyl)benzyl]-5-(1,1-dimethylethyl)]- **17.11**
 Cyclohexanamine, *N,N'*-[dithiobis[(5-ethyl-2,3-thiophenediyl)methylidyne]]bis- **13.63**
 Cyclohexanamine, *N*-[[5-methyl-2-(methylthio)-3-thienyl]methylene]- **13.64**
 Cyclohexane **17.13**
- Cyclohexane, cyclohexylidene- **2.109**
 Cyclohexane-*d*₁₂ **17.14**
 Cyclohexane, 1,4-dichloro-1,4-dinitroso-, (*E*)- **15.16**
 Cyclohexane, 1,4-dichloro-1,4-dinitroso-, (*Z*)- **15.17**
 Cyclohexane, ethylidene- **2.110**
 Cyclohexane, (hydroxymethylene)-, acetate **2.111**
 Cyclohexane, methylene- **2.112**, **2.101**, **2.153**
 1,3-Cyclohexanedione, 5,5-dimethyl- **17.15**
 Cyclohexanol, 2-(cyclohexylidene)- **2.113**
 Cyclohexanone, 2-cyclohexylidene- **2.114**
 Cyclohexanone, 5-methyl-2-(1-methylethenyl)- **2.115**
 Cyclohexene **2.116**, **2.45**, **2.123**, **2.263**
 Cyclohexene, 1-*tert*-butyl- **2.97**, **2.98**, **2.102**, **2.122**, **2.138**, **2.157**, **2.161**
 Cyclohexene, 1,2-dimethyl- **2.117**, **2.161**
 Cyclohexene, 1,3-dimethyl- **2.118**
 Cyclohexene, 1,4-dimethyl- **2.119**
 Cyclohexene, 2,3-dimethyl- **2.120**, **2.156**
 Cyclohexene, 4-(1-hydroxy-1-methylethyl)-1-methyl- **2.121**
 Cyclohexene, 1-methyl **2.122**, **2.11**, **2.21**, **2.120**, **2.153**, **2.161**, **2.236**, **2.237**, **2.238**, **2.239**, **2.243**, **2.309**, **2.326**, **2.327**
 Cyclohexene, 4-methyl- **2.123**
 Cyclohexene, 1-methyl-4-(1-methylethenyl)-, (*R*)- **2.124**, **13.8**, **13.52**, **13.97**, **13.119**, **13.136**
 Cyclohexene, 1-methyl-4-(1-methylethyl)- **2.125**
 2-Cyclohexene-1-thione, 3,5,5-trimethyl- **13.65**
 Cyclohexen-1-ol, 6,6-dimethyl-, acetate **2.126**
 2-Cyclohexen-1-one, 2-ethyl-3-methyl- **2.127**
 2-Cyclohexen-1-one, 2-hydroxy-3-methyl- **2.128**
 Cyclohexylamine **8.17**
 Cyclohexylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II) **12.130**
 Cyclohexylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) **12.131**
 4-Cyclohexylaminodiphenylamine **9.28**
 3-[(Cyclohexylamino)methylene]-5-ethyl-2-thiophenethione **13.170**
 1-Cyclohexylamino-4-phenylaminobenzene **9.28**
 (Cyclohexylidene)cyclohexane **2.109**
 2-(Cyclohexylidene)cyclohexanol **2.113**
 2-(Cyclohexylidene)cyclohexanone **2.114**
 2-(Cyclohexylidene)ethanol **2.216**
 Cyclohexylidenemethyl acetate **2.255**
N-Cyclohexylpiperidine **8.56**
 1,3-Cyclooctadiene **2.129**
 1,4-Cyclooctadiene, 6-hydroperoxy- **2.130**
 1,4-Cyclooctadiene, 6-methoxy- **2.131**
 1,4-Cyclooctadiene, 6-(trimethylsiloxy)- **2.132**
 1,5-Cyclooctadiene **2.133**, **2.137**
 2,5-Cyclooctadien-1-ol **2.134**
 Cyclooctane, ethylidene- **2.135**

- Cyclooctatetraene **2.136**
 1,3,5,7-Cyclooctatetraene **2.136**
 Cyclooctatetraene dibromide **2.32**
 Cyclooctene **2.137**
 Cyclooctene, 1-methyl- **2.138**
 2-(Cyclooctylidene)ethanol **2.217**
 Cyclooctyne **2.139**
 Cyclopenta[*b*][1]benzopyran, 2-phenyl- **5.8**
 Cyclopentadiene **2.140**
 1,3-Cyclopentadiene, 1,4-bis(4-chlorophenyl)- **2.141**
 1,3-Cyclopentadiene, 1,4-bis(4-fluorophenyl)- **2.142**
 1,3-Cyclopentadiene, 1,4-bis(4-methoxyphenyl)- **2.143**
 1,3-Cyclopentadiene, 1,4-bis(4-methylphenyl)- **2.144**
 1,3-Cyclopentadiene, 1-(4-chlorophenyl)-4-(4-methoxyphenyl)- **2.145**
 1,3-Cyclopentadiene, 1-(4-chlorophenyl)-4-phenyl- **2.146**
 1,3-Cyclopentadiene, 1,4-diphenyl- **2.147**
 1,3-Cyclopentadiene, 1-(4-methoxyphenyl)-4-phenyl- **2.148**
 1,3-Cyclopentadiene, 5-(1-methylethylidene)- **2.149**
 1,3-Cyclopentadiene, 5-(1-methylethylidene)-, endoperoxide **2.150**
 Cyclopentadienone, tetraphenyl- **3.60**
 Cyclopentane, (dicyclopropylmethylidene)- **2.151**
 Cyclopentane, ethylidene- **2.152**
 Cyclopentane, methylene- **2.153**
 Cyclopentanone, 2-cyclopentylidene- **2.154**
 Cyclopentanone, 2-(1-methylethenyl)- **2.155**
 Cyclopenta[*b*]quinoline, 4-methyl-1,2-diphenyl- **6.4**
 Cyclopentene **2.156**, **2.244**
 Cyclopentene, 1-*tert*-butyl- **2.157**
 Cyclopentene, 1,2-dimethyl- **2.158**
 Cyclopentene, 1,5-dimethyl- **2.159**
 Cyclopentene, 1-methoxy- **2.160**
 Cyclopentene, 1-methyl- **2.161**, **2.22**, **2.122**
 2-Cyclopentene-1-thione, 3-(4-methoxyphenyl)- **13.66**
 2-Cyclopenten-1-one, 2-acetoxy-3-methyl- **2.163**
 2-Cyclopenten-1-one, 3-acetoxy-2-methyl- **2.164**
 2-Cyclopenten-1-one, 2,3-dimethyl- **2.165**
 2-Cyclopenten-1-one, 2-ethyl-3-methyl- **2.166**
 2-Cyclopenten-1-one, 3-methoxy-2-methyl- **2.167**
 2-Cyclopenten-1-one, 3-methyl-2-pentyl- **2.162**
 2-Cyclopentylidenecyclopentanone **2.154**
 2-(Cyclopentylidene)ethanol **2.218**
 Cyclopropane, (cyclopropylcyclopropylidenemethyl)- **2.168**
 Cyclopropane, (dicyclopropylmethylidene)- **2.168**
 Cyclopropane, 1,1-di-1-propenyl- **2.333**, **2.334**
 Cyclopropane, 1,1',1''-(1-ethenyl-2-ylidene)tris- **2.169**
 Cyclopropane, (1-methylethylidene)- **2.170**
 Cyclopropane, 1-methyl-1-propenyl- (*E*) **2.49**
 Cyclopropane, (2-methyl-1-propenyl)- **2.335**
 Cyclopropanecarboxylic acid, 2,2-dimethyl-3-(2-methyl-1-propenyl), ester with 5-(phenylmethyl)-3-furanmethanol **5.76**
trans-2-Cyclopropyl-2-butene **2.49**
 (1-Cyclopropylethylidene)cyclobutane **2.94**
 2-(Cyclopropylidene)propane **2.170**
 2-Cyclopropyl-3-methyl-2-butene **2.50**
 (Cyclopropylmethylene)cyclobutane **2.95**
 1-Cyclopropyl-2-methyl-1-phenylpropene **3.113**
 1-Cyclopropyl-2-methylpropene **2.335**
 α -Cyclopropyl- β,β -trimethylstyrene **3.113**
 Cysteamine **13.93**
 Cysteine **13.67**, **13.50**, **13.69**
 L-Cysteine, *N*-acetyl- **13.69**
 L-Cysteine, *N*-acetyl-, negative ion **13.70**
 Cysteine, negative ion **13.68**
 5-S-Cysteinyl-dopa **4.71**
 Cytidine **16.18**
 Cytidine monophosphate **16.19**
 Cytidine 5'-monophosphate **16.19**
 5'-Cytidylic acid **16.19**
 Cytochrome a **10.10**
 Cytochrome a₃ **10.10**
 Cytochrome aa₃ **10.10**
 Cytochrome b **10.8**
 Cytochrome b₂ **10.8**
 Cytochrome C (ferro) **10.9**
 Cytochrome C oxidase **10.10**
 Cytosine **16.20**
 Daunomycin **4.79**
 Daunorubicin **4.79**
 1,6-Decadiene, 2,6,9-trimethyl-, (*E*)- **2.171**
 1,6-Decadiene, 2,6,9-trimethyl-, (*Z*)- **2.172**
 Decanedioic acid, bis(2,2,6,6-tetramethylpiperidin-4-yl)ester **8.78**
 Dccapreno- β -carotene **2.230**
 7-Dehydroandrosterone-3-acetate **16.5**
 Dehydrogenase, 3-hydroxybutyrate **10.11**
 Dehydrogenase, nicotinamide adenine dinucleotide (reduced) **10.12**
 Dehydrogenase, succinate **10.13**
 16-Dehydropregnolone-3-acetate **16.61**
 Dehydrostobadine **6.58**
 Deoxyadenosine monophosphate **16.21**
 2'-Deoxyadenosine 5'-monophosphate **16.21**
 Deoxyadenylic acid **16.21**
 2'-Deoxy-5'-adenylic acid **16.21**
 Deoxycytidine monophosphate **16.22**
 2'-Deoxycytidine-5'-monophosphate **16.22**
 Deoxycytidylic acid **16.22**

- 2'-Deoxy-5'-cytidylic acid **16.22**
 2'-Deoxyguanosine **16.37**
 2'-Deoxyguanosine 5'-monophosphate **16.23**
 Deoxyguanylic acid **16.23**
 Deoxyribonucleic acid **16.24**
 3'-Deoxythymidine 5'-monophosphate **16.25**
 Deoxythymidine 5'-monophosphate **16.25**
 Deuterium peroxide **14.6**
 Deuteroporphyrin, dimethyl ester **7.94**
 5-Deutero-1,3,5-triphenyl-2-pyrazoline **11.123**
 4,4'-Diacetamidostilbene-2,2'-disulfonate ion **3.106**
 2,3-Di(4-acetaminophenyl)-1,4-dioxene **2.185**
 3,5-Diacetyl-1,4-dihydro-2,6-dimethylpyridine **6.42**
 3,5-Diacetyl-1,4-dihdropyridine **6.41**
 3,6-Diaminoacridine **11.1**
 1,2-Diaminoanthraquinone **11.3, 11.4, 11.5, 11.8, 11.9, 11.10, 11.11, 11.13, 11.14**
 1,4-Diaminoanthraquinone **11.8**
 1,5-Diaminoanthraquinone **11.9**
 1,8-Diaminoanthraquinone **11.10**
 2,6-Diaminoanthraquinone **11.11**
 1,2-Diaminobenzene **9.25**
 4,4'-Diaminobiphenyl **9.23**
 1,5-Diaminobromo-4,8-dihydroxyanthraquinone **11.12**
 Diaminocercosporin **16.56**
 Diaminodiphosphopyridine nucleotide, reduced **16.42**
 1,4-Diamino-2-methoxyanthraquinone **11.13**
 3,6-Diamino-10-methylacridinium **11.2**
 3,7-Diaminophenotheniazinium **11.114**
 4,4'-Diaminostilbene-2,2'-disulfonate ion **3.107**
 2,5-Di-*tert*-amylhydroquinone **4.16**
 8,8'-Diapo- ψ -carotendioic acid, bis(6-*O*- β -D-glucopyranosyl- β -D-glucopyranosyl) ester **2.173**
 6,6'-Diapo- ψ , ψ -carotenedioic acid **2.174**
 6,6'-Diapo- ψ , ψ -carotenedioic acid, dimethyl ester **2.175**
 6,6'-Diapo- ψ , ψ -carotenedioic acid, dimethyl ester, 9-*cis*- **2.176**
 6,6'-Diapo- ψ , ψ -carotenedioic acid, monomethyl ester, 9-*cis*- **2.177**
 8,8'-Diapocarotenedioic acid **2.93**
 8,8'-Diapocarotenedioic acid, mono(6-*O*- β -D-glucopyranosyl- β -D-glucopyranosyl) ester **2.178**
 9-*cis*-6,6'-Diapo- ψ , ψ -carotenedioic acid **2.179**
 Diaquabis(acetylacetato)manganese(III) ion **12.9**
 2,3-Diazabicyclo[2.2.1]heptane, 2,3-dimethyl- **8.18**
 1,4-Diazabicyclo[2.2.2]octane **8.19**
 2,3-Diazabicyclo[2.2.2]oct-2-ene, 1,4-dichloro-, 2,3-dioxide **15.18**
 Diazene, 1-[4-*N*-(2-cyanoethyl)-*N*-ethylaminophenyl]-2-(4-methoxyphenyl)- **11.50**
 Diazene, 1-[4-*N*-(2-cyanoethyl)-*N*-ethylaminophenyl]-2-phenyl- **11.51**
 Diazene, 1-(4-diethylaminophenyl)-2-(2,4-dinitrophenyl) **11.52**
 Diazene, 1-(4-diethylaminophenyl)-2-(4-methoxyphenyl) **11.53**
 Diazene, 1-(4-diethylaminophenyl)-2-(3-nitrophenyl)- **11.54**
 Diazene, 1-(4-diethylaminophenyl)-2-(4-nitrophenyl)- **11.55**
 Diazene, 1-(4-diethylaminophenyl)-2-phenyl- **11.56, 11.90, 11.102**
 Diazene, diphenyl- **11.57**
 Diazene, 1-[4-*N*-ethyl-*N*-(2-hydroxyethyl)aminophenyl]- (4-methoxyphenyl)- **11.58**
 Diazene, 1-(4-methoxy-1-naphthyl)-2-phenyl- **11.59**
 Diazene, 1-(4-methylphenyl)-2-[1-(phenylaminocarbonyl)-2-oxopropyl]- **11.60**
 Diazene, 1-(4-nitrophenyl)-2-phenyl- **11.61**
 Diazene, 1-phenyl-2-[1-(phenylaminocarbonyl)-2-oxopropyl]- **11.62**
 Diazobis(2,4,6-trimethylphenyl)methane **11.19**
 Diazo(4-bromophenyl)phenylmethane **11.15**
 Diazodi(4-chlorophenyl)methane **11.17**
 Diazodi(4-methylphenyl)methane **11.18**
 Diazodiphenylmethane **11.16, 11.15, 11.17, 11.18, 11.20**
 9-Diazofluorene **11.63**
 Diazo(4-methoxyphenyl)phenylmethane **11.20**
 1,2,5,6-Dibenzanthracene **3.61**
 Dibenz[a,h]anthracene **3.61**
 Dibenzo[3,4:5,6][2]benzopyrano[7,8,1-*mna*]xanthene **3.20**
 Dibenzo[*fg,uv*]naphtho[2,3-*c*]pentaphene-9,19-dione, 1,2,3,4,11,12,13,14-octahydro- **3.62**
 Dibenzo[*a,j*]perylene **3.63**
 Dibenzo[*a,o*]perylene **3.67**
 Dibenzo[*a,o*]perylene, 7,16-diphenyl- **3.64**
 Dibenzo[*aj*]perylene-8,16-dione **3.68**
 1,2,7,8-Dibenzoperylene-3,9-quinone **3.68**
 Dibenzo[*de,g*]quinoline-2,9-diol, 5,6,6a,7-tetrahydro-1,10-dimethoxy-6-methyl- **8.20**
 Dibenzo[*de,g*]quinoline-2,9-diol, 5,6,6a,7-tetrahydro-1,2,9,10-tetramethoxy-6-methyl- **8.21**
 3,5-Dibenzoyl-1,4-dihydro-2,6-dimethylpyridine **6.43**
 4-(Dibenzoylmethylene)amino-3,*N,N*-triethylaniline **11.116**
N,N-Dibenzylhydroxylamine **15.11**
 Dibenzyl sulfide **13.71**
 2,5-Dibromofuran **5.27**
 (*E,E*)-1,4-Di(*tert*-butoxy)-1,3-butadiene **2.36**
 (*E,Z*)-1,4-Di(*tert*-butoxy)-1,3-butadiene **2.37**
 (*Z,Z*)-1,4-Di(*tert*-butoxy)-1,3-butadiene **2.38**
 3,5-Di-*tert*-butylcatechol **4.7**
 2,6-Di-*tert*-butyl-4-chlorophenol **4.110**
 2,6-Di-*tert*-butyl-*p*-cresol **4.102, 4.99, 4.116, 4.117, 4.118, 4.147**

- 2,5-Di-*tert*-butylfuran 5.28
 2,5-Di-*tert*-butylhydroquinone 4.14
 3,5-Di-*tert*-butyl-4-hydroxybenzyl alcohol 4.32
 2-(3',5'-Di-*tert*-butyl-2'-hydroxyphenyl)-5-chlorobenzotriazole 4.109
 3,5-Di-*tert*-butyl-4-hydroxyphenyl propionate 4.15
 3,5-Di-*tert*-butyl-4-hydroxytoluene 4.102, 4.99, 4.116, 4.117, 4.118, 4.147
 1,1-Di-*tert*-butyl-2-methoxyethene 2.353, 2.355
 2,4-Di-*tert*-butyl-5-methoxyphenol 4.100
 2,6-Di-*tert*-butyl-4-methoxyphenol 4.101
 2,6-Di-*tert*-butyl-4-methylanisole 3.29
 2,6-Di-*tert*-butyl-4-methylphenol 4.102, 4.99, 4.116, 4.117, 4.118, 4.147
 2,6-Di-*tert*-butylphenol 4.99, 2.280
 2',4'-Di-*tert*-butylphenyl 3,5-di-*tert*-butyl-4-hydroxybenzoate 4.38
 Dibutyl sulfide 13.72
 Di-2-butyl sulfide 13.73
 Di-*sec*-butyl sulfide 13.73
 Di-*tert*-butyl sulfide 13.74
 Di-*tert*-butylthioketene 13.165
 Di-*tert*-butylthioketone 13.117
 2,6-Dichloroanisole 3.37
 9,10-Dichloroanthracene 3.7
 1,3-Dichlorobenzene 3.35
 1,4-Dichlorobenzene 3.36
 1,4-Dichloro-2,3-diazabicyclo[2.2.2]oct-2-ene-2,3-dioxide 15.18
 5,5'-Dichloro-3,3-diethyl-2,2'-thiacarbocyanine bromide 11.163
cis-1,4-Dichloro-1,4-dinitrosocyclohexane 15.17
trans-1,4-Dichloro-1,4-dinitrosocyclohexane 15.16
 5,5'-Dichloro-1,1',3,3,3',3'-hexamethyltricarbocyanine, iodide 11.178
 Dichloromethane 17.16
 Dichloromethane-*d*₂ 17.17
 2,4-Dichlorophenol 4.113
 2,6-Dichlorophenol 4.114
 2,4-Dichlorophenoxy ion 4.159
 2,6-Dichlorophenoxy ion 4.160
 2,3-Di(3-chlorophenyl)-1,4-dioxene 2.186
 2,3-Di(4-chlorophenyl)-1,4-dioxene 2.187
 Dichloro[5,10,15,20-tetrakis(4-sulfonatophenyl)porphinatostannate(IV)] ion 7.58
 Dichloro(5,10,15,20-tetraphenylporphinate)tin(IV) 7.64
 4,4'-Dichlorothiobenzophenone 13.159
 5,5'-Dichloro-3,9,3'-triethyl-2,2'-thiacarbocyanine bromide 11.164
 5,5'-Dicyano-3,9,3'-triethyl-2,2'-thiacarbocyanine tetrafluoroborate 11.165
 3,5-Dicyclohexyloxycarbonyl-1,4-dihydro-2,6-dimethylpyridine 6.46
 Dicyclopentadienyiron 12.144
 Dicyclopentadienylnickel 12.165
 (Dicyclopropylmethyldiene)cyclobutane 2.96, 2.49, 2.50, 2.94, 2.95, 2.151, 2.168, 2.169, 2.333, 2.334, 2.335, 3.113
 (Dicyclopropylmethyldiene)cyclopentane 2.151
 (Dicyclopropylmethyldiene)cyclopropane 2.168
 1,1-Dicyclopropyl-2-methylpropene 2.334
 1,1-Dicyclopropylpropene 2.333
 3,4-Didehydro-9-ethenyl-14-ethyl-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-3-phorbinepropanoatomagnesate, hydrogen 7.37
 1,4-Di(4,4'-dimethoxyphenyl)-1,3-butadiene 2.35
 2,3-Di(4-dimethylaminophenyl)-1,4-dioxene 2.188
 Didodecylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II) 12.132
 2,5-Di-*sec*-dodecylhydroquinone 4.17
 4a,9a:9,10-Diethenoanthracene, 1,4,9,10-tetrahydro- 2.181
 4a,9a:9,10-Diethenoanthracene, 1,4,9,10-tetrahydro-5,8-dimethoxy- 2.180
 2,17-Diethenyl-1,10,19,22,23,24-hexahydro-3,7,13,18-tetramethyl-1,19-dioxobilane-8,12-dipropanoic acid 7.5, 7.6, 3.60
 2,17-Diethenyl-1,19,22,24-tetrahydro-3,7,13,18-tetramethyl-1,19-dioxobilane-8,12-dipropanoic acid 7.7
 7,12-Diethenyl-3,8,13,17-tetramethylporphine-2,18-dipropanoatozincate(II) 7.81
 7,12-Diethenyl-3,8,13,17-tetramethylporphine-2,18-dipropanoic acid 7.73
 7,12-Diethenyl-3,8,13,17-tetramethylporphine-2,18-dipropanoic acid, dimethyl ester 7.78
 3,5-Diethoxycarbonyl-1,4-dihydro-2,6-dimethylpyridine 6.47
 3,5-Diethoxycarbonyl-1,4-dihydro-2,6-diphenylpyridine 6.50
 3,5-Diethoxycarbonyl-1,4-dihydro-2,4,6-trimethylpyridine 6.51
 3,5-Diethoxycarbonyl-2,6-dimethylpyridine 6.52
 3,4-Diethoxycarbonylfuran 5.62
 1,1-Diethoxyethene 2.221
cis-1,2-Diethoxyethene 2.223, 2.222
(E)-1,2-Diethoxyethene 2.222
trans-1,2-Diethoxyethene 2.222
(Z)-1,2-Diethoxyethene 2.223, 2.222
 Diethylamine 8.27
 4-(Diethylamino)azobenzene 11.56, 11.53, 11.90, 11.102
 4-Diethylaminobenzaldehyde diphenylhydrazone 15.4
 7-(Diethylamino)coumarin 5.6
 4-(4'-Diethylamino-2',6'-dimethylphenyl)imino-3-(benzoylamino)-1-phenyl-2-pyrazolin-5-one 11.138

- 4-(4'-Diethylamino-2',6'-dimethylphenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one **11.140**
- 4-(4'-Diethylamino-2',6'-dimethylphenyl)imino-3-(2-methyl-2-propyl)-1-phenyl-2-pyrazolin-5-one **11.139**
- 4-(Diethylamino)-2',4'-dinitroazobenzene **11.52**
- 2-(Diethylamino)ethanol **8.23**
- Diethyl(7-aminoheptyl)amine **8.31**
- 2-(Diethylamino)-4-hydroxy-6-methylpyrimidine **16.68**
- 4-(Diethylamino)-4'-methoxyazobenzene **11.53**
- 1-(Diethylamino)-2-methylacetylene **8.87**
- 7-(Diethylamino)-4-methylcoumarin **5.7**
- 4-(4'-Diethylamino-2'-methylphenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one **11.144**
- 4-(4'-Diethylamino-2'-methylphenyl)imino-3-(2-methyl-2-propyl)-1-phenyl-2-pyrazolin-5-one **11.141**
- 4-(Diethylamino)-3'-nitroazobenzene **11.54**
- 4-(Diethylamino)-4'-nitroazobenzene **11.55**
- 4-(4'-Diethylaminophenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one **11.147**
- N,N*-Diethylbenzenemethanesulfenamide **13.35**
- 1,1'-Diethyl-4,4'-carbocyanine **11.42**
- 1,1'-Diethyl-2,2'-carbocyanine chloride **11.41**
- 1,1'-Diethyl-4,4'-carbocyanine toluenesulfonate **11.43**
- 1,1'-Diethyl-2,2'-cyanine iodide **11.45**
- Diethyl(2-cyanoethyl)amine **8.81**
- 3,3'-Diethyl-4,5,4',5'-dibenzo-2,2'-thiacarbocyanine toluenesulfonate **11.107**
- 3,3'-Diethyl-4,5,4',5'-dibenzo-2,2'-thiacyanine chloride **11.105**
- Diethyl 1,4-dihydro-2,4,6-trimethyl-3,5-pyridinedicarboxylate **6.51**
- 3,3'-Diethyl-5,5'-dimethoxy-2,2'-thiacarbocyanine toluenesulfonate **11.167**
- 1,2-Diethyl-1,2-dimethylhydrazine **8.33**
- N,N*-Diethyl-3,5-dimethyl-4-(4'-oxo-2',6'-dimethylcyclohexadienyliden)aminoaniline **11.49**
- Diethyl 2,6-dimethyl-3,5-pyridinedicarboxylate **6.52**
- Diethyl 2,4-dimethylpyrrole-3,5-dicarboxylate **6.66**
- 3,3'-Diethyl-8,10-dimethyl-2,2'-thiacarbocyanine toluenesulfonate **11.168**
- Diethyl disulfide **13.75**
- 3,3'-Diethyl-8,10-ethanediyi-2,2'-thiacarbocyanine toluenesulfonate **11.169**
- N,N*-Diethylethanolamine **8.23**
- Diethyl 3,4-furandicarboxylate **5.62**
- N,N*-Diethyl-1,7-heptanediamine **8.31**
- 2,17-Diethyl-1,10,19,22,23,24-hexahydro-3,7,13,18-tetramethyl-1,19-dioxobiline-8,12-dipropanoic acid **7.9**
- Diethyl(2-hydroxyethyl)amine **8.23**
- N,N*-Diethylhydroxylamine **15.23**
- O,N*-Diethylhydroxylamine **15.21**
- N,N*-Diethylhydroxylammonium ion **15.24**
- (*E,Z*)-4,5-Diethylidene-2,2-dimethyl-1,3-dioxolane **2.197**
- (*Z,Z*)-4,5-Diethylidene-2,2-dimethyl-1,3-dioxolane **2.198**
- Diethyl(2-methoxyethyl)amine **8.25**
- N,N*-Diethyl-3-methyl-4-[*N*-(phenylaminocarbonyl)pivaloylmethylene]aminoamine **11.111**
- 3,3'-Diethyl-2,2'-oxatricarbocyanine iodide **11.109**
- N,N*-Diethyl-4-[*N*-(phenylaminocarbonyl)pivaloylmethylene]aminoamine **11.112**
- 3,3'-Diethyl-8,9-(1,3-propanediyl)-2,2'-thiacarbocyanine iodide **11.170**
- Diethyl sulfide **13.76**
- 7,12-Diethyl-3,8,13,17-tetramethylporphine-2,18-dipropanoato cobaltate(II), dimethyl ester **7.86**
- 7,12-Diethyl-3,8,13,17-tetramethylporphine-2,18-dipropanoic acid, cobalt(II) **7.82**
- 7,12-Diethyl-3,8,13,17-tetramethylporphine-2,18-dipropanoic acid, dimethyl ester **7.85**
- 3,3'-Diethyl-2,2'-thiacarbocyanine toluenesulfonate **11.166**
- 3,3'-Diethyl-2,2'-thiadicarbocyanine iodide **11.173**
- 3,3'-Diethyl-2,2'-thiatricarbocyanine iodide **11.174**
- 5,7-Diethyltocol **4.45**
- 5,5'-Difluoro-1,1',3,3,3',3'-hexamethyltricarbocyanine, iodide **11.179**
- 2,5-Di-sec-hexadecylhydroquinone **4.18**
- 3,5-Dihexyloxycarbonyl-1,4-dihydro-2,6-dimethylpyridine **6.48**
- 2,4-Dihydro-4-(4-aminophenyl)imino-5-methyl-2-phenylpyrazol-3-one **11.133**
- 2,4-Dihydro-2-(4-bromophenyl)-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamylpyrazol-3-one **11.135**
- 2,4-Dihydro-2-(3-chlorophenyl)-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamylpyrazol-3-one **11.136**
- 2,4-Dihydro-2-(4-chlorophenyl)-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamylpyrazol-3-one **11.137**
- 2,4-Dihydro-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2-(3-methoxyphenyl)pyrazol-3-one **11.142**
- 2,4-Dihydro-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2-(3-methylphenyl)pyrazol-3-one **11.143**
- 2,4-Dihydro-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2-(4-nitrophenyl)pyrazol-3-one **11.145**
- 2,4-Dihydro-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2-phenylpyrazol-3-one **11.128**
- 2,4-Dihydro-4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2-(2,4,6-trichlorophenyl)pyrazol-3-one **11.146**
- 2,4-Dihydro-4-(4-diethylamino-2-methylphenyl)imino-5-methyl-2-phenylpyrazol-3-one **11.144**

- 2,4-Dihydro-4-(4-diethylaminophenyl)imino-5-methyl-2-phenylpyrazol-3-one **11.147**
- 2,4-Dihydro-4-(4-dimethylaminophenyl)imino-5-methyl-2-phenylpyrazol-3-one **11.150**
- 1,4-Dihydro-2,6-dimethyl-3,5-di(1-methylethoxycarbonyl)pyridine **6.49**
- 1,4-Dihydro-2,6-dimethyl-3,5-diphenylaminocarbonylpyridine **6.44**
- 2,3-Dihydro-4,5-dimethylfuran **5.30**
- 4,4'-(4,7-Dihydro-5,6-dimethylisobenzofuran-1,3-diy)bis(benzene-*p*-decanoate ion) **5.81**
- 2-[7-(1,3-dihydro-1,1-dimethyl-3-(sulfobutyl)benz[e]indol-2-ylidene]-1,3,5-heptatrienyl]-1,1-dimethyl-3-(sulfobutyl)benz[e]indolium, hydroxide, inner salt, Na salt **11.29**
- 2-[2-[3-[[1,3-dihydro-1,1-dimethyl-3-(3-sulfopropyl)benz[e]indol-2-ylidene]ethylidene]-2-[4-(ethoxycarbonyl)-1-piperidinyl]-1-cyclopenten-1-yl]ethenyl]-1,1-dimethyl-3-(3-sulfopropyl)benz[e]indolium, hydroxide, inner salt, compound with triethylamine **11.30**
- 2,3-Dihydro-1,4-dioxin **2.183, 2.122, 2.222, 2.223, 2.224, 3.101, 5.90**
- 4,5-Dihydro-1,5-diphenyl-3-(2-phenylethenyl)pyrazole **11.119**
- 2,4-Dihydro-4-[4-[ethyl(2-hydroxyethyl)amino]-2-methylphenyl]imino-5-methyl-2-phenylpyrazol-3-one **11.129**
- 1,4-Dihydro-1-ethyl-7-methyl-4-oxo-1,8-naphthyridine-3-carboxylate ion **16.46**
- 1,4-Dihydro-1-ethyl-7-methyl-4-oxo-1,8-naphthyridine-3-carboxylic acid **16.45**
- 2,3-Dihydrofuran **5.29, 7.59**
- 14,15-Dihydro-14-hydroxy-eburnamenine-14-carboxylic acid, methyl ester **8.100**
- 2,4-Dihydro-4-[4-[(2-hydroxyethyl)ethylamino]phenyl]imino-5-methyl-2-phenylpyrazol-3-one **11.130**
- 2,3-Dihydro-5-hydroxy-2,2,4,6,7-pentamethylbenzofuran **4.36**
- 2,3-Dihydro-5-hydroxy-2,4,6,7-tetramethylbenzofuran **4.37**
- 3,4-Dihydro-6-hydroxy-2,5,7,8-tetramethylbenzopyran-2-propionic acid **4.66**
- 3,4-Dihydrojasnone **2.162**
- Dihydrolipoate ion **13.115**
- Dihydroluteolin **4.59**
- 4,5-Dihydro-5-(4-methoxyphenyl)-3-[2-(4-methoxyphenyl)ethenyl]-1-phenylpyrazole **11.120**
- 2,3-Dihydro-5-methylfuran **5.31**
- 4,5-Dihydro-2-methylfuran **5.31**
- 2,4-Dihydro-4-[4-[(2-methylsulfamylethyl)ethylamino]-2-methylphenyl]imino-5-methyl-2-phenylpyrazol-3-one **11.78**
- 3,4-Dihydro-1-methyl-1-(4,8,12-trimethyltridecyl)-1-benzopyran-6-ol **4.48**
- Dihydronicotinamide adenine dinucleotide **16.48**
- 3,4-Dihydro-2,2,5,7,8-pentamethylbenzopyran-6-ol **4.49**
- Dihydroperoxy-2,6,10-trimethyl-2,6,10-dodecatriene **2.208**
- Dihydropyran **5.90**
- 2,3-Dihydropyran **5.90**
- 3,4-Dihydropyran **5.90**
- Dihydroquercetin **4.63**
- 1,4-Dihydro-2,4,6-trimethyl-3,5-diphenylaminocarbonylpyridine **6.45**
- 4,5-Dihydro-1,3,5-triphenylpyrazole **11.121, 11.122, 11.123, 11.124, 11.125, 11.126, 11.127**
- 1,8-Dihydroxy-9-anthrone **4.2**
- 1,8-Dihydroxy-9-anthrone, conjugate base **4.3**
- 1,2-Dihydroxybenzene **4.4**
- 1,3-Dihydroxybenzene **4.10**
- 1,4-Dihydroxybenzene **4.12**
- 1,4 Dihydroxybenzene ion(2-) **4.13**
- 2,2'-Dihydroxybiphenyl **4.128**
- 2,2'-Dihydroxybiphenyl, conjugate base **4.166**
- 2,5-Dihydroxybiphenyl **4.30**
- 2,5-Dihydroxybiphenyl, conjugate base **4.167**
- 2,3-Dihydroxy-1,4-butanedithiol (*R*^{*},*R*^{*})(±) **13.50**
- 2,3-Dihydroxy-1,4-butanedithiol (*R*^{*},*S*^{*}) **13.49**
- 3,3'-Dihydroxy-β-carotene-4,4'-dione **2.82**
- 3,4-Dihydroxycinnamic acid **4.68**
- 2,4-Di[2-hydroxy-4-(diethylamino)phenyl]squarylium **11.46, 12.66**
- 2,2'-Dihydroxydiethyl sulfide **13.164**
- 3,7-Dihydroxy-2-(3,4-dihydroxyphenyl)-1-benzopyran-4-one **4.75**
- 1,4-Dihydroxy-2,3-dimethylbenzene **4.22**
- 1,4-Dihydroxy-2,5-dimethylbenzene **4.23**
- 1,4-Dihydroxy-2,6-dimethylbenzene **4.24**
- Di(2-hydroxyethyl)-*tert*-butylamine **8.26**
- Di(2-hydroxyethyl)methylamine **8.46**
- 5,7-Dihydroxyflavone **4.73**
- Dihydroxylycopene **2.251**
- l*-3,4-Dihydroxy-α-(methylaminomethyl)benzyl alcohol **4.9**
- 1,4-Dihydroxy(methyl)benzene **4.29**
- 2,4-Dihydroxy-5-methylpyrimidine **16.79**
- 1,5-Dihydroxynaphthalene **4.82**
- 1,6-Dihydroxynaphthalene **4.83**
- 1,7-Dihydroxynaphthalene **4.84**
- 1,8-Dihydroxynaphthalene **4.85**
- 2,6-Dihydroxynaphthalene **4.86**
- 2,7-Dihydroxynaphthalene **4.87**
- 2,5-Di-(4-hydroxy-3-nitrophenyl)-2,5-dimethylpyrrolidin-1-oxyl **15.54**
- 3,4-Dihydroxyphenethylamine **4.5**
- 3-(3,4-Dihydroxyphenyl)-L-alanine **4.183**

- (*E*)-2-(3,4-Dihydroxyphenyl)-3,4-dihydro-1-benzopyran-3,5,7-triol **4.67**
 2,3-Di(4-hydroxyphenyl)-1,4-dioxene **2.189**
 2-(3,4-Dihydroxyphenyl)-3,5,7-trihydroxy-1-benzopyran-4-one **4.61**
 Diindeno[5,6-*a*:5',6'-*j*]perylene-8,17-dione, 1,2,3,10,11,12-hexahydro- **3.65**
 Diisobutyl sulfide **13.77**
 Diisopropylamine **8.29**
 Diisopropyl sulfide **13.78**
 Dimedone **17.15**
erythro-1,4-Dimercapto-2,3-butanediol **13.49**
threo-1,4-Dimercapto-2,3-butanediol **13.50**
 3,4-Dimercaptotoluene(2,2'-bipyridine)nickel(II) **12.153**
 3,4-Dimercaptotoluene(1,10-phenanthroline)nickel(II) **12.154**
 3,4-Dimercaptotoluene(1,10-phenanthroline)platinum(II) **12.155**
 Dimesna dianion **13.91**
 1,9:3,5-Dimethanocyclopenta[*d*]cyclopenta-[3,4][1,2]diazeto[1,2-*a*]pyridazine, decahydro- **8.22**
 Dimethirimol **16.66**
 9,10-Dimethoxyanthracene **3.8**
 1,2-Dimethoxybenzene **3.38**
 1,3-Dimethoxybenzene **3.39**
 1,4-Dimethoxybenzene **3.40**
 1,4-Dimethoxybenzonorbornene **3.73**
endo-2,3-Di(methoxycarbonyl)-7-adamantylenenorbornane **2.17**
exo-2,3-Di(methoxycarbonyl)-7-adamantylenenorbornane **2.18**
 1,3-Dimethoxy-4,6-di-*tert*-butylbenzene **3.28**
 3,4-Dimethoxy-*N,N*-dimethylaniline **9.5**
 1,4-Dimethoxy-9,10-diphenylanthracene **3.9**
 1,2-Dimethoxy-1,2-diphenylethene **3.101**
 5,5'-Dimethoxy-1,1',3,3,3',3'-hexamethyltricarbocyanine, iodide **11.180**
 5,6-Dimethoxyindole **6.9**
 2,3-Dimethoxy-5-methyl-1,4-benzenediol **4.21**
 2,3-Dimethoxy-5-methylhydroquinone **4.21**
 2,6-Dimethoxyphenol **4.115**
 2,6-Dimethoxyphenoxide ion **4.161**
 2,3-Di(3-methoxyphenyl)-1,4-dioxene **2.190**
 2,3-Di(4-methoxyphenyl)-1,4-dioxene **2.191**
 1,2-Dimethoxystilbene **3.101**
 4,4'-Dimethoxystilbene **3.100**
 α,β -Dimethoxystilbene **3.101**
 2,3-Dimethoxystyrychnidin-10-one **8.9**
 5,8-Dimethoxy-1,2,3,4-tetrahydro-9-isopropylidene-1,4-methanonaphthalene **2.257**
 4,4'-Dimethoxythiobenzophenone **13.160**
 11,17-Dimethoxy-18-[(3,4,5-trimethoxybenzoyl)oxy]yohimban-16-carboxylic acid methyl ester **8.98**
 4-(Dimethylamino)benzaldehyde **9.10**
p-Dimethylaminobenzaldehyde **9.10**
 4-(Dimethylamino)benzonitrile **9.22**
 4-(4'-Dimethylamino-3',5'-dimethylphenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one **11.149**
 2-Dimethylamino-5,6-dimethylpyrimidin-4-ol **16.69**
 1-[4-(Dimethylamino)phenylazo]-2-naphthol **11.96**
 5-[*p*-(Dimethylamino)phenyl]-3-[*p*-(dimethylamino)styryl]1-phenyl-2-pyrazoline **11.118**
 4-(4'-Dimethylaminophenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one **11.150**
 2-(4-Dimethylaminophenyl)-3-phenyl-1,4-dioxene **2.192**
 3-Dimethylamino-1-propanol **8.82**
 4-(4'-Dimethylamino-2',3',5',6'-tetramethylphenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one **11.151**
p-(Dimethylamino)thioanisole **13.6**
p-(*N,N*-Dimethylamino)- α,β,β -trimethylstyrene **3.115**
N,N-Dimethylaniline **9.6**
N,N-Dimethyl-*m*-anisidine **9.12**
N,N-Dimethyl-*o*-anisidine **9.11**
N,N-Dimethyl-*p*-anisidine **9.13**
 9,10-Dimethylanthracene **3.10**, **3.124**, **3.125**, **4.1**, **4.10**, **4.11**, **4.12**, **4.20**, **4.153**, **4.154**, **4.157**, **4.159**, **4.160**, **4.162**, **4.165**, **4.166**, **4.168**, **4.175**, **4.176**, **4.180**, **4.181**, **4.182**, **11.16**, **11.17**, **11.60**, **11.62**, **11.63**, **11.77**, **11.79**, **11.80**, **11.90**, **11.92**, **11.101**, **11.103**, **11.104**, **12.28**, **12.123**, **13.175**
 9,10-Dimethyl-1,2-benzanthracene **3.24**, **3.124**, **3.125**
 9,10-Dimethylbenz[a]anthracene **3.24**, **3.124**, **3.125**
 3,3-Dimethylbicyclo[2.2.1]heptane-2-thione **13.44**
 2,3-Dimethyl-1,3-butadiene **2.39**
N,N-Dimethyl-1-butanamine **8.14**
 2,3-Dimethyl-1-butene **2.45**
 2,3-Dimethyl-2-butene **2.51**, **2.38**, **2.50**, **2.52**, **2.53**, **2.56**, **2.103**, **2.117**, **2.140**, **2.168**, **2.170**, **2.184**, **2.185**, **2.186**, **2.187**, **2.188**, **2.189**, **2.190**, **2.191**, **2.192**, **2.193**, **2.194**, **2.195**, **2.196**, **2.223**, **2.225**, **2.303**, **2.310**, **2.332**, **2.334**, **3.10**, **3.11**, **3.28**, **3.66**, **3.125**, **4.3**, **4.100**, **5.34**, **7.5**, **7.65**, **8.83**, **9.1**, **13.86**, **13.145**
 2,3-Dimethyl-2-butene-*d*₁₂ **2.54**
cis-2,3-Dimethyl-2-butene-*d*₆ **2.53**
 (*E*)-2,3-Dimethyl-2-butene-*d*₆ **2.52**
trans-2,3-Dimethyl-2-butene-*d*₆ **2.52**
(*Z*)-2,3-Dimethyl-2-butene-*d*₆ **2.53**
 2,3-Dimethyl-2-butene-4-methoxyphenyl sulfide **13.18**
 2,3-Dimethyl-2-butene-4-methylphenyl sulfide **13.19**
 2,3-Dimethyl-2-butene phenyl sulfide **13.17**
 Dimethylcercosporin **16.57**
 2,2-Dimethylcyclohexa-3,5-dien-1-one **2.108**
 5,5-Dimethyl-1,3-cyclohexanedione **17.15**
 1,2-Dimethylcyclohexene **2.117**, **2.161**
 1,3-Dimethylcyclohexene **2.118**

1,4-Dimethylcyclohexene	2.119	5,7-Di(1-methylethyl)tocol	4.47
1,6-Dimethylcyclohexene	2.120, 2.156	<i>N,N</i> -Dimethylformamide	17.31
2,3-Dimethylcyclohexene	2.120, 2.156	6,6-Dimethylfulvene	2.149
6,6-Dimethylcyclohex-1-enyl acetate	2.126	6,6-Dimethylfulvene endoperoxide	2.150
1,2-Dimethylcyclopentene	2.158	2,4-Dimethylfuran	5.33
1,5-Dimethylcyclopentene	2.159	2,5-Dimethylfuran	5.34, 7.59, 9.28
2,3-Dimethyl-2-cyclopenten-1-one	2.165	Dimethyl 2,5-furandicarboxylate	5.61
2,3-Dimethyl-2,3-diazabicyclo[2.2.1]heptane	8.18	Dimethylglyoxime	15.19
1,2-Dimethyl-1,2-dibutylhydrazine	8.34	6,11-Dimethyl4,6,8,10,12-hexadecapentaene-2,3,14,15-tetraone (<i>all-E</i>)	2.234
2,4-Dimethyl-3,5-dicarbethoxypyrrole	6.66	2,5-Dimethyl-2,4-hexadiene	2.243, 13.87, 13.88
1,1-Dimethyl-2,2-di(2,2-dimethylpropyl)hydrazine	8.35	2,2'-Dimethyl-4,4',5,5',7,7'-hexahydroxymesonaphthodianthrone	16.60
1,2-Dimethyl-1,2-di(2,2-dimethylpropyl)hydrazine	8.36	Dimethylhomocoerdianthrone	3.95
1,1'-(1,2-Dimethyl-1,2-diethenyl)bisnaphthalene, (<i>E</i>)-	3.84	4-[2-(<i>N,N</i> -Dimethylhydrazono)ethylidene]-2,6-diphenylpyran	15.1
2,2'-(1,2-Dimethyl-1,2-diethenyl)bisnaphthalene, (<i>E</i>)-	3.85	2,7-Dimethyl-6-hydroperoxy-2,7-octadiene	2.279
(15 <i>E</i>)-Dimethyl-2,17-diethenyl-1,19,22,24-tetrahydro-3,7,13,18-tetramethyl-1,19-dioxobilin-8,12-dipropanoate	7.8	2,7-Dimethyl-7-hydroperoxy-2,5-octadiene	2.274
4,5-Dimethyl-2,3-dihydrofuran	5.30	2,3-Dimethylhydroquinone	4.22
4,4-Dimethyl-2,3-dihydropyran-2- <i>t</i>	5.96	2,5-Dimethylhydroquinone	4.23
4,4-Dimethyl-2,3-dihydropyran-3- <i>t</i>	5.94	2,6-Dimethylhydroquinone	4.24
5,6-Dimethyl-3,4-dihydropyran	5.91	2,3-Dimethyllidenebicyclo[2.2.1]heptane	2.8, 2.19, 2.140, 2.295, 2.296, 2.297, 2.357
(Z)-Dimethyl 7-[2-(dimethylamino)-2-oxoethyl]-8-ethyl-7,8-dihydro-3,7,12,17-tetramethylporphine-2,18-dipropionate	7.88	5,6-Dimethylidene-2-bicyclo[2.2.1]heptene	2.19
(Z)-Dimethyl 7-[2-(dimethylamino)-2-oxoethyl]-8-ethylidene-7,8-dihydro-3,7,12,17-tetramethylporphine-2,18-dipropionate	7.89	2,3-Dimethylidene-7-oxabicyclo[2.2.1]heptane	2.295
2,4-Di[2-methyl-4-(dimethylamino)phenyl]squarylium	11.47	5,6-Dimethylidene-7-oxa-2-bicyclo[2.2.1]heptene	2.297
1,1-Dimethyl-2,2-di(2-methylpropyl)hydrazine	8.37	1,3-Dimethylindole	6.10
1,2-Dimethyl-1,2-di(2-methylpropyl)hydrazine	8.38	2,3-Dimethylindole	6.11
1,2-Dimethyl-1,2-dipentylhydrazine	8.39	<i>N,N</i> -Dimethylisobut enylamine	8.83
1,2-Dimethyl-1,2-dipropylhydrazine	8.40	7,7-Dimethyl-2-methylenenorbornane	2.9
Dimethyldithiocarbamate ion	13.79	Dimethyl 4-methyl-1,3-naphthalenedipropionate	3.90
4,8-Dimethyl-4,8-dodecadiene	2.201	Dimethyl 3,3'-(4-methyl-1,3-naphthylene)dipropionate	3.90
(<i>E,E</i>)-4,8-Dimethyl-4,8-dodecadiene	2.202	2,4-Di[2-methyl-4-(<i>N</i> -methyl- <i>N</i> -octadecylamino)phenyl]squarylium	11.48
2,3-Dimethylenebicyclo[2.2.1]heptane	2.8, 2.19, 2.140, 2.295, 2.296, 2.297, 2.357	1,2-Dimethylnaphthalene	3.75
2,3-Dimethylenebicyclo[3.2.2]nonane	2.31	1,3-Dimethylnaphthalene	3.76
6,7-Dimethylenebicyclo[3.2.2]nonane	2.31	1,4-Dimethylnaphthalene	3.77
2,3-Dimethylenebicyclo[2.2.2]octane	2.33	1,5-Dimethylnaphthalene	3.75, 3.76, 3.77, 3.78, 3.79, 3.80, 3.81, 3.82, 3.83, 3.86
1,1-Dimethylethanol	17.62	1,6-Dimethylnaphthalene	3.78
2-[(1,1-Dimethylethoxy)methyl]furan	5.35	1,7-Dimethylnaphthalene	3.79
7-(1,1-Dimethylethyl)-3,4-dihydro-2,2-dimethyl-1-benzopyran-6-ol	4.54	1,8-Dimethylnaphthalene	3.80
2-(1,1-Dimethylethyl)-3,3-dimethyl-1-butene-1-thione	13.165	2,3-Dimethylnaphthalene	3.81
1,1-Dimethylethyl ethyl sulfide	13.56	2,6-Dimethylnaphthalene	3.82
1,1-Dimethylethyl methyl ether	17.18	2,7-Dimethylnaphthalene	3.83
4-(1,1-Dimethylethyl)phenol	4.121	1,1-Dimethyl-2-naphthalenethione	13.113
1,1-Dimethylethyl propyl sulfide	13.58	3,10-Dimethylnaphtho[1,2,3,4- <i>rst</i>]pentaphene-5,8-dione	3.95
		<i>N,N</i> -Dimethyl-4-nitrosoaniline	9.7
		2,7-Dimethyl-2,6-octadiene	2.277
		<i>cis</i> -2,6-Dimethyl-2,6-octadiene	2.276

- (E)-2,6-Dimethyl-2,6-octadiene 2.275
trans-2,6-Dimethyl-2,6-octadiene 2.275
(Z)-2,6-Dimethyl-2,6-octadiene 2.276
3,7-Dimethyl-1,6-octadien-3-ol 2.280, 2.398, 13.53, 13.54,
13.55, 13.143, 13.144
(E)-3,7-Dimethyl-2,6-octadien-1-ol 2.281
(E)-3,7-Dimethyl-2,7-octadien-1-ol 2.282
(Z)-3,7-Dimethyl-2,7-octadien-1-ol 2.283
2,6-Dimethyl-2,4,6-octatriene 2.285
3,7-Dimethyl-6-octen-1-ol 2.294
(E)-3,7-Dimethyl-2-octen-1-ol 2.292
(Z)-3,7-Dimethyl-2-octen-1-ol 2.293
3,3-Dimethyl-2-oxobutyric acid 17.7
2,4-Dimethyl-2-pentene 2.306
(E)-3,4-Dimethyl-2-penten-1-ol 2.316
(Z)-3,4-Dimethyl-2-penten-1-ol 2.317
3,4-Dimethyl-3-penten-2-one 2.324
2,4-Dimethylphenol 4.116
2,6-Dimethylphenol 4.117
3,4-Dimethylphenol 4.118
2,6-Dimethylphenoxy ion 4.162
2,3-Di(4-methylphenyl)-1,4-dioxene 2.193
2,2-Dimethyl-1-phenyl-1-propanethione 13.126
N,N-Dimethyl-1-piperidinamine 8.52
1,2-Dimethylpiperidine 8.57
2,6-Dimethylpiperidine 8.58
(1,2-Dimethyl-1-propenyl)benzene 3.123, 2.96
1-(1,2-Dimethyl-1-propenyl)-4-methoxybenzene 3.118,
3.115, 3.123
2,2-Dimethylpropyl methyl ether 17.19
Di(1-methylpropyl) sulfide 13.73
2,6-Dimethylpyran-4-thione 13.139
1,3-Dimethylpyrazolo[1,2-*a*]benzotriazole 11.132
1,3-Dimethylpyrazolo[1',2':2,3][1,2,3]triazolo[4,5-*a*]phenazin-4-i um 6.39
2,5-Dimethylpyrrole 6.60
N,N-Dimethyl-1-pyrrolidinamine 8.95
5,5-Dimethyl-1-pyrroline-1-oxyl 15.56
cis- α , α' -Dimethylstilbene 3.103
trans- α , α' -Dimethylstilbene 3.102, 3.103
 β , β -Dimethylstyrene 3.114
Dimethyl sulfide 13.80, 13.81
Dimethyl sulfoxide 13.81
1,1'-(7,13-Dimethyl-1,4,8,12-tetraazacyclopentadeca-
4,7,12,15-tetraene-6,14-diyl)bis[ethanoato]nickel(II)
12.133
4,4'-Dimethyl-3,3',5,5'-tetracarbethoxy-2,2'-
dipyrromethane 6.65
8,13-Dimethyl-2,2,19,19-tetramethoxy-4,6,8,10,12,14,16-
eicosahptaene-3,18-dione (*all-E*) 2.211
6,11-Dimethyl-2,2,15,15-tetramethoxy-4,6,8,10,12-
hexadecapentaene-3,14-dione (*all-E*) 2.233
Dimethyl 3,7,12,17-tetramethylporphine-2,18-dipropanone
7.94
2,2-Dimethyl-3-thiahexane 13.58
2,2-Dimethyl-3-thiapentane 13.56
4,4'-Dimethylthiobenzophenone 13.161
3,3-Dimethylthiocamphor 13.45
5,7-Dimethyltocol 4.51
N,N-Dimethyl-*p*-toluidine 9.19
7,7-Dimethyl-2-(trimethylsiloxy)-norborna-2,5-diene 2.1
7,7-Dimethyl-2-(trimethylsiloxy)norborn-2-ene 2.5, 2.7,
2.24
(E)-2,6-Dimethyl-1,6-undecadiene 2.359
(Z)-2,6-Dimethyl-1,6-undecadiene 2.360
1-(β , β -Dimethylvinyl)naphthalene 3.87
(E)-2,3-Di(α -naphthyl)-2-butene 3.84
(E)-2,3-Di(β -naphthyl)-2-butene 3.85
(Z)-2,3-Di(α -naphthyl)-2-butene 3.84
(Z)-2,3-Di(β -naphthyl)-2-butene 3.85
2,4-Dinitrophenol 4.123
2,5-Dinitrophenol 4.124
2,6-Dinitrophenol 4.125
2,4-Dinitrophenoxide ion 4.164
4,4'-Di-*tert*-octyldiphenylamine 9.18
2,5-Di-*sec*-octylhydroquinone 4.19
2,3-Dioxabicyclo[2.2.1]hept-5-ene, 7-(1-
methylenelidene)- 2.150
2,3-Dioxabicyclo[2.2.2]oct-5-ene 2.182
3,6-Dioxacyclohexene 2.183, 2.122, 2.222, 2.223, 2.224,
3.101, 5.90
3,5-Dioxacyclopentene 2.199
Dioxane 17.20
1,4-Dioxane 17.20
1,6-Dioxapyrene 3.94
13,14-Dioxatricyclo[8.2.1.1^{4,7}]tetradeca-4,6,10,12-tetraen
5.77
1,4-Dioxene 2.183, 2.122, 2.222, 2.223, 2.224, 3.101, 5.90
1,4-Dioxene, 2-(4-cyanophenyl)-3-(4-
dimethylaminophenyl)- 2.184
1,4-Dioxene, 2,3-di(4-acetaminophenyl)- 2.185
1,4-Dioxene, 2,3-di(3-chlorophenyl)- 2.186
1,4-Dioxene, 2,3-di(4-chlorophenyl)- 2.187
1,4-Dioxene, 2,3-di(4-dimethylaminophenyl)- 2.188
1,4-Dioxene, 2,3-di(4-hydroxyphenyl)- 2.189
1,4-Dioxene, 2,3-di(3-methoxyphenyl)- 2.190
1,4-Dioxene, 2,3-di(4-methoxyphenyl)- 2.191
1,4-Dioxene, 2-(4-dimethylaminophenyl)-3-phenyl- 2.195
1,4-Dioxene, 2,3-di(4-methylphenyl)- 2.193
1,4-Dioxene, 2,3-diphenyl- 2.194, 8.19
1,4-Dioxene, 2-(4-methoxyphenyl)-3-phenyl- 2.195
1,4-Dioxene, 2,3,5,6-tetraphenyl- 2.196

p-Dioxene 2.183, 2.122, 2.222, 2.223, 2.224, 3.101, 5.90
1,3-Dioxolane, 4,5-diethylidene-2,2-dimethyl-, (*E,Z*) 2.1

- 1,3-Dioxolane, 4,5-diethylidene-2,2-dimethyl-, (Z,Z) **2.198**
- 1,3-Dioxolane, 2-(2-furanyl)- **5.36**
- 2-[2-(1,3-Dioxolanyl)]furan **5.36**
- 1,3-Dioxole **2.199**
- Dioxygen anion **14.14**
- exo,exo*-2,3-Dioxymethyl-7-adamantylidenenorbornane **2.10**
- Di(1,2,2,6,6-pentaamethyl-4-piperidinol) 1,10-decanedioate **8.74**
- 2,5-Di-*tert*-pentylhydroquinone **4.16**
- Diphenylamine **9.16**
- 4-[N-Di(phenylaminocarbonyl)methylene]amino-3,N,N-triethylaniline **11.115**
- 2-[2-[2-(Diphenylamino)-3-[3-(4-methoxy-4-oxobutyl)naphtho[2,3-*d*]thiazol-2-ylidene]ethylidene]-1-cyclopenten-1-yl]ethenyl]-3-(4-methoxy-4-oxobutyl)naphtho[2,3-*d*]thiazolium, perchlorate **11.108**
- 4-(4'-Diphenylaminophenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one **11.152**
- 9,10-Diphenylanthracene **3.11, 4.50, 11.78, 11.128, 11.129, 11.130, 11.135, 11.136, 11.137, 11.142, 11.143, 11.144, 11.145, 11.146, 11.147, 13.76**
- 2,5-Diphenyl-3,4-benzofuran **5.83, 2.51, 3.60, 3.96, 3.98, 4.49, 4.54, 4.184, 4.185, 6.39, 7.5, 7.78, 8.19, 10.23, 10.34, 10.35, 10.36, 10.38, 10.42, 11.33, 11.37, 11.38, 11.70, 11.71, 11.72, 11.73, 11.75, 11.76, 11.109, 11.174, 11.178, 11.179, 11.180, 11.181, 11.182, 11.183, 11.184, 12.20, 12.21, 12.22, 12.23, 12.84, 12.138, 13.63, 13.64, 13.108, 13.169, 13.170, 13.171, 15.1, 17.30**
- 1,4-Diphenyl-1,3-butadiene **2.40, 2.35**
- (E)-2,3-Diphenyl-2-butene **3.102, 3.103**
- (Z)-2,3-Diphenyl-2-butene **3.103**
- 1,4-Diphenyl-1,3-cyclopentadiene **2.147**
- Diphenyldiazene **11.57**
- 1,3-Diphenyl-5,6-dimethylisobenzofuran **5.82**
- 1,2-Diphenyl-3,6-dioxacyclohexene **2.194, 8.19**
- 2,3-Diphenyl-1,4-dioxene **2.194, 8.19**
- 2,6-Diphenyl-4-(2,6-diphenyltelluropyran-4-ylidene)methyltelluropyrylium **11.162**
- Diphenyldiselenide **13.83**
- Diphenyl disulfide **13.84**
- Diphenylditelluride **13.85**
- 4,5-Diphenyl-1,3-dithiane-2-thione **13.89**
- 4,5-Diphenyl 1,3-dithiole-2-thione **13.89**
- 2,3-Diphenylfuran **5.37**
- 2,5-Diphenylfuran **5.38, 4.50, 5.25, 5.44, 5.47, 5.50, 5.83**
- 3,4-Diphenylfuran **5.39**
- α,α -Diphenylfurfuryl alcohol **5.67**
- 1,2-Diphenylindene **3.70**
- 1,3-Diphenylisobenzofuran **5.83, 2.51, 3.60, 3.96, 3.98, 4.49, 4.54, 4.184, 4.185, 6.39, 7.5, 7.78, 8.19, 10.23, 10.34, 10.35, 10.36, 10.38, 10.42, 11.33, 11.37, 11.38, 11.70, 11.71, 11.72, 11.73, 11.75, 11.76, 11.109, 11.174, 11.178, 11.179, 11.180, 11.181, 11.182, 11.183, 11.184, 12.20, 12.21, 12.22, 12.23, 12.84, 12.138, 13.63, 13.64, 13.108, 13.169, 13.170, 13.171, 15.1, 17.30**
- 1,2-Diphenyl-4-methylcyclopenta[*b*]quinoline **6.4**
- 2,5-Diphenyl-4-methyloxazole **6.32**
- 1,1-Diphenyl-2-methylpropene **3.46**
- Diphenyl nitrone **15.10**
- 2,5-Diphenyloxazole **6.31**
- (E)-2,3-Diphenyloxirane **17.49**
- trans*-Diphenyloxirane **17.49**
- N,N'*-Diphenyl-*p*-phenylenediamine **9.29**
- 2,6-Diphenylpyran-4-thione **13.140**
- Diphenyl sulfide **13.34, 13.16, 13.23, 13.26, 13.31**
- 3,4-Diphenylsydnone **5.102**
- (E)-2,3-Diphenylthiirane **13.151**
- trans*-Diphenylthiirane **13.151**
- 2,6-Diphenylthiopyran-4-thione **13.173**
- 9,18-Diphenyltribenzo[*a,f,j*]perylene **3.129**
- 4-(*N*-Dipivaloylmethylene)amino-3,N,N-triethylaniline **11.67**
- 3,8,3',10'-Di(1,3-propanediyl)-2,2'-thiacarbocyanine iodide **11.156**
- Dipropylamine **8.86**
- Dipropyl sulfide **13.82**
- Diselenide, diphenyl- **13.83**
- Disilirane, 1,1,2,2-tetrakis(2,4,6-trimethylphenyl)- **17.21**
- Dismutase, superoxide **10.14**
- Disodium [*N,N'*-ethylenebis(5-sulfosalicylideneiminato)]nickelate(II) **12.143**
- Disodium [*N,N'*-*o*-phenylenebis(5-sulfosalicylideneiminato)]nickelate(II) **12.167**
- Disodium [*N,N'*-propylenebis(5-sulfosalicylideneiminato)]nickelate(II) **12.169**
- Disperse Blue 1 **11.14**
- Disperse Blue 14 **11.7**
- Disperse blue polyester **11.12**
- Disperse bright pink **11.13**
- Disperse Orange **11.6**
- Disperse Orange 11 **11.6**
- Disperse Red 11 **11.13**
- Disperse Red 15 **11.5**
- Disperse red 2S **11.5**
- Disperse violet K **11.8**
- Disulfide, diphenyl- **13.84**
- Ditelluride, diphenyl- **13.85**
- Di(2,2,6,6-tetramethyl-4-piperidinol) 1,10-decanedioate **8.78**
- 1,5-Dithiacyclooctane **13.86**
- 1,5-Dithiacyclooctane 1-oxide **13.87**

- 1,4-Dithiane **13.88**
 1,3-Dithiane-2-thione, 4,5-diphenyl- **13.89**
 Dithiazanine iodide **11.173**
 2,2'-Dithiobisethanesulfonate ion **13.91**
N,N'-[Dithiobis[(5-ethyl-2,3-thiophenediyi)methylidyne]]biscyclohexanamine **13.63**
 1,5-Dithiocane **13.86**
 1,5-Dithiocane 1-oxide **13.87**
 Dithioerythritol **13.49**
 1,2-Dithiolane-3-pentanoic acid **13.90**, *3.125*
 Dithiothreitol **13.50**
 Docosahexaenoic acid **2.200**
 Docosanoic acid **17.22**
 4,8-Dodecadiene, 4,8-dimethyl- **2.201**
 4,8-Dodecadiene, 4,8-dimethyl-, (*E,E*)- **2.202**
 4,8-Dodecadiene, hydroperoxy-4,8-dimethyl- **2.203**
 (*E,E*)-8,10-Dodecadienol **2.204**
 Dodecane **17.23**
 Dodecanoic acid **17.24**
 2,4,6,8,10-Dodecapentaene-1,12-dione, 1,12-bis(3-hydroxy-1,2,2-trimethylcyclopentyl)-4,7-dimethyl-[1*R*-[1*α*[2*E,4E,6E,8E,10E,12(1*R*^{*},3*R*^{*})],3*α*]-] **2.205**, **2.206**
 2,4,6,8,10-Dodecapentaene-1,12-dione, 1,12-bis(4-hydroxy-1,2,2-trimethylcyclopentyl)-4,7-dimethyl-[1*R*-[1*α*[2*E,4E,6E,8E,10E,12(1*R*^{*},4*R*^{*})],4*β*]-] **2.207**
 Dodecapreno- β -carotene **2.358**
 2,6,10-Dodecatriene, dihydroperoxy-2,6,10-trimethyl- **2.208**
 2,6,10-Dodecatriene, hydroperoxy-2,6,10-trimethyl- **2.209**
 2,6,10-Dodecatriene, 2,6,10-trimethyl- **2.210**
 Dodecylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II) **12.134**
 Dodecylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) **12.135**
 4-Dodecyloxy-2-hydroxybenzophenone **4.42**
 Dodecyl sulfate, sodium salt **17.25**
 Dodecyltrimethylammonium chloride **17.26**
 Dopa-melanin **4.78**
 Dopamine **4.5**
 Doxorubicin **4.80**
 Dregamine **8.101**
 Durohydroquinone monoethyl ether **4.126**
 Ebselen **13.40**
 E.C. 1.1.1.30 **10.11**
 E.C. 1.3.99.1 **10.13**
 E.C. 1.6.99.3 **10.12**
 E.C. 1.9.3.1 **10.10**
 E.C. 1.15.1.1 **10.14**
 E.C. 3.4.4.4 **10.33**
 E.C. 4.2.1.1 **10.7**
 Echinone **2.88**
 Ehrlich's Reagent **9.10**
 4,6,8,10,12,14,16-Eicosahptaene-3,18-dione, 8,13-dimethyl-2,2,19,19-tetramethoxy- (*all-E*) **2.211**
 Eicosanoic acid **17.27**
 5,8,11,14-Eicosatetraenoic acid **2.212**
 5,8,11,14-Eicosatetraenoic acid, phenyl ester **2.214**
 Elsinochrome A **16.10**
 Elsinochrome B **16.9**
 emi-Crocin **2.178**
 3,6-Endoperoxycyclohexene **2.182**
 Eosin dianion **11.65**
 1,4-Epidioxynaphthalene-1-propanoic acid, 4-methyl- **3.91**
 1,4-Epidioxynaphthalene-1-propanoic acid, 4-methyl-anion **3.92**
 Epinephrine **4.9**
 Ergosta-5,7,22-trien-3-ol (3*β*) **16.26**
 Ergosta-5,7,22-trien-3-ol acetate **16.27**
 Ergosterol **16.26**
 Ergosterol acetate **16.27**
 Ergothioneine **13.104**
 Ergothioneine ion **13.105**
 Eriodictyol **4.59**
 Estra-1,3,5(10),8-tetraen-17-one, 3-methoxy- **16.28**
 1,3,5(10),9(11)-Estratetraen-17-one, 3-(acetoxy)- **16.29**
 1,3,5(10),9(11)-Estratetraen-17-one, 3-(benzyloxy)- **16.30**
 1,3,5(10),9(11)-Estratetraen-17-one, 3-hydroxy- **16.31**
 1,3,5(10),9(11)-Estratetraen-17-one, 3-methoxy- **16.32**, **16.29**, **16.30**, **16.31**, **16.33**
 1,3,5(10),9(11)-Estratetraen-17-one, 3-(toluenesulfonyloxy)- **16.33**
 1,2-Ethandiylidenebis[3,5-di(1,1-dimethylethyl)-2,5-cyclohexadien-4-one] **17.12**
 Ethanone, iodo- **17.28**
 (Z,Z)-3,3'-(1,2-Ethanediylbis(iminomethylidyne))bis[5-ethyl-2-thiophenethione] **13.171**
 2,2'-(1,2-Ethanediylbis(nitrilodecylidyne))bis[4-methylphenolato]nickel(II) **12.136**
 2,2'-(1,2-Ethanediylbis(nitriloethylidyne))bis[phenolato]nickel(II) **12.137**
 [[3,3]-(1,2-Ethanediylbis(nitrilomethylidyne))bis[5-ethylthiophenethionate]]copper(II) **12.138**
 2,2'-(1,2-Ethanediylbis(nitrilomethylidyne))bis[phenolato]cobalt(II) **12.139**
 2,2'-(1,2-Ethanediylbis(nitrilomethylidyne))bis[phenolato]nickel(II) **12.140**
 1,2-Ethanediylidenebis[3-[2-hydroxy-5-methyl-3-(1,1-dimethylethyl)benzyl]-5-(1,1-dimethylethyl)-2,5-cyclohexadien-4-one] **17.11****

- 3,7,4,6-Ethanediylidenepentaleno[1,6-*cd*]pyridazine, 3,7a-dichloro-3*a*,4,5,5*a*,6,7,7*a*,7*b*-octahydro-, 1,2-dioxide **15.20**
- Ethanesulfonate ion, 2,2'-dithiobis- **13.91**
- Ethanesulfonic acid, 2-mercaptop-, ion(1-) **13.92**
- Ethanethiol, 2-amino- **13.93**
- Ethanethiol, 2-[(3-aminopropyl)amino]- **13.94**
- Ethanethiol, 2-hydroxy-, **13.95**
- Ethanethiol, 2-hydroxy-, negative ion **13.96**
- Ethanol **17.29**
- Ethanol, 2-cyclododecylidene- **2.215**
- Ethanol, 2-cyclohexylidene- **2.216**
- Ethanol, 2-cyclooctylidene- **2.217**
- Ethanol, 2-cyclopentylidene- **2.218**
- Ethanol, 2-(diethylamino)- **8.23**
- Ethanol, 2,2'-(1,1-dimethylethyl)imino]bis- **8.26**
- Ethanol, 1-(2-methylene-3,5-cyclohexadienylidene)- **2.219**
- Ethanol, 2-(4-methylphenyl)thio- **13.97**
- Ethanol, 1-tricyclo[3.3.1.1^{3,7}]decylidene-, acetate **2.220**
- Ethanone, 1-(3,4-dihydro-6-methyl-2*H*-pyran-5-yl)- **5.88**
- Ethanone, 1-(4-hydroxyphenyl)-, ion(1-) **4.151**
- Ethene, 1,1-bis(4-methoxyphenyl)- **3.100**
- Ethene, 1,1-diethoxy- **2.221**
- Ethene, 1,2-diethoxy-, (*E*)- **2.222**
- Ethene, 1,2-diethoxy-, (*Z*)- **2.223, 2.222**
- Ethene, ethoxy- **2.224**
- Ethene, tetraethoxy- **2.225**
- Ethene, 1,1,2-tricyclopropyl- **2.169**
- Ethene, 1,1,2-triethoxy- **2.226**
- Ethenylbenzene **3.108**
- 9-Ethenyl-14-ethyl-13-formyl-21-(methoxycarbonyl)-4,8,18-trimethyl-20-oxo-3-phorbinepropanoic acid, 3,7,11,15-tetramethyl-2-hexadecenyl ester **7.27**
- 9-Ethenyl-14-ethyl-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-3-phorbinepropanoic acid, 3,7,11,15-tetramethyl-2-hexadecenyl ester **7.26**
- 8-Ethenyl-13-ethyl-3,7,12,17-tetramethylporphine-2,18-dipropanoic acid **7.90**
- 8-Ethenyl-13-ethyl-3,7,12,17-tetramethylporphine-2,18-dipropanoic acid, dimethyl ester **7.91**
- 7-Ethenyl-12-(1-hydroxyethyl)-3,8,13,17-tetramethylporphine-2,18-dipropanoic acid **7.92**
- 7-Ethenyl-3,8,13,17-tetramethylporphine-2,18-dipropanoic acid **7.93**
- Ether, methyl 1-methylpropenyl **2.55**
- Ethirimol **16.67**
- 5-Ethoxycarbonyl-3,4-dihydro-6-methylpyran **5.98**
- 5-Ethoxycarbonyl-3,4-dihydro-6-phenylpyran **5.100**
- 2-(4-Ethoxycarbonylphenyl)-1,8-dihydropyrrolo[3',2':3,4]cyclopenta[1,2-*b*]pyridine **6.70**
- 2-[(4-Ethoxycarbonyl)phenyl]furan **5.40**
- Ethoxyethene **2.224**
- 1-Ethoxy-2-methyl-1-propene **2.336**
- Ethyl alcohol **17.29**
- Ethylamine, *N,N*-diethyl- **8.24**
- Ethylamine, *N,N*-diethyl-2-methoxy- **8.25**
- Ethylamine, 3,4-dihydroxyphenyl- **4.5**
- Ethylamine, 1,1-dimethyl-*N,N*-bis(2-hydroxyethyl)- **8.26**
- Ethylamine, *N*-ethoxy- **15.21**
- Ethylamine, *N*-ethyl- **8.27**
- Ethylamine, *N*-ethyl-*N*-ethoxy- **15.22**
- Ethylamine, *N*-ethyl-*N*-hydroxy- **15.23**
- Ethylamine, *N*-ethyl-*N*-hydroxy-, conjugate acid **15.24**
- Ethylamine, 1-methyl- **8.28**
- Ethylamine, 1-methyl-*N,N*-(1-methylethyl)- **8.29**
- Ethylamine[2,2'-thiobis(3,4-dimethylphenolato)]nickel(II) **12.141**
- Ethylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) **12.142**
- Ethyl 4-aminobenzoate **9.9**
- Ethyl *p*-aminobenzoate **9.9**
- Ethylbenzene **3.41**
- Ethyl 5-bromo-2-furoate **5.55**
- 2-Ethyl-1,3-butadiene **2.41**
- Ethyl β -apo-8'-carotenoate **2.85**
- 6-Ethyl-3,4-dihydropyran-5-carboxylic acid ethyl ester **5.101**
- Ethyl 4-(dimethylamino)benzoate **9.21**
- Ethyl *p*-dimethylaminobenzoate **9.21**
- Ethyl disulfide **13.75**
- [*N,N'*-Ethylenebis(5-sulfosalicylideneiminato)]nickelate(II) disodium salt **12.143**
- Ethylene-propylene-ethylenenorbornene terpolymer **17.30**
- 3-Ethyl-2-[7-(3-ethyl-2-benzothiazolylidene)-1,3,5-heptatrienyl]benzothiazolium iodide **11.174**
- 3-Ethyl-2-[5-(3-ethyl-2-benzothiazolylidene)-1,3-pentadienyl]benzothiazolium iodide **11.173**
- 3-Ethyl-2-[7-(3-ethyl-2-benzoxazolylidene)-1,3,5-heptatrienyl]benzoxazolium iodide **11.109**
- 1-Ethyl-2-[(1-ethyl-2-quinolinylidene)methyl]quinolinium iodide **11.45**
- 1-Ethyl-4-[3-(1-ethyl-4-quinolinylidene)-1-propenyl]quinolinium **11.42**
- 1-Ethyl-2-[3-(1-ethyl-2-quinolinylidene)-1-propenyl]quinolinium chloride **11.41**
- Ethyl 2-furoate **5.56**
- 4-[*N*-Ethyl-*N*-(2-hydroxyethyl)amino]-4'-methoxyazobenzene **11.58**
- Ethylidenecycloheptane **2.100**
- Ethylidenecyclohexane **2.110**
- Ethylidenecyclooctane **2.135**
- Ethylidenecyclopentane **2.152**
- Ethyl iodide **17.28**
- Ethyl linolenate **2.268**

- 2-Ethyl-3-methyl-2-cyclohexen-1-one **2.127**
 2-Ethyl-3-methyl-2-cyclopenten-1-one **2.166**
 3-Ethyl-2-penten-1-ol **2.318**
 Ethyl phytol ether **2.236**
 Ethyl sulfide **13.76**
 Ethyl vinyl ether **2.224**
 Ethynediamine, tetraethyl- **8.30**
 Etiobilirubin-IV γ **7.4**
 Etiobiliverdin-IV γ **7.3**
 Eucatropine **8.76**
 Fatty acid methyl esters from peanut oil **17.42**
 Fatty acid methyl esters from soybean oil **17.43**
 Fatty acid methyl esters from sunflower oil **17.44**
 Ferric acetylacetone **12.174**
 Ferrocene **12.144**
 Ferrocytocrome C **10.9**
 Ferrocytocrome C oxygen oxidoreductase **10.10**
 Ferulic acid **4.70**
 Fisetin **4.75**
 Flavanone, 4',5,7-trihydroxy- **4.72**
 Flavocytochrome b₂ (Fe³⁺) **10.8**
 Flavone, 5,7-dihydroxy- **4.73**
 Flavone, 3-hydroxy- **4.74**
 Flavone, 3,3',4',7-tetrahydroxy- **4.75**
 Fluorocene, 9-diazo- **11.63**
 Fluorene, 9-(fluoren-9-ylidene)- **3.66**
 Fluorene, 9-(phenylsulfonyl)-, anion **13.98**
 9-(Fluoren-9-ylidene)fluorene **3.66**
 Fluorescein, 2',7'-dibromo-4'-(hydroxymercuri)-, disodium salt **11.64**
 Fluorescein dianion, 2',4',5',7'-tetrabromo- **11.65**
 Fluorescein dianion, 3,4,5,6-tetrachloro-2',4',5',7'-tetraido- **11.66**
 1-Fluoro-4-(butylthio)benzene **13.11**
 2-(4-Fluoro-3-nitrophenyl)-2,5,5-trimethylpyrrolidin-1-oxyl **15.55**
 4-Fluorophenoxy ion **4.165**
 1-(4-Fluorophenylazo)-2-naphthol **11.97**
 7-Fluoro-2-phenyl-1,2-benziselenazol-3-one **13.37**
 3-(4-Fluorophenyl)furan **5.41**
 4-Fluorophenyl methyl sulfide **13.21**
p-Fluorophenyl methyl sulfide **13.21**
 3-(4-Fluorophenyl)thio-1-propanol **13.134**
 4'-Fluoropivalothiophenone **13.128**
p-Fluorothioanisole **13.21**
 Formamide, *N,N*-dimethyl- **17.31**
 Fucoxanthin **2.67**
 Fullerene-C₆₀ **17.32**
 Fullerene-C₇₀ **17.33**
 2-Furaldehyde **5.9**
 2-Furaldehyde, 5-(2-hydroxyethoxy)methyl- **5.10**
 2-Furaldehyde, 5-(hydroxymethyl)- **5.11**
 2-Furancarboxaldehyde, 5-(hydroxymethyl)- **5.12**
 2-Furancarboxaldehyde, 5-methyl **5.13**

- 2-Furancarboxylic acid **5.54**
 2-Furancarboxylic acid, 5-bromo-, ethyl ester **5.55**
 2-Furancarboxylic acid, ethyl ester **5.56**
 2-Furancarboxylic acid, methyl ester **5.57**
 3-Furancarboxylic acid, 4,5-dihydro-2-methyl-, methyl ester **5.58**
2,5-Furandicarboxaldehyde 5.59
2,5-Furandicarboxylic acid, diethyl ester 5.60
2,5-Furandicarboxylic acid, dimethyl ester 5.61
3,4-Furandicarboxylic acid, diethyl ester 5.62
2,5-Furandimethanol 5.18, 10.2, 10.3, 10.23, 10.24, 10.26, 13.104, 13.106, 16.37, 16.64, 17.77
2-Furanitrile 5.63
2-Furanitrile, 5-methyl- 5.64
2-Furanmethamine, *N*-methyl- 5.65
2-Furanmethanediol, diacetate 5.66
2-Furanmethanol, α,α -diphenyl- 5.67
2-Furanmethanol, α -(diphenylmethyl)- 5.68
2-Furanmethanol, α -methyl- 5.69
2-Furanmethanol, α -phenyl- 5.70
2-Furanmethanol, α -(2-phenylethyl)- 5.71
2-Furanmethanol, α -(phenylmethyl)- 5.72
2-Furanmethanol, α -(3-phenylpropyl)- 5.73
3-Furanmethanol, 5-(phenylmethyl)- 5.74
3-Furanmethanol, 5-(phenylmethyl)-, acetate 5.75
3-Furanmethanol, 5-(phenylmethyl)-, ester with 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylic acid 5.76
2-Furanone, 3-(1-methylethylidene)- 2.227
[2,2](2,5)Furanophane 5.77
Furfural 5.9
Furfural diacetate 5.66
Furfuryl alcohol 5.78, 2.51, 4.1, 4.12, 4.40, 4.41, 4.92, 4.93, 4.106, 4.107, 4.108, 4.112, 4.113, 4.115, 4.121, 4.129, 4.130, 4.131, 4.135, 4.137, 4.138, 4.139, 4.140, 4.146, 4.150, 4.151, 4.155, 4.156, 4.157, 4.158, 4.159, 4.161, 4.163, 4.164, 4.169, 4.170, 4.171, 4.172, 4.173, 4.174, 4.176, 4.177, 4.178, 4.179, 4.182, 5.9, 5.10, 5.11, 5.12, 5.13, 5.14, 5.18, 5.32, 5.34, 5.35, 5.36, 5.46, 10.20, 10.23, 10.27, 10.34, 13.76, 13.108, 13.109, 13.111
Furfurylamine 5.79
Furfuryl methyl ether 5.43
Furo[2,3-*h*][1]benzopyran-2-one 5.1
Furo[3,2-*g*][1]benzopyran-7-one 5.86
Furo[3,2-*g*][1]benzopyran-7-one, 9-methoxy- 5.87
2-Furoic acid 5.54
2-Furonitrile 5.63
Furyl methyl ketone 5.15
Galangin 4.65
Gallium, tetranaphtho[2,3-*b*:2',3'-*g*;2'',3''-*l*:2'',3''-*q*]porphyratobis(trihexylsilanolato)- 7.25
Gentian Violet 11.44
Geraniol 2.281
Glaucine 8.21
Glucose 16.34
Glutamylcysteinylglycine, negative ion 13.100
 γ -L-Glutamyl-L-cysteinylglycine 13.99
Glutathione 13.99
Glutathione, negative ion 13.100
Glycero-3-phosphocholine 17.53
Glycine 10.15
Glycine, glycyl- 10.16
Glycine, glycylglycyl- 10.17
Glycine, glycylglycylglycyl- 10.18
Glycine, glycyl-L-histidyl- 10.19
Glycine, histidyl- 10.20
Glycine, *N*-(2-mercaptop-1-oxopropyl)- 13.101
Glycine, *N*-(2-mercaptop-1-oxopropyl)-, negative ion 13.102
Glycine, tryptophyl- 10.21
Glycine, L-tryptophylglycyl- 10.22
Glycine, L-tyrosyl- 4.76
Glycylglycine 10.16
Glycylglycylglycine 10.17
Glycylglycylglycylglycine 10.18
Glycylglycyl-L-histidine 10.28
Glycyldhistidine 10.27
Glycyl-L-histidylglycine 10.19
Glycylmethionine 13.109
Glycyltryptophan 10.41
Glycyltyrosine 4.189
Glycyl-L-tyrosyl-L-alanine 4.1
Gold(I) methanethiol triethylphosphine complex 12.159
Gold(I) tetra-*O*-acetylglucose-1-thiol triethylphosphine complex 12.170
Guaiacol 4.129
Guanine 16.35
Guanosine 16.36
Guanosine, 2'-deoxy- 16.37
Guanosine monophosphate 16.38
Guanosine 5'-monophosphate 16.38
5'-Guanylic acid 16.38
Helianthrene 3.67
Hematoporphyrin derivative 7.14
Hematoporphyrin-human serum albumin complex 7.15
Hematoporphyrin IX 7.72
1,6-Heptadiene-3,5-dione, 1,7-bis(4-hydroxy-3-methoxyphenyl)- 4.77
Heptane 17.34
1,7-Heptanediamine, *N,N*-diethyl- 8.31
3,5-Heptanedione, 4-[[2-ethyl-4-(diethylamino)phenyl]imino]-2,2,6,6-tetramethyl- 11.67
1-Heptene 2.228
cis-2-Heptene 2.229

- (Z)-2-Heptene 2.229
 Heterocoerdianthrone 3.68
 Hexaamminecobalt(III) ion 12.145
 $(all-E)$ -1,3,5,7,9,11,13,15,17,19,21,23,25-
 Hexacosatridecaene, 3,7,11,16,20,24-hexamethyl-
 1,26-bis(2,6,6-trimethyl-1-cyclohexyl- 2.230
 2,4,6,8,10,12,14-Hexadecaheptaene-1,16-dione, 1,16-bis(3-
 hydroxy-1,2,2-trimethylcyclopentyl)-6,11-dimethyl-
 [1R-[1 α [2E,4E,6E,8E,10E,12E,14E,16(1R*,3S*)],3 β]-
 2.231
 2,4,6,8,10,12,14-Hexadecaheptaene-1,16-dione, 1,16-bis(4-
 hydroxy-1,2,2-trimethylcyclopentyl)-6,11-dimethyl-
 [1R-[1 α [2E,4E,6E,8E,10E,12E,14E,16(1R*,4S*)],4 β]-
 2.232
 Hexadecanoic acid 17.37
 4,6,8,10,12-Hexadecapentaene-3,14-dione, 6,11-dimethyl-
 2,2,15,15-tetramethoxy- ($all-E$) 2.233
 4,6,8,10,12-Hexadecapentaene-2,3,14,15-tetraone, 6,11-
 dimethyl- ($all-E$) 2.234
 2,6,10,14-Hexadecatetraene, 2,6,10,14-tetramethyl- 2.235
 2-Hexadecene, 1-ethoxy-3,7,11,15-tetramethyl- 2.236
 2-Hexadecene, 1-methoxy-3,7,11,15-tetramethyl- 2.237
 2-Hexadecen-1-ol, 3,7,11,15-tetramethyl- 2.238
 2-Hexadecen-1-ol, 3,7,11,15-tetramethyl-, acetate 2.239
 1,5-Hexadiene 2.240
 2,4-Hexadiene, 2,5-dimethyl- 2.243, 13.87, 13.88
 cis-2,trans-4-Hexadiene 2.242
 (E,E)-2,4-Hexadiene 2.241
 (E,Z)-2,4-Hexadiene 2.242
 trans,trans-2,4-Hexadiene 2.241
 3,4,5,6,7,8-Hexahydro-1-benzopyran-2-one 2.92
 1,2,3,10,11,12-Hexahydroindeno[5,6-a:5',6'-j]perylene-
 8,17-dione 3.65
 Hexahydro-1,2-dimethylpyridazine 8.91
 cis-(-)-2,3,4,4 α ,5,9 b -Hexahydro-2,8-dimethylpyrido[4,3-
 b]indole 6.56
 1,2,3,8,9,10-Hexahydro-3,3,5,6,8,8-hexamethylbenzo[1,2-
 b:4,3-b']dipyran 5.2
 2,3,4,7,8,9-Hexahydro-2,2,5,7,7,10-hexamethylbenzo[1,2-
 b:4,5-b']dipyran 5.3
 Hexahydropyrazolo[1,2- a][1,2]diazepine 8.88
 3,4,6,7,9,10-Hexahydro-3,3,6,6-tetramethyl-1,8-
 acridinedione 6.1
 Hexamethoxybenzene 3.42
 1,2,3,4,5,6-Hexamethylbicyclo[2.2.0]hexa-2,5-diene 2.29
 Hexamethylbicyclo[2.2.0]hexa-2,5-diene 2.29
 Hexamethyl Co α .Co β -dicyano-7-de(carboxymethyl)-7,8-
 didehydrocobyminate 16.16
 Hexamethyldewarbenzene 2.29
 Hexamethylenedithiocarbamate 13.1
 Hexamethylenetetramine 8.32
 1,3,3,1',3',3'-Hexamethyl-8,10-indolo-2,2'-carbocyanine
 iodide 11.40
 1,1',3,3,3',3'-Hexamethylindotricarbocyanine iodide
 11.182
 1,1',3,3,3',3'-Hexamethylindotricarbocyanine perchlorate
 11.183
 2,6,10,15,19,23-Hexamethyl-2,6,10,14,18,22-
 tetracosahexaene ($all-E$) 2.346
 1,1',3,3,3',3'-Hexamethyltricarbocyanine, fluoride 11.181
 1,1',3,3,3',3'-Hexamethyltricarbocyanine, iodide 11.182
 1,1',3,3,3',3'-Hexamethyltricarbocyanine, perchlorate
 11.183
 Hexanoic acid 17.38
 1-Hexanol, 6-(4-methylphenyl)thio- 13.103
 Hexaphenylisobenzofuran 5.84
 2-Hexene 2.244, 2.116
 cis-3-Hexene 2.246
 (E)-3-Hexene 2.245
 trans-3-Hexene 2.245
 (Z)-3-Hexene 2.246
 Histamine 6.5
 Histidine 10.23, 4.10, 4.11, 4.12, 4.20, 4.35, 4.111, 4.153, 4.154,
 4.155, 4.156, 4.157, 4.162, 4.175, 4.176, 4.177, 4.178, 5.81,
 6.20, 6.22, 6.26, 6.29, 10.29, 10.36
 Histidine, β -alanyl-1-methyl- 10.25
 Histidine, glycyl- 10.27
 L-Histidine, β -alanyl- 10.24
 L-Histidine, N-(4-amino-1-oxobutyl)- 10.26
 L-Histidine, glycylglycyl- 10.28
 Histidinyl- β -alanine 10.3
 Histidylglycine 10.20
 Homocarnosine 10.26
 Hydrazine, 1,2-bis(1-methylethylidene)- 15.3, 2.40
 Hydrazine, 1,2-diethyl-1,2-dimethyl- 8.33
 Hydrazine, 1,2-dimethyl-1,2-dibutyl- 8.34
 Hydrazine, 1,1-dimethyl-2,2-di(2,2-dimethylpropyl)- 8.35
 Hydrazine, 1,2-dimethyl-1,2-di(2,2-dimethylpropyl)- 8.36
 Hydrazine, 1,1-dimethyl-2,2-di(2-methylpropyl)- 8.37
 Hydrazine, 1,2-dimethyl-1,2-di(2-methylpropyl)- 8.38
 Hydrazine, 1,2-dimethyl-1,2-dipentyl- 8.39
 Hydrazine, 1,2-dimethyl-1,2-dipropyl- 8.40
 Hydrazine, tetramethyl- 8.41
 Hydrazine, tetra(methyl- d_3)- 8.42
 Hydrazine, tetra(2-methylpropyl)- 8.43
 Hydrazine, 1,1,2-trimethyl-2-(2,2-dimethylpropyl)- 8.44
 Hydrazine, 1,1,2-trimethyl-2-(2-methylpropyl)- 8.45
 Hydrazine hydrate 14.7
 Hydrogen azide 14.3
 Hydroperoxide, 2,5-cyclooctadien-1-yl- 2.130
 Hydroperoxide, 1,1,6-trimethyl-2,5-heptadienyl- 2.274
 6-Hydroperoxy-1,4-cyclooctadiene 2.130
 Hydroperoxy-2,6-dimethyl-2,6-octadiene 2.278
 6-Hydroperoxy-2,7-dimethyl-2,7-octadiene 2.279
 7-Hydroperoxy-2,7-dimethyl-2,5-octadiene 2.274

5-Hydroperoxy-4-methyl-3-octene	2.286	Hydroxyethylvinyldeuteroporphyrin	7.92
Hydroperoxy-2,6,10-trimethyl-2,6,10-dodecatriene	2.209	3-Hydroxyflavone	4.74
Hydroquinone	4.12	5-Hydroxyindole	6.12
Hydroquinone, chloro-	4.20	Hydroxylamine, <i>N,N</i> -dibenzyl-	15.11
Hydroquinone, 2,5-di- <i>tert</i> -butyl-	4.14	Hydroxylamine, <i>N,N</i> -diethyl-	15.23
Hydroquinone, 2,5-di(2,2-dimethylpropyl)-	4.16	Hydroxylamine, <i>N,N</i> -diethyl-, conjugate acid	15.24
Hydroquinone, 2,5-di- <i>sec</i> -dodecyl-	4.17	4-Hydroxy-3-methoxycinnamic acid	4.70
Hydroquinone, 2,5-di- <i>sec</i> -hexadecyl-	4.18	17-Hydroxy-4-methyl-4-androsten-3-one	16.4
Hydroquinone, 2,3-dimethoxy-5-methyl-	4.21	2-Hydroxy-3-methyl-2-cyclohexen-1-one	2.128
Hydroquinone, 2,3-dimethyl-	4.22	4-Hydroxy-6-methyl-2-(dimethylamino)pyrimidine	16.70
Hydroquinone, 2,5-dimethyl-	4.23	5-(Hydroxymethyl)furan-2-carboxaldehyde	5.11
Hydroquinone, 2,6-dimethyl-	4.24	5-(Hydroxymethyl)furfural	5.11
Hydroquinone, 2,5-di- <i>sec</i> -octyl-	4.19	4-Hydroxymethyl-2-(phenylmethyl)furan	5.74
Hydroquinone, trimethyl-	4.31	3-Hydroxy- <i>N</i> -methylpiperidinc	8.70
Hydroquinone dianion	4.13	4-Hydroxy- <i>N</i> -methylpiperidine	8.71
Hydroquinone dimethyl ether	3.40	<i>N</i> -[5-Hydroxy-8-[(2-(methylsulfonyl)-4-nitrophenyl]azo]-1-naphthalenyl-1,3-benzenedisulfonamide, conjugate base	11.85
Hydroxide- <i>d</i> ion	14.8	<i>N</i> -[5-Hydroxy-8-[(2-(methylsulfonyl)-4-nitrophenyl]azo]-1-naphthalenyl-1,3-benzenedisulfonamide, conjugate dibase	11.86
4'-Hydroxyacetophenone	4.93	2-Hydroxynaphthalene	4.89
<i>p</i> -Hydroxyacetophenone	4.93	5-Hydroxy-1,4-naphthoquinone	4.90
Hydroxyaluminum(III) tetra(1-phenyl)-2,3-naphthalocyanine	7.23	6-Hydroxy-1,4-naphthoquinone	4.91
2-Hydroxybenzoate ion	4.35	4-Hydroxy-4-(2-naphthyl)-2,2,6,6-tetramethylpiperidine-1-oxyl	15.31
4-Hydroxybenzonitrile	4.112	2-Hydroxy-4-octyloxybenzophenone	4.43
2-Hydroxybiphenyl	4.141	6-Hydroxy-2,2,5,7,8-pentamethylchroman	4.49
2-Hydroxybiphenyl, conjugate base	4.180	4-Hydroxy-1,2,2,6,6-pentamethylpiperidine	8.72
4-Hydroxybiphenyl, conjugate base	4.181	1-(4-Hydroxyphenylazo)-2-naphthol	11.98
Hydroxybis(dimethylglyoximato)pyridinecobalt(II)	12.146	4-Hydroxy-4-(2-phenylethynyl)-2,2,6,6-tetramethylpiperidine- <i>N</i> -oxyl	15.32
Hydroxybis(dimethylglyoximato)triethylaminecobalt(II)	12.147	<i>N</i> -[(2-Hydroxyphenyl)methylene]benzenamine, <i>N</i> -oxide	15.7
Hydroxybis(dimethylglyoximato)triphenylphosphinecobalt(II)	12.148	2-Hydroxyphenyl phenyl nitrone	15.7
Hydroxybis(diphenylglyoximato)pyridinecobalt(II)	12.149	3-(<i>p</i> -Hydroxyphenyl)propionic acid	4.34
Hydroxybis(diphenylglyoximato)triphenylphosphinecobalt(II)	12.150	<i>N</i> -(3-Hydroxypropyl)- <i>N,N</i> -dimethylamine	8.82
D(-)-β-Hydroxybutyrate dehydrogenase	10.11	8-Hydroxyquinoline	6.73
2-(2'-Hydroxy-3'-chloro-5'- <i>tert</i> -butylphenyl)benzotriazole	4.97	6-Hydroxy-2,5,7,8-tetramethylchroman-2-carboxylate ion	4.44
2-Hydroxycinnamic acid	4.69	3-(6-Hydroxy-2,5,7,8-tetramethylchroman-2-yl)propionic acid	4.66
<i>o</i> -Hydroxycinnamic acid	4.69	4-Hydroxy-2,2,6,6-tetramethylpiperidine	8.77, 8.19
6-Hydroxy-1,4-cyclooctadiene	2.134	4-Hydroxy-2,2,6,6-tetramethylpiperidine-1-oxyl	15.33
4-Hydroxy-5,6-dimethyl-2-(dimethylamino)pyrimidine	16.69	4-Hydroxy-2,2,6,6-tetramethylpiperidine- <i>N</i> -oxyl	15.33
3-Hydroxy-3,4-dimethylpentyl <i>p</i> -methylphenyl sulfide	13.119	<i>p</i> -Hydroxythioanisole	13.120
4-(1-Hydroxy-2,2-dimethyl)propyl-2-methoxyphenol	4.127	3-Hydroxytyramine	4.5
2-(2'-Hydroxy-3',5'-di- <i>tert</i> -pentylphenyl)benzotriazole	4.96	<i>dl</i> -Hyoscyamine	8.1
3-Hydroxydiphenylamine	4.142	Hypericin	16.60
3-Hydroxy-1,3,5(10),9(11)-estratetraen-17-one	16.31	Imidazole	6.6
5-(2-Hydroxyethoxy)methylfuran-2-carboxaldehyde	5.10	Imidazole, 4-methyl-	6.7
2-Hydroxyethylsulfide ion	13.96	Imidazole-4-ethanamine	6.5
<i>N</i> -(2-Hydroxyethyl)-2,2,6,6-tetramethylpiperidine	8.68		

- Imidazole-4-ethanaminium, α -carboxy-2,3-dihydro-*N,N,N*-trimethyl-2-thioxo-, *S*, inner salt **13.104**
- Imidazole-4-ethanaminium, α -carboxy-2,3-dihydro-*N,N,N*-trimethyl-2-thioxo-, ion **13.105**
- Imidazole-4-propionic acid, α -amino-2,3-dihydro-2-thioxo-, (*S*) **13.106**
- Imidazo[1,2-*a*]pyrazin-3-one, 4-methyl-6-[4-[2-[3-carboxy-4-(6-hydroxy-3-xanthenon-9-yl)phenylthiocarbamylethoxy]phenyl]- **16.39**
- Imidazo[1,2-*a*]pyrazin-3-one, 2-methyl-6-(4-methoxyphenyl)- **16.40**
- Imidazo[1,2-*a*]pyrazin-3-one, 2-methyl-6-phenyl- **16.41**
- 5-Iminodaunomycin **4.81**
- Indene **3.69**
- Indene, 1,2-diphenyl- **3.70**
- Indene, 1-methyl-2-phenyl- **3.71**
- Indene, 3-methyl-2-phenyl- **3.71**
- Indene, 1,1'-spirobis[5,6-di(methoxycarbonyl)-3,3-dimethyl- **3.98**
- Indene-1,3-dione, 2-(2-quinolylidene)- **11.68**
- Indigo **11.69**
- Indole **6.8**
- Indole, 5,6-dimethoxy- **6.9**
- Indole, 1,3-dimethyl- **6.10**
- Indole, 2,3-dimethyl- **6.11**
- Indole, 5-hydroxy- **6.12**
- Indole, 5-methoxy- **6.13**
- Indole, 5-methoxy-1,2,3-trimethyl- **6.14**
- Indole, 1-methyl- **6.15**
- Indole, 3-methyl- **6.16**
- Indole, 1-methyl-3-(β -methoxyvinyl)- (*E*) **2.247**, **2.248**
- Indole, 1-methyl-3-(β -methoxyvinyl)- (*Z*) **2.248**
- Indole, 3-methyl-2-phenyl- **6.17**
- Indole, 1-(phenylsulfonyl)-3-(β -methoxyvinyl)- (*E*) **2.247**, **2.249**
- Indole, 1-(phenylsulfonyl)-3-(β -methoxyvinyl)- (*Z*) **2.249**
- Indole, 1,2,3-trimethyl- **6.18**
- Indole, 1,2,3-trimethyl-5-nitro- **6.19**
- Indole-3-acetate ion **6.20**
- Indole-3-acetic acid **6.21**
- Indole-3-butyrate ion **6.22**
- Indole-3-butyrlic acid **6.23**
- 3-Indolebutyric acid **6.23**
- Indole-3-carbinol **6.24**
- Indole-3-carboxylate ion **6.26**
- Indole-2-carboxylic acid **6.25**
- Indole-3-carboxylic acid **6.27**
- 3-Indolemethanol **6.24**
- Indole-3-propanamide, α -amino- **10.43**
- Indole-3-propionamide **6.28**
- Indole-3-propionate ion **6.29**
- Indole-3-propionic acid **6.30**
- Indolium, 2-[7-[4-bromo-1-(1,3,3-trimethyl-2-indol-2-ylidene)]-1,3,5-heptatrienyl]-1,3,3-trimetetrafluoroborate **11.70**
- Indolium, 5-chloro-1-[7-(5-chloro-1,3-dihydro)-1,3,3-trimethylindol-2-ylidene)-1,3,5-heptatrienyl]-, iodide **11.178**
- Indolium, 2-[7-[4-chloro-1-(1,3,3-trimethyl-2-indol-2-ylidene)]-1,3,5-heptatrienyl]-1,3,3-trimethyl-tetrafluoroborate **11.71**
- Indolium, 1-[7-(1,3-dihydro)-1,3,3,5-tetramethylindol-2-ylidene]-1,3,5-heptatrienyl]-, iodide **11.184**
- Indolium, 1-[7-(1,3-dihydro)-1,3,3-trimethylindol-2-ylidene]-1,3,5-heptatrienyl]-, fluoride **11.181**
- Indolium, 1-[7-(1,3-dihydro)-1,3,3-trimethylindol-2-ylidene)-1,3,5-heptatrienyl]-, iodide **11.182**
- Indolium, 1-[7-(1,3-dihydro)-1,3,3-trimethylindol-2-ylidene)-1,3,5-heptatrienyl]-, perchlorate **11.183**
- Indolium, 5-fluoro-1-[7-(5-fluoro-1,3-dihydro)-1,3,3-trimethylindol-2-ylidene)-1,3,5-heptatrienyl]-, iodide **11.179**
- Indolium, 2-[7-[4-iodo-1-(1,3,3-trimethyl-2-indol-2-ylidene)]-1,3,5-heptatrienyl]-1,3,3-trimethyl-, iodide **11.72**
- Indolium, 2-[7-[4-iodo-1-(1,3,3-trimethyl-2-indol-2-ylidene)]-1,3,5-heptatrienyl]-1,3,3-trimethyl-, tetrafluoroborate **11.73**
- Indolium, 5-methoxy-1-[7-(5-methoxy-1,3-dihydro)-1,3,3-trimethylindol-2-ylidene)-1,3,5-heptatrienyl]-, iodide **11.180**
- Indolium, 2-[7-(1,3,3-trimethyl-2-indol-2-ylidene)-1-[4-(2,2-dimethoxyethyl)-1,3,5-heptatrienyl]-1,3,3-trimethyl-, tetrafluoroborate **11.74**
- Indolium, 2-[7-(1,3,3-trimethyl-2-indol-2-ylidene)-1-[4-(1-piperidino)-1,3,5-heptatrienyl]-1,3,3-trimethyl-bis(tetrafluoroborate) **11.75**
- Indolium, 2-[7-(1,3,3-trimethyl-2-indol-2-ylidene)-1-[4-(1,3,3-trimethyl-2-indolylidene)-2-propenyl]-1,3,3-heptatrienyl]-1,1,3-trimethyl-, tetrafluoroborate **11.76**
- 5-Indolol **6.12**
- Inosine diphosphate, ester with 1,4-dihydro-1- β -D-ribofuranosyl-3-pyridinecarboxamide **16.42**
- Iodide ion **14.9**
- Iodine **14.10**
- Iodoethane **17.28**
- 4-Iodophenoxy ion **4.168**
- 1-(4-Iodophenylazo)-2-naphthol **11.99**
- 4-Iodo-2,2,6,6-tetramethylpiperidin-1-oxyl **15.42**
- 2-[7-(4-Iodo-1,3,3-trimethyl-2-indol-2-ylidene)-1,3,5-heptatrienyl]-1,3,3-trimethylindolium iodide **11.72**
- 2-[7-(4-Iodo-1,3,3-trimethyl-2-indol-2-ylidene)-1,3,5-heptatrienyl]-1,3,3-trimethylindolium tetrafluoroborate **11.73**
- β -Ionone **2.61**

- IR 125 **11.29**
 IR 132 **11.108**
 IR 140 **11.33**
 IR 144 **11.30**
 Iridium(I) bromocarbonylbis(triphenylphosphine) **12.119**
 Iridium(I) carbonylchlorobis(triphenylphosphine) **12.123**
 Iridium(I) carbonyliodobis(triphenylphosphine) **12.125**
 Iron(III) (acetato)mesoporphyrin IX, dimethyl ester **7.74**
 Iron(III) (acetato)5,10,15,20-tetraphenylporphyrin **7.47**
 Iron(III) bis(diisopropylidithiocarbamate) **12.48**
 Iron(III) (chloro)5,10,15,20-tetraphenylporphyrin **7.49**
 Iron(II) 5,10,15,20-tetraphenylporphyrin, μ -oxobis- **7.69**
 Iron(III) (thiocyanato-S)5,10,15,20-tetraphenylporphyrin
 7.70
 Iron(III) tris(acetylacetone) **12.174**
 Isobenzofuran **5.80**
 Isobenzofuran, 1,3-bis[4-(11-carboxynonyl)phenyl]-4,7-dihydro-5,6-dimethyl-, dianion **5.81**
 Isobenzofuran, 5,6-dimethyl-1,3-diphenyl- **5.82**
 Isobenzofuran, 1,3-diphenyl- **5.83**, *2.51, 3.60, 3.96, 3.98, 4.49, 4.54, 4.184, 4.185, 6.39, 7.5, 7.78, 8.19, 10.23, 10.34, 10.35, 10.36, 10.38, 10.42, 11.33, 11.37, 11.38, 11.70, 11.71, 11.72, 11.73, 11.75, 11.76, 11.109, 11.174, 11.178, 11.179, 11.180, 11.181, 11.182, 11.183, 11.184, 12.20, 12.21, 12.22, 12.23, 12.84, 12.138, 13.63, 13.64, 13.108, 13.169, 13.170, 13.171, 15.1, 17.30*
 Isobenzofuran, hexaphenyl- **5.84**
 Isobenzofuran, 1,3,4,7-tetraphenyl- **5.85**
 Isobutylamine **8.85**
 Isobutylene **2.337**
 2-Isobutylideneadamantane **2.355**
 Isobutyl sulfide **13.77**
 (-)-Isocaryophyllene. **2.34**
 Isodicyclopentadiene **2.254**
 Isoindole, 1,1,3-triphenyl-, 2-oxide **15.25**
 Isoprene **2.42**
 Isopropanol **17.61**
 Isopropyl alcohol **17.61**
 Isopropylamine **8.28**
 1-Isopropylamino-4-phenylaminobenzene **9.30**
 6-Isopropyl-3,4-dihydropyran-5-carboxylic acid ethyl ester
 5.99
 2-Isopropylideneadamantane **2.353**
 2-(Isopropylidene)cyclopentanone **2.155**
 Isopropylidenecyclopropane **2.170**
 3-Isopropylidene-2-furanone **2.227**
 1-Isopropyl-4-methyl-1,3-cyclohexadiene **2.105**
 4-Isopropyl-1-methylcyclohexene **2.125**
 Isopropyl sulfide **13.78**
 Isopsoralen **5.1**
 Isostilbene **3.99**
 Isozeaxanthin **2.81**
 Juglone **4.90**
 Kaempferol **4.64**
 Ketene diethyl acetal **2.221**
 2-Ketoglutaric acid **17.51**
 Kryptocyanine cation **11.42**
 Lactate dehydrogenase (cytochrome) **10.8**
 Lauric acid **17.24**
 Lead(II) bis(*O,O'*-diphenylidithiophosphate) **12.70**
 Lead(II) bis[2,2'-thiobis[*O,O'*-di(4-*tert*-butylphenyl)dithiophosphate]] **12.113**
 Lead(II) 2,2'-methylenebis[*O,O'*-di(4-*tert*-butylphenyl)dithiophosphate] **12.157**
 Lecithin **17.53**
 L-L Leucine, L-methionyl- **13.107**
 Leucomalachite Green **11.77**
 Limonene **2.124, 13.8, 13.52, 13.97, 13.119, 13.136**
 Linalool **2.280, 2.338, 13.53, 13.54, 13.55, 13.143, 13.144**
 Linoleic acid **2.264**
 Linolenic acid **2.267**
 Linolenic acid, phenyl ester **2.270**
 Lipoic acid **13.90, 3.125**
 Luciferase **10.29**
 Luminofor Red-Violet 440RT **11.121, 11.122, 11.123, 11.124, 11.125, 11.126, 11.127**
 Luminol **9.33**
 Luminor 8 **11.121, 11.122, 11.123, 11.124, 11.125, 11.126, 11.127**
 2,6-Lupetidine **8.58**
 Lutein **2.75**
 (*all-E*)-Lutein **2.75**
all-trans-Lutein **2.75**
 Luteolin **4.60**
 Lycopene **2.250**
 Lycopene, (*all-E*)- **2.250**
 (*all-E*)-Lycopene **2.250**
all-trans-Lycopene **2.250**
 Lycopene, dihydroxy- **2.251**
 Lysine decarboxylase (*B. cadaveris*) **10.30**
 Lysozyme **10.30**
 LysRSSR **10.30**
 Magenta dye 9 **11.154**
 Magnesium(II) tetraphenylporphyrin **7.65**
 Magnesium(II) 5,10,15,20-tetraphenylporphyrin **7.65**
 Malonic acid **17.39**
 Manganese(III) (acetato)5,10,15,20-tetraphenylporphyrin
 7.48
 Manganese(II) bis(diisopropylidithiocarbamate) **12.49**
 Manganese(II) chloride hexahydrate **12.151**
 Manganese(II) chloride tetrhydrate **12.152**
 Manganese(III) mesoporphyrin diethyl ester **7.84**
 Manganese(III) 5,10,15,20-tetraphenylporphyrin **7.66**
 Manganese(III) tris(acetylacetone) **12.175**

- Melanin (from DOPA) **4.78**
 Menadione **16.44**
 Menaquinone **16.44**
p-Menth-1-ene **2.125**
 Merbromin **11.64**
 2-Mercaptoethanesulfonate ion **13.92**
 2-Mercaptoethanol **13.95**
 2-Mercaptoethanol, negative ion **13.96**
 (*S*)-1-(3-Mercapto-2-methyl-1-oxopropyl)-L-proline **13.130**
 2-Mercaptopropionylglycine **13.101**
 2-Mercaptopropionylglycine, negative ion **13.102**
N-(2-Mercaptopropionyl)glycine **13.101**
 Mercurochrome **11.64**
 Merocyanine 540 **11.39**
 Mesitol **4.148**
 Mesobilirubin IX **7.9**
 Mesodiphenylbenzheianthrene **3.129**
 Mesodiphenylhelianthrene **3.64**
 Mesoporphyrin di[4-(diphenylmethylaminocarbonyl-2-nitrophenylmethyl)ester] **7.83**
 Mesoporphyrin diester with *N*-benzhydryl-4-bromomethyl-3-nitrobenzamide **7.83**
 Mesoporphyrin IX, dimethyl ester **7.85**
 Methacrylic acid **2.252**
 Methanediol, 2-furanyl-, diacetate **5.66**
 Methanesulfonamide, *N*-[2-[4-(1,5-dihydro-3-methyl-5-oxo-1-phenylpyrazol-4-ylidene)amino]-methylphenyl]ethylamino]ethyl- **11.78**
 Methanethione, bis(4-chlorophenyl)- **13.159**
 Methanethione, bis(4-methylphenyl)- **13.161**
 Methanethione, (4-chlorophenyl)phenyl- **13.158**
 Methanethione, (4-methoxyphenyl)phenyl- **13.162**
 Methanethione, (2,2,6,6-tetramethylcyclohexylidene)-
13.166
 2,5-Methano-1,3-dithiole[4,5-*d*][1,3]dithiin, dihydro-2,3a,5-trimethyl-7-methylene- **2.253**
 4,7-Methanoindene, 4,5,6,7-tetrahydro- **2.254**
 Methanol **17.40**
 Methanol, cyclohexylidene-, acetate **2.255**
 Methanol, 1-tricyclo[3.3.1.1^{3,7}]decylidene-, acetate **2.256**,
2.126
 1,4-Methanonaphthalene, 5,8-dimethoxy-1,2,3,4-tetrahydro-9-isopropylidene- **2.257**
 1,4-Methanonaphthalene, 5,6,7,8-tetrachloro-1,2,3,4-tetrahydro-9-isopropylidene- **2.258**
 1,4-Methanonaphthalene, 5,6,7,8-tetrafluoro-1,2,3,4-tetrahydro-9-isopropylidene- **2.259**
 1,4-Methanonaphthalene, 1,2,3,4-tetrahydro- **3.72**
 1,4-Methanonaphthalene, 1,2,3,4-tetrahydro-5,8-dimethoxy- **3.73**
 1,4-Methanonaphthalene, 1,2,3,4-tetrahydro-9-isopropylidene- **2.260**
 5,8-Methano-[1,2,4]triazolo[1,2-*a*]pyridazine-1,3-dione,
 tetrahydro-2-methyl- **17.41**
 Methenamine **8.32**
 Methionine **13.108**, **4.30**, **4.76**, **4.92**, **4.152**, **4.153**, **4.157**, **4.158**,
4.160, **4.165**, **4.166**, **4.168**, **4.180**, **4.181**, **4.182**, **4.187**, **4.188**,
10.21, **10.36**, **10.41**
 Methionine, glycyl- **13.109**
 Methionine, *S*-methyl-*N*-(phenylmethyl)carbonyl-, methylene
 ester **13.110**
 Methionine methyl ester **13.111**
 L-Methionyl-L-leucine **13.107**
 Methone **17.15**
 Methoxsalen **5.87**
 9-Methoxyanthracene **3.12**
 Methoxybenzene **3.43**
 2-Methoxybicyclo[2.2.1]cycloheptene **2.20**
 2-Methoxybiphenyl **3.59**
o-Methoxybiphenyl **3.59**
 2-Methoxy-2-butene **2.55**
 1-Methoxy-4-(butylthio)benzene **13.12**
 2-(Methoxycarbonyl)phenoxide ion **4.41**
N-Methoxycarbonyltryptamine **10.35**
 6-Methoxy-1,4-cyclooctadiene **2.131**
 1-Methoxycyclopentene **2.160**
 3-Methoxy-4,6-di-*tert*-butylphenol **4.100**
 2-Methoxy-*N,N*-dimethylaniline **9.11**
 3-Methoxy-*N,N*-dimethylaniline **9.12**
N-Methoxy-*N,N*-dimethylaniline **9.13**
 4-Methoxy-2,6-dimethylphenol **4.132**
 3-Methoxyestra-1,3,5(10),8-tetraen-17-one **16.28**
 3-Methoxy-1,3,5(10),9(11)-estratetraen-17-one **16.32**,
16.29, **16.30**, **16.31**, **16.33**
 2-Methoxyfuran **5.42**
 9-Methoxyfuro[3,2-*g*][1]benzopyran-7-one **5.87**
 5-Methoxyindole **6.13**
 3-Methoxy-2-methyl-2-cyclopenten-1-one **2.167**
 5-Methoxymethyl-2-(dimethoxymethyl)furan **5.32**
 5-(Methoxymethyl)furan-2-carboxaldehyde **5.12**
 5-Methoxy-4-[2-(methylsulfonyl)-4-nitrophenyl]azo-1-naphthoxide ion **11.88**
 1-Methoxy-2-nitrobenzene **3.44**
 2-Methoxynorbornene **2.20**
 2-Methoxyphenol **4.129**
 3-Methoxyphenol **4.130**
 4-Methoxyphenol **4.131**
 2-Methoxyphenoxide ion **4.169**
 3-Methoxyphenoxide ion **4.170**
 4-Methoxyphenoxide ion **4.171**
 2-(4-Methoxyphenoxy)methylidene)adamantane **2.352**
 1-Methoxy-4-(phenylazo)naphthalene **11.59**
 1-(4-Methoxyphenylazo)-2-naphthol **11.100**
 4-(4-Methoxyphenylazo)-1-naphthol **11.89**

- 7-Methoxy-2-phenyl-1,2-benziselenazol-3-one **13.38**
 3-(4-Methoxyphenyl)-2-cyclopentene-1-thione **13.66**
 3-(4-Methoxyphenyl)-1,5-diphenyl-2-pyrazoline **11.124**
 5-(4-Methoxyphenyl)-1,3-diphenyl-2-pyrazoline **11.125**
 2-(4-Methoxyphenyl)furan **5.44**
 3-(4-Methoxyphenyl)furan **5.45**
 4-(4'-Methoxyphenyl)imino-3-methyl-1-phenyl-2-pyrazolin-5-one **11.148**
 4-Methoxyphenyl methyl sulfide **13.22**
 1-(4-Methoxyphenyl)-4-phenyl-1,3-cyclopentadiene **2.148**
 2-(4-Methoxyphenyl)-3-phenyl-1,4-dioxene **2.195**
 (4-Methoxyphenyl)phenylmethanethione **13.162**
 4-Methoxyphenyl phenyl sulfide **13.23**
 1-(4-Methoxyphenyl)-4,4,8,8-tetramethyl-2,3,4,5,7,8,9,10-octahydpyrrolo[4,3,2-*m,n*]acridine-10-one **6.68**
 1-Methoxy-4-(phenylthio)benzene **13.23**
 3-(4-Methoxyphenyl)thio-1-propanol **13.135**
 4'-Methoxypivalothiophenone **13.129**
 3-Methoxypsoralen **5.87**
 6-Methoxyquinoline **6.74**
 4-Methoxy-2,3,5,6-tetramethylphenol **4.133**
p-Methoxythioanisole **13.22**
 4-Methoxythiobenzamide **13.154**
 4-Methoxythiobenzophenone **13.162**
 5-Methoxy-1,2,3-trimethylindole **6.14**
 4-Methoxy-2,3,6-trimethylphenol **4.134**
m-Methoxy- α,β,β -trimethylstyrene **3.117**
p-Methoxy- α,β,β -trimethylstyrene **3.118, 3.115, 3.123**
 Methyl acetal of oxidized octaethylpurpurin ethyl ester **7.16**
 2-Methylacetophenone enol **2.219**
 Methyl alcohol **17.40**
 Methylamine, *N,N*-bis(2-hydroxyethyl)- **8.46**
 Methylamine, *N,N*-dimethyl- **8.47**
 1-Methylamino-9,10-diphenylanthracene **9.20**
 Methyl *tert*-amyl ether **17.19**
N-Methylaniline **9.14**
 9-Methylanthracene **3.13**
 Methyl arachidonate **2.213**
 [4-Methyl-1,2-benzenedithiolato](2,2'-bipyridine)nickel(II) **12.153**
 [4-Methyl-1,2-benzenedithiolato](1,10-phenanthroline)nickel(II) **12.154**
 [4-Methyl-1,2-benzenedithiolato](1,10-phenanthroline)platinum(II) **12.155**
 Methyl benzoate **3.57**
 Methyl-1,4-benzoquinone **16.13**
 α -Methylbenzyl alcohol **3.55**
 Methylbixin **2.176**
trans-Methylbixin **2.175**
 Methyl 4-bromophenyl sulfide **13.7**
 2-Methyl-1,3-butadiene **2.42**
 2-Methyl-2-butene **2.56, 2.122, 2.161, 2.307, 2.309, 2.310, 3.110, 15.2, 15.13, 15.14, 15.15**
 (*E*)-2-Methyl-2-butenoic acid **2.58**
 3-Methyl-2-buten-1-ol **2.60**
 Methyl *tert*-butyl ether **17.18**
 Methyl 4-*tert*-butylphenyl sulfide **13.20**
 Methyl butyl sulfide **13.57**
 1-Methyl-4-(butylthio)benzene **13.13**
 2-Methyl-6-[4-[2-[3-carboxy-4-(6-hydroxy-3-xanthenon-9-yl)phenylthiocarbamylethoxy]-phenyl]imidazo[1,2-*a*]pyrazin-3-one **16.39**
 Methylcercosporin **16.58**
 4-Methyl-2-(3-chlorophenyl)-5-phenyloxazole **6.33**
 4-Methyl-2-(4-chlorophenyl)-5-phenyloxazole **6.34**
 Methyl 3-chlorophenyl sulfide **13.14**
 3-Methyl-2-[7-(5-chloro-1,3,3-trimethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)benzothiazolium iodide **11.34**
 4-Methyl-4-cholest-3-one **2.90**
 1-Methylcyclobutene **2.98**
 1-Methylcycloheptene **2.102**
 1-Methyl-1,4-cyclohexadiene **2.107**
 3-Methyl-1,2-cyclohexanedione, 2-enol **2.128**
 1-Methylcyclohexene **2.122, 2.11, 2.21, 2.120, 2.153, 2.161, 2.236, 2.237, 2.238, 2.239, 2.243, 2.309, 2.326, 2.327**
 4-Methylcyclohexene **2.123**
 1-Methylcyclooctene **2.138**
 1-Methylcyclopentene **2.161, 2.22, 2.122**
 Methyl 3,5-di-*tert*-butyl-4-hydroxybenzoate **4.39**
N-Methyldiethanolamine **8.46**
 5-Methyl-2,3-dihydrofuran **5.31**
 Methyl 4,5-dihydro-2-methyl-3-furancarboxylate **5.58**
 4-Methyl-2,3-dihydropyran-2-*t* **5.97**
 4-Methyl-2,3-dihydropyran-3-*t* **5.95**
 4-Methyl-2,3-dihydropyran-4-*d* **5.92**
 4-Methyl-2,3-dihydropyran-4-*t* **5.93**
 6-Methyl-3,4-dihydropyran-5-carboxylic acid ethyl ester **5.98**
 Methyl 4-(1,1-dimethylethyl)phenyl sulfide **13.20**
 2-Methyl-4,6-dinitrophenoxide ion **4.173**
 4-Methyl-2,6-dinitrophenoxide ion **4.174**
 Methylenecyclopropylcyclobutenylidene- **2.96, 2.49, 2.50, 2.94, 2.95, 2.151, 2.168, 2.169, 2.333, 2.334, 2.335, 3.113**
 2-Methylenebicyclo[2.2.1]heptane **2.11, 2.9, 2.13, 2.14, 2.21, 2.23**
 5-Methylenebicyclo[2.2.1]hept-2-ene **2.22, 2.6, 2.11, 2.21, 2.153**
 2,2'-Methylenebis(6-*tert*-butyl-4-methylphenol) **4.136**
 2,2'-Methylenebis[*O,O'*-di(4-*tert*-butylphenyl)dithiophosphate]cobalt(II) **12.156**
 2,2'-Methylenebis[*O,O'*-di(4-*tert*-butylphenyl)dithiophosphate]lead(II) **12.157**

- 2,2'-Methylenebis[*O,O'*-di(4-*tert*-butylphenyl)dithiophosphate]nickel(II) 12.158
- Methylene Blue cation 11.113
- Methylene chloride 17.16
- Methylene-*d*₂ chloride 17.17
- Methylenecycloheptane 2.101
- Methylenecyclohexane 2.112, 2.101, 2.153
- Methylenecyclopentane 2.153
- 2-Methylenenorbornane 2.11, 2.9, 2.13, 2.14, 2.21, 2.23
- endo*-2-Methylenenorbornane-3-*d* 2.13
- exo*-2-Methylenenorbornane-3-*d* 2.14
- 5-Methylene-2-norbornene 2.22, 2.6, 2.11, 2.21, 2.153
- Methyl esters of fatty acids from peanut oil 17.42
- Methyl esters of fatty acids from soybean oil 17.43
- Methyl esters of fatty acids from sunflower oil 17.44
- Methyl 4-fluorophenyl sulfide 13.21
- 3-Methyl-2-[7-(5-fluoro-1,3,3-trimethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)benzothiazolium iodide 11.35
- 2-Methylfuran 5.46
- 5-Methylfuran-2-carboxaldehyde 5.13
- 5-Methyl-2-furanitrile 5.64
- 5-Methylfurfural 5.13
- α-Methylfurfuryl alcohol 5.69
- N*-Methylfurfurylamine 5.65
- Methyl 2-furoate 5.57
- 3-Methylheptane 17.35
- 4-Methylheptane 17.36
- Methylhydroquinone 4.29
- 4-Methylimidazole 6.7
- 2,2'-Methylenimino diethanol 8.46
- 1-Methylindole 6.15
- 3-Methylindole 6.16
- 2-Methyl-5-isopropylthiacyclopent-2-ene-1-oxide 13.144
- Methyl linoleate 2.265, 4.46, 4.50, 4.53
- Methyl linolenate 2.269
- 4-(Methylmercapto)pheno 13.120
- (E)-1-Methyl-3-(β-methoxylvinyl)indole 2.247, 2.248
- (Z)-1-Methyl-3-(β-methoxylvinyl)indole 2.248
- 2-Methyl-6-(4-methoxyphenyl)imidazo[1,2-*a*]pyrazin-3-one 16.40
- 4-Methyl-2-(4-methoxyphenyl)-5-phenyloxazole 6.35
- Methyl 4-methoxyphenyl sulfide 13.22
- 3-Methyl-2-[7-(5-methoxy-1,3,3-trimethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)benzothiazolium iodide 11.36
- (R)-1-Methyl-4-(1-methylethenyl)cyclohexene 2.124, 13.8, 13.52, 13.97, 13.119, 13.136
- 1-Methyl-4-(1-methylethyl)-1,3-cyclohexadiene 2.105
- 2-Methyl-5-(1-methylethyl)-1,3-cyclohexadiene 2.106
- 4-Methyl-2-(4-methylphenyl)-5-phenyloxazole 6.36
- Methyl 3-methylphenyl sulfide 13.24
- Methyl 4-methylphenyl sulfide 13.25
- 4-Methyl-3-(4-methylphenyl)sydnone 5.103
- Methyl 1-methylpropenyl ether 2.55
- N*-[(5-Methyl-2-(methylthio)-3-thienyl)methylene]cyclohexanamine 13.64
- 1-Methylnaphthalene 3.86
- 4-Methyl-1,3-naphthalenedipropionic acid 3.89
- 4-Methyl-1-naphthalenepropionate ion 3.91
- 4-Methyl-1-naphthalenepropionate ion endoperoxide 3.91
- 4-Methyl-1-naphthalenepropionic acid endoperoxide 3.91
- 2-Methyl-1,4-naphthoquinone 16.44
- 3,3'-(4-Methyl-1,3-naphthylene)dipropionic acid 3.89
- Methyl 2-naphthyl ketone 3.1
- 4-Methyl-2-nitrophenoxyde ion 4.175
- 1-(4-Methyl-2-nitrophenoxy)-2-naphthol 11.101
- 4-Methyl-2-(4-nitrophenyl)-5-phenyloxazole 6.37
- 2-Methyl-2-nitrosopropane 15.50
- 9-Methyl-6-nitro-2,3,4,9-tetrahydrocarbazole 6.3
- 2-Methylnorborna-2,5-diene 2.6
- 2-Methylnorborn-2-ene 2.21, 2.23
- 4-Methyl-4-octene 2.289
- cis*-4-Methyl-4-octene 2.290
- (E)-4-Methyl-4-octene 2.290
- trans*-4-Methyl-4-octene 2.291
- (Z)-4-Methyl-4-octene 2.291
- Methyl oleate 2.272
- 4-Methyl-1,3-pentadiene 2.300
- 2-Methyl-2-pentene 2.307, 2.66, 2.86, 2.92, 2.122, 2.243, 2.250, 2.256, 2.311, 2.312, 2.343, 3.71, 4.32, 4.102, 4.147, 13.87, 15.1
- 3-Methyl-2-pentene 2.308
- cis*-3-Methyl-2-pentene 2.310
- cis*-4-Methyl-2-pentene 2.312, 2.116
- (E)-3-Methyl-2-pentene 2.309
- (E)-4-Methyl-2-pentene 2.311
- trans*-3-Methyl-2-pentene 2.309
- trans*-4-Methyl-2-pentene 2.311
- (Z)-3-Methyl-2-pentene 2.310
- (Z)-4-Methyl-2-pentene 2.312, 2.116
- 2-Methyl-2-penten-4-ol 2.323
- 4-Methyl-3-penten-2-ol 2.323
- (E)-3-Methyl-2-penten-1-ol 2.319
- (Z)-3-Methyl-2-penten-1-ol 2.320
- 3-Methyl-2-pentyl-2-cyclopenten-1-one 2.162
- 4-Methylphenol 4.135
- 10-Methylphenothiazine 13.123
- N*-Methylphenothiazine 13.123
- 4-Methylphenoxyde ion 4.172
- 2-(4-Methylphenoxy)methylideneadamantane 2.354
- 1-(4-Methylphenylazo)-2-naphthol 11.102
- 4-(4-Methylphenylazo)-1-naphthol 11.90

- 1-(4-Methylphenylazo)-3-(phenylaminocarbonyl)-2-naphthol **11.80**
- 1-(4-Methylphenylazo)-1-(phenylaminocarbonyl)-2-propanone **11.60**
- 3-Methyl-1-phenyl-2-butene **3.45**
- N*-Methyl-4-phenyl-2,3-dihydropyridinium **6.53**
- 3-(4-Methylphenyl)-1,5-diphenyl-2-pyrazoline **11.126**
- 5-(4-Methylphenyl)-1,3-diphenyl-2-pyrazoline **11.127**
- 2-(4-Methylphenyl)furan **5.47**
- 3-(4-Methylphenyl)furan **5.48**
- 2-Methyl-6-phenylimidazo[1,2-*a*]pyrazin-3-one **16.41**
- 1-Methyl-2-phenylindene **3.71**
- 3-Methyl-2-phenylindene **3.71**
- 3-Methyl-2-phenylindole **6.17**
- 2-Methyl-*N*-phenylmethyleno-2-propanamine *N*-oxyl **15.46**
- 4-Methylphenyl phenyl nitrone **15.8**
- 4-Methylphenyl phenyl sulfide **13.26**
- 3-(4-Methylphenyl)-4-phenylsydnone **5.106**
- 2-Methyl-1-phenyl-1-propene **3.114**
- N*-Methyl-4-phenylpyridinium **6.55**
- Methyl phenyl sulfide **13.29**
- 2-Methyl-3-phenylsulfinyl-2-butene **13.53**
- 3-Methyl-4-phenylsydnone **5.105**
- 3-(4-Methylphenyl)sydnone **5.104**
- N*-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine **8.94**
- 1-Methyl-4-(phenylthio)benzene **13.26**
- 4-(4-Methylphenyl)thio-1-butanol **13.52**
- 2-(4-Methylphenyl)thioethanol **13.97**
- 6-(4-Methylphenyl)thio-1-hexanol **13.103**
- 5-(4-Methylphenyl)thio-1-pentanol **13.118**
- 3-(4-Methylphenyl)thio-1-propanol **13.136**
- Methyl phosphite **17.73, 17.74**
- Methyl phytol ether **2.237**
- 2-Methylpiperidine **8.59**
- N*-Methylpiperidine **8.60**
- 1-Methyl-3-piperidinol **8.70**
- 1-Methyl-4-piperidinol **8.71**
- N*-Methyl-4-piperidinol **8.71**
- 2-Methyl-2-propaneamine **8.15**
- 2-Methyl-2-propanol **17.62**
- 2-Methylpropene **2.337**
- 2-Methylpropenoic acid **2.252**
- 1,1'-(2-Methyl-1-propenylidene)bisbenzene **3.46**
- (2-Methylpropylidene)tricyclo[3.3.1.1^{3,7}]decane **2.355**
- 2-Methylpropyl methyl ether **17.18**
- 1-Methylpropyl propyl sulfide **13.59**
- 1-Methylpyridinium **6.54**
- 4-(*N*-Methylpyridinium)-*tert*-butylnitronate **15.45**
- α -(1-Methyl-4-pyridyl)-*tert*-butyl nitronate **15.45**
- 1-Methylpyrrole **6.64**
- N*-Methylpyrrole **6.64**
- Methyl salicylate **4.40**
- Methyl salicylate, conjugate base **4.41**
- Methylselenobenzene **13.27**
- 2-(Methylseleno)-*N*-phenylbenzamide **13.2**
- Methyl stearate **16.43**
- cis*- α -Methylstilbene **3.105**
- (*E*)- α -Methylstilbene **3.104**
- trans*- α -Methylstilbene **3.104**
- (*Z*)- α -Methylstilbene **3.105**
- cis*- β -Methylstyrene **3.120**
- (*E*)- β -Methylstyrene **3.119**
- trans*- β -Methylstyrene **3.119**
- (*Z*)- β -Methylstyrene **3.120**
- Methyl sulfoxide **13.80, 13.81**
- Methyl sulfoxide **13.81**
- Methyltellurobenzene **13.28**
- 9-Methyl-2,3,4,9-tetrahydrocarbazole **6.2**
- 3-Methyl-2-[7-(1,3,3,5-tetramethyl-2-indolylidene)-1-(1,3,5-heptatrienyl)benzothiazolium, iodide **11.38**
- 2-Methylthiacyclohex-2-ene-1-oxide **13.143**
- 1-Methyl-3-(1-thiaethyl)benzene **13.24**
- 3-Methyl-4-thiaheptane **13.59**
- Methylthiirane **13.152**
- p*-Methylthioanisole **13.25**
- 4-Methylthiobenzamide **13.155**
- Methylthiolato(triethylphosphino)gold(I) **12.159**
- 2-(Methylthio)-*N*-phenylbenzamide **13.3**
- N*-Methylthiourea **13.176**
- 3-Methyl-2-[7-(1,3,3,5-trimethyl-2-indolylidene)-1,3,5-heptatrienyl]benzothiazolium, iodide **11.37**
- m*-Methyl- α , β , β -trimethylstyrene **3.121**
- p*-Methyl- α , β , β -trimethylstyrene **3.122, 3.109, 3.111, 3.112, 3.117, 3.118, 3.123**
- 5-Methyluracil **16.79**
- Mitomycin C **16.7**
- Monohydroperoxides of 4,8-dimethyl-4,8-dodecadiene **2.203**
- Monohydroperoxides of *trans*-2,6-dimethyl-2,6-octadiene **2.278**
- Monohydroperoxy-3,7-dimethyl-2,6-octadien-1-ol **2.284**
- Monohydroperoxygeraniol **2.284**
- Monohydroxyethyl vinyl deuteroporphyrin **7.92**
- Morpholine, 4-[(phenylmethyl)thio]- **13.112**
- 4-Morpholineethanesulfonic acid **17.45**
- Muscalure **2.349**
- Myristic acid **17.67**
- NADH dehydrogenase **10.12**
- Nagarse **10.32**
- Nalidixic acid **16.45**
- Nalidixic acid, anion **16.46**
- Naphthacene **3.124, 3.12, 7.53, 8.72, 8.77, 9.18, 15.46**

- 5,12-Naphthacenedione, 8-acetyl-10[(3-amino-2,3,6-trideoxy-hexopyranosyloxy]-tetrahydro-6,8,11-trihydroxy-1-methoxy- **4.79**
- 5,12-Naphthacenedione, 10-[(3-amino-2,3,6-trideoxy- α -L-*lyxo*-hexopyranosyl)oxy]-7,8,9,10-tetrahydro-6,8,11-trihydroxy-8-(hydroxyacetyl)-1-methoxy- **4.80**
- 12-Naphthaceneone, 8-acetyl-10[(3-amino-2,3,6-trideoxyhexopyranosyloxy]-tetrahydro-6,7,11-trihydroxy-5-imino-1-methoxy- **4.81**
- 2-Naphthalenamine **9.24**
- Naphthalene **3.74**
- Naphthalene, 2-acetyl- **3.1**
- Naphthalene, 2-amino- **9.24**
- Naphthalene, 1,5-dihydroxy- **4.82**
- Naphthalene, 1,6-dihydroxy- **4.83**
- Naphthalene, 1,7-dihydroxy- **4.84**
- Naphthalene, 1,8-dihydroxy- **4.85**
- Naphthalene, 2,6-dihydroxy- **4.86**
- Naphthalene, 2,7-dihydroxy- **4.87**
- Naphthalene, 1,2-dimethyl- **3.75**
- Naphthalene, 1,3-dimethyl- **3.76**
- Naphthalene, 1,4-dimethyl- **3.77**
- Naphthalene, 1,5-dimethyl- **3.75, 3.76, 3.77, 3.78, 3.79, 3.80, 3.81, 3.82, 3.83, 3.86**
- Naphthalene, 1,6-dimethyl- **3.78**
- Naphthalene, 1,7-dimethyl- **3.79**
- Naphthalene, 1,8-dimethyl- **3.80**
- Naphthalene, 2,3-dimethyl- **3.81**
- Naphthalene, 2,6-dimethyl- **3.82**
- Naphthalene, 2,7-dimethyl- **3.83**
- Naphthalene, 1,1'-(1,2-dimethyl-1,2-diethenyl)bis-, (*E*)- **3.84**
- Naphthalene, 2,2'-(1,2-dimethyl-1,2-diethenyl)bis-, (*E*)- **3.85**
- Naphthalene, 2,2'-(1,2-dimethyl-1,2-diethenyl)bis-, (*Z*)- **3.84, 3.85**
- Naphthalene, 1-methoxy-4-(phenylazo)- **11.59**
- Naphthalene, 1-methyl- **3.86**
- Naphthalene, 1-(2-methyl-1-propenyl)- **3.87**
- Naphthalene, 1,2,3,4,5,6,7,8-octahydro- **2.261, 13.112**
- 2-Naphthalenecarboxamide, 4-[[4-(aminocarbonyl)phenyl]azo]-3-hydroxy-*N*-(2-methoxyphenyl)- **11.79**
- 2-Naphthalenecarboxamide, 3-hydroxy-4-[(4-methylphenyl)azo]-*N*-phenyl- **11.80**
- 1,5-Naphthalenediol **4.82**
- 1,6-Naphthalenediol **4.83**
- 1,7-Naphthalenediol **4.84**
- 1,8-Naphthalenediol **4.85**
- 2,6-Naphthalenediol **4.86**
- 2,7-Naphthalenediol **4.87**
- 1,4-Naphthalenedipropionate ion **3.88**
- 3,3'-(1,4-Naphthalene)dipropionate ion **3.88**
- 1,3-Naphthalenedipropionic acid, 4-methyl- **3.89**
- 1,3-Naphthalenedipropionic acid, 4-methyl-, dimethyl ester **3.90**
- 1-Naphthalenepropionate ion, 4-methyl- **3.91**
- 1-Naphthalenepropionate ion, 4-methyl-, endoperoxide **3.92**
- 1-Naphthalenepropionic acid, 4-methyl-, endoperoxide **3.93**
- 2-Naphthalenthione, 1,2-dihydro-1,1-dimethyl- **13.113**
- 2-Naphthalenthione, 1,2-dihydro-1,1,3-trimethyl- **13.114**
- 1-Naphthalenone, 4-[[2,6-dimethyl-4-(diethylamino)phenyl]imino]-2-(2-methoxy-5-fluorosulfonylphenyl)aminocarbonyl- **11.81**
- 1-Naphthalenone, 4-[[4-(diphenylamino)phenyl]imino]- **11.82**
- 1-Naphthalenone, 4-[(*N*-methyl-*N*-phenyl)amino]imino- **11.83**
- 1-Naphthalenone, 4-[[4-(phenylamino)phenyl]imino]- **11.84**
- Naphthalocyanine, bis(tribenzylsiloxy)silicon **7.17**
- Naphthalocyanine, bis(*tri-n*-hexyloxysiloxy)silicon **7.18**
- Naphthalocyanine, bis(trihexylsiloxy)silicon **7.18**
- Naphthalocyanine, bis(trihexylsiloxy)tin **7.19**
- Naphthalocyanine, bis(triisobutylsiloxy)silicon **7.20**
- Naphthalocyanine, 1,6,10,15,19,24,28,33-octabutoxy-, palladium(II) **7.21**
- Naphthalocyanine, 2,11,20,29-tetrakis(1,1-dimethylethyl) **7.22**
- Naphthalocyanine, 1,10,19,28-tetraphenyl-, hydroxyaluminum(III) **7.23**
- Naphthalocyanine, trihexylsiloxyaluminum **7.24**
- Naphthalocyanine, trihexylsiloxygallium **7.25**
- Naphtho[1,8-*b*:5,4-*b'*,*c*']dipyran **3.94**
- 1-Naphthol **4.88**
- 1-Naphthol, 5-[[3-(aminosulfonyl)phenyl]sulfonylamino]-4-[2-(methylsulfonyl)-4-nitrophenylazo]-, conjugate base **11.85**
- 1-Naphthol, 5-[[3-(aminosulfonyl)phenyl]sulfonylamino]-4-[2-(methylsulfonyl)-4-nitrophenylazo]-, conjugate dibase **11.86**
- 1-Naphthol, 4-(4-chlorophenylazo)- **11.87**
- 1-Naphthol, 5-methoxy-4-[2-(methylsulfonyl)-4-nitrophenylazo]-, conjugate base **11.88**
- 1-Naphthol, 4-(4-methoxyphenylazo)- **11.89**
- 1-Naphthol, 4-(4-methylphenylazo)- **11.90**
- 1-Naphthol, 4-(4-nitrophenylazo)- **11.91**
- 1-Naphthol, 4-phenylazo- **11.92, 11.87, 11.89, 11.90, 11.91**
- 2-Naphthol **4.89**
- 2-Naphthol, 1-(4-aminophenylazo)- **11.93**
- 2-Naphthol, 1-(4-bromophenylazo)- **11.94**
- 2-Naphthol, 1-(4-chlorophenylazo)- **11.95**
- 2-Naphthol, 1-[4-(dimethylamino)phenylazo]- **11.96**
- 2-Naphthol, 1-(4-fluorophenylazo)- **11.97**

- 2-Naphthol, 1-(4-hydroxyphenylazo)- 11.98
 2-Naphthol, 1-(4-iodophenylazo)- 11.99
 2-Naphthol, 1-(4-methoxyphenylazo)- 11.100
 2-Naphthol, 1-(4-methyl-2-nitrophenylazo)- 11.101
 2-Naphthol, 1-(4-methylphenylazo)- 11.102
 2-Naphthol, 1-(4-methylphenylazo)-3-(phenylaminocarbonyl)- 11.80
 2-Naphthol, 1-(4-nitrophenylazo)- 11.103
 2-Naphthol, 1-phenylazo- 11.104, 11.94, 11.95, 11.96, 11.97, 11.98, 11.99, 11.100, 11.102, 11.103
 4-Naphthol, 1-(4-methylphenylazo)- 11.90
 4-Naphthol, 1-phenylazo- 11.91
 α -Naphthol 4.88
 β -Naphthol 4.89
 Naphtho[1,2,3,4-*rst*]pentaphene-5,8-dione, 3,10-dimethyl-3.95
 1,4-Naphthoquinone, 5-hydroxy- 4.90
 1,4-Naphthoquinone, 6-hydroxy- 4.91
 1,4-Naphthoquinone, 2-methyl- 16.44
 1,4-Naphthoquinone *N*-methyl-*N*-phenylhydrazone 11.83
 Naphtho[1,2-*d*]thiazolium, 1-ethyl-2-[(1-ethylnaphtho[1,2-*d*]thiazol-2-ylidene)methyl]-, chloride 11.105
 Naphtho[1,2-*d*]thiazolium, 1-ethyl-2-[(1-ethylnaphtho[1,2-*d*]thiazol-2-ylidene)methyl]-1-but enyl-, bromide 11.106
 Naphtho[1,2-*d*]thiazolium, 1-ethyl-2-[(1-ethylnaphtho[1,2-*d*]thiazol-2-ylidene)-1-propenyl]-, toluenesulfonate 11.107
 Naphtho[2,3-*d*]thiazolium, 2-[2-[2-(diphenylamino)-3-[3-(4-methoxy-4-oxobutyl)naphtho[2,3-*d*]thiazol-2-ylidene]ethylidene]1-cyclopenten-1-yl]ethenyl]-3-(4-methoxy-4-oxobutyl)-, perchlorate 11.108
 2-Naphthylamine 9.24
 2,2'-[1,8-Naphthylenebis(nitrilomethylidyne)]-bis[phenolato]nickel(II) 12.160
 1,8-Naphthyridine-3-carboxylic acid, 1,4-dihydro-1-ethyl-7-methyl-4-oxo- 16.45
 1,8-Naphthyridine-3-carboxylic acid, 1,4-dihydro-1-ethyl-7-methyl-4-oxo-, anion 16.46
 Naringenin 4.72
 Neoalloocimene 2.285
 Neurosporene 2.70
 $(all-E)$ -Neurosporene 2.70
 $(all-trans)$ -Neurosporene 2.70
 Nickel(I), bis(1,2-diphenyl-1,2-ethanedithionato)-, tetrabutylammonium salt 12.65
 Nickel(II), bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenethionato]- 12.21
 Nickel(II), bis[1,2-di(2-bromophenyl)-1,2-ethenedithiolato]- 12.25
 Nickel(II), bis[1,2-di(2-chlorophenyl)-1,2-ethenedithiolato]- 12.33
 Nickel(II), bis[1,2-di(4-chlorophenyl)-1,2-ethenedithiolato]- 12.34
 Nickel(II), bis[1,2-di(4-ethylphenyl)-1,2-ethenedithiolato]- 12.42
 Nickel(II), 3,4-dimercaptotoluene(2,2'-bipyridine) 12.153
 Nickel(II), 3,4-dimercaptotoluene(1,10-phenanthroline) 12.154
 Nickel(II) acetate 12.161
 Nickel(II) ammine[2,2'-thiobis(4-*tert*-octyl)phenolate] 12.2
 Nickel(II) aniline[2,2'-thiobis(4-*tert*-octyl)phenolate] 12.3
 Nickel(II) aquo[2,2'-thiobis(3,4-dimethyl)phenolate] 12.4
 Nickel(II) aquo[2,2'-thiobis(4-*tert*-octyl)phenolate] 12.5
 Nickelate(1-), bis[1,2-di(4-trifluoromethylphenyl)-1,2-ethanedithionato]-, tetrabutylammonium salt 12.75
 Nickel(II) bis(acetylacetone) 12.10
 Nickel(II) bis(acetylacetone) dihydrate 12.11
 Nickel(II) bis[2-(*N*-(*p*-anilinophenyl)formimidoyl)phenol] 12.106
 Nickel(II) bis[1,3-bis(2-pyridylimino)isoindoline] 12.45
 Nickel(II) bis(5-bromosalicylaldehyde)⁺2H₂O 12.14
 Nickel(II) bis[*O*-butyl-3,5-di-*tert*-butyl-4-hydroxybenzylphosphonate] 12.16
 Nickel(II) bis[2-(*N*-butylformimidoyl)-4-bromophenol] 12.17
 Nickel(II) bis[2-(*N*-butylformimidoyl)-4-methoxyphenol] 12.18
 Nickel(II) bis[2-(*N*-butylformimidoyl)phenol] 12.19
 Nickel(II) bis[2-(*N*-sec-butylformimidoyl)phenol] 12.105
 Nickel(II) bis[2-(*N*-*tert*-butylformimidoyl)phenol] 12.57
 Nickel(II) bis[*p*-*tert*-butylphenylsalicylate] 12.58
 Nickel(II) bis[2-(*N*-cyclohexylformimidoyl)phenol] 12.24
 Nickel(II) bis(dibutyldithiocarbamate) 12.28
 Nickel(II) bis[*O,O'*-di(4-*tert*-butylphenyl)dithiophosphate] 12.32
 Nickel(II) bis(dicyclohexylthiophosphinate) 12.35
 Nickel(II) bis[1,2-di(2,4-dichlorophenyl)-1,2-ethenedithiolato]- 12.37
 Nickel(II) bis[1,2-di(3,4-dichlorophenyl)-1,2-ethenedithiolato]- 12.38
 Nickel(II) bis[1,2-di(dimethylaminophenyl)-1,2-ethenedithiolato]- 12.39
 Nickel(II) bis(diethylthiocarbamate) 12.40
 Nickel(II) bis(*O,O'*-diethylthiophosphate) 12.44
 Nickel(II) bis(*O,O'*-diethylphosphorodithiolato)- 12.44
 Nickel(II) bis(diisopropylthiocarbamate) 12.50
 Nickel(II) bis[diisopropylthiophosphate] 12.52
 Nickel(II) bis(*O,O'*-diisopropylthiophosphate) 12.52
 Nickel(II) bis(*O,O'*-diisopropylphosphorodithiolato)- 12.52
 Nickel(II) bis[3,5-diisopropylsalicylate] 12.60
 Nickel(I) bis[1,2-di(4-methoxyphenyl)-1,2-ethanedithionato]-, tetrabutylammonium salt 12.53
 Nickel(II) bis[1,2-di(4-methoxyphenyl)-1,2-ethenedithiolato]- 12.54
 Nickel(II) bis(dimethylthiocarbamate) 12.56
 Nickel(II) bis(1,2-dimethyl-1,2-ethenedithiolato)- 12.15

- Nickel(II) bis[*O,O'*-di(4-methylphenyl)dithiophosphate] 12.63
- Nickel(II) bis(diphenyldithiocarbamate) 12.64
- Nickel(II) bis(*O,O'*-diphenyldithiophosphate) 12.71
- Nickel(II) bis[1,2-diphenyl-1,2-ethenedithiolato]- 12.66
- Nickel(II) bis(*O,O'*-diphenyl phosphorodithioato)- 12.71
- Nickel(II) bis[dithioacetylacetone] 12.74
- Nickel(II) bis(dithiobenzil) 12.66
- Nickel(II) bis[dithiobiacyl] 12.15
- Nickel(II) bis(dithiohexafluorobiacyl) 12.85
- Nickel(I) bis[1,2-di(4-trifluoromethylphenyl)-1,2-ethenedithiolato]-, tetrabutylammonium salt 12.75
- Nickel(II) bis[1,2-di(4-trifluoromethylphenyl)-1,2-ethenedithiolato]- 12.76
- Nickel(I) bis[1,2-di(3,4,5-trimethoxyphenyl)-1,2-ethanedithionato]-, tetrabutylammonium salt 12.77
- Nickel(II) bis[1,2-di(3,4,5-trimethoxyphenyl)-1,2-ethenedithiolato]- 12.79
- Nickel(II) bis[2-(*N*-dodecylformimidoyl)phenol] 12.82
- Nickel(II) bis[*O*-ethyl-3,5-di-*tert*-butyl-4-hydroxybenzylphosphonate] 12.83
- Nickel(II) bis[2-(formimidoyl)phenol] 12.96
- Nickel(II) bis[hydrotris(1-pyrazolyl)borate] 12.86
- Nickel(II) bis(2'-hydroxyacetophenone oxime) 12.87
- Nickel(II) bis[2'-hydroxy-4'-methylacetophenone oxime] 12.93
- Nickel(II) bis[2-hydroxy-5-methylbenzophenone] 12.94
- Nickel(II) bis[2'-hydroxy-4'-methylundecanophenone oxime] 12.95
- Nickel(II) bis[4-imino-2-pentene-2-ol] 12.97
- Nickel(II) bis[2-(*N*-isopropylformimidoyl)phenol] 12.101
- Nickel(II) bis(isopropylxanthate) 12.98
- Nickel bis(mesitylenesulfonate) 12.118
- Nickel(II) bis(5-methoxysalicylaldehyde)²H₂O 12.92
- Nickel(II) bis(1-methylamino-2-methylinocycloheptatriene) 12.102
- Nickel(II) bis[*O*-(1-methylethyl) carbonodithionato]- 12.98
- Nickel(II) bis[1-(4'-methylphenyl)amino-2-(4'-methylphenyl)iminocycloheptatriene] 12.104
- Nickel(II) bis(4-methylphenyldithiocarbamate) 12.103
- Nickel(II) bis[2-*N*-phenylaminoformimidoylphenol] 12.90
- Nickel(II) bis(phenyldithiocarbamate) 12.107
- Nickel(II) bis[2-(*N*-phenylformimidoyl)phenol] 12.110
- Nickel(II) bis(salicylaldehyde) dihydrate 12.88
- Nickel(II) bis(salicylaldehyde oxime) 12.89
- Nickel(II) bis[2,2'-thiobis[*O,O'*-di(4-*tert*-butylphenyl)dithiophosphate]] 12.114
- Nickel(II) bis[2,2'-thiobis(4-*tert*-octyl)phenolate] 12.116
- Nickel(II) bis[μ -toluene-3,4-dithiolato] tetrabutylammonium salt 12.100
- Nickel(I) bis[3,4,6-trichloro-1,2-benzenedithiolato]-, tetrabutylammonium salt 12.117
- Nickel(II) butylamine[2,2'-thiobis(3,4-dimethyl)phenolate] 12.120
- Nickel(II) butylamine[2,2'-thiobis(4-*tert*-octyl)phenolate] 12.122
- Nickel chloride 12.162
- Nickel chloride hexahydrate 12.163
- Nickel(II) cyclohexylamine[2,2'-thiobis(3,4-dimethyl)phenolate] 12.130
- Nickel(II) cyclohexylamine[2,2'-thiobis(4-*tert*-octyl)phenolate] 12.131
- Nickel(II) deuteroporphyrin 7.95
- Nickel(II) deuteroporphyrin bis(piperidine) 7.96
- Nickel(II) dibutyldithiocarbamate 12.28
- Nickel(II) dichloride 12.162
- Nickel(II) dichloride hexahydrate 12.163
- Nickel(II) didodecylamine[2,2'-thiobis(3,4-dimethyl)phenolate] 12.132
- Nickel dimethyldithiocarbamate 12.56
- Nickel(II) dodecylamine[2,2'-thiobis(3,4-dimethyl)phenolate] 12.134
- Nickel(II) dodecylamine[2,2'-thiobis(4-*tert*-octyl)phenolate] 12.135
- Nickel(II) ethylamine[2,2'-thiobis(3,4-dimethyl)phenolate] 12.141
- Nickel(II) ethylamine[2,2'-thiobis(4-*tert*-octyl)phenolate] 12.142
- Nickel(II) 2,2'-[ethylenebis(nitrilodecylidyne)]di-*p*-cresol 12.136
- Nickel(II) 2,2'-[ethylenebis(nitrioloethylidyne)]diphenol 12.137
- Nickel(II) 2,2'-[ethylenebis(nitrilomethylidyne)]diphenol 12.140
- Nickel(I) ion 12.164
- Nickel(II) mesoporphyrin 7.87
- Nickel(II) 2,2'-methylenebis[*O,O'*-di(4-*tert*-butylphenyl)dithiophosphate] 12.158
- Nickel(II) 2,2'-[1,8-naphthylenebis(nitrilomethylidyne)]diphenol 12.160
- Nickelocene 12.165
- Nickelous acetate 12.161
- Nickel(II) 2,2'-[*o*-phenylenebis(nitrilomethylidyne)]diphenol 12.166
- Nickel(II) propylamine[2,2'-thiobis(4-*tert*-octyl)phenolate] 12.168
- Nickel(II) protoporphyrin 7.80
- Nickel(II) protoporphyrin, dimethyl ester 7.79
- Nickel(II) tetraphenylporphyrin 7.67
- Nickel(II) 5,10,15,20-tetraphenylporphyrin 7.67
- Nickel(II) triethanolamine[2,2'-thiobis(4-*tert*-octyl)phenolate] 12.171
- Nicotinamide adenine dinucleotide 16.47
- Nicotinamide adenine dinucleotide, reduced 16.48
- Nicotinamide adenine dinucleotide phosphate 16.49

Nicotinamide-adenine dinucleotide phosphate, reduced	
16.50	
Nicotinamide hypoxanthine dinucleotide, reduced	16.42
Nicotinamide mononucleotide, reduced	16.51
Nicotinamide ribose monophosphate, reduced	16.51
Nicotine	8.93
Nitrite ion	14.11
2-Nitroanisole	3.44
4-Nitroazobenzene	11.61
Nitrobenzene	3.47
nitrone 3	15.25
2-Nitrophenol	4.137
3-Nitrophenol	4.138
4-Nitrophenol	4.139
2-Nitrophenoxide ion	4.176
3-Nitrophenoxide ion	4.177
4-Nitrophenoxide ion	4.178
1-(4-Nitrophenylazo)-2-naphthol	11.103
4-(4-Nitrophenylazo)-1-naphthol	11.91
4-(<i>p</i> -Nitrophenylazo)-1-naphthol	11.91
7-Nitro-2-phenyl-1,2-benziselenazol-3-one	13.39
2-(4-Nitrophenyl)furan	5.49
4-Nitrophenyl phenyl sulfide	13.31
1-Nitro-4-(phenylthio)benzene	13.31
4-Nitroso- <i>N,N</i> -dimethylaniline	9.7
4-Nitrothiobenzamide	13.156
Nitroxide, bis[4-(1,1-dimethylethyl)phenyl]-	15.26
Nitroxide, bis[4-(1-methyl-1-phenylethyl)phenyl]-	15.27
2,4,6,8-Nonatetraenoic acid, 9-(4-methoxy-2,3,6-trimethylphenyl)-3,7-dimethyl-, ethyl ester, (<i>all-E</i>)-	2.262
1-Nonene	2.263
Nopadiene	2.25
Noradrenaline	4.6
Norbixin	2.179
<i>trans</i> -Norbixin	2.174
Norbornadiene	2.4
Norepinephrine	4.6
1,6,10,15,19,24,28,33-Octabutoxynaphthalocyaninatopalladium(II)	7.21
1,4,8,11,15,18,22,25-Octabutoxyphthalocyanine	7.42
Octa(3,6-butoxy)phthalocyanine	7.42
9,12-Octadecadienoic acid, methyl ester	2.265, 4.46, 4.50, 4.53
9,12-Octadecadienoic acid, phenyl ester	2.266
(<i>Z,Z</i>)-9,12-Octadecadienoic acid	2.264
Octadecanoic acid	17.66
6,9,12,15-Octadecatetraenoic acid, methyl ester	2.213
9,12,15-Octadecatrienoic acid, ethyl ester (<i>all-Z</i>)-	2.268
9,12,15-Octadecatrienoic acid, methyl ester	2.269
9-Octadecenoic acid	2.271
9-Octadecenoic acid, methyl ester	2.272
9-Octadecenoic acid, phenyl ester	2.273
Octadecyl 3-(3',5'-di- <i>tert</i> -butyl-4'-hydroxyphenyl)propionate	4.33
2,5-Octadiene, 7-hydroperoxy-2,7-dimethyl-	2.274
2,6-Octadiene, 2,6-dimethyl-, (<i>E</i>)	2.275
2,6-Octadiene, 2,6-dimethyl-, (<i>Z</i>)	2.276
2,6-Octadiene, 2,7-dimethyl-	2.277
2,6-Octadiene, hydroperoxy-2,6-dimethyl-	2.278
2,7-Octadiene, 6-hydroperoxy-2,7-dimethyl-	2.279
1,6-Octadien-3-ol, 3,7-dimethyl-	2.280, 2.338, 13.53, 13.54, 13.55, 13.143, 13.144
2,6-Octadien-1-ol, 3,7-dimethyl-, (<i>E</i>)	2.281
2,7-Octadien-1-ol, 3,7-dimethyl-, (<i>E</i>)-	2.282
2,7-Octadien-1-ol, 3,7-dimethyl-, (<i>Z</i>)-	2.283
2,7-Octadien-1-ol, 6-hydroperoxy-3,7-dimethyl-	2.284
Octaethylidihydropurpurin ethyl ester	7.28
2,3,10,11,16,17,24,25-Octaethyl-5,6,7,8,19,20,21,22-octadehydro[26]porphyrin-(2.4.2.4)	7.100
2,3,7,8,12,13,17,18-Octaethyl-5-porphinepropenoic acid ethyl ester	7.97
Octaethyl[22]porphyrin-(2.2.2.2) (di- <i>trans</i>)	7.98
Octaethylporpurin ethyl ester	7.30
1,2,5,6,6a,6b,6c,6d-Octahydro-1,6- <i>o</i> -benzenobenzo[1,3]cyclopropa[1,2,3- <i>cd</i>]cyclopropa[<i>gh</i>]pentalene	2.181
1,2,5,6,6a,6b,6c,6d-Octahydro-9,12-dimethoxy-1,6- <i>o</i> -benzenobenzo[1,3]cyclopropa[1,2,3- <i>cd</i>]cyclopropa[<i>gh</i>]pentalene	2.180
1,2,3,4,11,12,13,14-Octahydrodinaphtho[2,3- <i>a</i> :2',3'- <i>j</i>]perylene-9,19-dione	3.62
1,2,5,6,6a,6b,6c,6d-Octahydro-1,6[1',2']-4-methyl-3,5-dioxo-1,2,4-triazolo[1,3]cyclopropa[1,2,3- <i>cd</i>]cyclopropa[<i>gh</i>]pentalene	17.69
1,2,5,6,6a,6b,6c,6d-Octahydro-1,6[1',2']-4-methyl-3,5-dioxo-1,2,4-triazolo[1,3]cyclopropa[1,2,3- <i>cd</i>]cyclopropa[<i>gh</i>]pentalene	17.70
Octahydropyridazino[1,2- <i>a</i>]pyridazine	8.92
Octalin	2.261, 13.112
$\Delta^{9,10}$ -Octalin	2.261, 13.112
(<i>all-E</i>)-3,7,11,15,20,24,28,32-Octamethyl- 1,34-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)-1,3,5,7,9,11,13,15,17,19,21,23,25,27,29,31,33-tritriacontaheptadecaene	2.358
1,1',3,3,3',3',5,5'-Octamethyltricarbocyanine, iodide	11.184
Octanoate ion, 6,8-dimercapto-	13.115
2,4,6-Octatriene, 2,6-dimethyl-	2.285
3-Octene, 5-hydroperoxy-4-methyl-	2.286
4-Octene, (<i>E</i>)-	2.287
4-Octene, (<i>Z</i>)-	2.288
4-Octene, 4-methyl-	2.289
4-Octene, 4-methyl-, (<i>E</i>)-	2.290
4-Octene, 4-methyl-, (<i>Z</i>)-	2.291

- cis*-4-Octene 2.288
(E)-4-Octene 2.287
trans-4-Octene 2.287
(Z)-4-Octene 2.288
 2-Octen-1-ol, 3,7-dimethyl-, (*E*)- 2.292
 2-Octen-1-ol, 3,7-dimethyl-, (*Z*)- 2.293
 6-Octen-1-ol, 3,7-dimethyl- 2.294
 Oleic acid 2.271
 Ommochrome 16.52
 $\Delta^{1,6}$ -Oxabicyclo[4.4.0]decen-3-one 2.92
 7-Oxabicyclo[2.2.1]heptane, 2,3-bis(methylene)- 2.295
 7-Oxabicyclo[2.2.1]heptane, 2,3,5,6-tetrakis(methylene)- 2.296
 7-Oxabicyclo[2.2.1]hept-2-ene, 5,6-bis(methylene)- 2.297
 Oxadisilirane, 2,2,3,3-tetrakis(2,6-diethylphenyl)- 17.46
 Oxadisilirane, 2,2,3,3-tetrakis(2,6-diisopropylphenyl)- 17.47
 Oxalate ion 17.48
 1-Oxa-4-thiacyclohexane 13.177
 1,4-Oxathiane 13.177
 2,2'-Oxatricarbocyanine, 3,3'-diethyl-, iodide 11.109
exo,exo-2,3-(1 Oxatrimethylene)-7-adamantylidenenorbornane 2.12
 Oxazole, 2,5-diphenyl- 6.31
 Oxazole, 2,5-diphenyl-4-methyl- 6.32
 Oxazole, 4-methyl-2-(3-chlorophenyl)-5-phenyl- 6.33
 Oxazole, 4-methyl-2-(4-chlorophenyl)-5-phenyl- 6.34
 Oxazole, 4-methyl-2-(4-methoxyphenyl)-5-phenyl- 6.35
 Oxazole, 4-methyl-2-(4-methylphenyl)-5-phenyl- 6.36
 Oxazole, 4-methyl-2-(4-nitrophenyl)-5-phenyl- 6.37
 Oxirane, 2,3-diphenyl-, (*E*)- 17.49
 μ -Oxobis[5,10,15,20-tetraphenylporphinatoiron(II)] 7.69
 5'-Oxo-3',4'-diethyl-5-methyl-1',5'-dihydro-(2,2')-dipyrromethene 7.101
 5'-Oxo-3'-ethyl-5,4'-dimethyl-1',5'-dihydro-(2,2')-dipyrromethene 7.107
 5'-Oxo-4-ethyl-3,5-dimethyl-1',5'-dihydro-(2,2')-dipyrromethene 7.108
 5'-Oxo-4-ethyl-3'-ethenyl-3,5,4'-trimethyl-1',5'-dihydro-(2,2')-dipyrromethene 7.106
 5'-Oxo-3'-ethyl-4',3,5-trimethyl-1',5'-dihydro-(2,2')-dipyrromethene 7.103
 α -Oxoglutaric acid 17.51
 Oxolane 17.68
 2-Oxo-3-phenylpropionic acid 17.60
 4-Oxo-2,2,6,6-tetramethylpiperidinoxy free radical 15.44
 5'-Oxo-3',4',4-triethyl-3,5-dimethyl-1',5'-dihydro-(2,2')-dipyrromethene 7.102
 5'-Oxo-4',4,5-trimethyl-3'-ethyl-1',5'-dihydro-(2,2')-dipyrromethene 7.104
 5'-Oxo-4'-vinyl-4-ethyl-3',3,5-trimethyl-1',5'-dihydro-(2,2')-dipyrromethene 7.105
- Oxygen 14.12
 Oxygen-18 14.13
 Oxygen, lowest excited singlet state 17.64
 Palladium(II) bis[2'-hydroxy-4'-*tert*-butyloctadecanophenone oxime] 12.91
 Palmitic acid 17.37
 Pantetheine-adenosine triphosphate 16.17
 Para Red 11.103
 P-422 (8 conjugated bonds) 2.298
 P-438 (9 conjugated bonds) 2.299
 PDS 8.79
 Penicillamine 13.116
 Pentacene 3.96
 Pentachlorophenol 4.140
 Pentachlorophenoxy ion 4.179
 1,3-Pentadiene, 4-methyl- 2.300
 1,3-Pentadiene, 1-phenyl-, (*E,E*)- 2.301
(E)-1,3-Pentadiene 2.328
(Z)-1,3-Pentadiene 2.329
 Pentadienylium, 1,1,5,5-tetrakis[4-(diethylamino)phenyl]- 11.110
 Pentamethoxybenzene 3.48
N,N,2,4,6-Pentamethylaniline 9.15
 1,3,3,7,7-Pentamethylbicyclo[2.2.1]heptane-2-thione 13.
 Pentamethylene sulfide 13.145
 1,2,2,6,6-Pentamethylpiperidine 8.61
 1,2,2,6,6-Pentamethylpiperidin-4-ol 8.72
 Pentanamide, 2-[(4-diethylamino)-2-methylphenyl]imino-4,4-dimethyl-3-oxo-*N*-phenyl- 11.111
 Pentanamide, 2-[(4-diethylamino)phenyl]imino-4,4-dimethyl-3-oxo-*N*-phenyl- 11.112
 Pentane 17.50
 Pentane, 3-(methoxymethylene)-2,2,4,4-tetramethyl- 2.355
n-Pentane 17.50
 Pentanedioic acid, 2-oxo- 17.51
 2,4-Pentanedione iron(III) derivative 12.174
 2,4-Pentanedione nickel(II) derivative 12.10
 3-Pantanethione, 2,2,4,4-tetramethyl- 13.117
 Pentanoic acid 17.79
 1-Pentanol, 5-(4-methylphenyl)thio- 13.118
 3-Pentanol, 3,4-dimethyl-1-[(4-methylphenyl)thio]- 13.1
 1-Pentene 2.302
 1-Pentene, 2,3,4-trimethyl- 2.303
 2-Pentene, (*E*)- 2.304
 2-Pentene, (*Z*)- 2.305
 2-Pentene, 2,4-dimethyl- 2.306
 2-Pentene, 2-methyl- 2.307, 2.66, 2.86, 2.92, 2.122, 2.243, 2.250, 2.256, 2.311, 2.312, 2.343, 3.71, 4.32, 4.102, 4.147, 13.87, 15.1
 2-Pentene, 3-methyl- 2.308
 2-Pentene, 3-methyl-, (*E*)- 2.309

- 2-Pentene, 3-methyl-, (*Z*)- **2.310**
 2-Pentene, 4-methyl-, (*E*)- **2.311**
 2-Pentene, 4-methyl-, (*Z*)- **2.312, 2.116**
 2-Pentene, 2,3,4-trimethyl- **2.313**
 2-Pentene, 2,4,4-trimethyl- **2.314**
cis-2-Pentene **2.305**
(E)-2-Pentene **2.304**
trans-2-Pentene **2.304**
(Z)-2-Pentene **2.305**
 4-Pentenoic acid **2.315**
 2-Penten-1-ol, 3,4-dimethyl-, (*E*)- **2.316**
 2-Penten-1-ol, 3,4-dimethyl-, (*Z*)- **2.317**
 2-Penten-1-ol, 3-ethyl- **2.318**
 2-Penten-1-ol, 3-methyl-, (*E*)- **2.319**
 2-Penten-1-ol, 3-methyl-, (*Z*)- **2.320**
 2-Penten-1-ol, 3,4,4-trimethyl-, (*E*)- **2.321**
 2-Penten-1-ol, 3,4,4-trimethyl-, (*Z*)- **2.322**
 2-Penten-4-ol, 2-methyl- **2.323**
 3-Penten-2-ol, 4-methyl- **2.323**
 3-Penten-2-one, 3,4-dimethyl- **2.324**
 Perinaphthenone **3.97**
 Permanax 45 **6.72**
 Perspex **17.57**
 3,10-Perylenedione, 4,9-dihydroxy-1,12-bis(2-hydroxypropyl)-2,6,7,11-tetramethoxy- **16.53**
 Perylo[1,12-*def*]-1,3-dioxepin-5,11-dione, 6-amino-12-hydroxy-8,9-bis(2-hydroxypropyl)-7,10-dimethoxy- **16.54**
 Perylo[1,12-*def*]-1,3-dioxepin-5,11-dione, 6,12-dihydroxy-8,9-bis(2-hydroxypropyl)-7,10-dimethoxy- **16.55**
 Perylo[1,12-*def*]-1,3-dioxepin-5,12-dione, 6,11-diamino-8,9-bis(2-hydroxypropyl)-7,10-dimethoxy- **16.56**
 Perylo[1,12-*def*]-1,3-dioxepin-6,11-dione, 8,9-bis(2-hydroxypropyl)-5,7,10,12-tetramethoxy- **16.57**
 Perylo[1,12-*def*]-1,3-dioxepin-6,11-dione, 5-hydroxy-8,9-bis(2-hydroxypropyl)-7,10,12-trimethoxy- **16.58**
 Phenalenone **3.97**
 Phenalen-1-one **3.97**
 1-Phenanthrenecarboxylic acid, 1,2,3,4,4a,4b,5,6,10,10a-decahydro-1,4a-dimethyl-7-(1-methylethyl)-, [1*R*-(1*α*,4*α**β*,4*β**α*,10*αα*)]- **2.325**
 Phenanthro[1,10,9,8-*opqra*]perylene-7,14-dione, 1,3,4,6,8,13-hexahydroxy-10,11-dialkyl- **16.59**
 Phenanthro[1,10,9,8-*opqra*]perylene-7,14-dione, 1,3,4,6,8,13-hexahydroxy-10,11-dimethyl- **16.60**
 Phenethylamine **8.48**
 α-Phenethylfurfuryl alcohol **5.71**
 Phenol **4.92**
 Phenol, 4-acetyl- **4.93**
 Phenol, 4-acetyl-2,6-bis(1,1-dimethylethyl)- **4.94**
 Phenol, 2-acetyl-5-dodecyloxy- **4.42**
 Phenol, 4-amino- **4.95**
 Phenol, 2-(benzotriazol-2-yl)-4,6-bis(1,1-dimethylpropyl)- **4.96**
 Phenol, 2-(benzotriazol-2-yl)-6-chloro-4-(1,1-dimethylethyl)- **4.97**
 Phenol, 2-benzyl- **4.98**
 Phenol, 2,6-bis(1,1-dimethylethyl)- **4.99, 2.280**
 Phenol, 2,4-bis(1,1-dimethylethyl)-5-methoxy- **4.100**
 Phenol, 3,5-bis(1,1-dimethylethyl)-4-methoxy- **4.101**
 Phenol, 2,6-bis(1,1-dimethylethyl)-4-methyl- **4.102, 4.99, 4.116, 4.117, 4.118, 4.147**
 Phenol, 2,6-bis(1,1-dimethylethyl)-4-phenyl- **4.103**
 Phenol, 2,6-bis(1,1-dimethylethyl)-4-(phenylmethyl)- **4.104**
 Phenol, 4-bromo-2,6-bis(1,1-dimethylethyl)- **4.105**
 Phenol, 2-*tert*-butyl- **4.120**
 Phenol, 4-*tert*-butyl- **4.121**
 Phenol, 2-chloro- **4.106**
 Phenol, 3-chloro- **4.107**
 Phenol, 4-chloro- **4.108**
 Phenol, 2-(5-chlorobenzotriazol-2-yl)-4,6-bis(1,1-dimethylethyl)- **4.109**
 Phenol, 4-chloro-2,6-di(1,1-dimethylethyl)- **4.110**
 Phenol, 2-chloro-4-hydroxy- **4.20**
 Phenol, 4-chloro-3-hydroxy- **4.11**
 Phenol, 3-chloro-5-methoxy- **4.111**
 Phenol, 4-cyano- **4.112**
 Phenol, 2,4-dichloro- **4.113**
 Phenol, 2,6-dichloro- **4.114**
 Phenol, 2,6-dimethoxy- **4.115**
 Phenol, 2,6-dimethoxy-, ion(1-)- **4.161**
 Phenol, 2,4-dimethyl- **4.116**
 Phenol, 2,6-dimethyl- **4.117**
 Phenol, 3,4-dimethyl- **4.118**
 Phenol, 4-(1,1-dimethylethoxy)-2,6-bis(1,1-dimethylethyl)- **4.119**
 Phenol, 3-(1,1-dimethylethyl)-4-methoxy- **4.122**
 Phenol, 2,4-dinitro- **4.123**
 Phenol, 2,5-dinitro- **4.124**
 Phenol, 2,6-dinitro- **4.125**
 Phenol, 4-ethoxy-2,3,5,6-tetramethyl- **4.126**
 Phenol, 4-(1-hydroxy-2,2-dimethyl)propyl-2-methoxy- **4.127**
 Phenol, 2-(2-hydroxyphenyl)- **4.128**
 Phenol, 2-methoxy- **4.129**
 Phenol, 3-methoxy- **4.130**
 Phenol, 4-methoxy- **4.131**
 Phenol, 4-methoxy-2,6-dimethyl- **4.132**
 Phenol, 4-methoxy-2,3,5,6-tetramethyl- **4.133**
 Phenol, 4-methoxy-2,3,6-trimethyl- **4.134**
 Phenol, 4-methyl- **4.135**
 Phenol, 2,2'-methylenebis(6-*tert*-butyl-4-methyl- **4.136**
 Phenol, 4-(methylthio)- **13.120**

- Phenol, 2-nitro- 4.137
 Phenol, 3-nitro- 4.138
 Phenol, 4-nitro- 4.139
 Phenol, pentachloro- 4.140
 Phenol, 2-phenyl- 4.141
 Phenol, 3-(phenylamino)- 4.142
 Phenol, 2-[phenylimino)methyl]-, *N*-oxide 15.7
 Phenol, 2,3,5,6-tetrafluoro- 4.143
 Phenol, 4-(1,1,3,3-tetramethylbutyl)-, salicylate 4.144
 Phenol, 2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)-] 4.145
 Phenol, 2,4,6-tri-*tert*-butyl- 4.147
 Phenol, 2,4,6-trichloro- 4.146
 Phenol, 2,4,6-trimethyl- 4.148
 Phenol, 2,4,6-triphenyl- 4.149
 Phenolate ion 4.150
 Phenothiazine 13.121
 Phenothiazine, 2-chloro-10-dimethylaminopropyl- 13.122
 Phenothiazine, 10-methyl- 13.123
 Phenothiazine, 10-(12-sulfonatododecyl)- 13.124
 Phenothiazine-10-propanamine, 2-chloro-*N,N*-dimethyl- 13.122
 Phenothiazinium, 3,7-bis(dimethylamino)- 11.113
 Phenothiazinium, 3,7-diamino- 11.114
 12-(10'-Phenothiazinyl)dodecyl-1-sulfonate ion 13.124
 Phenoxazine 6.38
 Phenoxide ion 4.150
 Phenoxide ion, 4-acetyl- 4.151
 Phenoxide ion, 2-benzyl- 4.152
 Phenoxide ion, 4-bromo- 4.153
 Phenoxide ion, 4-bromo-2,6-dimethyl- 4.154
 Phenoxide ion, 2-chloro- 4.155
 Phenoxide ion, 3-chloro- 4.156
 Phenoxide ion, 4-chloro- 4.157
 Phenoxide ion, 4-cyano- 4.158
 Phenoxide ion, 2,4-dichloro- 4.159
 Phenoxide ion, 2,6-dichloro- 4.160
 Phenoxide ion, 2,6-dimethoxy- 4.161
 Phenoxide ion, 2,6-dimethyl- 4.162
 Phenoxide ion, 4-(1,1-dimethylethyl)- 4.163
 Phenoxide ion, 2,4-dinitro- 4.164
 Phenoxide ion, 4-fluoro- 4.165
 Phenoxide ion, 4-hydroxy-, conjugate base 4.13
 Phenoxide ion, 2-(2-hydroxyphenyl)- 4.166
 Phenoxide ion, 5-hydroxy-2-phenyl- 4.167
 Phenoxide ion, 4-iodo- 4.168
 Phenoxide ion, 2-methoxy- 4.169
 Phenoxide ion, 3-methoxy- 4.170
 Phenoxide ion, 4-methoxy- 4.171
 Phenoxide ion, 4-methyl- 4.172
 Phenoxide ion, 2-methyl-4,6-dinitro- 4.173
 Phenoxide ion, 4-methyl-2,6-dinitro- 4.174
 Phenoxide ion, 4-methyl-2-nitro- 4.175
 Phenoxide ion, 2-nitro- 4.176
 Phenoxide ion, 3-nitro- 4.177
 Phenoxide ion, 4-nitro- 4.178
 Phenoxide ion, pentachloro- 4.179
 Phenoxide ion, 2-phenyl- 4.180
 Phenoxide ion, 4-phenyl- 4.181
 Phenoxide ion, 2,4,6-trichloro- 4.182
 2-(Phenoxy methylidene)adamantane 2.350, 2.351, 2.352, 2.354
 Phenylalanine 10.31
 L-Phenylalanine, 3,4-dihydroxy- 4.183
 o-Phenylanisole 3.59
 9-Phenylanthracene 3.14
 Phenyl arachidonate 2.214
 1-Phenylazo-2-naphthol 11.104, 11.94, 11.95, 11.96, 11.97, 11.98, 11.99, 11.100, 11.102, 11.103
 4-Phenylazo-1-naphthol 11.92, 11.87, 11.89, 11.90, 11.91
 1-(Phenylazo)-1-(phenylaminocarbonyl)-2-propanone 11.62
 2-Phenyl-1,2-benzisoselenazol-3-one 13.40
 2-Phenyl-1,2-benzisothiazol-3-one 13.41
 2-Phenylbenzo[*b*]cyclopenta[*e*]pyran 5.8
 Phenyl benzyl nitrone 15.12
 α -Phenyl-*N*-benzyl nitrone 15.12
 4-Phenylbutylamine 8.49
 Phenyl-*N*-*tert*-butyl nitrone 15.46
 2-Phenylcyclopenta[*b*][1]benzopyran 5.8
 4-Phenyl-2,6-di-*tert*-butylphenol 4.103
 6-Phenyl-3,4-dihydropyran-5-carboxylic acid ethyl ester 5.100
 Phenyl disulfide 13.84
 2,2'-(1,2-Phenylenebis(nitrilomethylidyne))-bis[phenolato]nickel(II) 12.166
 [*N,N'*-*o*-Phenylenebis(5-sulfosalicylideneiminato)]nickelate(II) disodium salt 12.167
m-Phenylenediamine, *N,N,N',N'*-tetramethyl- 9.27
o-Phenylenediamine 9.25
o-Phenylenediamine, *N,N,N',N'*-tetramethyl- 9.26
p-Phenylenediamine, *N*-cyclohexyl-*N'*-phenyl- 9.28
p-Phenylenediamine, *N,N'*-diphenyl- 9.29
p-Phenylenediamine, *N*-(1-methylethyl)-*N'*-phenyl- 9.3
p-Phenylenediamine, *N,N,N',N'*-tetramethyl- 9.31
 1-Phenylethanol 3.55
 2-Phenylethylamine 8.48
 Phenyl fluorenyl nitrone 15.6
 2-Phenylfuran 5.50
 3-Phenylfuran 5.51
 α -Phenylfurfuryl alcohol 5.70
 Phenylhydrazine 9.32
 Phenylhydroquinone 4.30

- Phenyl linoleate **2.266**
 Phenyl linolenate **2.270**
 5-(Phenylmethyl)-3-furanyl methyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate **5.76**
 Phenyl 4-methylphenyl nitrone **15.9**
 4-[(Phenylmethyl)thio]morpholine **13.112**
 1-[(Phenylmethyl)thio]piperidine **13.125**
 Phenyl 9,12-octadecadienoate **2.266**
 Phenyl 9-octadecenoate **2.273**
 Phenyl oleate **2.273**
 (E,E)-1-Phenyl-1,3-pentadiene **2.301**
 4-Phenylphenolate anion **4.181**
 2-Phenylphenoxyde **4.180**
 4-Phenylphenoxyde ion **4.181**
 N-Phenyl-2-(phenylmethyl)selenobenzamide **13.4**
 N-Phenyl-2-(phenylmethyl)thiobenzamide **13.5**
 (E)-1-Phenylpropene **3.119**
 (Z)-1-Phenylpropene **3.120**
 3-Phenylpropylamine **8.50**
 α -(3-Phenylpropyl)furfuryl alcohol **5.73**
 β -Phenyl- α -pyrrolidinylbenzeneethanol **8.2**
 Phenylpyruvic acid **17.60**
 Phenyl sulfide **13.34, 13.16, 13.23, 13.26, 13.31**
 (E)-2-Phenylsulfinyl-2-butene **13.54**
 (Z)-2-Phenylsulfinyl-2-butene **13.55**
 1-Phenylsulfinyl-2-methylpropene **2.338**
 9-(Phenylsulfonyl)fluorene anion **13.98**
 9-(Phenylsulfonyl)fluoren-9-yl anion **13.98**
 (E)-1-(Phenylsulfonyl)-3-(β -methoxylvinyl)indole **2.247, 2.249**
 (Z)-1-(Phenylsulfonyl)-3-(β -methoxylvinyl)indole **2.249**
 3-Phenylsydnone **5.107**
 1-Phenyl-4,4,8,8-tetramethyl-2,3,4,5,7,8,9,10-octahdropyrrolo[4,3,2-m,n]acridine-10-one **6.69**
 4-Phenylthiobzenophenone **13.163**
 3-Phenylthio-1-propanol **13.137**
 Pheophytin *a* **7.26**
 Pheophytin *b* **7.27**
 Phleochrome **16.53**
 20-Phorbinecarboxylic acid, 3,4-didhydro-3,4,8,9,13,14,18,19-octaethyl-18,19-dihydro-, ethyl ester **7.28**
 20-Phorbinecarboxylic acid, 3,4-didehydro-3,4,8,9,13,14,18,19-octaethyl-18,19-dihydro-, ethyl ester, dichlorotin(IV) **7.29**
 20-Phorbinecarboxylic acid, 3,4,20,21-tetradehydro-3,4,8,9,13,14,18,19-octaethyl-18,19-dihydro-, ethyl ester **7.30**
 20-Phorbinecarboxylic acid, 3,4,20,21-tetradehydro-4,9,14,19-tetraethyl-18,19-dihydro-3,8,13,18-tetramethyl-, ethyl ester, zinc(II) **7.31**
 3-Phorbinepropanoic acid, 9-acetyl-14-ethyl-13,14-dihydro-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, 3,7,11,15-tetramethyl-2-hexadecenyl ester **7.32**
 3-Phorbinepropanoic acid, 9-acetyl-14-ethylidene-13,14-dihydro-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, 3,7,11,15-tetramethyl-2-hexadecenyl ester **7.33**
 3-Phorbinepropanoic acid, 3,4-didehydro-4,9-diethenyl-14-ethyl-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, 3,7,11,15-tetramethyl-2-hexadecenyl ester, magnesium complex **7.34, 7.35**
 3-Phorbinepropanoic acid, 3,4-didehydro-9-ethenyl-14-ethyl-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, 2-amino-2-(methoxycarbonyl)ethyl ester, magnesium(II), **7.36**
 3-Phorbinepropanoic acid, 3,4-didehydro-9-ethenyl-14-ethyl-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, magnesium(II) **7.37**
 3-Phorbinepropanoic acid, 3,4-didehydro-9-ethenyl-14-ethyl-21-(methoxycarbonyl)-4,8,13,18-tetramethyl-20-oxo-, 3,7,11,15-tetramethyl-2-hexadecenyl ester **7.38, 7.39**
 Phosphatidylcholine **17.53**
 Phosphatidylethanolamine **17.52**
 5-O-Phosphono- β -D-ribofuranosyl-3-pyridinecarboxamide **16.51**
 Phosphoric acid, mono(2-aminoethyl) mono(2,3-dihydroxypropyl) ester **17.52**
 Phosphoric acid, mono(2-dimethylammonioethyl) mono(2,3-dihydroxypropyl) ester **17.53**
 Phthalazine-1,4-dione, 5-amino-2,3-dihydro- **9.33**
 Phthalocyanine, bis(tribenzylsiloxy)silicon **7.40**
 Phthalocyanine, bis(tripropylsiloxy)silicon **7.41**
 Phthalocyanine, 1,4,8,11,15,18,22,25-octabutoxy- **7.42**
 Phthalocyanine, sulfo-, chloroaluminum(III) **7.43**
 Phthalocyanine, sulfo-, zinc(II) **7.44**
 Phthalocyanine, tetracarboxy-, copper(II) **7.45**
 Phthalocyanine, 2,9,16,23-tetra(1,1-dimethylethyl)- **7.46**
 Phycol-400 **17.54**
 15-*cis*-Phytoene **2.74**
 15-(Z)-Phytoene **2.74**
 15-*cis*-Phytofluene **2.73**
 Phytol **2.238**
 Phytyl acetate **2.239**
 Pigment Red 1 **11.103**
 Pigment Red 3 **11.101**
 Pinacyanol chloride **11.41**
 α -Pinene **2.326, 13.103, 13.118, 13.137**
 β -Pinene **2.327**
 2-Pipecoline **8.59**
 Piperazine **8.51**
 1-Piperidinamine, *N,N*-dimethyl- **8.52**
 Piperidine **8.53**

- Piperidine, 4-amino-2,2,6,6-tetramethyl- 8.54
 Piperidine, 1-butoxy-2,2,6,6-tetramethyl- 8.55
 Piperidine, 1-cyclohexyl- 8.56
 Piperidine, 1,2-dimethyl- 8.57
 Piperidine, 2,6-dimethyl- 8.58
 Piperidine, 2-methyl- 8.59
 Piperidine, *N*-methyl- 8.60
 Piperidine, *N*-methyl-4-hydroxy- 8.71
 Piperidine, 1,2,2,6,6-pentamethyl- 8.61
 Piperidine, 1-[(phenylmethyl)thio]- 13.125
 Piperidine, 1-(1-pyrrolidinyl)- 8.62
 Piperidine, 2,2,6,6-tetramethyl- 8.63
 Piperidine, 2,2,6,6-tetramethyl-4,4'-(1,6-hexanediaminyl)bis- 8.64
 Piperidine, 2,2,6,6-tetramethyl-4,4'-(1,6-hexanediaminyl)bis-, polymer 8.65
 1-Piperidineethanol, diester with [[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]butylpropanedioic acid 8.66
 1-Piperidineethanol, 4-hydroxy-2,2,6,6-tetramethyl-, polymer with butanedioic acid 8.67
 1-Piperidineethanol, 2,2,6,6-tetramethyl- 8.68
 1-Piperidineethanol, 2,2,6,6-tetramethyl-, acetate (ester) 8.69
 Piperidin-1-ol, 2,2,6,6-tetramethyl- 15.28
 Piperidin-3-ol, 1-methyl- 8.70
 Piperidin-4-ol, 1-methyl- 8.71
 Piperidin-4-ol, 1,2,2,6,6-pentamethyl- 8.72
 Piperidin-4-ol, 1,2,2,6,6-pentamethyl-, diester with [[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]butylpropanedioic acid 8.73
 Piperidin-4-ol, 1,2,2,6,6-pentamethyl-, diester with 1,10-decanedioic acid 8.74
 Piperidin-4-ol, 1,2,2,6,6-pentamethyl-, triester with nitrilo-triacetic acid 8.75
 Piperidin-4-ol, 1,2,2,6-tetramethyl-, phenylacetate ester 8.76
 Piperidin-4-ol, 2,2,6,6-tetramethyl- 8.77, 8.19
 Piperidin-4-ol, 2,2,6,6-tetramethyl-, diester with 1,10-decanedioic acid 8.78
 Piperidin-4-ol, 2,2,6,6-tetramethyl-, ester with copolymer of styrene + methacrylic acid 8.79
 Piperidin-4-ol, 2,2,6,6-tetramethyl-, triester with nitrilo-triacetic acid 8.80
 1-Piperidinyloxy, 4-amino-2,2,6,6-tetramethyl- 15.29
 1-Piperidinyloxy, 4-azido-2,2,6,6-tetramethyl- 15.30
 1-Piperidinyloxy, 4,4'-(1,10-dioxo-1,10-decanediyl)bis(oxy)[2,2,6,6-tetramethyl- 15.37
 1-Piperidinyloxy, 4,4'-(1,6-dioxo-1,6-hexanediyil)bis(oxy)[2,2,6,6-tetramethyl- 15.38
 1-Piperidinyloxy, 4-hydroxy-4-(2-naphthyl)-2,2,6,6-tetramethyl-, 15.31
 1-Piperidinyloxy, 4-hydroxy-4-(2-phenylethynyl)-2,2,6,6-tetramethyl- 15.32
 1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl- 15.33
 1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, benzoate 15.34
 1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, butyrate 15.35
 1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, crotonate 15.36
 1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, diester with 1,10-decanedioic acid 15.37
 1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, diester with 1,6-hexanedioic acid 15.38
 1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, diester with terephthalic acid 15.39
 1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, methacrylate 15.40
 1-Piperidinyloxy, 4-hydroxy-2,2,6,6-tetramethyl-, pentanoate 15.41
 1-Piperidinyloxy, 4-iodo-2,2,6,6-tetramethyl- 15.42
 1-Piperidinyloxy, 4,4'-(1,4-phenylenebis(carbonyloxy))bis[2,2,6,6-tetramethyl- 15.39
 1-Piperidinyloxy, 2,2,6,6-tetramethyl- 15.43
 1-Piperidinyloxy, 2,2,6,6-tetramethyl-4-[(2-methyl-1-oxo-2-propenyl)oxy]- 15.40
 1-Piperidinyloxy, 2,2,6,6-tetramethyl-4-(2-oxo-1-butenyl)oxy- 15.36
 1-Piperidinyloxy, 2,2,6,6-tetramethyl-4-(1-oxobutoxy)- 15.35
 4-Piperidone, 2,2,6,6-tetramethyl-1-oxyl- 15.44
 (*E*)-Piperylene 2.328
 (*Z*)-Piperylene 2.329
 Pivalothiophenone 13.126
 Pivalothiophenone, 4'-chloro- 13.127
 Pivalothiophenone, 4'-fluoro- 13.128
 Pivalothiophenone, 4'-methoxy- 13.129
 Platinum(II) bis[1,2-di(4-ethylphenyl)-1,2-ethenedithiolato]- 12.43
 Platinum(II) 3,4-dimercaptotoluene(1,10-phenanthroline) 12.155
 Poly(*N,N'*-Bis[4-(2,2,6,6-tetramethylpiperidinyl)]-1,6-hexanediamine) 8.65
 Polybutadiene 2.330
cis-Poly(butadiene) 2.330
(Z)-Poly(butadiene) 2.330
 Polycarbonate 17.55
 Polyester of succinic acid and 4-hydroxy-1-(2-hydroxyethyl)-2,2,6,6-tetramethylpiperidine 8.67
 Poly(ethylene glycol) 17.56
 Polyisoprene 2.331
cis-Poly(isoprene) 2.331
(Z)-Poly(isoprene) 2.331

- Poly(methyl methacrylate) **17.57**
- Polystyrene **17.58**
- Polystyrene-substituted with 4-(2,2,6,6-tetramethylpiperidinyl) alkanoate **8.79**
- Poly[[6-[(1,1,3,3-tetramethylbutyl)amino]-1,3,5-triazine-2,4-diyl][(2,2,6,6-tetramethyl-4-piperidinyl)imino]-1,6-hexanediy[(2,2,6,6-tetramethyl-4-piperidinyl)imino]] **8.16**
- Porphinatozinc(II) **7.71**
- Porphine, (acetato)5,10,15,20-tetraphenyl-, iron(III) **7.47**
- Porphine, (acetato)5,10,15,20-tetraphenyl-, manganese(III) **7.48**
- Porphine, 2,3-dihydro-5,10,15,20-tetraphenyl- **7.50**
- Porphine, 2,3-dihydro-5,10,15,20-tetraphenyl-, cadmium(II) **7.51**
- Porphine, 2,3-dihydro-5,10,15,20-tetraphenyl-, copper(II) **7.52**
- Porphine, 2,3-dihydro-5,10,15,20-tetraphenyl-, zinc(II) **7.53**
- Porphine, 7,8,17,18-tetrahydro-5,10,15,20-tetraphenyl-, (*E*) **7.54**
- Porphine, 5,10,15,20-tetrakis(4-iodophenyl)-, copper(II) **7.55**
- Porphine, 5,10,15,20-tetrakis(4-methoxyphenyl)-, copper(II) **7.56**
- Porphine, tetrakis(4-sulfonatophenyl)- **7.57**
- Porphine, 5,10,15,20-tetrakis(4-sulfonatophenyl)- **7.57**
- Porphine, 5,10,15,20-tetrakis(4-sulfonatophenyl)-, dichlorotin(IV) **7.58**
- Porphine, 5,10,15,20-tetraphenyl- **7.59**
- Porphine, 5,10,15,20-tetraphenyl-, cadmium(II) **7.60**
- Porphine, 5,10,15,20-tetraphenyl-, chloroaluminum(III) **7.61**
- Porphine, 5,10,15,20-tetraphenyl-, cobalt(II) **7.62**
- Porphine, 5,10,15,20-tetraphenyl-, copper(II) **7.63**
- Porphine, 5,10,15,20-tetraphenyl-, dichlorotin(IV) **7.64**
- Porphine, 5,10,15,20-tetraphenyl-, magnesium(II) **7.65**
- Porphine, 5,10,15,20-tetraphenyl-, manganese(III) **7.66**
- Porphine, 5,10,15,20-tetraphenyl-, nickel(II) **7.67**
- Porphine, 5,10,15,20-tetraphenyl-, zinc(II) **7.68**
- Porphine, zinc(II) **7.71**
- Porphine-2,18-dipropanoic acid, 7,12-bis(1-hydroxyethyl)-3,8,13,17-tetramethyl- **7.72**
- Porphine-2,18-dipropanoic acid, 7,12-diethenyl-3,8,13,17-tetramethyl- **7.73**
- Porphine-2,18-dipropanoic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, acetatoferrate(III), dimethyl ester **7.74**
- Porphine-2,18-dipropanoic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, cadmium(II) **7.75**
- Porphine-2,18-dipropanoic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, cobalt(II) **7.76**
- Porphine-2,18-dipropanoic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, copper(II) **7.77**
- Porphine-2,18-dipropanoic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, dimethyl ester **7.78**
- Porphine-2,18-dipropanoic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, dimethyl ester, nickel(II) **7.79**
- Porphine-2,18-dipropanoic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, nickel(II) **7.80**
- Porphine-2,18-dipropanoic acid, 7,12-diethenyl-3,8,13,17-tetramethyl-, zinc(II) **7.81**
- Porphine-2,18-dipropanoic acid, 7,12-diethyl-3,8,13,17-tetramethyl-, cobalt(II) **7.82**
- Porphine-2,18-dipropanoic acid, 7,12-diethyl-3,8,13,17-tetramethyl-, di[4-(diphenylmethylaminocarbonyl-2-nitrophenylmethyl] ester **7.83**
- Porphine-2,18-dipropanoic acid, 7,12-diethyl-3,8,13,17-tetramethyl-, diethyl ester, manganese(III) **7.84**
- Porphine-2,18-dipropanoic acid, 7,12-diethyl-3,8,13,17-tetramethyl-, dimethyl ester **7.85**
- Porphine-2,18-dipropanoic acid, 7,12-diethyl-3,8,13,17-tetramethyl-, dimethyl ester, cobalt(II) **7.86**
- Porphine-2,18-dipropanoic acid, 7,12-diethyl-3,8,13,17-tetramethyl-, nickel(II) **7.87**
- Porphine-2,18-dipropanoic acid, 7-[2-(dimethylamino)-2-oxoethyl]-8-ethyl-7,8-dihydro-3,7,12,17-tetramethyl-, dimethyl ester, (*Z*) **7.88**
- Porphine-2,18-dipropanoic acid, 7-[2-(dimethylamino)-2-oxoethyl]-8-ethylidene-7,8-dihydro-3,7,12,17-tetramethyl-, dimethyl ester **7.89**
- Porphine-2,18-dipropanoic acid, 8-ethenyl-13-ethyl-3,7,12,17-tetramethyl- **7.90**
- Porphine-2,18-dipropanoic acid, 8-ethenyl-13-ethyl-3,7,12,17-tetramethyl-, dimethyl ester **7.91**
- Porphine-2,18-dipropanoic acid, 7-ethenyl-12-(1-hydroxyethyl)-3,8,13,17-tetramethyl- **7.92**
- Porphine-2,18-dipropanoic acid, 7-ethenyl-3,8,13,17-tetramethyl- **7.93**
- Porphine-2,18-dipropanoic acid, 3,7,12,17-tetramethyl-, dimethyl ester **7.94**
- Porphine-2,18-dipropanoic acid, 3,7,12,17-tetramethyl-, nickel(II) **7.95**
- Porphine-2,18-dipropanoic acid, 3,7,12,17-tetramethyl-, nickel(II), bis(piperidine) **7.96**
- 5-Porphinepropenoic acid, octaethyl-, ethyl ester **7.97**
- [22]Porphyrin-(2.2.2.2), octaethyl-, (di-*trans*) **7.98**
- [26] Porphyrin **7.99**
- [26] Porphyrin-(2.4.2.4), 2,3,10,11,16,17,24,25-octaethyl-5,6,7,8,19,20,21,22-octadehydro- **7.100**
- PPO **6.31**
- Pregna-5,16-dien-20-one, 3-(acetoxy)- (β) **16.61**
- Pregnenolone **16.62**
- Pregn-5-en-20-one, 3-hydroxy-, (β) **16.62**
- Proflavine **11.1**
- L-Proline, 1-(3-mercaptop-2-methyl-1-oxopropyl)-, (*S*)- **13.130**

- 2-Propanamine, 2-methyl-*N*-(1-methyl-4-pyridylmethylene)-, *N*-oxyl **15.45**
- 2-Propanamine, 2-methyl-*N*-phenylmethylene-, *N*-oxyl **15.46**
- 2-Propanamine, 2-methyl-*N*-(4-pyridylmethylene)-, *N,N*-dioxyl- **15.47**
- 2-Propanamine, 2-methyl-*N*-(4-pyridylmethylene)-, *N*-oxyl **15.48**
- 2-Propanamine, 2-methyl-*N*-(2-sulfonatophenyl)methylene-, *N*-oxyl **15.49**
- Propane, 1-(1,1-dimethylethylthio)- **13.58**
- Propane, 2-methyl-2-nitroso- **15.50**
- Propane, 1-methyl-1-(propylthio)- **13.59**
- Propanediamide, 2-[[4-diethylamino)-2-ethylphenyl]imino]-*N,N'*-diphenyl- **11.115**
- Propanedioic acid, [[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]butyl-, bis[2-(2,2,6,6-tetramethyl-1-piperidinyl)ethyl] ester **8.66**
- 1,3-Propanediol, 2-amino-2-(hydroxymethyl)- **17.59**
- 1,3-Propanedione, 2-[[2-ethyl-4-(diethylamino)phenyl]imino]-1,3-diphenyl- **11.116**
- Propanenitrile, 3-(diethylamino)- **8.81**
- 1-Propanethiol, 3-amino-2-hydroxy- **13.131**
- 1-Propanethione, 1-(4-chlorophenyl)-2,2-dimethyl- **13.127**
- 1-Propanethione, 1-(4-fluorophenyl)-2,2-dimethyl- **13.128**
- 1-Propanethione, 1-(4-methoxyphenyl)-2,2-dimethyl- **13.129**
- Propanoic acid, 2-oxo-3-phenyl- **17.60**
- 1-Propanol, 3-(3-chlorophenyl)thio- **13.132**
- 1-Propanol, 3-(4-chlorophenyl)thio- **13.133**
- 1-Propanol, 3-(dimethylamino)- **8.82**
- 1-Propanol, 3-(4-fluorophenyl)thio- **13.134**
- 1-Propanol, 3-(4-methoxyphenyl)thio- **13.135**
- 1-Propanol, 3-(4-methylphenyl)thio- **13.136**
- 1-Propanol, 3-(phenylthio)- **13.137**
- 2-Propanol **17.61**
- 2-Propanol, 1-amino-3-thio- **13.131**
- 2-Propanone **17.2**
- 2-Propanone, (1-methylethylidene)hydrazone **15.3, 2.40**
- 2-Propanone, 1-(4-methylphenylazo)-1-(phenylaminocarbonyl)- **11.60**
- 2-Propanone, 1-(phenylazo)-1-(phenylaminocarbonyl)- **11.62**
- 2-Propanone azine **15.3, 2.40**
- Propenal **2.1**
- 1-Propen-1-amine, *N,N*,2-trimethyl- **8.83**
- 1-Propene, 1-anisyl- **3.116**
- Propene, 1,1-bis(cyclobutyl)-2-methyl- **2.332**
- Propene, 1,1-bis(cyclopropyl)- **2.333**
- Propene, 1,1-bis(cyclopropyl)-2-methyl- **2.334**
- Propene, 1-cyclopropyl-2-methyl- **2.335**
- Propene, 1-ethoxy-2-methyl- **2.336**
- Propene, 2-methyl- **2.337**
- Propene, 2-methyl-1-naphthalenyl- **3.87**
- Propene, 2-methyl-1-phenyl- **3.114**
- Propene, 2-methyl-1-(phenylsulfinyl)- **2.338**
- 2-Propenoic acid, 3-(2-hydroxyphenyl)- **4.69**
- p*-Propenylanisole **3.116**
- 4,4'-(1,3-Propenyl)bis[2,6-di(1,1-dimethylethyl)pyrylium **11.159**
- Propylamine **8.84**
- Propylamine, 2-methyl- **8.85**
- Propylamine, *N*-propyl- **8.86**
- Propylamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) **12.168**
- [*N,N'*-Propylenebis(5-sulfosalicylideneiminato)]nickelate(II) disodium salt **12.169**
- Propylene sulfide **13.152**
- Propyn-1-amine, *N,N*-diethyl- **8.87**
- Protease Type VII **10.32**
- Protochlorophyll **7.39**
- Protochlorophyllide **7.37**
- Protochlorophyllide 2-amino-2-(methoxycarbonyl)ethyl ester **7.36**
- Protopheophytin **7.38**
- Protoporphyrin IX **7.73**
- Protoporphyrin IX, dimethyl ester **7.78**
- Pseudoisocyanine **11.45**
- Psoralen **5.86**
- Psoralen, 8-methoxy- **5.87**
- Pulegone **2.115**
- Purine, 6-amino- **16.63**
- Purin-2,6,8-trione, 7,9-dihydro-, ion(1-) **16.64**
- Pyran-4-*d*, 2,3-dihydro-4-methyl- **5.92**
- Pyran-4-*d*, 3,4-dihydro-4-methyl- **5.92**
- Pyran-4-*t*, 2,3-dihydro-4-methyl- **5.93**
- Pyran-4-*t*, 3,4-dihydro-4-methyl- **5.93**
- Pyran-5-*t*, 2,3-dihydro-4,4-dimethyl- **5.94**
- Pyran-5-*t*, 3,4-dihydro-4,4-dimethyl- **5.94**
- Pyran-5-*t*, 2,3-dihydro-4-methyl- **5.95**
- Pyran-5-*t*, 3,4-dihydro-4-methyl- **5.95**
- Pyran-6-*t*, 2,3-dihydro-4,4-dimethyl- **5.96**
- Pyran-6-*t*, 3,4-dihydro-4,4-dimethyl- **5.96**
- Pyran-6-*t*, 2,3-dihydro-4-methyl- **5.97**
- Pyran-6-*t*, 3,4-dihydro-4-methyl- **5.97**
- Pyran, 5-benzoyl-3,4-dihydro-6-phenyl- **5.89**
- Pyran, 3,4-dihydro- **5.90**
- Pyran, 3,4-dihydro-5,6-dimethyl- **5.91**
- Pyran, 5,6-dihydro-2,3-dimethyl- **5.91**
- Pyran, 3,4-dihydro-6-methyl-5-acetyl- **5.88**
- Pyran, 5-ethoxycarbonyl-3,4-dihydro-6-methyl- **5.98**
- Pyran, 5-ethoxycarbonyl-3,4-dihydro-6-phenyl- **5.100**
- Pyran-5-carboxylic acid, 3,4-dihydro-6-methyl-, ethyl ester **5.98**

- Pyran-5-carboxylic acid, 3,4-dihydro-6-(1-methylethyl)-, ethyl ester **5.99**
- Pyran-5-carboxylic acid, 3,4-dihydro-6-phenyl-, ethyl ester **5.100**
- Pyran-5-carboxylic acid, 6-ethyl-3,4-dihydro-, ethyl ester **5.101**
- Pyran-4-thione **13.138**
- Pyran-4-thione, 2,6-dimethyl- **13.139**
- Pyran-4-thione, 2,6-diphenyl- **13.140**
- Pyrazole, 4,5-dihydro-5-[4-(diethylamino)phenyl]-3-[2-[4-(diethylamino)phenyl]ethenyl]-1-phenyl- **11.117**
- Pyrazole, 4,5-dihydro-5-[4-(dimethylamino)phenyl]-3-[2-[4-(dimethylamino)phenyl]ethenyl]-1-phenyl- **11.118**
- Pyrazole, 4,5-dihydro-1,5-diphenyl-3-(2-phenylethenyl)- **11.119**
- Pyrazole, 4,5-dihydro-5-(4-methoxyphenyl)-3-[2-(4-methoxyphenyl)ethenyl]-1-phenyl- **11.120**
- Pyrazole, 3,4,4,5-tetramethyl- **15.3**
- Pyrazole dye 2 **11.155**
- Pyrazole dye 3 **11.22**
- Pyrazole dye 4 **11.24**
- Pyrazole dye 5 **11.25**
- Pyrazole dye 6 **11.21**
- Pyrazole dye 7 **11.23**
- Pyrazole-3-selone, 4-(aminomethylene)-2,4-dihydro-5-methyl-2-phenyl-, (Z)- **13.141**
- Pyrazole-3-thione, 4-(aminomethylene)-2,4-dihydro-5-methyl-2-phenyl- **13.142**
- 2-Pyrazoline, 3-(4-chlorophenyl)-1,5-diphenyl- **11.121**
- 2-Pyrazoline, 5-(4-chlorophenyl)-1,3-diphenyl- **11.122**
- 2-Pyrazoline, 5-deutero-1,3,5-triphenyl- **11.123**
- 2-Pyrazoline, 1,5-diphenyl-3-styryl- **11.119**
- 2-Pyrazoline, 3-(4-methoxyphenyl)-1,5-diphenyl- **11.124**
- 2-Pyrazoline, 5-(4-methoxyphenyl)-1,3-diphenyl- **11.125**
- 2-Pyrazolinone, 5-[*p*-methoxyphenyl]-3-[*p*-methoxystyryl]-1-phenyl- **11.120**
- 2-Pyrazoline, 3-(4-methylphenyl)-1,5-diphenyl- **11.126**
- 2-Pyrazoline, 5-(4-methylphenyl)-1,3-diphenyl- **11.127**
- 2-Pyrazoline, 1,3,5-triphenyl- **11.121, 11.122, 11.123, 11.124, 11.125, 11.126, 11.127**
- 3-Pyrazolinone, 4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2-phenyl- **11.128**
- 3-Pyrazolinone, 4-[4-[ethyl(2-hydroxyethyl)amino]-2-methylphenyl]imino-5-methyl-2-phenyl- **11.129**
- 3-Pyrazolinone, 4-[4-[(2-hydroxyethyl)ethylamino]phenyl]imino-5-methyl-2-phenyl- **11.130**
- 3-Pyrazolinone, 4-[4-[(2-methylsulfamylethyl)ethylamino]-2-methylphenyl]imino-5-methyl-2-phenyl- **11.78**
- Pyrazolo[1,2-*a*]benzotriazole **11.131**
- Pyrazolo[1,2-*a*]benzotriazole, 1,3-dimethyl- **11.132**
- Pyrazolo[1,2-*a*]benzotriazol-4-ium, hydroxide, inner salt **11.131**
- Pyrazolo[1,2-*a*][1,2]diazepine, hexahydro- **8.88**
- Pyrazol-3-one, 4-[(4-aminophenyl)imino]-2,4-dihydro-5-methyl-2-phenyl- **11.133**
- Pyrazol-3-one, 4-[(4-amino-2,3,5,6-tetramethylphenyl)imino]-2,4-dihydro-5-methyl-2-phenyl- **11.134**
- Pyrazol-3-one, 2-(4-bromophenyl)-4-(4-diethylamino-2-methylphenyl)-2,4-dihydro-imino-5-methylcarbamyl- **11.135**
- Pyrazol-3-one, 2-(3-chlorophenyl)-4-(4-diethylamino-2-methylphenyl)-2,4-dihydro-imino-5-methylcarbamyl- **11.136**
- Pyrazol-3-one, 2-(4-chlorophenyl)-4-(4-diethylamino-2-methylphenyl)-2,4-dihydro-imino-5-methylcarbamyl- **11.137**
- Pyrazol-3-one, 4-[[4-(diethylamino)-2,6-dimethylphenyl]imino]-2,4-dihydro-5-(benzoylamino)-2-phenyl- **11.138**
- Pyrazol-3-one, 4-[[4-(diethylamino)-2,6-dimethylphenyl]imino]-2,4-dihydro-5-(1,1-dimethylethyl)-2-phenyl- **11.139**
- Pyrazol-3-one, 4-[[4-(diethylamino)-2,6-dimethylphenyl]imino]-2,4-dihydro-5-methyl-2-phenyl- **11.140**
- Pyrazol-3-one, 4-[[4-(diethylamino)-2-methylphenyl]imino]-2,4-dihydro-5-(1,1-dimethylethyl)-2-phenyl- **11.141**
- Pyrazol-3-one, 4-(4-diethylamino-2-methylphenyl)imino-2,4-dihydro-5-methylcarbamyl-2-(3-methoxyphenyl)- **11.142**
- Pyrazol-3-one, 4-(4-diethylamino-2-methylphenyl)imino-2,4-dihydro-5-methylcarbamyl-2-(3-methylphenyl)- **11.143**
- Pyrazol-3-one, 4-[[4-(diethylamino)-2-methylphenyl]imino]-2,4-dihydro-5-methyl-2-phenyl- **11.144**
- Pyrazol-3-one, 4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2,4-dihydro-2-(4-nitrophenyl)- **11.145**
- Pyrazol-3-one, 4-(4-diethylamino-2-methylphenyl)imino-5-methylcarbamyl-2,4-dihydro-2-(2,4,6-trichlorophenyl)- **11.146**
- Pyrazol-3-one, 4-[[4-(diethylamino)phenyl]imino]-2,4-dihydro-5-methyl-2-phenyl- **11.147**
- Pyrazol-3-one, 2,4-dihydro-4-[(4-methoxyphenyl)imino]-5-methyl-2-phenyl- **11.148**
- Pyrazol-3-one, 4-[[4-(dimethylamino)-3,5-dimethylphenyl]imino]-2,4-dihydro-5-methyl-2-phenyl- **11.149**
- Pyrazol-3-one, 4-[[4-(dimethylamino)phenyl]imino]-2,4-dihydro-5-methyl-2-phenyl- **11.150**
- Pyrazol-3-one, 4-[[4-(dimethylamino)-2,3,5,6-tetramethylphenyl]imino]-2,4-dihydro-5-methyl-2-phenyl- **11.151**
- Pyrazol-3-one, 4-[[4-(diphenylamino)phenyl]imino]-2,4-dihydro-5-methyl-2-phenyl- **11.152**

- Pyrazol-3-one, 4,4'-(2,3,5,6-tetramethyl-1,4-phenylene)dinitrilo]bis[2,4-dihydro-5-methyl-2-phenyl- 11.153
- Pyrazolo[1,2-*a*]pyrazole, tetrahydro- 8.89
- Pyrazolo[1,2-*a*]pyrazole, tetrahydro-2,2,6,6-tetramethyl- 8.90
- Pyrazolo[5,1-*c*]-1,2,4-triazole, 7-[4-(*N*-ethyl- [*N*-(2-methylsulfonylamino)ethyl]amino- 2-methylphenylimino]-6-methyl-3-[4-[3-[(3-butyl-4-hydroxyphenoxy(dodecyl)methoxy)-carbonylamino)phenyl]propyl]- 11.154
- Pyrazolo[5,1-*c*]-1,2,4-triazole-6-carboxylic acid, 7-[[4-(diethylamino)-2-methylphenyl]imino]-3-phenyl-, ethyl ester 11.155
- Pyrazolo[1',2':2,3][1,2,3]triazolo[4,5-*a*]phenazin-4-iium, 1,3-dimethyl- 6.39
- Pyridazine, hexahydro-1,2-dimethyl- 8.91
- Pyridazine, octahydro- 8.92
- Pyridine 6.40
- Pyridine, 3,5-diacetyl-1,4-dihydro- 6.41
- Pyridine, 3,5-diacetyl-1,4-dihydro-2,6-dimethyl- 6.42
- Pyridine, 3,5-dibenzoyl-1,4-dihydro-2,6-dimethyl- 6.43
- Pyridine, 3-(1-methyl-2-pyrrolidinyl)- 8.93
- Pyridine, 1,2,3,6-tetrahydro-1-methyl-4-phenyl- 8.94
- 3-Pyridinecarboxamide, 1,4-dihydro-1-(5-*O*-phosphono-β-D-ribofuranosyl- 16.51
- Pyridine-3,5-dicarboxamide, 1,4-dihydro-2,6-dimethyl-, *N,N*-diphenyl- 6.44
- Pyridine-3,5-dicarboxamide, 1,4-dihydro-2,4,6-trimethyl-, *N,N*-diphenyl- 6.45
- Pyridine-3,5-dicarboxylic acid, 1,4-dihydro-2,6-dimethyl-, dicyclohexyl ester 6.46
- Pyridine-3,5-dicarboxylic acid, 1,4-dihydro-2,6-dimethyl-, diethyl ester 6.47
- Pyridine-3,5-dicarboxylic acid, 1,4-dihydro-2,6-dimethyl-, dihexyl ester 6.48
- Pyridine-3,5-dicarboxylic acid, 1,4-dihydro-2,6-dimethyl-, di(1-methylethyl) ester 6.49
- Pyridine-3,5-dicarboxylic acid, 1,4-dihydro-2,6-diphenyl-, diethyl ester 6.50
- Pyridine-3,5-dicarboxylic acid, 1,4-dihydro-2,4,6-trimethyl-, diethyl ester 6.51
- Pyridine-3,5-dicarboxylic acid, 2,6-dimethyl-, diethyl ester 6.52
- Pyridinium, 2,3-dihydro-1-methyl-4-phenyl- 6.53
- Pyridinium, 4-[(1,1-dimethylethyl)imino]methyl]-1-methyl- 15.45
- Pyridinium, 1-methyl- 6.54
- Pyridinium, 1-methyl-4-phenyl- 6.55
- Pyrido[2,1-*b*]benzothiazolium, 4-[(2,3-dihydro)pyrido[2,1-*b*]benzothiazol-4-yl)methylene]-1,2,3,4-tetrahydro-, iodide 11.156
- Pyrido[4,3-*b*]indole, *N*-acetyl-2,3,4,4*a*,5,9*b*-hexahydro-2,8-dimethyl- 6.57
- Pyrido[4,3-*b*]indole, 2,3,4,4*a*,5,9*b*-hexahydro-2,8-dimethyl-*cis*(-), 6.56
- Pyrido[4,3-*b*]indole, 1,2,3,4-tetrahydro-2,8-dimethyl- 6.58
- α-(4-Pyridyl)-*tert*-butyl nitrone 15.48
- α-(4-Pyridyl 1-oxide)-*N*-*tert*-butylnitrone 15.47
- 2-Pyrimidinamine, *N,N*,4,5-tetramethyl-6-(phenylmethoxy)- 16.65
- Pyrimidin-4-ol, 5-butyl-2-(dimethylamino)-6-methyl- 16.66
- Pyrimidin-4-ol, 5-butyl-2-(ethylamino)-6-methyl- 16.67
- Pyrimidin-4-ol, 2-(diethylamino)-6-methyl- 16.68
- Pyrimidin-4-ol, 2-(dimethylamino)-5,6-dimethyl- 16.69
- Pyrimidin-4-ol, 2-(dimethylamino)-6-methyl- 16.70
- 4-Pyrimidinone, 5-butyl-2-(dimethylamino)-6-methyl- 16.66
- 4-Pyrimidinone, 5-butyl-2-(ethylamino)-6-methyl- 16.67
- 4-Pyrimidinone, 2-(diethylamino)-6-methyl- 16.68
- 4-Pyrimidinone, 2-(dimethylamino)-5,6-dimethyl- 16.69
- 4-Pyrimidinone, 2-(dimethylamino)-6-methyl- 16.70
- Pyrocatechol 4.4
- Pyrrole 6.59
- Pyrrole, 3,4-dihydro-2,2,3-trimethyl-, 1-oxide 15.51
- Pyrrole, 3,4-dihydro-3,5,5-trimethyl-, 1-oxide 15.52
- Pyrrole, 2,5-dimethyl- 6.60
- Pyrrole, 1-(1,1-dimethylethyl)- 6.61
- Pyrrole, 2-(1,1-dimethylethyl)- 6.62
- Pyrrole, 3-(1,1-dimethylethyl)- 6.63
- Pyrrole, 1-methyl- 6.64
- Pyrrole-2,4-dicarboxylic acid, 5,5'-methylenebis[3-methyl- tetraethyl ester 6.65
- Pyrrole-3,5-dicarboxylic acid, 2,4-dimethyl-, diethyl ester 6.66
- 1-Pyrrolidinamine, *N,N*-dimethyl- 8.95
- Pyrrolidine 8.96
- 1-Pyrrolidinyloxy, 3-cyano-2,2,5,5-tetramethyl-4-nitro- 15.53
- 1-Pyrrolidinyloxy, 2,5-di-(4-hydroxy-3-nitrophenyl)-2,5-dimethyl- 15.54
- 1-Pyrrolidinyloxy, 2-(4-fluoro-3-nitrophenyl)-2,5,5-trimethyl- 15.55
- 1-(1-Pyrrolidinyl)piperidine 8.62
- 1-Pyrrolidine-1-oxyl, 5,5-dimethyl- 15.56
- Pyrrolo[2,3,4-*k*,*l*]acridine-10-one, 1-(4-bromophenyl)-2,3,5,7,8,9-hexahydro-4,4,8,8-tetramethyl- 6.67
- Pyrrolo[2,3,4-*k*,*l*]acridine-10-one, 2,3,5,7,8,9-hexahydro-1-(4-methoxyphenyl)-4,4,8,8-tetramethyl- 6.68
- Pyrrolo[2,3,4-*k*,*l*]acridine-10-one, 2,3,5,7,8,9-hexahydro-4,4,8,8-tetramethyl-1-phenyl- 6.69
- Pyrrolo[3',2':3,4]cyclopenta[1,2-*b*]pyridine, 2-(4-ethoxycarbonylphenyl)-1,8-dihydro- 6.70
- Pyrrol-2-one, 3,4-diethyl-1,5-dihydro-5-[(5-methylpyrrol-2-yl)methylene]- 7.101

- Pyrrol-2-one, 3,4-diethyl-5-[(4-ethyl-3,5-dimethylpyrrol-2-yl)methylene]-1,5-dihydro- **7.102**
- Pyrrol-2-one, 5-[(3,5-dimethylpyrrol-2-yl)methylene]-4-ethyl-1,5-dihydro-3-methyl- **7.103**
- Pyrrol-2-one, 5-[(4,5-dimethylpyrrol-2-yl)methylene]-4-ethyl-1,5-dihydro-3-methyl- **7.104**
- Pyrrol-2-one, 3-ethenyl-5-[(4-ethyl-3,5-dimethylpyrrol-2-yl)methylene]-1,5-dihydro-4-methyl- **7.105**
- Pyrrol-2-one, 4-ethenyl-5-[(4-ethyl-3,5-dimethylpyrrol-2-yl)methylene]-1,5-dihydro-3-methyl- **7.106**
- Pyrrol-2-one, 4-ethyl-1,5-dihydro-3-methyl-5-[(5-methylpyrrol-2-yl)methylene]- **7.107**
- Pyrrol-2-one, 5-[(4-ethyl-3,5-dimethylpyrrol-2-yl)methylene]-1,5-dihydro- **7.108**
- Pyrylium, 2,6-bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)selenopyran-4-ylidene]-3-propenyl]- **11.157**
- Pyrylium, 2,6-bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)telluropyan-4-ylidene]-3-propenyl]- **11.158**
- Pyrylium, 4,4'-(1,3-propenyl)bis[2,6-di(1,1-dimethylethyl)- **11.159**
- PZ 25 **13.41**
- PZ 51 **13.40**
- Quadricyclane **17.63**
- Quadricyclo[2.2.1.0^{2,6}.0^{3,5}]heptane **17.63**
- Quercetin **4.61**
- Quinaldine Blue **11.41**
- Quinoline **6.71**
- Quinoline, 1,2-dihydro-2,2,4-trimethyl-, homopolymer **6.72**
- Quinoline, 2-(1,3-dioxoindolidene)- **11.68**
- Quinoline, 8-hydroxy- **6.73**
- Quinoline, 6-methoxy- **6.74**
- Quinolinium, 1-ethyl-2-[(1-ethyl-2-quinolinylidene)methyl]-, iodide **11.45**
- Quinolinium, 1-ethyl-2-[3-(1-ethyl-2-quinolinylidene)-1-propenyl]-, chloride **11.41**
- Quinolinium, 1-ethyl-4-[3-(1-ethyl-4-quinolinylidene)-1-propenyl]- **11.42**
- Quinolinium, 1-ethyl-4-[3-(1-ethyl-4-quinolylidene)-1-propenyl]-, toluenesulfonate **11.43**
- 8-Quinolinol **6.73**
- Quinone **16.12**
- Quinophthalone **11.68**
- Quinuclidine **8.97**
- Reserpine **8.98**
- Resorcinol **4.10**
- Resorcinol dimethyl ether **3.39**
- 11-*cis*-Retinal **2.339**
- 11-(*Z*)-Retinal **2.339**
- 13-*cis*-Retinal **2.340**
- 13-(*Z*)-Retinal **2.340**
- (*all-E*)-Retinal **2.341**
- all-trans*-Retinal **2.341**
- Retinoic acid **2.342**
- (*all-E*)-Retinoic acid **2.342**
- all-trans*-Retinoic acid **2.342**
- (*all-E*)-Retinol **2.343**
- all-trans*-Retinol **2.343**
- Retinol acetate **2.344**
- Retinyl acetate **2.344**
- Rhodopsin **16.71**
- Riboflavin **16.72**
- RO 10-9359 **2.262**
- Rose Bengal dianion **11.66**
- Rubellin A **16.8**
- Rubidomycin **4.79**
- Rubrene **3.125, 3.8, 3.10, 3.24, 3.124, 5.83, 7.5, 7.7, 8.6, 8.19, 8.24, 8.27, 9.1, 9.6, 9.14, 9.16, 14.1, 14.2, 14.7, 15.44**
- Rubrene-2,3,8,9-tetracarboxylate ion **3.126, 3.88**
- Rutin **4.58**
- Saccharose **16.75**
- Salcomine **12.139**
- Salicylate ion **4.35**
- Sarcina phytoene (3 conjugated bonds) **2.74**
- Sarcina phytofluene **2.73**
- 9,10-Secococholesta-5,7,10(19)-trien-3-ol (3β) **2.361**
- Selenopyriium, 2,6-bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)selenopyran-4-ylidene]-3-propenyl]- **11.160**
- Silane, (bicyclo[2.2.1]hepta-2,5-dien-2-yloxy)trimethyl- **2.7**
- Silane, (bicyclo[2.2.1]hept-2-en-2-yloxy)trimethyl- **2.24**
- Silane, (2,5-cyclooctadien-1-yloxy)trimethyl- **2.132**
- Silane, (7,7-dimethylbicyclo[2.2.1]hepta-2,5-dien-2-yloxy)trimethyl- **2.5**
- Silane, (7,7-dimethylbicyclo[2.2.1]hept-2-en-2-yloxy)trimethyl- **2.5, 2.7, 2.24**
- Silicon, [2,3-*b*:2',3'-*g*:2'',3''-*l*:2'',3''-*q*]porphyrazinatobis(tribenzylsilanolato)- **7.40**
- Silicon, [2,3-*b*:2',3'-*g*:2'',3''-*l*:2'',3''-*q*]porphyrazinatobis(tripropylsilanolato)- **7.41**
- Silicon, tetranaphtho[2,3-*b*:2',3'-*g*:2'',3''-*l*:2'',3''-*q*]porphyrazinatobis(tribenzylsilanolato)- **7.17**
- Silicon, tetranaphtho[2,3-*b*:2',3'-*g*:2'',3''-*l*:2'',3''-*q*]porphyrazinatobis(trihexylsilanolato)- **7.18**
- Silicon, tetranaphtho[2,3-*b*:2',3'-*g*:2'',3''-*l*:2'',3''-*q*]porphyrazinatobis(triisobutylsilanolato)- **7.20**
- Singlet oxygen **17.64**
- Sitosteryl acetate **16.73**
- Skatole **6.16**
- Sodium dodecyl sulfate **17.25**
- Sodium lauryl sulfate **17.25**
- Solvent Yellow 14 **11.104, 11.94, 11.95, 11.96, 11.97, 11.98, 11.99, 11.100, 11.102, 11.103**
- Soybean oil **17.65, 2.62, 4.50**

- Spermidine **8.10**
 Spermine **8.11**
 Spheroidene **2.68**
(all-E)-Spheroidene **2.68**
(all-trans)-Spheroidene **2.68**
 Spheroindenone **2.69**
(all-E)-Spheroindenone **2.69**
(all-trans)-Spheroindenone **2.69**
 Spirilloxanthin **2.71**
(all-E)-Spirilloxanthin **2.71**
(all-trans)-Spirilloxanthin **2.71**
 2,2'-Spirobi[1-benzopyran]-6,6'-diol, 7,7'-bis(1,1-dimethylethyl)-3,3',4,4'-tetrahydro-4,4',4'-tetramethyl- **4.184**
 2,2'-Spirobi[1-benzopyran]-6,6'-diol, 3,3',4,4'-tetrahydro-4,4',4',7,7'-hexamethyl- **4.185**
 1,1'-Spirobi[indene]-5,5',6,6'-tetracarboxylic acid, 2,2',3,3'-tetrahydro-3,3,3',3'-tetramethyl-, tetramethyl ester **3.98**
 2,2'-Spirobis[7-*tert*-butyl-3,4-dihydro-6-hydroxy-4,4-dimethylbenzopyran] **4.184**
 2,2'-Spirobis[2-(7-*tert*-butyl-6-hydroxy-4,4-dimethyl-1-benzopyran)] **4.184**
 1,1'-Spirobis[2,3-dihydro-5,6-di(methoxycarbonyl)-3,3-dimethylindene] **3.98**
 2,2'-Spirobis[3,4-dihydro-6-hydroxy-4,4,7-trimethyl-1-benzopyran] **4.185**
 Spirocyclopropane[4,7]methanoindene, 4,5,6,7-tetrahydro- **2.345**
 Squalene **2.346**
 Squarylium Dye 1a **11.46**, **12.66**
 Squarylium Dye 3a **11.32**
 Squarylium Dye 1b **11.47**
 Squarylium Dye 1c **11.48**
 Stearic acid **17.66**
 Stigmasta-5,22-dien-3-ol acetate (3β) **16.74**
 Stigmasta-5-en-3-ol acetate (3β) **16.73**
 Stigmasteryl acetate **16.74**
cis-Stilbene **3.99**
 Stilbene, 4,4'-dimethoxy- **3.100**
 Stilbene, α,β -dimethoxy- **3.101**
 Stilbene, α,α' -dimethyl-, (*E*)- **3.102**, **3.103**
 Stilbene, α,α' -dimethyl-, (*Z*)- **3.103**
 Stilbene, α -methyl-, (*E*)- **3.104**
 Stilbene, α -methyl-, (*Z*)- **3.105**
 Stilbene (*Z*)- **3.99**
(Z)-Stilbene **3.99**
 Stilbene-2,2'-disulfonate ion, 4,4'-diacetamido- **3.106**
 Stilbene-2,2'-disulfonate ion, 4,4'-diamino- **3.107**
(E)-Stilbene oxide **17.49**
 Stobadine **6.56**
 Strychnidin-10-one **8.99**
 Strychnine **8.99**
l-Strychnine **8.99**
 Styrene **3.108**
 Styrene, 3-chloro- α,β,β -trimethyl- **3.109**
 Styrene, 4-chloro- α,β,β -trimethyl- **3.110**, **3.121**, **3.122**
 Styrene, 3-cyano- α,β,β -trimethyl- **3.111**
 Styrene, 4-cyano- α,β,β -trimethyl- **3.112**
 Styrene, α -cyclopropyl- β,β -trimethyl- **3.113**
 Styrene, β,β -dimethyl- **3.114**
 Styrene, 4-(dimethylamino)- α,β,β -trimethyl- **3.115**
 Styrene, 4-methoxy- β -methyl- **3.116**
 Styrene, 3-methoxy- α,β,β -trimethyl- **3.117**
 Styrene, 4-methoxy- α,β,β -trimethyl- **3.118**, **3.115**, **3.123**
 Styrene, β -methyl-, (*E*)- **3.119**
 Styrene, β -methyl-, (*Z*)- **3.120**
 Styrene, 3, α,β,β -tetramethyl- **3.121**
 Styrene, 4, α,β,β -tetramethyl- **3.122**, **3.109**, **3.111**, **3.112**, **3.117**, **3.118**, **3.123**
 Styrene, α,β,β -trimethyl- **3.123**, **2.96**
 Subtilisin B **10.32**
 Subtilisin BPN' **10.32**
 Subtilisin Novo **10.32**
 Succinate dehydrogenase **10.13**
 Sucrose **16.75**
 Sudan I **11.104**, **11.94**, **11.95**, **11.96**, **11.97**, **11.98**, **11.99**, **11.100**, **11.102**, **11.103**
 Sulfinylbismethane **13.81**
 10-(12-Sulfonatododecyl)phenothiazine **13.124**
 2-Sulfonatophenyl-*N*-*tert*-butylnitron **15.49**
 Superoxide dismutase **10.14**
 Superoxide ion **14.14**
 Superoxide radical anion **14.14**
 Sydnone, 3,4-diphenyl- **5.102**
 Sydnone, 4-methyl-3-(4-methylphenyl)- **5.103**
 Sydnone, 3-(4-methylphenyl)- **5.104**
 Sydnone, 3-methyl-4-phenyl- **5.105**
 Sydnone, 3-(4-methylphenyl)-4-phenyl- **5.106**
 Sydnone, 3-phenyl- **5.107**
 Tangeretin **4.62**
 Taxifolin **4.63**
 Telluropyrylum, 2,6-bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)telluropyran-4-ylidene]-3-propenyl]- **11.161**
 Telluropyrylum, 2,6-diphenyl-4-(2,6-diphenyltelluropyran-4-ylidene)methyl- **11.162**
 TEMPAMIN **15.29**
 [1,1',3',1''-Terphenyl]-2'-ol, 5'-phenyl- **4.149**
 α -Terpinene **2.105**
 Terpinolene **2.121**
 Tetra-*O*-acetylglucose-1-thiolato(triethylphosphino)gold(I) **12.170**
 1,4,5,8-Tetraaminoanthraquinone **11.14**

- 29,30,31,32-
Tetraazapentacyclo[24.2.1.1^{4,7}.1^{12,15}.1^{18,21}]dotriaconta-1,3,5,7,9,10,11,13,15(31),16,18,20,26(29),27-pentadecaen-22,24-diyne, 5,6,13,14,19,20,27,28-octaethyl- 7.100
- 25,26,27,28-
Tetraazapentacyclo[20.2.1.1^{4,7}.1^{10,13}.1^{16,19}]octacosa-1(25),2,4,6,8,10(27),11,13,15,17,19,21,23-tridecaene, 5,6,11,12,17,18,23,24-octaethyl- 7.98
- Tetrabenzofulvalene 3.66
- 2',4',5',7'-Tetrabromofluorescein dianion 11.65
- Tetrabutylammonium bis[1,2-diphenyl-1,2-ethenedithiolato]nickelate(I) 12.65
- Tetrabutylammonium bis[1,2-di(4-trifluoromethylphenyl)-1,2-ethenedithiolato]nickelate(I) 12.75
- 2,11,20,29-Tetra-*tert*-butylnaphthalocyanine 7.22
- Tetra(6-*tert*-butyl)-2,3-naphthalocyanine 7.22
- Tetra-*tert*-butylphthalocyanine 7.46
- Tetracene 3.124, 3.12, 7.53, 8.72, 8.77, 9.18, 15.46
- Tetracene, 5,6,11,12-tetraphenyl- 3.125, 3.8, 3.10, 3.24, 3.124, 5.83, 7.5, 7.7, 8.6, 8.19, 8.24, 8.27, 9.1, 9.6, 9.14, 9.16, 14.1, 14.2, 14.7, 15.44
- Tetracene-2,3,8,9-tetracarboxylate, 5,6,11,12-tetraphenyl- 3.126, 3.88
- 5,6,7,8-Tetrachloro-1,2,3,4-tetrahydro-9-isopropylidene-1,4-methanonaphthalene 2.258
- 3,4,5,6-Tetrachloro-2',4',5',7'-tetraiodofluorescein dianion 11.66
- 4,6,8,10,12,14,16,18,20-Tetracosanonaene-3,22-dione, 6,10,15,19-tetramethyl-2,2,23,23-tetramethoxy- (*all-E*) 2.347
- 4,6,8,10,12,14,16,18,20-Tetracosanonaene-2,3,22,23-tetraone, 6,10,15,19-tetramethyl- (*all-F*) 2.348
- Tetracyclo[3.2.0.0^{2,7}.0^{4,6}]heptane 17.63
- Tetracyclone 3.60
- Tetradecanoic acid 17.67
- 2,9,16,23-Tetra(1,1-dimethylethyl)phthalocyanine 7.46
- Tetraethoxyethene 2.225
- 2,3,5,6-Tetrafluorophenol 4.143
- 5,6,7,8-Tetrafluoro-1,2,3,4-tetrahydro-9-isopropylidene-1,4-methanonaphthalene 2.259
- Tetraglycine 10.18
- 7,8,11,12-Tetrahydro- ψ -carotene 2.72
- 1,2,3,4-Tetrahydro-5,8-dimethoxy-1,4-methanonaphthalene 3.73
- Tetrahydrofuran 17.68
- 1,2,3,4-Tetrahydro-9-isopropylidene-1,4-methanonaphthalene 2.260
- 4,5,6,7-Tetrahydro-4,7-methanoindene 2.254
- 1,2,3,4-Tetrahydro-1,4-methanonaphthalene 3.72
- 2,3,6,7-Tetrahydro-9-methyl[1]benzopyran[6,7,8-*i,j*]quinolizin-11-one 8.4
- Tetrahydro-2-methyl-5,8-methano-[1,2,4]triazolo[1,2-*a*]pyridazine-1,3-dione 17.41
- Tetrahydropyrazolo[1,2-*a*]pyrazole 8.89
- Tetrahydropyrrole 8.96
- 4,5,6,7-Tetrahydrospirocyclopropane[4,7]methanoindene 2.345
- Tetrahydro-2,2,6,6-tetramethylpyrazolo[1,2-*a*]pyrazole 8.90
- Tetrahydrothiophene 13.168
- Tetrahydrothiopyran 13.145
- 3,3',4',7-Tetrahydroxyflavone 4.75
- 2,2,3,3-Tetrakis(2,6-diethylphenyl)oxadisilirane 17.46
- 2,2,3,3-Tetrakis(2,6-diisopropylphenyl)oxadisilirane 17.47
- 5,10,15,20-Tetrakis(4-iodophenyl)porphine, copper(II) 7.55
- 5,10,15,20-Tetrakis(4-methoxyphenyl)porphinatocopper(II) 7.56
- Tetrakis(4-sulfonatophenyl)porphine 7.57
- 5,10,15,20-Tetrakis(4-sulfonatophenyl)porphine 7.57
- 1,1,2,2-Tetrakis(2,4,6-trimethylphenyl)disilirane 17.21
- 1,1,2,2-Tetramesityldisilirane 17.21
- 1,2,3,4-Tetramethoxybenzene 3.49
- 1,2,3,5-Tetramethoxybenzene 3.50
- 1,2,4,5-Tetramethoxybenzene 3.51, 3.42, 3.48, 3.50
- 2,2,6,6-Tetramethyl-4-aminopiperidine-*N*-oxyl 15.29
- N,N*,2,4-Tetramethylaniline 9.17
- (*all-E*)-3,7,12,16-Tetramethyl-1,18-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)-1,3,5,7,9,11,13,15,17-octadecanonaene 2.66, 2.183
- 4-(1,1,3,3-Tetramethylbutyl)phenyl salicylate 4.144
- 2,2,5,5-Tetramethyl-3-cyano-4-nitropyrrolidin-1-oxyl 15.53
- 2,2,4,4-Tetramethyl-1,3-cyclobutanedithione 13.60
- 2,2,4,4-Tetramethylcyclobutanethione 13.61
- (2,2,6,6-Tetramethylcyclohexylidene)methanethione 13.166
- 2,2,6,6-Tetramethylcyclohexylthioketene 13.166
- Tetramethylene sulfide 13.168
- Tetramethylethylene 2.51, 2.38, 2.50, 2.52, 2.53, 2.56, 2.103, 2.117, 2.140, 2.168, 2.170, 2.184, 2.185, 2.186, 2.187, 2.188, 2.189, 2.190, 2.191, 2.192, 2.193, 2.194, 2.195, 2.196, 2.223, 2.225, 2.303, 2.310, 2.332, 2.334, 3.10, 3.11, 3.28, 3.66, 3.125, 4.3, 4.100, 5.34, 7.5, 7.65, 8.83, 9.1, 13.86, 13.145
- 2,6,10,14-Tetramethyl-2,6,10,14-hexadecatetraene 2.235
- 3,7,11,15-Tetramethyl-2-hexadecen-1-ol 2.238
- Tetramethylhydrazine 8.41
- Tetra(methyl-*d*₃)hydrazine 8.42
- 2,2,6,6-Tetramethyl-4-hydroxypiperidine 8.77, 8.19
- 2,3,5,6-Tetramethylidene-7-oxabicyclo[2.2.1]heptane 2.296
- 2,3,5,6-Tetramethylidene-7-oxa-2-bicyclo[2.2.1]heptane monoendoperoxide 2.357
- 4,4'-[2,3,5,6-Tetramethylphenyl]benzene-3-methyl-1-phenyl-2-pyrazolin-5-one 11.153
- N,N,N',N'*-Tetramethyl-*m*-phenylenediamine 9.27

- N,N,N',N'-Tetramethyl-o-phenylenediamine* **9.26**
N,N,N',N'-Tetramethyl-p-phenylenediamine **9.31**
2,2,6,6-Tetramethylpiperidine **8.63**
2,2,6,6-Tetramethylpiperidine-N-oxyl **15.43**
2,2,6,6-Tetramethylpiperidin-1-ol **15.28**
2,2,6,6-Tetramethylpiperidin-4-ol **8.77, 8.19**
2,2,6,6-Tetramethyl-4-piperidinol N-oxyl **15.33**
2,2,6,6-Tetramethyl-4-piperidinol-1-oxyl benzoate **15.34**
2,2,6,6-Tetramethyl-4-piperidinol-1-oxyl butyrate **15.35**
2,2,6,6-Tetramethyl-4-piperidinol-1-oxyl crotonate **15.36**
2,2,6,6-Tetramethyl-4-piperidinol-1-oxyl methacrylate
15.40
2,2,6,6-Tetramethyl-4-piperidinol-1-oxyl pentanoate **15.41**
2,2,6,6-Tetramethyl-4-piperidone N-oxyl **15.44**
3,7,12,17-Tetramethylporphine-2,18-dipropanoic acid,
dimethyl ester **7.94**
Tetra(2-methylpropyl)hydrazine **8.43**
3,4,4,5-Tetramethylpyrazole **15.3**
3, α , β -Tetramethylstyrene **3.121**
4, α , β -Tetramethylstyrene **3.122, 3.109, 3.111, 3.112, 3.117,**
3.118, 3.123
6,10,15,19-Tetramethyl-4,6,8,10,12,14,16,18,20-
tetracosanonaene-2,3,22,23-tetraone (all-E) **2.348**
6,10,15,19-Tetramethyl-2,2,23,23-tetramethoxytetracos-
4,6,8,10,12,14,16,18,20-nonaene-3,22-dione (all-E)
2.347
3,3,6,6-Tetramethylthiacyclohept-4-ene **13.150**
3,3,6,6-Tetramethylthiacyclohept-4-yne **13.147**
3,3,6,6-Tetramethylthiacyclohept-4-yne 1,1-dioxide **13.148**
3,3,6,6-Tetramethylthiacyclohept-4-yne 1-oxide **13.149**
2,2,4,4-Tetramethyl-3-thioxocyclobutanone **13.62**
Tetramethyl-o-xylylene **2.104**
Tetranaphtho[2,3-b:2',3'-g:2'',3''-l:2''',3'''-q]porphyrazine,
1,6,10,15,19,24,28,33-octabutoxy-, palladium(II)
7.21
Tetranaphtho[2,3-b:2',3'-g:2'',3''-l:2''',3'''-q]porphyrazine,
2,11,20,20-tetrakis(1,1-dimethylethyl)- **7.22**
Tetraphene **3.23**
trans-Tetraphenylbacteriochlorin **7.54**
Tetraphenylchlorin **7.50**
Tetraphenylcyclopentadienone **3.60**
Tetraphenyl-p-dioxin **2.196**
Tetraphenylfuran **5.52**
2,3,4,5-Tetraphenylfuran **5.52**
1,3,4,7-Tetraphenylisobenzofuran **5.85**
5,6,11,12-Tetraphenylnaphthacene **3.125, 3.8, 3.10, 3.24,**
3.124, 5.83, 7.5, 7.7, 8.6, 8.19, 8.24, 8.27, 9.1, 9.6, 9.14,
9.16, 14.1, 14.2, 14.7, 15.44
[1,10,19,28-
Tetraphenylnaphthalocyanine]hydroxyaluminum(III)
7.23
5,10,15,20-Tetraphenylporphinatocadmium(II) **7.60**
5,10,15,20-Tetraphenylporphinato(chloro)aluminum(III)
ion **7.61**
5,10,15,20-Tetraphenylporphinatocobalt(II) **7.62**
5,10,15,20-Tetraphenylporphinatocupper(II) **7.63**
5,10,15,20-Tetraphenylporphinatomagnesium(II) **7.65**
5,10,15,20-Tetraphenylporphinatonickel(II) **7.67**
5,10,15,20-Tetraphenylporphinato(thiocyanato-S)iron(III)
7.70
5,10,15,20-Tetraphenylporphinatozinc(II) **7.68**
5,10,15,20-Tetraphenylporphine **7.59**
meso-Tetraphenylporphine **7.59**
5,6,11,12-Tetraphenyltetracene **3.125, 3.8, 3.10, 3.24, 3.124,**
5.83, 7.5, 7.7, 8.6, 8.19, 8.24, 8.27, 9.1, 9.6, 9.14, 9.16, 14.1,
14.2, 14.7, 15.44
Tetraphenyltetrahydroporphine **7.54**
2,2'-Thiacarbocyanine, 5,5'-dichloro-3,3-diethyl-, bromide
11.163
2,2'-Thiacarbocyanine, 5,5'-dichloro-3,9,3'-triethyl-,
bromide **11.164**
2,2'-Thiacarbocyanine, 5,5'-dicyano-3,9,3'-triethyl-,
tetrafluoroborate **11.165**
2,2'-Thiacarbocyanine, 3,3'-diethyl-, toluenesulfonate
11.166
2,2'-Thiacarbocyanine, 3,3'-diethyl-5,5'-dimethoxy-,
toluenesulfonate **11.167**
2,2'-Thiacarbocyanine, 3,3'-diethyl-8,10-dimethyl-,
toluenesulfonate **11.168**
2,2'-Thiacarbocyanine, 3,3'-diethyl-8,10-ethanediyl-,
toluenesulfonate **11.169**
2,2'-Thiacarbocyanine, 3,3'-diethyl-8,9-(1,3-propanediyl)-,
iodide **11.170**
2,2'-Thiacarbocyanine, 3,9,3'-triethyl-, bromide **11.171**
2,2'-Thiacarbocyanine, 3,3',9-triethyl-5,5'-dimethoxy-,
toluenesulfonate **11.172**
Thiacycloheptane **13.146**
Thiacyclohexane **13.145**
Thiacyclohex-2-ene, 2-methyl-, 1-oxide **13.143**
Thiacyclopent-2-ene, 2-methyl-, 1-oxide **13.144**
2,2'-Thiadicarbocyanine, 3,3'-diethyl-, iodide **11.173**
3-Thiaheptane **13.82**
2-Thiahexane **13.57**
Thiane **13.145**
2,2'-Triaticarbocyanine, 3,3'-diethyl-, iodide **11.174**
Thiepane **13.146**
Thiepin, 4,5-didehydro-2,3,6,7-tetrahydro-3,3,6,6-
tetramethyl- **13.147**
Thiepin, 4,5-didehydro-2,3,6,7-tetrahydro-3,3,6,6-
tetramethyl-, 1,1-dioxide **13.148**
Thiepin, 4,5-didehydro-2,3,6,7-tetrahydro-3,3,6,6-
tetramethyl-, 1-oxide **13.149**
Thiepin, 2,3,6,7-tetrahydro-3,3,6,6-tetramethyl- **13.150**
Thiirane, 2,3-diphenyl-, (E) **13.151**
Thiirane, methyl- **13.152**

- Thioanisole 13.29
 Thiobenzamide 13.153, 13.154, 13.155, 13.156, 13.157
 Thiobenzamide, 4-chloro- 13.153
 Thiobenzamide, 4-methoxy- 13.154
 Thiobenzamide, 4-methyl- 13.155
 Thiobenzamide, 4-nitro- 13.156
 Thiobenzamide, 4-(trifluoromethyl)- 13.157
 Thiobenzophenone, 4-chloro- 13.158
 Thiobenzophenone, 4,4'-dichloro- 13.159
 Thiobenzophenone, 4,4'-dimethoxy- 13.160
 Thiobenzophenone, 4,4'-dimethyl- 13.161
 Thiobenzophenone, 4-methoxy- 13.162
 Thiobenzophenone, 4-phenyl- 13.163
 2,2'-Thiobis[4-(1,1,3,3-tetramethylbutyl)phenol] 4.145
 [2,2'-Thiobis(4-(1,1,3,3-tetramethylbutyl)phenolato)](*n*-butylamine)nickel(II) 12.122
 Thiocamphenilone 13.44
 Thiocamphor 13.47
 Thioctic acid 13.90, 3.125
 2,2'-Thiodiethanol 13.164
 Thiofenchone 13.46
 Thioketene, di-*tert*-butyl- 13.165
 Thioketene, 2,2,6,6-tetramethylcyclohexyl- 13.166
 Thiola 13.101
 2-Thiol-L-histidine 13.106
 2-Thiol-L-histidine betaine 13.104
 2-Thiol-L-histidine betaine, anion 13.105
 Thionine cation 11.114
 Thionorcamphor 13.43
 Thiophene 13.167
 Thiophene, 2,3-dihydro-5-methyl-2-(1-methylethyl)-, 1-oxide 13.144
 Thiophene, tetrahydro- 13.168
 2,5-Thiophenedithione, 3,4-bis[(cyclohexylamino)methylene]dihydro-, (*Z,Z*) 13.169
 Thiophene-3-methanamine, 5-ethyl-2-thio-*N,N'*-(ethanediyl)bis- 13.171
 2-Thiophenethione, 3-[(cyclohexylamino)methylene]-5-ethyl-, (*Z*)- 13.170
 2-Thiophenethione, 3,3'-[1,2-ethanediyl]bis(iminomethylidyne)]bis[5-ethyl-, (*Z,Z*) 13.171
 Thiopropionylglycine 13.101
 Thiopropionylglycine, negative ion 13.102
 Thiopyran, 3,4-dihydro-6-methyl-, 1-oxide 13.143
 Thiopyran-4-thione 13.172
 Thiopyran-4-thione, 2,6-diphenyl- 13.173
 Thiopyrylium, 2,6-bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)selenopyran-4-ylidene]-3-propenyl]- 11.175
 Thiopyrylium, 2,6-bis(1,1-dimethylethyl)-4-[1-[2,6-bis(1,1-dimethylethyl)thiopyran-4-ylidene]-3-propenyl]- 11.176
 Thiosinamine 13.175
 Thiourea 13.174
 Thiourea, allyl- 13.175
 Thiourea, methyl- 13.176
 4-Thiouridine 16.76
 1,4-Thioxane 13.177
 α -Thujene 2.30
 Thymidine 16.77
 Thymidine monophosphate 16.78
 Thymidine 5'-monophosphate 16.78
 5'-Thymidylic acid 16.78
 Thymine 16.79
 Tiglic acid 2.58
 Tin, tetrานaphtho[2,3-*b*:2',3'-*g*:2'',3''-*l*:2'',3''-*q*]porphyratinatobis(trihexylsilanolato)- 7.19
 Tin(IV) dichloro{octaethylidihydropurpurin ethyl ester} 7.29
 Tin(IV) (dichloro)5,10,15,20-tetraphenylporphyrin 7.64
 Tinuvin 144 8.73
 Tinuvin 622 8.67
 Tinuvin 770 8.78
 Tocol 4.48
 α -Tocopherol 4.50
 β -Tocopherol 4.52
 γ -Tocopherol 4.53
 δ -Tocopherol 4.46
 α -Tocopherol ethyl ether 5.5
 α -Tocopherol hydroquinone 4.28
 β -Tocopherol hydroquinone 4.26
 γ -Tocopherol hydroquinone 4.27
 α -Tocopherol methyl ether 5.4
 α -Tocopheryl acetate 16.11
 Toluene 3.127
 Toluene-*d*₈ 3.128
 Toluene-*d*₈ 3.128
 α -Toluenethiol 13.42
 Toluidine Red 11.101
 p -Toluquinone 16.13
 3-Tosyloxy-1,3,5(10),9(11)-estratetraen-17-one 16.33
 1,2,4-Triazolobenzo[1,3]cyclopropa[1,2,3-*cd*]cyclopropa[*gh*]pentalene, 1,2,5,6,6*a*,6*b*,6*c*,6*d*-octahydro-1,6[1',2']-4-methyl-3,5-dioxo- 17.69
 1,2,3-Triazolo[1,2-*a*]benzotriazole 11.177
 1,2,4-Triazolo[1,3]cyclopropa[1,2,3-*cd*]cyclopropa[*gh*]pentalene, 1,2,5,6,6*a*,6*b*,6*c*,6*d*-octahydro-1,6[1',2']-4-methyl-3,5-dioxo- 17.70
 Tribenzo[*a,f,j*]perylene, 9,18-diphenyl- 3.129
 Tribenzyllamine 8.6
 Tributylamine 8.13

- 2,4,6-Tri-*tert*-butylphenol 4.147
 Tributyl phosphite 17.71
 Tricarbocyanine, 5,5'-dichloro-1,1',3,3,3',3'-hexamethyl-, iodide 11.178
 Tricarbocyanine, 5,5'-difluoro-1,1',3,3,3',3'-hexamethyl-, iodide 11.179
 Tricarbocyanine, 5,5'-dimethoxy-1,1',3,3,3',3'-hexamethyl-, iodide 11.180
 Tricarbocyanine, 1,1',3,3,3',3'-hexamethyl-, fluoride 11.181
 Tricarbocyanine, 1,1',3,3,3',3'-hexamethyl-, iodide 11.182
 Tricarbocyanine, 1,1',3,3,3',3'-hexamethyl-, perchlorate 11.183
 Tricarbocyanine, 1,1',3,3,3',3',5,5'-octamethyl-, iodide 11.184
 2,4,6-Trichlorophenol 4.146
 2,4,6-Trichlorophenoxyde ion 4.182
cis-9-Tricosene 2.349
(*Z*)-9-Tricosene 2.349
 Tricyclo[3.3.1.1^{3,7}]decane, 2-(3-chlorophenoxy)methylene)- 2.350
 Tricyclo[3.3.1.1^{3,7}]decane, 2-(4-chlorophenoxy)methylene)- 2.351
 Tricyclo[3.3.1.1^{3,7}]decane, 2-(hydroxymethylene)-, acetate 2.2
 Tricyclo[3.3.1.1^{3,7}]decane, 2-(methoxyphenoxy)methylene)- 2.352
 Tricyclo[3.3.1.1^{3,7}]decane, (1-methylethylene)- 2.353
 Tricyclo[3.3.1.1^{3,7}]decane, 2-(methylphenoxy)methylene)- 2.354
 Tricyclo[3.3.1.1^{3,7}]decane, (2-methylpropylidene)- 2.355
 Tricyclo[3.3.1.1^{3,7}]decane, 2-(phenoxy)methylene)- 2.350, 2.351, 2.352, 2.354
 Tricyclo[3.3.1.1^{3,7}]decane, tricyclo[3.3.1.1^{3,7}]decylidene- 2.356, 13.112
 Tricyclo[3.1.0.0^{2,6}]hex-3-ene 2.3
 Tricyclohexyl phosphite 17.72
 1,1,2-Tricyclopropylethene 2.169
 4-[1,5,5-Tri(4-diethylaminophenyl)pentadienylidene]anilinium perchlorate 11.110
 Triethanolamine[2,2'-thiobis[4-(1,1,3,3-tetramethylbutyl)]phenolato]nickel(II) 12.171
 1,1,2-Triethoxyethene 2.226
 Triethylamine 8.24
 3,9,3'-Triethyl-4,5,4',5'-dibenzo-2,2'-thiacarbocyanine bromide 11.106
 3,9,3'-Triethyl-5,5'-dimethoxy-2,2'-thiacarbocyanine toluenesulfonate 11.172
 Triethylenediamine 8.19
O,N,N-Triethylhydroxylamine 15.22
 Triethyl phosphite 17.71, 17.72, 17.73
 3,9,3'-Triethyl-2,2'-thiacarbocyanine bromide 11.171
 4-(Trifluoromethyl)thiobenzamide 13.157
 Triglycine 10.17
 Trihexylsiloxyaluminum naphthalocyanine 7.24
 Trihexylsiloxygallium naphthalocyanine 7.25
 4',5,7-Trihydroxyflavanone 4.72
 3,5,7-Trihydroxy-2-(4-hydroxyphenyl)-2-benzopyran-4-one 4.64
 1,2,3-Trimethoxybenzene 3.52
 1,2,4-Trimethoxybenzene 3.53, 3.51
 1,3,5-Trimethoxybenzene 3.54
 Trimethylamine 8.47
N,N,4-Trimethylaniline 9.19
 1,3,3-Trimethylbicyclo[2.2.1]heptane-2-thione 13.46
 1,7,7-Trimethylbicyclo[2.2.1]heptane-2-thione 13.47
 4,6,6-Trimethylbicyclo[3.1.1]hept-3-ene-2-thione 13.48
 3,5,5-Trimethyl-2-cyclohexene-1-thione 13.65
 4-(2,6,6-Trimethyl-1-cyclohexen-1-yl)-3-butene-2-one 2.61
(*E*)-2,6,9-Trimethyl-1,6-decadiene 2.171
(*Z*)-2,6,9-Trimethyl-1,6-decadiene 2.172
 1,1,2-Trimethyl-2-(2,2-dimethylpropyl)hydrazine 8.44
 2,6,10-Trimethyl-2,6,10-dodecatriene 2.210
 Trimethylhydroquinone 4.31
 1,2,3-Trimethylindole 6.18
 2-[7-(1,3,3-Trimethyl-2-indol-2-ylidene)-1-[4-(2,2-dimethoxyethyl)-1,3,5-heptatrienyl]-1,1,3-trimethylindolium, tetrafluoroborate 11.74
 2-[7-(1,3,3-Trimethyl-2-indol-2-ylidene)-1-[4-(1-piperidinio)-1,3,5-heptatrienyl]-1,1,3-trimethylindolium, bis(tetrafluoroborate) 11.75
 2-[7-(1,3,3-Trimethyl-2-indol-2-ylidene)-1-[4-[3-(1,3,3-trimethyl-2-indolylidene)-2-propenyl]-1,3,5-heptatrienyl]-1,1,3-trimethylindolium, tetrafluoroborate 11.76
 1,5,7-Trimethyl-3-methylidene-2,6,9,10-tetrathiaadamantane 2.253
 1,1,2-Trimethyl-2-(2-methylpropyl)hydrazine 8.45
 1,1,3-Trimethyl-2-naphthalenethione 13.114
 1,2,3-Trimethyl-5-nitroindole 6.19
 2,7,7-Trimethylnorborn-2-ene 2.23, 2.9, 2.11
 2,3,4-Trimethyl-1-pentene 2.303
 2,3,4-Trimethyl-2-pentene 2.313
 2,4,4-Trimethyl-2-pentene 2.314
(*E*)-3,4,4-Trimethyl-2-penten-1-ol 2.321
(*Z*)-3,4,4-Trimethyl-2-penten-1-ol 2.322
 2,4,6-Trimethylphenol 4.148
 Trimethyl phosphite 17.73, 17.74
 2,4,4-Trimethyl- Δ^1 -pyrroline-*N*-oxide 15.52
 4,5,5-Trimethyl- Δ^1 -pyrroline-*N*-oxide 15.51
 6-(Trimethylsiloxy)-1,4-cyclooctadiene 2.132
 2-(Trimethylsiloxy)norborna-2,5-diene 2.7
 2-(Trimethylsiloxy)norborn-2-ene 2.24
 α,β,β -Trimethylstyrene 3.123, 2.96

1,3,3-Trimethyl-2-[7-[1,3,3-trimethylindolidene]heptatrienyl]indolium perchlorate	11.183	Ubiquinol 10	4.25
8,9,11-Trioxatetracyclo[4.2.2.1.0 ^{5,6}]undec-5-ene, 2,3-bis(methylene)-	2.357	1,6-Undecadiene, 2,6-dimethyl-, (<i>E</i>)-	2.359
Tri(1,2,2,6,6-pentamethyl-4-piperidinol) nitrilotriacetate	8.75	1,6-Undecadiene, 2,6-dimethyl-, (<i>Z</i>)-	2.360
Triphenylamine	9.8	Undecanoic acid	17.76
1,1,3-Triphenylisoindole <i>N</i> -oxyl	15.25	Urate ion	16.64
Triphenyl nitrone	15.5	Urea	17.77
2,4,6-Triphenylphenol	4.149	Urea, allyl-	17.78
Triphenylphosphine	17.75	Uric acid anion	16.64
1,3,5-Triphenyl-2-pyrazoline	11.121, 11.122, 11.123, 11.124, 11.125, 11.126, 11.127	Uridine	16.80
Triphosphopyridine nucleotide	16.49	Valeric acid	17.79
Tris(acetylacetonato)chromate(III)	12.172	Vaska's complex	12.123
Tris(acetylacetonato)cobalt(III)	12.173	Veratrole	3.38
Tris(acetylacetonato)iron(III)	12.174	Vincamine	8.100
Tris(acetylacetonato)manganese(III)	12.175	Vinylacetic acid	2.59
Tris(2,2'-bipyridine)cobalt(II) ion	12.176	2-Vinylfuran	5.53
Tris(2,2'-bipyridine)ruthenium(II) dichloride	12.177	4-Vinylprotochlorophyll	7.34
Tris(hydroxymethyl)aminomethane	17.59	4-Vinylprotopheophytin	7.35
Tris(1,10-phenanthroline)cobalt(II) ion	12.178	Violaxanthin	2.80
Tri(2,2,6,6-tetramethyl-4-piperidinol) nitrilotriacetate	8.80	Vitamin A	2.343
(all- <i>E</i>)-1,3,5,7,9,11,13,15,17,19,21,23,25,27,29,31,33-Triacontaheptadecaene, 3,7,11,15,20,24,28,32-octamethyl-1,34-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)-	2.358	Vitamin A acid	2.342
Trolox C anion	4.44	Vitamin B ₂	16.72
Troxerutin	4.57	Vitamin D ₃	2.361
Trypaflavine cation	11.2	Vobasan-17-oic acid, 19,20-dihydro-3-oxo-, methyl ester, (20 α)	8.101
Trypsin	10.33	WR-1065	13.94
Tryptamine	10.34	Xanthophyll	2.75
Tryptamine, <i>N</i> -methoxycarbonyl-	10.35	Xanthotoxin	5.87
Tryptophan	10.36	2,4-Xylenol	4.116
Tryptophan, <i>N</i> -acetyl-	10.37	2,6-Xylenol	4.117
Tryptophan, <i>N</i> -acetyl-, methyl ester	10.38	3,4-Xylenol	4.118
Tryptophan, glycyl-	10.41	Zeaxanthin	2.79
L-Tryptophan, ethyl ester	10.40	Zinc(II), bis[3-(cyclohexylimino)methyl]-5-ethyl-2-thiophenethionato]-	12.23
Tryptophan, methyl ester	10.42	Zinc(II) bis(acetylacetone)	12.13
L-Tryptophan, L-alanyl-	10.39	Zinc(II) bis(dibutylthiocarbamate)	12.29
Tryptophanamide	10.43	Zinc(II) bis(diisopropylthiocarbamate)	12.51
L-Tryptophanamide, <i>N</i> -acetyl-	10.44	Zinc(II) bis(<i>O,O'</i> -diphenylthiophosphate)	12.72
Tryptophylglycine	10.21	Zinc(II) bis[2,2'-thiobis[<i>O,O'</i> -di(4- <i>tert</i> -butylphenyl)dithiophosphate]]	12.115
L-Tryptophylglycylglycine	10.22	Zinc(II) etiopurpurin ethyl ester	7.31
Tyramine	4.186	Zinc(II) porphyrin	7.71
Tyrosine	4.187	Zinc(II) protoporphyrin	7.81
Tyrosine, glycyl-	4.189	Zinc(II) protoporphyrin IX	7.81
L-Tyrosine, <i>N</i> -acetyl-, ethyl ester	4.188	Zinc(II) sulfophthalocyanine	7.44
Tyrosine, methyl ester	4.190	Zinc(II) tetraphenylchlorin	7.53
L-Tyrosylglycine	4.76	Zinc(II) tetraphenylporphyrin	7.68
Ubiquinol 0	4.21	Zinc(II) 5,10,15,20-tetraphenylporphyrin	7.68