

Estimation of the Thermodynamic Properties of Hydrocarbons at 298.15 K

Eugene S. Domalski and Elizabeth D. Hearing¹

Chemical Thermodynamics Division, Center for Chemical Physics, National Institute of Standards and Technology, Gaithersburg, Maryland 20899

Received September 11, 1986; revised manuscript received February 2, 1988

An estimation method developed by S.W. Benson and coworkers, for calculating the thermodynamic properties of organic compounds in the gas phase, has been extended to the liquid and solid phases for hydrocarbon compounds at 298.15 K. The second order approach which includes nearest neighbor interactions has been applied to the condensed phase. A total of 1311 comparisons are made between experimentally determined values and those calculated using additive group values. Of the 559 comparisons given for the enthalpy of formation ($\Delta_f H^\circ$) in the gas, liquid, and solid phases, the average difference (residual), without regard to sign, is 2.6 kJ/mol. The average differences for 390 comparisons for the heat capacity (C_p°) and 352 comparisons for the entropy (S°) in the three phases are 1.9 and 2.3 J/mol·K, respectively. The good agreement between experimental and calculated values shows that the Benson group additivity approach to the estimation of thermodynamic properties of organic compounds is applicable to the liquid and solid phases as well as the gas phase. Appendices provide example calculations of the thermodynamic properties of selected hydrocarbon compounds, total symmetry numbers, and methyl repulsion corrections. Most of the 144 references listed offer an indication of the activity in the development of estimation methods for calculating thermodynamic properties since 1931.

Key words: condensed phase; enthalpy of formation; entropy; estimation methods; gas phase; heat capacity; hydrocarbons; thermodynamic properties.

Contents

1. Introduction.....	1638	Appendix C. Methyl Repulsion Corrections for Branched Hydrocarbons	1651
2. Discussion of Results.....	1639	Appendix D. Example Calculations of Total Symmetry Number for Some Hydrocarbon Compounds	1655
2.1. <i>n</i> -Alkanes.....	1639		
2.2. Substituted Alkanes	1640		
2.3. Linear Alkenes	1641		
2.4. Substituted Alkenes	1641		
2.5. Alkynes.....	1641		
2.6. Aromatic Hydrocarbons	1641		
2.7. Cycloalkanes and Related Hydrocarbons	1642	Table 1. Additive group contribution values for hydrocarbons at 298.15 K.....	1658
3. Symmetry Numbers and Entropy	1642	Table 2. Ring strain corrections to be applied to cycloalkanes and related hydrocarbons.....	1660
4. Summary and Conclusions	1642	Table 3. <i>n</i> -Alkane hydrocarbons in the gas phase at 298.15 K	1661
5. Acknowledgments	1643	Table 4. <i>n</i> -Alkane hydrocarbons in the liquid phase at 298.15 K	1661
6. References	1643	Table 5. <i>n</i> -Alkane hydrocarbons in the solid phase at 298.15 K	1662
Appendix A. Example Calculations of Enthalpy of Formation, Heat capacity, and Entropy at 298.15 K for Some Hydrocarbon Compounds	1648	Table 6. Substituted alkane hydrocarbons (tertiary carbon) in the gas phase at 298.15 K	1662
Appendix B. Comparison of Calculated Thermodynamic Properties in the Gas Phase with Benson's Values at 298.15 K	1650	Table 7. Substituted alkane hydrocarbons (tertiary carbon) in the liquid phase at 298.15 K	1663
		Table 8. Substituted alkane hydrocarbons (tertiary carbon) in the solid phase at 298.15 K	1664

¹Present address: 2247 Regina Drive, Clarksburg, MD 20871.

©1988 by the U. S. Secretary of Commerce on behalf of the United States. This copyright is assigned to the American Institute of Physics and the American Chemical Society.

Reprints available from ACS; see Reprints List at back of issue.

Table 9.	Substituted alkane hydrocarbons (quaternary carbon) in the gas phase at 298.15 K	1664
Table 10.	Substituted alkane hydrocarbons (quaternary carbon) in the liquid phase at 298.15 K	1664
Table 11.	Substituted alkane hydrocarbons (quaternary carbon) in the solid phase at 298.15 K	1665
Table 12.	Linear alkene hydrocarbons in the gas phase at 298.15 K	1665
Table 13.	Linear alkene hydrocarbons in the liquid phase at 298.15 K	1666
Table 14.	Linear alkene hydrocarbons in the solid phase at 298.15 K	1667
Table 15.	Substituted alkene hydrocarbons in the gas phase at 298.15 K	1668
Table 16.	Substituted alkene hydrocarbons in the liquid phase at 298.15 K	1669
Table 17.	Substituted alkene hydrocarbons in the solid phase at 298.15 K	1669
Table 18.	Alkyne hydrocarbons in the gas phase at 298.15 K	1670
Table 19.	Alkyne hydrocarbons in the liquid phase at 298.15 K	1670
Table 20.	Alkyne hydrocarbons in the solid phase at 298.15 K	1670
Table 21.	Aromatic hydrocarbons in the gas phase at 298.15 K	1670
Table 22.	Aromatic hydrocarbons in the liquid phase at 298.15 K	1672
Table 23.	Aromatic hydrocarbons in the solid phase at 298.15 K	1674
Table 24.	Cycloalkane and related hydrocarbons in the gas phase at 298.15 K	1675
Table 25.	Cycloalkane and related hydrocarbons in the liquid phase at 298.15 K	1676
Table 26.	Cycloalkane and related hydrocarbons in the solid phase at 298.15 K	1678

1. Introduction

For many years, thermodynamicists have been active in the development of numerical methods which correlate molecular structure and sub-structure with a corresponding energy contribution to a thermodynamic property. Some indication of this activity can be found by examining the 144 references at the end of this paper which primarily cover estimation methods and also identify some important reference sources of evaluated thermodynamic properties of hydrocarbon and related compounds. A variety of estimation schemes have emerged which demonstrate that good agreement can be achieved between calculated and experimentally measured thermodynamic properties. Of the estimation schemes which have been developed, the one put together by S.W. Benson and coworkers (58BEN/BUS, 68BEN, 69BEN/CRU, 69SHA, 71SHA, 73EIG/GOL, 76BEN, 77LUR/BEN, 77SHA/GOL, and 77STE/GOL) has the most desirable characteristics and appears

to be generally acceptable to scientists within and between the disciplines of physical chemistry and chemical engineering. The attractive features of the Benson approach consist of simple additivity, clarity of notation, second order character, i.e., inclusion of nearest-neighbor interactions, ease of application, and satisfactory agreement between the estimated thermodynamic property and its experimentally determined value. Initially, focus was directed toward estimation of thermodynamic properties of organic compounds in the gas phase. The development of numerical methods oriented toward estimation of thermodynamic properties in the condensed phase has also been reported (69SHA, 71SHA, 77LUR/BEN), but primary attention has usually been placed upon gas phase processes. The estimation procedure developed by Benson and coworkers has been adopted into CHETAH, the ASTM Chemical Thermodynamic and Energy Release Evaluation Program (74SEA/FRE), for the estimation of thermodynamic properties of organic compounds in the gas phase and for the classification of chemical compounds or compositions according to whether they are likely to be impact sensitive.

The purpose of this paper is to extend the Benson group contribution values to the liquid and solid phases for hydrocarbon compounds, and when possible, to provide improved group contribution values for the gas phase. The thermodynamic properties at 298.15 K considered are: enthalpy of formation, heat capacity, and entropy. The approach taken in the evaluation of data and in the development of group values consisted of calculating and examining group increments in a systematic manner. This procedure was followed by the selection of group values which appeared to give minimum residuals between the calculated and experimental values. The development of groups and group values followed the path from alkanes to alkenes, alkynes, aromatic hydrocarbons, cycloalkanes, and other hydrocarbon derivatives. A global least squares, least sums, or regression-type fit of all the group values was not carried out because of differences in the quality of the data and because of the minimal amount of data available for certain group values. Computations were performed using a conventional desk-top electronic calculator. It is expected that the results we obtained are approximately equivalent to what would have been derived from a weighted regression analysis.

A companion document, 89DOM/HEA, is in preparation which describes a database and algorithm for calculating the enthalpy of formation, heat capacity, and entropy at 298.15 K of 1311 hydrocarbon compounds in the gas, liquid, and solid phases. The database includes the group contribution values used by the algorithm, the corresponding experimental and calculated data for the enthalpy of formation, heat capacity, and entropy of the hydrocarbon compounds, and the differences or residuals which result. The document, 89DOM/HEA, is also available on diskette and contains an interactive program which allows the user to carry out the calculation of the enthalpy of formation, heat capacity, or entropy for any hydrocarbon compound based on the assembly of the

corresponding group contribution values in a particular phase. Also calculated are the entropy of formation, $\Delta_f S^\circ$, Gibbs energy of formation, $\Delta_f G^\circ$, and the natural logarithm of the equilibrium constant, $\ln K_f$.

The thermodynamic data used as the data-base for this study were obtained from the following reference sources: 53ROS/PIT, 69STU/WES, 70COX/PIL, 77PED/RYL, 84DOM/EVA, 86MAR, and 86PED/NAY. Differences between the enthalpy of formation, heat capacity, and entropy for the gas and liquid phases, and liquid and solid phases are related to the corresponding transition properties. For example, the difference between the enthalpies of formation at 298.15 K for the gas and liquids phases equals the enthalpy of vaporization at 298.15 K for that hydrocarbon. Similarly, the difference between the enthalpy of formation liquid and solid phases gives the transition property equal to the sum of the enthalpies of fusion and the enthalpies of the first-order crystal-crystal transitions at 298.15 K. In the tables containing data on hydrocarbon compounds in the solid phase, the experimental values for $\Delta_f H^\circ$ were sometimes derived from a combination of $\Delta_f H^\circ$ (liquid), ΔH (fusion), and any ΔH (transition).

2. Discussion of Results

Group contribution values for hydrocarbons at 298.15 K are given for the enthalpy of formation, $\Delta_f H^\circ$, heat capacity, C_p° , and entropy, S° , for the gas, liquid, and solid phases in Table 1 in joules. The superscript symbol, $^\circ$, indicates that the substance is in its standard state. For a gaseous compound, the standard thermodynamic properties are for the pure compound at a pressure of 0.1 MPa (1 bar) and in a hypothetical state in which it exhibits ideal gas behavior. The heat capacity and enthalpy of the compound are those of the real gas extrapolated to zero pressure. For a liquid or solid compound, the standard thermodynamic properties are for the pure compound in the liquid or solid phase, respectively, under a pressure of 0.1 MPa (1 bar).

Ring strain corrections which are applicable to cycloalkanes and related compounds are shown in Table 2 and are also in joules. The joule is related to the thermochemical calorie through the definition: one thermochemical calories equals 4.1840 joules.

Recommendations by the IUPAC (82COX/ANG) to reference the standard state pressure as 100,000 Pa (1 bar) rather than 101,325 Pa (1 atmosphere) were examined. Calculations indicated that numerical changes to group contribution values were negligible when compared to the uncertainties the values themselves. Systematic differences imposed by a change in standard state pressure to 100,000 Pa were well below 0.4 kJ/mol (0.1 kcal/mol) for $\Delta_f H^\circ$ and 0.4 J/mol·K (0.1 cal/mol·K) for C_p° and S° in most instances. Hence, no corrections to accommodate this change were deemed necessary.

Example calculations which give details of the additivity of the group values in the estimation of $\Delta_f H^\circ$, C_p° , and S° are provided for several hydrocarbon compounds in Appendix A. Appendix B shows a comparison of gas

phase property data calculated from this study with that of 76BEN. Overall the residuals found between the two calculated values are small. Comparison to experimental values in other tables indicates that improvement takes place about half of the time.

2.1. *n*-Alkanes

Tables 3, 4, and 5 show the agreement between 122 experimental and calculated values for $\Delta_f H^\circ$, C_p° , and S° at 298.15 K for *n*-alkanes. In the gas phase, 50 comparisons were made between the experimental and calculated values for $\Delta_f H^\circ$, C_p° , and S° at 298.15 K; here, residuals do not exceed 1.5 kJ/mol for $\Delta_f H^\circ$ or 1.5 J/mol·K for C_p° or S° . In the liquid phase, 39 comparisons showed residuals for $\Delta_f H^\circ$ of 1.6 kJ/mol or less and residuals for S° of 0.4 J/mol·K or less. Residuals for liquid C_p° result in only 3 of 14 exceeding 2.0 J/mol·K. In the solid phase, 17 comparisons between experimental and calculated values for $\Delta_f H^\circ$ show 8 residuals below 2.0 kJ/mol. The development of group values for C_p° and S° in the solid phase for *n*-alkanes presented considerable difficulty. Data for C_p° and S° in the solid phase were not plentiful and some of the results reported are of dubious quality. For example, some (10) residuals for C_p° and S° in Table 5 appear to be unusually large, while others are quite satisfactory. In particular, the data on C_p° and S° for crystalline *n*-octadecane reported by 67MES/GUT and for crystalline *n*-hexacosane reported by 76AND/MAR were of high quality as were the extrapolated values for crystalline *n*-hexadecane derived from the liquid phase data of 67MES/GUT. However, data on C_p° and S° for solid phase *n*-alkanes reported before 1950 were considered unreliable because low temperature measurements did not extend below the boiling point of liquid nitrogen (~80 K). As a result of using the latter data, large residuals, shown in Table 5, were obtained for C_p° and S° for solid *n*-alkanes. In some of the reported studies, sample impurities may have introduced errors into the experimental data for C_p° and S° . Additional experimental studies on solid *n*-alkanes in the C_{20} to C_{40} range are needed to improve the reliability of estimation of thermodynamic properties using group contribution values.

Although separate values for odd and even carbon containing homologues were developed for the calculation of values for the group $C-(C)_n(H)$, for $\Delta_f H$ in the solid phase for *n*-alkanes, this did not result in an improvement in the estimated values when compared to experimental data. Therefore, the separate values were discarded in preference for the single value listed in Table 1, $C-(C)_n(H)$. -29.41 kJ/mol Effects from odd and even numbers of carbon atoms in the data for C and S° in *n*-alkanes were even less apparent and, hence, only a single value appears in Table 1 for the $C-(C)_n(H)$ group. It should also be mentioned that the value for the methyl group does not change, except for the physical phase, no matter to what structural entity it is attached, i.e., $C-(H)-(C)$ — $C-(H)-(X)$, where X is another carbon atom neighbor, such as C_a , C_b , or C_c .

2.2. Substituted Alkanes

Tables 6 through 11 compare the experimental and calculated values for substituted or branched alkanes for $\Delta_f H^\circ$, C_p° , and S° at 298.15 K. During the evaluation of data on the enthalpies of formation of substituted alkanes, the Benson approach was modified for the treatment of gauche isomers. A new treatment was developed which corrects for the repulsive interaction of hydrogen atoms on methyl groups attached to tertiary or quaternary carbon atoms. A group value was assigned to a tertiary or quaternary carbon along with a group value for the number of methyl groups attached to the tertiary or quaternary carbon. For example, $\Delta_f H^\circ$ for 2-methylpropane required 3 methyl repulsion correction values, 2-methylbutane required 2 methyl repulsion correction values, 3-methyl-pentane required 1 methyl repulsion correction value, and 3-ethyl-pentane needed no correction of this kind. $\Delta_f H^\circ$ for the gas phase for these molecules is shown in the sample calculations below.


A similar rationale was applied to the methyl repulsion correction for a quaternary carbon atom. Molecules with quaternary carbon atoms which corresponded to the examples above were 2,2-dimethylpropane, 2,2-dimethylbutane, 3,3-dimethylpentane, 3-ethyl-3-methylpentane, and 3,3-diethylpentane and required 4, 3, 2, 1, and 0 methyl repulsion correction values, respectively, in the calculation of $\Delta_f H^\circ$ at 298.15 K. In addition, different methyl repulsion correction values were required for quaternary carbon atoms which had a tertiary carbon atom adjacent to it and another such correction when an alkane had two quaternary carbon atoms in its structure. These additional methyl repulsion corrections have been developed for $\Delta_f H^\circ$ in the gas and liquid phases, but not for the solid phase due to a lack of precise data on solid hydrocarbon compounds having the appropriate structures. The introduction of the methyl repulsion correction values was made because of the significant reduction observed in the differences in the experimental and calculated values for hydrocarbon compounds having tertiary and/or quaternary carbon atoms and because it appears simpler in both concept and application to the gauche correction developed earlier. Some further examples of the methyl repulsion corrections are provided in Appendix C.

Although the notion of methyl repulsion interactions in the arrays of data on C_p° and S° for substituted alkanes was examined carefully, correction terms did not emerge which brought about a reduction in the magnitude of the residuals for these hydrocarbons.

Among the substituted alkanes shown in Tables 6 through 11 for all three phases, 215 comparisons were made between experimental and calculated values. In the gas phase, out of a total of 117 comparisons, 101 show residuals of less than or equal to 4.0 kJ/mol for $\Delta_f H^\circ$ and less than or equal to 4.0 J/mol-K for C_p° and S° . In the liquid phase, 77 comparisons are listed. Of these, 42 show residuals of less than or equal to 4.0 kJ/mol for $\Delta_f H^\circ$ and less than or equal to 4.0 J/mol-K for C_p° and S° . In the solid phase, out of 19 comparisons for $\Delta_f H^\circ$, 17

are less than or equal to 4.0 kJ/mol. Comparisons of C_p° and S° for solid substituted alkanes are virtually nonexistent. Experimental data are needed on key substituted alkanes to assist in the estimation of other family members.

2-Methylpropane




Enthalpy of formation (gas phase)

No. of groups	Group	Group value	Sum (kJ/mol)
3	C-(C)(H) ₃	-42.26	-126.78
1	C-(C) ₂ (H)	-1.17	-1.17
3	-CH ₃ corr	-2.26	-6.78
-134.73			

Compare with experimental value: -134.18 kJ/mol (51PRO/MAR, 86PED/NAY).

Compare with estimated value from 76 BEN: -135.98 kJ/mol.

2-Methylbutane




Enthalpy of formation (gas phase)

No. of groups	Group	Group value	Sum (kJ/mol)
3	C-(C)(H) ₃	-42.26	-126.78
1	C-(C) ₂ (H) ₂	-20.63	-20.63
1	C-(C) ₃ (H)	-1.17	-1.17
2	-CH ₃ corr	-2.26	-4.52
-153.10			

Compare with experimental value: -153.68 kJ/mol (70GOO, 86PED/NAY).

Compare with estimated value from 76BEN: -153.26 kJ/mol.

3-Methylpentane




Enthalpy of formation (gas phase)

No. of groups	Group	Group value	Sum (kJ/mol)
3	C-(C)(H) ₃	-42.26	-126.78
2	C-(C) ₂ (H) ₂	-20.63	-41.26
1	C-(C) ₃ (H)	-1.17	-1.17
1	-CH ₃ corr	-2.26	-2.26
-171.47			

Compare with experimental value: -172.09 kJ/mol (41PRO/ROS, 86PED/NAY).

Compare with estimated value from 76BEN: -170.54 kJ/mol.

3-Ethylpentane

Enthalpy of formation (gas phase)

No. of groups	Group	Group value	Sum (kJ/mol)
3	C-(C)(H) ₃	-42.26	-126.78
3	C-(C) ₂ (H) ₂	-20.63	-61.89
1	C-(C) ₃ (H)	-1.17	-1.17
			-189.84

Compare with experimental value: -189.62 kJ/mol (41PRO/ROS 2, 86PED/NAY).

Compare with estimated value from 76BEN: -187.82 kJ/mol.

2.3. *n*-Alkenes

Tables 12, 13, and 14 compare 154 experimental and calculated values for $\Delta_f H^\circ$, C_p° , and S° at 298.15 K for linear alkenes in the gas, liquid and solid phases. In the gas phase, 73 out of 83 comparisons show residuals of 2.0 kJ/mol or less for $\Delta_f H^\circ$ and 2.0 J/mol·K or less for C_p° and S° . In the liquid phase, 40 out of 54 comparisons show residuals of 2.0 kJ/mol or less for $\Delta_f H^\circ$ and 2.0 J/mol·K or less for C_p° and S° . In the solid phase, experimental data were unavailable for C_p° and S° on linear alkenes, suggesting the need for additional research. Data for $\Delta_f H^\circ$ on solid linear alkene are derived from data on the liquid phase and data on enthalpies of fusion and transition. Here, 15 out of 17 comparisons show residuals of less than 2.0 kJ/mol. The residual for $\Delta_f H^\circ$ of 1-hexadecene in the solid phase listed in Table 14 is large, 17.2 kJ/mol. Because the $\Delta_f H^\circ$ residuals listed for most of the other linear alkenes are below 2.0 kJ/mol, it appears that the group contribution values are satisfactory. Low temperature heat capacity studies were performed on seven 1-olefins by 57MCC/FIN. The presence of a first order crystal-crystal transition in the temperature range between 200 K and the triple point of 1-decene and linear alkenes with higher carbon numbers was present, but not in 1-hexadecene, however, a lambda transition was observed. Also significant is the presence of a higher than normal impurity (1.3 mol%) in the 1-hexadecene sample studied by 57MCC/FIN. It is possible that the impurity in the 1-hexadecene has obscured a crystal-crystal transition near the lambda transition which is about the magnitude of the residual we find in Table 14, 17.2 kJ/mol. A careful phase transition study on a good quality sample would answer the question.

Attempts to extrapolate the low temperature data on C_p° and S° reported by 57MCC/FIN for 1-alkenes in the solid phase through the melting point to 298.15 K produced group values with large uncertainties and poor reliability, hence, such group values are not listed.

2.4. Substituted Alkenes

Tables 15, 16, and 17 show the agreement between 118 experimental and calculated values for substituted alkenes in the gas, liquid, and solid phases, respectively, for $\Delta_f H^\circ$, C_p° , and S° at 298.15 K. In the gas phase, 62 comparisons were made. Of these, 43 agreed within 4.0 kJ/mol or less for $\Delta_f H^\circ$ and within 4.0 J/mol·K or less for C_p° and S° . In the liquid phase, 47 comparisons were made of which 16 residuals were 4.0 kJ/mol or less for $\Delta_f H^\circ$ and 12 residuals were 4.0 J/mol·K or less for C_p° and S° . As with the data on solid linear alkenes, data are limited on solid substituted alkenes. Of the available data on $\Delta_f H^\circ$, 5 out of 9 residuals were 4.0 kJ/mol or less.

We have examined the available data on the enthalpies of formation at 298.15 K for branched alkenes to determine whether a methyl repulsion correction could be derived which would be the counterpart of gauche alkene correction discussed in 69BEN/CRU. Our examination showed that a similar correction was needed for alkene groups which involved tertiary and quaternary carbon atoms. A cis correction for alkenes was also developed in the gas, liquid, and solid phases for $\Delta_f H^\circ$, but for C_p° and S° in the gas phase only.

2.5. Alkynes

Tables 18, 19, and 20 give comparisons between 61 experimental and calculated values for alkynes in the gas, liquid, and solid phases, respectively, for $\Delta_f H^\circ$, C_p° , and S° at 298.15 K. In the gas phase, 41 comparisons were made of which 36 were 2.0 kJ/mol or less for $\Delta_f H^\circ$ and 2.0 J/mol·K or less for C_p° and S° . In the liquid phase, 18 comparisons were made of which 11 were 2.0 kJ/mol or less for $\Delta_f H^\circ$ and 2.0 J/mol·K or less for C_p° and S° . The solid phase is limited to 2 values of $\Delta_f H^\circ$.

The experimental values for C_p° and S° for the alkynes in the gas phase were those calculated by 45WAG/KIL, using statistical thermodynamic procedures, and the methods established by 46TAY/WAG for calculating the regularities in gaseous thermodynamic functions for a homologous series. Those values were tabulated by 53ROS/PIT and 69STU/WES. The good agreement between the experimental and calculated values was expected because of the group additive features in both sets of values. Data for C_p° and S° for alkynes in the liquid phase were limited and those for the solid phase were absent.

2.6. Aromatic Hydrocarbons

Tables 21, 22, and 23 show the agreement between experimental and calculated values for aromatic hydrocarbons in the gas, liquid, and solid phases, respectively, for $\Delta_f H^\circ$, C_p° , and S° at 298.15 K. A total of 368 comparisons were made. In the gas phase, 145 out of 168 comparisons had residuals of 4.0 kJ/mol or less for $\Delta_f H^\circ$, or 4.0 J/mol·K or less for C_p° and S° . In the liquid phase, 80 out of 103 comparisons had residuals of 4.0 kJ/mol or less for $\Delta_f H^\circ$, or 4.0 J/mol·K or less for C_p° and S° . In the

solid phase, 57 out of 97 comparisons had residuals of 4.0 kJ/mol or less for $\Delta_f H^\circ$, or 4.0 J/mol·K or less for C_p° and S° .

The experimental values for C_p° and S° for benzene in the solid phase at 298.15 K which appeared in Table 23, were obtained by linear extrapolation of the data on the solid phase from the melting point to room temperature. Note that separate corrections were applicable for ortho substitution in the gas phase for C_p° and S° for naphthalene.

The experimental enthalpies of formation, heat capacities, and entropies of 1-methylnaphthalene and 2-methylnaphthalene in the gas, liquid, and solid phases have been based upon the data of 56MIL/BEC, 57MCC/FIN, 60SPE/ROS, and 74SAB/CHA. Although actual differences of 2.97, -6.39, and -0.58 kJ/mol can be calculated between the experimental enthalpies of formation of 1-methylnaphthalene and 2-methylnaphthalene for the gas, liquid, and solid phases, respectively, no correction to the calculated values has been applied because the experimental uncertainties in $\Delta_f H^\circ$ for an isomer can be difficult to evaluate and the corresponding uncertainties for the calculated differences in $\Delta_f H^\circ$ between isomers can become large.

2.7. Cycloalkanes and Related Hydrocarbons

The corrections listed in Table 2 are applicable to cycloalkanes and related compounds reflect the magnitude of the internal ring strain imposed for a given size ring. Note that somewhat different ring corrections were applied to substituted versus unsubstituted values for cyclopentane, cyclopentene, and cyclohexane.

Tables 24, 25, and 26 give comparisons between the experimental and calculated values for cycloalkanes, substituted cycloalkanes, and related compounds in the gas, liquid, and solid phases, respectively, for $\Delta_f H^\circ$, C_p° , and S° at 298.15 K. A total of 273 comparisons were made. Among the three phases, unsubstituted cycloalkanes showed equivalence between the experimental and calculated values, i.e., zero residual, since the ring corrections are derived from the differences which are found between the experimental and "strain-free" values. A total of 129 comparisons were made which gave zero residuals. For substituted cycloalkanes, a total 144 comparisons were made. In the gas phase, 44 out of 66 comparisons had residuals of 4.0 kJ/mol or less for $\Delta_f H^\circ$ and 4.0 J/mol·K or less for C_p° and S° . In the liquid phase, 50 out of 68 comparisons showed residuals of 4.0 kJ/mol or less for $\Delta_f H^\circ$, or 4.0 J/mol·K or less for C_p° and S° .

3. Symmetry Numbers and Entropy

In the calculation of the entropy of a gaseous molecule, it is important that proper symmetry contributions imposed by the presence of certain structural features are accounted for. Molecular symmetry is not a factor in the additive summation of group values in the

calculation of the enthalpy of formation or heat capacity, but it does influence the calculation of the entropy of a compound in the gas phase in a manner which is not simply additive. In order to implement the contribution to the entropy of a compound because of symmetry features, the actual or real entropy, S° , is divided into two parts:

- (1) intrinsic entropy, $S^\circ(\text{group})$, which is the entropy contribution determined by adding the group values for the entropy of the structural parts of the compound; and,
- (2) symmetry-related entropy, $S^\circ(\text{sym})$, which is the entropy contribution determined from the symmetry features of the compound as a whole.

Hence, $S^\circ = S^\circ(\text{group}) + S^\circ(\text{sym})$. The symmetry-related entropy is in turn divided into two parts:

- (1) entropy due to optical isomerism in the molecule, $S^\circ(\text{opt})$; and,
- (2) entropy from the total symmetry number of the molecule, $S^\circ(\text{tot})$, which is a reflection of how symmetrical the molecule is.

It therefore follows that: $S^\circ(\text{sym}) = S^\circ(\text{opt}) + S^\circ(\text{tot})$, or $S^\circ(\text{sym}) = R \ln N(\text{opt}) + R \ln N(\text{tot})$, where R is the gas constant, $N(\text{opt})$ is the number of optical isomers, $N(\text{tot})$ is the total symmetry number, and \ln is the natural logarithm. Note that $R \ln N(\text{opt})$ is a positive quantity and that $R \ln N(\text{tot})$ is a negative quantity in the summation of the actual entropy, S° .

One of the ways in which a molecule may exhibit optical isomerism occurs when its mirror image cannot be superimposed upon itself as a result of simple molecular rotations. Such mirror images are called enantiomers and are said to be chiral. The two enantiomers have identical physical properties except for their interaction with transmitted polarized light. Other elements of optical isomerism require consideration in the calculation of $R \ln N(\text{opt})$, and discussions can be found in textbooks on organic chemistry and stereoisomerism.

A discussion of how to calculate the total symmetry number as well as several sample calculations for selected hydrocarbon compounds is given in Appendix D. Any of the tables which show gas phase comparisons between experimental and calculated entropies, also contains a column entry on the extreme right giving the value for $N(\text{tot})$ for that hydrocarbon molecule.

4. Summary and Conclusions

A series of group contribution values has been developed which permits the estimation of the enthalpy of formation, heat capacity, and entropy at 298.15 K for hydrocarbon compounds in the gas, liquid, and solid-phases. The set of group values which has been pro-

vided in the gas phase for $\Delta_f H^\circ$, C_p° , and S° at 298.15 K are comparable in accuracy of prediction to the group values developed by 58BEN/BUS, 68BEN, 69BEN/CRU, and 76BEN. This comparison is shown in Appendix B for the gas phase. The present set of group values have the advantage of including experimental data on hydrocarbon compounds which were reported over the period 1975-1987. The evaluation of thermodynamic data on gaseous hydrocarbon compounds also provided group values which have internal consistency with the corresponding values for the liquid and solid phases.

A total of 1311 comparisons have been made between an experimentally determined value and one calculated using the additive group contribution values reported here for $\Delta_f H^\circ$, C_p° , and S° at 298.15 K. From the total, 559 comparisons are listed for $\Delta_f H^\circ$ among the three phases and have an average residual, calculated without regard to sign, of 2.6 kJ/mol. Similarly, the average residuals obtained for the 390 comparisons made for C_p° and the 352 comparisons made for S° are 1.9 and 2.3 J/mol·K, respectively. Here also, the average residuals were calculated without regard to sign, and the unusually large residuals found for C_p° and S° for five *n*-alkanes in the solid phase in Table 5 have been omitted. The average residuals calculated for $\Delta_f H^\circ$, C_p° , and S° are comparable to what are often reported for corresponding experimental uncertainties. These results demonstrate the general applicability of the Benson second order group additivity approach to the estimation of thermodynamic properties in the liquid and solid phases.

An approach has been developed which corrects for the repulsive interactions of hydrogen atoms on methyl groups in branched hydrocarbon compounds. Different methyl repulsion correction values are needed for branching to a tertiary carbon, a quaternary carbon, a tertiary or quaternary carbon atom adjacent to a quaternary carbon atom, and for two non-adjacent quaternary carbon atoms. These corrections are applicable not only to branched hydrocarbon groups which are present in saturated hydrocarbons, but to those present in unsaturated, alicyclic, or aromatic hydrocarbon compounds.

Plans are underway to develop the Benson approach of group value additivity to the estimation of $\Delta_f H^\circ$, C_p° , and S° at 298.15 K for organic compounds containing the elements CHO, CHN, CHNO, CHS, and CHSO in the gas, liquid, and solid phases.

5. Acknowledgments

The authors thank Mr. Barry L. Jacobs for his assistance in the early stages of development of this research project and Mrs. Wendy L. King for her assistance in implementing format and numerical changes to the manuscript. The authors are grateful to the NIST Office of Standard Reference Data for the project's financial support and to Mrs. Connie I Seymour and Miss Mary E. Trapane of that office for manuscript preparation for publication.

6. References

- 31SMI/AND Smith, R. H., and Andrews, D. H., "Thermal Energy Studies. I. Phenyl Derivatives of Methane, Ethane, and Some Related Compounds", *J. Am. Chem. Soc.* **53**, 3644-3660 (1931).
- 32PAR/HUF Parks, G. S., and Huffman, H. M., "The Free Energies of Some Organic Compounds", *Am. Chem. Soc. Monograph Ser. (The Chem. Catalog Book Co., Inc.)*, New York, 1932.
- 40AST Aston, J. G., "Some Observations on the Thermodynamics of Hydrocarbons and Related Compounds", *Chem. Rev.* **27**, 59-73 (1940)
- 40AST/MES Aston, J. G., and Messerly, G. H., "The Heat Capacity, Entropy, Heats of Fusion and Vaporization and the Vapor Pressure of *n*-Butane", *J. Am. Chem. Soc.* **62**, 1917-1933 (1940).
- 40PAR Parks, G. S., "Some Free-Energy Data for Typical Hydrocarbons Containing Six or More Carbon Atoms", *Chem. Rev.* **27**, 75-83 (1940).
- 40PIT Pitzer, K. S., "Chemical Equilibria, Free Energies, and Heat Contents for Gaseous Hydrocarbons", *Chem. Rev.* **27**, 39-57 (1940).
- 40PIT 2 Pitzer, K. S., "The Vibration Frequencies and Thermodynamic Functions of Long Chain Hydrocarbons", *J. Phys. Chem.* **8**, 711-720 (1940).
- 40ROS Rossini, F. D., "Heats of Formation of Gaseous Hydrocarbons", *Chem. Rev.* **27**, 1-16 (1940).
- 40WIL Wilson, E. B., Jr., "The Present Status of the Statistical Method of Calculating Thermodynamic Functions", *Chem. Rev.* **27**, 17-38 (1940).
- 41DOB Dobratz, C. J., "Heat Capacities of Organic Vapors", *Ind. Eng. Chem.* **33**, 759-762 (1941).
- 41PRO/ROS Prosen, E. J., and Rossini, F. D., "Heats of Isomerization of the Five Hexanes", *J. Research NBS* **27**, 289-310 (1941).
- 41PRO/ROS 2 Prosen, E. J., and Rossini, F. D., "Heats of Isomerization of the Nine Heptanes", *J. Research NBS* **27**, 519-528 (1941).
- 43STU/MAY Stull, D. R., and Mayfield, F. D., "Heat Capacities of Hydrocarbon Gases", *Ind. Eng. Chem.* **39**, 517-550 (1943).
- 44AND/BEY Andersen, J. W., Beyer, G. H., and Watson, K. M., "Thermodynamic Properties of Organic Compounds. Estimation from Group Contributions", *Nat. Petrol. News, Tech. Sect.* **36**, R476-R484 (July 5, 1944).
- 44GUT/SPI Guthrie, G. B., Jr., Spitzer, R. W., and Huffman, H. M., "Thermal Data. XVIII. The Heat Capacity, Heat of Fusion, Entropy, and Free Energy of Ethylbenzene", *J. Am. Chem. Soc.* **66**, 2120-2121 (1944).
- 44PRO/ROS Prosen, E. J., and Rossini, F. D., "Heats of Combustion of Eight Normal Paraffin Hydrocarbons in the Liquid State", *J. Research NBS* **33**, 255-272 (1944).
- 45GUT/PIT Guttman, L., and Pitzer, K. S., "trans-2-Butene. The Heat Capacity, Heats of Fusion and Vaporization, and Vapor Pressure. The Entropy and Barrier to Internal Rotation", *J. Am. Chem. Soc.* **67**, 324-327 (1945).
- 45JOH/PRO Johnson, W. H., Prosen, E. J., and Rowan, E. D., "Heats of Combustion and Isomerization of the Eight C_8H_{10} Alkylbenzenes", *J. Research NBS* **35**, 141-146 (1945).
- 45PRO/ROS Prosen, E. J., and Rowan, E. D., "Heats of Isomerization of the 18 Octanes", *J. Research NBS* **34**, 163-174 (1945).
- 45PRO/ROS ? Prosen, E. J., and Rowan, E. D., "Heats of Combustion of the Paraffin Hydrocarbons at 25°C.", *J. Research NBS* **34**, 263-269 (1945).

- 45WAG/KIL Wagman, D. D., Kilpatrick, J. E., Pitzer, K. S., and Rossini, F. D., "Heats, Equilibrium Constants, and Free Energies of Formation of the Acetylenic Hydrocarbons through the Pentynes, to 1,500 K", *J. Research NBS* **35**, 467-496 (1945).
- 46AST/FIN Aston, J. G., Fink, H. L., Bestul, A. B., Pace, E. L., and Szasz, G. J., "The Heat Capacity and Entropy, Heats of Fusion and Vaporization and Vapor Pressure of 1-Butene", *J. Am. Chem. Soc.* **68**, 52-62 (1946).
- 46TAY/WAG Taylor, W. J., Wagman, D. D., Williams, M. G., Pitzer, K. S., and Rossini, F. D., "Heats, Equilibrium Constants, and Free Energies of Formation of the Alkylbenzenes", *J. Research NBS* **37**, 95-122 (1946).
- 47BRE/THO Bremner, J. G. M., and Thomas, G. D., "The Calculation of Resonance Energies from Thermal Data", *Trans. Faraday Soc.* **43**, 338-341 (1947).
- 47BRE/THO 2 Bremner, J. G. M., and Thomas, G. D., "The Extension of Thermodynamic Values from the Aliphatic to the Aromatic Series", *Trans. Faraday Soc.* **43**, 779-785 (1947).
- 47JOH/PRO Johnson, W. H., Prosen, E. J., and Rossini, F. D., "Heats of Combustion of Six Nonanes", *J. Research NBS* **38**, 419-422 (1947).
- 48BRE/THO Bremner, J. G. M., and Thomas, G. D., "The Standard Free Energies of Formation of Gaseous Organic Compounds: A Group Method of Summation", *Trans. Faraday Soc.* **44**, 230-238 (1948).
- 49FRA Franklin, J. L., "Prediction of Heat and Free Energies of Organic Compounds", *Ind. Eng. Chem.* **41**, 1070-1076 (1949).
- 49KLA Klages, F., "Improvement in the Additive Calculation of Heats of Combustion and the Mesomeric Energy from Heats of Combustion", *Ber.* **82**, 358-375 (1949).
- 49SOU/MAT Souders, M., Jr., Matthews, C. S., and Hurd, C. O., "Relationship of Thermodynamic Properties to Molecular Structure. Heat Capacities and Heat Contents of Hydrocarbon Vapors", *Ind. Eng. Chem.* **41**, 1037-1048 (1949).
- 49SOU/MAT 2 Souders, M., Jr., Matthews, C. S., and Hurd, C. O., "Entropy and Heat of Formation of Hydrocarbon Vapors", *Ind. Eng. Chem.* **41**, 1048-1056 (1949).
- 51PRO/MAR Prosen, E. J., Maron, F. W., and Rossini, F. D., "Heats of Combustion and Isomerization of Ten C_4 Hydrocarbons", *J. Research NBS* **46**, 106-112 (1951).
- 51VAN/CHE Van Krevelen, D. W., and Chermin, H. A. G., "Estimation of the Free Enthalpy (Gibbs Free Energy) of Formation of Organic Compounds from Group Contributions", *Chem. Eng. Sci.* **1**, 66-80 (1951).
- 52SCO/DOU Scott, D. W., Douslin, D. R., Gross, M. E., Oliver, G. D., and Huffman, H. F., "2,2,3,3-Tetramethylbutane: Heat Capacity, Heats of Transition, Fusion, and Sublimation, Vapor Pressure, Entropy and Thermodynamic Functions", *J. Am. Chem. Soc.* **74**, 883-887 (1952).
- 53RAT/GWI Rathjens, G. W., Jr., and Gwinn, W. D., "Heat Capacities and Entropy of Cyclobutane", *J. Am. Chem. Soc.* **75**, 5629-5633 (1953).
- 53ROS/PIT Rossini, F. D., Pitzer, K. S., Arnett, R. L., Braun, R. M., and Pimentel, G. C., "Selected Values of Physical and Thermodynamic Properties of Hydrocarbons and Related Compounds" (Carnegie Press, Pittsburgh, Pa. 1953).
- 54FIN/GRO Finke, H. L., Gross, M. E., Waddington, G., and Huffman, H. M., "Low Temperature Thermal Data for the Nine Normal Paraffin Hydrocarbons from Octane to Hexadecane", *J. Am. Chem. Soc.* **76**, 333-341 (1954).
- 55STAY/KIL Taylor, R. D., and Kilpatrick, J. E., "Entropy, Heat Capacity, and Heats of Transition of 1,3,5-Trimethylbenzene", *J. Chem. Phys.* **23**, 1232-1235 (1955).
- 56HAN Handrick, G. R., "Heats of Combustion of Organic Compounds", *Ind. Eng. Chem.* **48**, 1366-1374 (1956).
- 56LAI Laidler, K. J., "A System of Molecular Thermochemistry for Organic Gases and Liquids", *Can. J. Chem.* **34**, 626-648 (1956).
- 56MIL/BEC Milligan, D. E., Becker, E. D., and Pitzer, K. S., "Thermodynamic Functions of Alkylnaphthalenes from 298 to 1500 K", *J. Am. Chem. Soc.* **78**, 2707-2711 (1956).
- 57GAM Gambill, W. R., "You Can Predict Heat Capacities", *Chem. Eng.* **64**, May, 263-268 (1957).
- 57GAM 2 Gambill, W. R., "You Can Predict Heat Capacities", *Chem. Eng.* **64**, June, 243-248 (1957).
- 57GAM 3 Gambill, W. R., "Predict Liquid Heat Capacities", *Chem. Eng.* **64**, July, 263-268 (1957).
- 57GAM 4 Gambill, W. R., "Equations Give Liquid Heat Capacities", *Chem. Eng.* **64**, August, 257-258 (1957).
- 57GAM 5 Gambill, W. R., "Generalized Gas Heat Capacities", *Chem. Eng.* **64**, October, 283-288 (1957).
- 57JOH/HUA Johnson, A. I., and Huang, C. J., "Estimation of Ideal-Gas Heat Capacities of Hydrocarbons", *Can. J. Technol.* **34**, 405-409 (1957).
- 57MCC/FIN McCullough, J. P., Fink, H. L., Gross, M. E., Messerly, J. F., and Waddington, G., "Low Temperature Calorimetric Studies of Seven 1-Olefins: Effect of Orientational Disorder in the Solid State", *J. Phys. Chem.* **61**, 289-301 (1957).
- 58BEN/BUS Benson, S. W., and Buss, J. H., "Additive Rules for the Estimation of Molecular Properties. Thermodynamic Properties", *J. Chem. Phys.* **29**, 546-572 (1958).
- 58GRE/ROS Greenshields, J. B., and Rossini, F. D., "Molecular Structure and Properties of Hydrocarbons and Related Compounds", *J. Phys. Chem.* **62**, 271-280 (1958).
- 59ALL Allen, T. L., "Bond Energies and the Interactions between Next-Nearest Neighbors. I. Saturated Hydrocarbons, Diamond, Sulfanes, S₈, and Organic Sulfur Compounds", *J. Chem. Phys.* **31**, 1039-1049 (1959).
- 59OTH/ZUD Othmer, D. F., and Zudkevitch, D., "Correlating Latent Heats and Entropies of Vaporization with Temperature", *Ind. Eng. Chem.* **51**, 791-796 (1959).
- 59SCH Schaefer, J. R., "Estimation of the Heat and Entropy of Fusion of Some Polyhydrocarbons", *J. Polymer Sci.* **38**, 549-552 (1959).
- 60SPE/ROS Speros, D. M., and Rossini, F. D., "Heats of Combustion and Formation of Naphthalene, the two Methylnaphthalenes, cis- and trans-Decahydro-naphthalene, and Related Compounds", *J. Phys. Chem.* **64**, 1723-1727 (1960).
- 60TAT/PAP Tatevskii, V. M., and Papulov, Yu. G., "Relationship between the Energy of Formation of a Molecule from its Atoms and its Structure. I. Energy of a Molecule as the Sum of Energies of its Chemical Bonds", *Zhur. Fiz. Khim.* **34**, 241-258 (1960).
- 60TAT/PAP 2 Tatevskii, V. M., and Papulov, Yu. G., "Relationship between the Energy of Formation of a Molecule from its Atoms and its Structure. II. Energy of a Molecule as the Sum of Energies of Interaction between Pairs of Atoms", *Zhur. Fiz. Khim.* **34**, 489-504 (1960).
- 60TAT/PAP 3 Tatevskii, V. M., and Papulov, Yu. G., "Relationship between the Structure of a Molecule and its Energy of Formation from its Atoms. III. Other Methods of Considering the Relationship between the Energy and Structure of a Molecule", *Zhur. Fiz. Khim.* **34**, 708-715 (1960).
- 61MCC/GOO McCullough, J. P., and Good, W. D., "Correlation of the Heat of Formation Data for Organic Sulfur Compounds", *J. Phys. Chem.* **65**, 1430-1432 (1961).


- 61ROC/ROS Rockenfeller, J. D., and Rossini, F. D., "Heats of Combustion, Isomerization, and Formation of Selected C₆, C₈, and C₁₀ Monoolefin Hydrocarbons", *J. Phys. Chem.* **65**, 267-272 (1961).
- 61TAT/BEN Tatevskii, V. M., Benderskii, V. A., and Yarovoi, S. S., "Rules and Methods for Calculating the Physicochemical Properties of Paraffinic Hydrocarbons", translated from the Russian by M. F. Mullins and edited by B. P. Mullins, (Pergamon Press, New York, Oxford, London, Paris, 1961).
- 62COX Cox, J. D., "A Bond Energy Scheme for Aliphatic and Benzenoid Compounds", *Tetrahedron* **18**, 1337-1350 (1962).
- 62SKI Skinner, H. A., "An Examination of Allen's Empirical Bond-Energy Scheme and its Application to Paraffins, Cycloalkanes, Olefins, Alkyl Alcohols and Bromides, and Amines", *J. Chem. Soc.* 4396-4408 (1962).
- 63BON Bondi, A., "Heat of Sublimation of Molecular Crystals. A Catalog Molecular Structure Increments", *J. Chem. Eng. Data* **8**, 371-381 (1963).
- 63FIS Fishtine, S. H., "Reliable Latent Heats of Vaporization", *Ind. Eng. Chem.* **55**, 47-56 (1963).
- 63SKI/PIL Skinner, H. A., and Pilcher, G., "Bond Energy Term Values in Hydrocarbons and Related Compounds", *Quart. Rev.* **17**, 264-288 (1963).
- 65FIN/MES Finke, H. L., Messerly, J. F., and Todd, S. S., "Thermodynamic Properties of n-Propyl-, n-Butyl-, and n-Decyl-substituted Cyclohexane from 10 to 370 K", *J. Phys. Chem.* **69**, 2094-2100 (1965).
- 65RIH/DOR Rihani, D. N., and Doraiswamy, L. K., "Estimation of Heat Capacity of Organic Compounds from Group Contributions", *Ind. Eng. Chem. Fundamentals* **4**, 17-21 (1965).
- 65VER/DOR Verma, K. K., and Doraiswamy, L. K., "Estimation of Heats of Formation of Organic Compounds", *Ind. Eng. Chem. Fundamentals* **4**, 389-396 (1965).
- 66BON Bondi, A., "Estimation of the Heat Capacity of Liquids", *Ind. Eng. Chem. Fundamentals* **5**, 442-449 (1966).
- 66NEL/SUN Nelander, B., and Sunner, S., "On the Interpretation of Ring Strain", *J. Chem. Phys.* **44**, 2476-2480 (1966).
- 66SOM/ZWO Somayajulu, G. R., and Zwolinski, B. J., "Generalized Treatment of Alkanes", *Trans. Faraday Soc.* **62**, 2327-2340 (1966).
- 66SOM/ZWO 2 Somayajulu, G. R., and Zwolinski, B. J., "Isomeric Variation Procedures for Physicochemical Properties of Alkanes", *J. Phys. Chem.* **70**, 3498-3501 (1966).
- 66VIT Vitvitskii, A. I., "Problems concerning the Calculation of the Heats of Formation of Aliphatic Hydrocarbons", *Teor. Eksp. Khim.* **2**, 392-398 (1966).
- 67BRA/THO Bradford, M. L., and Thodos, G., "Latent Heats of Vaporization of Hydrocarbons", *J. Chem. Eng. Data* **12**, 373-376 (1967).
- 67JAN Janz, G. J., "Thermodynamic Properties of Organic Compounds", (Academic Press, New York, 1967).
- 67MES/GUT Messerly, J. F., Guthrie, G. B., Todd, S. S., and Finke, H. L., "Low Temperature Thermal Data for n-Pentane, n-Heptadecane, and n-Octadecane, Revised Thermodynamic Functions for the n-Alkanes, C₅-C₁₈", *J. Chem. Eng. Data* **12**, 338-346 (1967).
- 67ROD Rodgers, A. S., "Estimation of the Heats of Formation of Chlorofluorocarbons", *J. Phys. Chem.* **71**, 1996-2000 (1967).
- 67VIT Vitvitskii, A. I., "Calculation of the Heats of Formation of Aliphatic Hydrocarbons", *Zhur. Org. Khim.* **3**, 1354-1358 (1967).
- 68BEN Benson, S. W., "Thermochemical Kinetics" (J. Wiley & Sons, Inc., New York, 1968)
- 68LAC/SKI Lacher, J. R., and Skinner, H. A., "Bond Energy Additivity and Bond Interactions in Fluoro-halogenated Hydrocarbons", *J. Chem. Soc. A*, 1034-1038 (1968).
- 68MOF Moffat, J. B., "Correlation of Some Thermodynamic Properties of Nitrites", *J. Chem. Eng. Data* **13**, 36-38 (1968).
- 68RIH Rihani, D. N., "How to Estimate Heats of Formation", *Hydrocarbon Process.* **47**, No. 3, 137-144 (1968).
- 68RIH/DOR Rihani, D. N., and Doraiswamy, L. K., "Estimation of the Ideal Gas Entropy of Organic Compounds", *Ind. Eng. Chem. Fundamentals* **7**, 375-380 (1968).
- 68SCH Schotte, W., "Prediction of Heats of Formation", *J. Phys. Chem.* **72**, 2422-2431 (1968).
- 69BEN/CRU Benson, S. W., Cruickshank, F. R., Golden, D. M., Haugen, G. R., O'Neal, H. E., Rodgers, A. S., Shaw, R., and Walsh, R., "Additive Rules for the Estimation of Thermochemical Properties", *Chem. Rev.* **69**, 269-324 (1969).
- 69GOO/SMI Good, W. D., and Smith, N. K., "Enthalpies of Combustion of Toluene, Benzene, Cyclohexane, Cyclohexene, Methylcyclopentane, 1-Methylcyclopentene, and n-Hexane", *J. Chem. Eng. Data* **14**, 102-106 (1969).
- 69STU/WES Stull, D. R., Westrum, E. F., Jr., and Sinke, G. C., "The Chemical Thermodynamics of Organic Compounds" (J. Wiley & Sons, Inc., New York, 1969).
- 69SHA Shaw, R., "Heat Capacities of Liquids. Estimation of Heat Capacity at Constant Pressure and 25°C using Additivity Rules", *J. Chem. Eng. Data* **14**, 461-465 (1969).
- 70COX/PIL Cox, J. D., and Pilcher, G., "Thermochemistry of Organic and Organometallic Compounds" (Academic Press, London, 1970).
- 70GOO Good, W. D., "The Enthalpies of Combustion of the Isomeric Pentanes", *J. Chem. Thermodynam.* **2**, 237-244 (1970).
- 70JOS Joshi, R. M., "A Critical Evaluation of Bond Energy/Group Contribution Methods for Calculating the Heat of Formation: Development of a New Generalized Scheme. Part I. Alkanes", *J. Macromol. Sci.-Chem.* **A4**, 1819-1839 (1970).
- 71JOS Joshi, R. M., "Bond Energy/Group Contribution Methods of Calculating the Standard Heat of Formation: Development of a New Generalized Bond-Energy Scheme for Monomers and Polymers. Part II. Hydrocarbons", *J. Macromol. Sci.-Chem.* **A5**, 687-734 (1971).
- 71SHA Shaw, R., "Heat of Formation of Nitroaromatics. Group Additivity for Solids", *J. Phys. Chem.* **75**, 4047-4049 (1971).
- 72JOS Joshi, R. M., "Bond energy/Group Contribution Methods of Calculating the Standard Heat of Formation: Development of a New Generalized Bond-Energy Scheme for Monomers and Polymers Part III. Alicyclic Hydrocarbons", *J. Macromol. Sci.-Chem.* **A6**, 595-629 (1972).
- 72SOM/ZWO Somayajulu, G. R., and Zwolinski, B. J., "Generalized Treatment of Alkanes. Part 2", *J. Chem. Soc. Faraday Trans. 2*, 68, 1971-1987 (1972).
- 73EIG/GOL Eigenmann, H. K., Golden, D. M., and Benson, S. W., "Revised Group Additivity Parameters for the Enthalpies of Formation of Oxygen Containing Organic Compounds", *J. Phys. Chem.* **77**, 1687-1691 (1973).
- 73KIS/SUG Kishimoto, K., Suga, H., Syuzo, S., "Calorimetric Study of the Glassy State VIII. Heat Capacity and Relaxation Phenomena of Isopropylbenzene", *Bull. Chem. Soc. Japan* **46**, 3020-3031 (1973).

- 74HER Herndon, W. C., "Thermochemical Parameters for Benzenoid Hydrocarbons", *Thermochim. Acta* **8**, 225-237 (1974).
- 74JOS Joshi, R. M., "Bond Energy/Group Contribution Methods of Calculating the Standard Heat of Formation: Development of a New Generalized Bond-Energy Scheme for Monomers and Polymers. Part IV. Halocarbons", *J. Macromol. Sci. -Chem.* A8, 861-885 (1974).
- 74MAN Mansson, M., "A Calorimetric Study of peri-Strain in 1,8-Dimethylnaphthalene and 1,4,5,8-Tetramethyl-naphthalene", *Acta Chem. Scand.* 28B, 677-680 (1974).
- 74SAB/CHA Sabbah, R., Chastel, R., and Laffitte, M., "Etude Thermochimique des Methylnaphthalenes", *Thermochim. Acta* 10, 353-358 (1974).
- 74SEA/FRE Seaton, W. H., Freedman, E., and Treweek, D. N., American Society for Testing and Materials, ASTM Data Series Publication DS 51, "CHETAH - The ASTM Chemical Thermodynamic and Energy Release Evaluation Program" (ASTM, 1916 Race St., Philadelphia, PA 19103, 1974).
- 74SOM/ZWO Somayajulu, G. R., and Zwolinski, B. J., "Generalized Treatment of Alkanes. Part 3. Triatomic Additivity", *J. Chem. Soc., Faraday Trans. 2*, 70, 967-972 (1974).
- 74SOM/ZWO 2 Somayajulu, G. R., and Zwolinski, B. J., "Generalized Treatment of Alkanes. Part 4. Triatomic Additivity Applications to Substituted Alkanes", *J. Chem. Soc., Faraday Trans. 2*, 70, 973-993 (1974).
- 74SOM/ZWO 3 Somayajulu, G. R., and Zwolinski, B. J., "Generalized Treatment of Aromatic Hydrocarbons. Part I. Triatomic Additivity to Parent Aromatic Hydrocarbons", *J. Chem. Soc., Faraday Trans. 2*, 71, 1928-1941 (1974).
- 75JOS Joshi, R. M., "Bond Energy/Group Contribution Methods of Calculating the Standard Heat of Formation: Development of a New Generalized Bond-Energy Scheme for Monomers and Polymers. Part V. Oxygen Compounds", *J. Macromol. Sci. -Chem.* A9, 1309-1383 (1975).
- 75SHA Shaw, R., "Thermochemistry of Hydrazo, Azo, and Azoxy Groups", Chapter 3 in *The Chemistry of the Hydrazo, Azo, and Azoxy Groups*, Part 1, edited by Saul Patai (John Wiley & Sons, Inc., Chichester, 1975).
- 76AND/MAR Andon, R. J. L., and Martin, J. F., "Thermodynamic Properties of Hexacosane", *J. Chem. Thermodynamics* 8, 1159-1166 (1976).
- 76BEN Benson, S. W., "Thermochemical Kinetics", Second Edition, (J. Wiley & Sons, Inc., New York, 1976).
- 77LUR/BEN Luria, M., and Benson, S. W., "Heat Capacities of Liquid Hydrocarbons. Estimation of Heat Capacities at Constant Pressure as a Temperature Function, using Additivity Rules", *J. Chem. Eng. Data* 22, 90-100 (1977).
- 77PED/RYL Pedley, J. B., and Rylance, J., "Sussex - Computer Analysed Thermochemical Data: Organic and Organometallic Compounds", (University of Sussex, School of Molecular Sciences, Falmer, Brighton, U.K., 1977).
- 77REI/PRA Reid, R. C., Prausnitz, J. M., and Sherwood, T. K., "The Properties of Gases and Liquids", Third Edition, (McGraw-Hill Book Co., 1977).
- 77SHA/GOL Shaw, R., Golden, D. M., and Benson, S. W., "Thermochemistry of Some Six-Membered Cyclic and Polycyclic Compounds Related to Coal", *J. Phys. Chem.* 81, 1716-1729 (1977).
- 77STE/GOL Stein, S. E., Golden, D. M., and Benson, S. W., "Predictive Scheme for Thermochemical Properties of Polycyclic Aromatic Hydrocarbons", *J. Phys. Chem.* 81, 314-317 (1977).
- 78SEL/STR Sellers, P., Stridh, G., and Sunner, S., "An Examination of the Constancy of the CH₂ Increment in Homologous Series", *J. Chem. Eng. Data*, 23, 250-255 (1978).
- 78SHA Shaw, R., "Thermochemistry of Acetylenes", Chapter 3 in *The Chemistry of the Carbon-Carbon Triple Bond*, Part 1, edited by Saul Patai (John Wiley & Sons, Inc., Chichester, 1978).
- 79DAV Davidson, I. M. T., "Enthalpies of Formation of Silanes", *J. Organometal. Chem.* 170, 365-372 (1979).
- 79JOS Joshi, R. M., "Bond Energy Scheme for Estimating Heats of Formation of Monomers and Polymers. Part VI. Sulfur Compounds", *J. Macromol. Sci. -Chem.* A13, 1015-144 (1979).
- 80DUC/GRU Ducros, M., Gruson, J. F., and Sannier, H., "Estimation des Enthalpies de Vaporisation des Composes Organiques Liquides. Partie 1. Applications aux Alcanes, Cycloalcanes, Alcenes, Alcanes, Thiols, Chloro et Bromoalcanes, Nitriles, Esters, Acides et Aldehydes", *Thermochim. Acta* 36, 39-65 (1980).
- 81HOL/FIN Holmes, J. L., Fingas, M., and Lossing, F. P., "Towards a General Scheme for Estimating the Heats of Formation of Organic Ions in the Gas Phase. Part I. Odd-electron Cations", *Can. J. Chem.* 59 80-93 (1981).
- 81LEV Levanova, S. V., "Use of Additivity Schemes for Calculating Thermodynamic Functions of Haloalkanes", *Izv. Vyssh. Uchebn. Zaved. Khim. Khim. Tekhnol.* 24, 1213-1216 (1981).
- 81ONE/RIN O'Neal, H. E., and Ring, M. A., "An Additivity Scheme for the Estimation of Heats of Formation, Entropies, and Heat Capacities of Silanes, Polysilanes, and their Alkyl Derivatives", *J. Organometal. Chem.* 213, 419-434 (1981).
- 81STE/BAR Stein, S. E., and Barton, B. D., "Chemical Thermodynamics of Polyaromatic Compounds containing Heteroatoms and Five-Membered Rings", *Thermochim. Acta* 44, 265-281 (1981).
- 82COX/ANG Cox, J. D., Angus, S., Armstrong, G. T., Freeman, R. D., Laffitte, M., Schneider, G. M., Somsen, G., Alcock, C. B., and Gilles, P. W., (Task Group), IUPAC, Phys. Chem. Div., Comm. on Thermodynamics, "Notation for States and Processes, Significance of the Word Standard in Chemical Thermodynamics, and Remarks on and Remarks on Commonly Tabulated Forms of Thermodynamic Functions", *Pure & Appl. Chem.* 54, 1239-1250 (1982).
- 82CHU/NGU Chu, J. Y., Tam T. Nguyen, and King, K. D., "Group Additivity Parameters for the Estimation of Thermochemical Properties of Gaseous and Liquid Nitriles", *J. Phys. Chem.* 86, 443-447 (1982).
- 82JOS Joshi, R. M., "Bond Energy Scheme for Estimating Heats of Formation of Monomers and Polymers. Part VII. Nitrogen Compounds", *J. Macromol. Sci. -Chem.* A18, 861-911 (1982).
- 82KIZ/LEB Kizin, A. N., and Lebedev, Yu. A., "Calculation of Enthalpies of Formation of Polysubstituted Aliphatic Compounds in the Solid Phase", *Doklady Akad. Nauk SSSR* 262, 914-917 (1982).
- 82WAG/EVA Wagman, D. D., Evans, W. H., Parker, V. B., Schumm, R. H., Halow, I., Bailey, S. M., Churney, K. L., and Nuttall, R. N., "The NBS Tables of Chemical Thermodynamic Properties. Selected Values for Inorganic and C1 and C2 Organic Substances in SI Units", *J. Phys. & Chem. Ref. Data* 11, Supplement No. 2, 392 pp. (1982).

- 83AND/ANT Andreevskii, D. N., Antonova, Z. A., and Radyuk, V. I., "Increments in the Thermodynamic Properties for Halogenoalkanes", *Zhur. Fiz. Khim.* **57**, 837-841 (1983).
- 83AND/ANT 2 Andreevskii, D. N. and Antonova, Z. A., "Estimation of the Thermodynamic Properties of Polyhalogenoethanes and Polyhalogenoethylenes. II. Entropies and Heat Capacities", *Zhur. Fiz. Khim.* **57**, 2202-2205 (1983).
- 83KOH Kohli, R., "Heat Capacity and Thermodynamic Properties of Alkali Metal Compounds. II. Estimation of the Thermodynamic Properties of Cesium and Rubidium Zirconates", *Thermochim. Acta* **65**, 285-293 (1983).
- 83KOH Kohli, R., "Heat Capacity and Thermodynamic Properties of Alkali Metal Compounds. II. Estimation of the Thermodynamic Properties of Cesium and Rubidium Zirconates", *Thermochim. Acta* **65**, 285-293 (1983).
- 83LAV/SKO Lavygin, I. A., and Skorokhodov, I. I., "An Additive Method for the Estimation of the Heats of Vaporization of Organosilicon Compounds", *Zhur. Fiz. Khim.* **57**, 1044-1045 (1983).
- 84DOM/EVA Domalski, E. S., Evans, W. H., and Hearing, E. D., "Heat Capacities and Entropies of Organic Compounds in the Condensed Phase", *J. Phys. & Chem. Ref. Data* **13**, Supplement No. 1, 286 pp., (1984).
- 85HIS/BEN Hisham, M. W. M. and Benson, S. W., "Empirical Relations between the Heats of Formation of the Different Valence States of Solid Polyvalent Metal Oxides", *J. Phys. Chem.* **89**, 1905-1909 (1985).
- 85HIS/BEN 2 Hisham, M. W. M. and Benson, S. W., "Correlations of Heats of Formation of Different Valency States of Solid, Polyvalent Binary Metal Compounds", *J. Phys. Chem.* **89**, 3417-3425 (1985).
- 86CAR Cardozo, R. L., "Prediction of the Enthalpy of Combustion of Organic Compounds", *AIChE Journal* **32**, 844-847 (1986).
- 86HIS/BEN Hisham, M. W. M. and Benson, S. W., "Thermochemistry of Inorganic Solids. 3. Enthalpies of Formation of Solid Metal Oxyhalide Compounds", *J. Phys. Chem.* **90**, 885-888 (1986).
- 86MAR Marsh, K. N., *et al.*, "Selected Values of Hydrocarbons and Related Compounds", Thermodynamics Research Center, Texas A & M University, College Station, TX (loose-leaf data sheets, extant 1986).
- 86PED/NAY Pedley, J. B., Naylor, R. D., and Kirby, S. P., "Thermochemical Data of Organic Compounds", Second Edition, (Chapman and Hall, London and New York, 1986).
- 87HIS/BEN Hisham, M. W. M. and Benson, S. W., "Thermochemistry of Inorganic Solids. 5. Empirical Relations among Enthalpies of Formation of Oxides, Carbonates, Sulfates, Hydroxides, and Nitrates", *J. Chem Eng. Data* **32**, 243-247 (1987).
- 87CHI/ANN Chickos, J. S., Annunziata, R., Ladon, L. H., Hyman, A. S., and Liebman, J. F., "Estimating Heats of Sublimation of Hydrocarbons. A Semiempirical Approach", *J. Org. Chem.* **51**, 4311-4313 (1987).
- 87REI/PRA Reid, R. C., Prausnitz, J. M., and Poling, B. E., "The Properties of Gases and Liquids", Fourth Edition, (McGraw-Hill Book Co., 1987).
- 89DOM/HEA Domalski, E. S., and Hearing, E. D., "Thermodynamic Properties of Organic Compounds at 298.15 K. I. Estimation Algorithm and Database for over 1300 Hydrocarbon Compounds", NBS Technical Note (in preparation), 1989.

Appendix A. Example Calculations of Enthalpy of Formation, Heat Capacity, and Entropy at 298.15 K for Some Hydrocarbon Compounds

n-Hexane




Enthalpy of formation (gas phase)

No. of groups	Group	Group value	Sum (kJ/mol)
2	C-(C)(H) ₃	-42.26	-84.52
4	C-(C) ₂ (H) ₂	-20.63	-82.52
			-167.04

Compare with experimental value: -167.11 kJ/mol (44PRO/ROS, 69GOO/SMI, 86PED/NAY).

2-Methylhexane




Enthalpy of formation (liquid phase)

No. of groups	Group	Group value	Sum (kJ/mol)
3	C-(C)(H) ₃	-47.61	142.83
3	C-(C) ₂ (H) ₂	-25.73	-77.19
1	C-(C) ₃ (H)	-4.77	-4.77
2	-CH ₃ corr	-2.18	-4.36
			-229.15

Compare with experimental value: -229.49 kJ/mol (45PRO/ROS 2, 86PED/NAY).

2,2,4,4-Tetramethylpentane




Enthalpy of formation (liquid phase)

No. of groups	Group	Group value	Sum (kJ/mol)
6	C-(C)(H) ₃	-47.61	-285.66
1	C-(C) ₂ (H) ₂	-25.73	-25.73
2	C-(C) ₄	17.99	35.98
6	-CH ₃ , corr	-0.64	-3.84
			-279.25

Compare with experimental value: -279.99 kJ/mol (47JOH/PRO, 86PED/NAY)

***trans*-4,4-Dimethylpentene-2**




Enthalpy of formation (liquid phase)

No. of groups	Group	Group value	Sum (kJ/mol)
3	C-(C)(H) ₃	-47.61	-142.83
1	C-(C ₆)(H) ₃	-47.61	-47.61
2	Cd-(C)(H)	31.05	62.10
1	C-(C ₆)(C) ₃	20.79	20.79
3	-CH ₃ corr	-4.39	-13.17
			-120.72

Compare with experimental value: -121.71 kJ/mol (61ROC/ROS 86PED/NAY).

Isopropylbenzene




Enthalpy of formation (solid phase)

No. of groups	Group	Group value	Sum (kJ/mol)
2	C-(C)(H) ₃	-46.74	-93.48
1	C-(C) ₂ (C _B)(H)	-3.50	-3.50
1	C _B -(C)(C _B) ₂	13.90	13.90
5	C _B -(H)(C _B) ₂	6.53	32.65
2	-CH ₃ corr.	-2.34	-4.68
			-55.11

Compare with experimental value: -48.45 kJ/mol (45JOH/PRO, 73KIS/SUG).


1-Octyne



Heat capacity (gas phase)

No. of groups	Group	Group value	Sum (kJ/mol)
1	C-(C)(H) ₃	25.73	25.73
4	C-(C) ₂ (H) ₂	22.89	91.56
1	C-(C ₂)(C)(H) ₂	20.97	20.97
1	C _r -(C)	13.22	13.22
1	C _r -(H)	22.55	22.55
			174.03


Compare with experimental value: 173.97 J/mol·K (69STU/WES).

1,3,5-Trimethylbenzene

Heat capacity (liquid phase)

No. of groups	Group	Group value	Sum (kJ/mol)
3	C-(C _B)(H) ₃	36.48	109.44
3	C _B -(H)(C _B) ₂	22.68	68.04
3	C _B -(C)(C _B) ₂	10.10	30.30
207.78			


Compare with experimental value: 209.33 J/mol·K (55TAY/KIL, 84DOM/EVA).

n-Butylcyclohexane

Heat capacity (liquid phase)

No. of groups	Group	Group value	Sum (kJ/mol)
1	C-(C)(H) ₃	36.48	36.48
8	C-(C) ₂ (H) ₂	30.42	243.36
1	C-(C) ₃ (H)	21.38	21.38
1	Ring corr.	-26.21	-26.21
275.01			


Compare with experimental value: 271.04 J/mol·K (65FIN/MES, 84DOM/EVA).

1,2-Diphenylethane

Heat capacity (solid phase)

No. of groups	Group	Group value	Sum (kJ/mol)
10	C _B -(H)(C _B) ₂	20.13	201.30
2	C _B -(C)(C _B) ₂	-23.26	-46.52
2	C-(C _B)(C)(H) ₂	49.38	98.76
253.54			


Compare with experimental value: 253.55 J/mol·K (31SMI/AND, 84DOM/EVA).

n-Hexadecane

Entropy (gas phase)

No. of groups	Group	Group value	Sum (kJ/mol)
2	C-(C)(H) ₃	127.32	254.64
14	C-(C) ₂ (H) ₂	39.16	548.24
1	-R ln[N(tot)]	-24.03	-24.03
778.85			


Compare with experimental value: 778.31 J/mol·K (69STU/WES)

2,2-Dimethylhexane

Entropy (gas phase)


No. of groups	Group	Group value	Sum (kJ/mol)
4	C-(C)(H) ₃	127.32	509.28
3	C-(C) ₂ (H) ₂	39.16	117.48
1	C-(C) ₄	-149.49	-149.49
1	-R ln[N(tot)]	-45.67	-45.67
431.60			

Compare with experimental value: 431.20 J/mol·K (69STU/WES).

Ethylbenzene**Entropy (liquid phase)**

No. of groups	Group	Group value	Sum (kJ/mol)
1	C-(C)(H) ₃	83.30	83.30
1	C-(C)(C _B)(H) ₂	47.40	47.40
1	C _B -(C)(C _B) ₂	-19.50	-19.50
5	C _B -(H)(C _B) ₂	28.87	144.35
255.55			

Compare with experimental value: 255.01 J/mol·K (44GUT/SPI, 84DOM/EVA).

Naphthalene**Entropy (solid phase)**

No. of groups	Group	Group value	Sum (kJ/mol)
8	C _B -(H)(C _B) ₂	22.75	182.00
2	C _{BF} -(C _{BF})(C _B) ₂	-6.00	-12.00
170.00			

Compare with experimental value: 167.40 J/mol·K (57MCC/FIN, 84DOM/EVA).

Appendix B. Comparison of Calculated Thermodynamic Properties in the Gas Phase with Benson's Values at 298.15 K**Enthalpy of formation**

Compound	$\Delta_f H^\circ$ (This work)	$\Delta_f H^\circ$ (76BEN)	Residual kJ/mol
propane	-105.15	-105.98	0.83
2-methylbutane	-153.10	-153.26	0.16
<i>trans</i> -2-hexene	-53.39	-54.02	0.63
<i>p</i> -xylene	18.00	15.98	2.02
<i>n</i> -butylbenzene	-12.17	-12.18	0.01

Heat capacity

Compound	C_p° (This work)	C_p° (76BEN)	Residual J/mol·K
<i>n</i> -hexane	143.02	143.85	-0.83
2,5-dimethylhexane	188.86	187.61	-1.25
<i>cis</i> -3-hexene	122.17	123.85	-1.68
ethylbenzene	129.14	129.29	-0.15
cyclopentane	83.01	87.86	-4.85

Entropy

Compound	S° (This work)	S° (76BEN)	Residual J/mol·K
<i>n</i> -pentane	348.09	348.69	-0.60
2-methylpropane	291.82	294.68	-2.86
1-hexene	383.28	386.81	3.53
benzene	269.20	268.78	0.42
methylcyclohexane	344.36	343.34	1.02


Comparison of the small data set in Appendix B to corresponding experimental values for these compounds shows that residual for $\Delta_f H^\circ$ are less than 4 kJ/mol and those for C_p° and S° are below 4 J/mol·K. An improvement on the part of our group contribution values for the gas phase over those of Benson *et al.*, is demonstrated about half of the time.

Appendix C. Methyl Repulsion Corrections for Branched Hydrocarbons

Methyl Repulsion Corrections in Branched Alkane Hydrocarbons with Quaternary Carbon Atoms

Several examples of application of the methyl repulsion corrections to the calculation of $\Delta_f H^\circ$ in the gas phase for hydrocarbons which contain quaternary carbon atoms are given below.

2,2-Dimethylpropane




Enthalpy of formation (gas phase)

No. of groups	Group	Group value	Sum (kJ/mol)
4	C-(C)(H) ₃	-42.26	-169.04
1	C-(C) ₄	19.20	19.20
4	Quat. CH ₃ corr	-4.56	-18.24
-168.08			

Compare with experimental value: -167.99 kJ/mol (70GOO, 86PED/NAY).

Compare with estimated value from 76BEN: -168.62 kJ/mol.

2,2-Dimethylbutane




Enthalpy of formation (gas phase)

No. of groups	Group	Group value	Sum (kJ/mol)
4	C-(C)(H) ₃	-42.26	-169.04
1	C-(C) ₂ (H) ₂	-20.63	-20.63
1	C-(C) ₄	19.20	19.20
3	Quat. CH ₃ corr	-4.56	-13.68
-184.15			

Compare with experimental value: -186.10 kJ/mol (41PRO/ROS, 86PED/NAY).

Compare with estimated value from 76BEN: -182.55 kJ/mol

3,3-Dimethylpentane




Enthalpy of formation (gas phase)

No. of groups	Group	Group value	Sum (kJ/mol)
4	C-(C)(H) ₃	-42.26	-169.04
2	C-(C) ₂ (H) ₂	-20.63	-41.26
1	C-(C) ₄	19.20	19.20
2	Quat. CH ₃ corr	-4.56	-9.12
-200.22			

Compare with experimental value: -201.17 kJ/mol (41PRO/ROS, 86PED/NAY).

Compare with estimated value from 76BEN: -196.48 kJ/mol.

3-Ethyl-3-methylpentane




Enthalpy of formation (gas phase)

No. of groups	Group	Group value	Sum (kJ/mol)
4	C-(C)(H) ₃	-42.26	-169.04
3	C-(C) ₂ (H) ₂	-20.63	-61.89
1	C-(C) ₄	19.20	19.20
1	Quat. CH ₃ corr	-4.56	-4.56
-216.29			

Compare with experimental value: -214.85 kJ/mol (45PRO/ROS, 86PED/NAY).

Compare with estimated value from 76BEN: -210.41 kJ/mol

3,3-Diethylpentane



Enthalpy of formation (gas phase)

No. of groups	Group	Group value	Sum (kJ/mol)
4	C-(C)(H) ₃	-42.26	-169.04
4	C-(C) ₂ (H) ₂	-20.63	-82.52
1	C-(C) ₄	19.20	19.20
-232.36			

Compare with experimental value: -232.34 kJ/mol (47JOH/PRO, 86PED/NAY).

Compare with estimated value from 76BEN: -224.35 kJ/mol.

Comparison of the methyl repulsion correction and Benson's gauche — correction in branched alkane hydrocarbons (gas phase)

Compound name and formula	Gauche isomers (76BEN)	Benson method (76 BEN) Δ_fH° (kJ/mol)			Methyl repulsion method (This work) Δ_fH° (kJ/mol)		
		expt	calc	resd	expt	calc	resd
2-Methylpropane	0	-134.18	-135.98	1.80	-134.18	-134.73	0.55
2,2-Dimethylpropane	0	-167.99	-168.62	0.63	-167.99	-168.08	0.09
2-Methylbutane	1	-153.63	-153.26	-0.37	-153.63	-153.10	-0.53
3-Methylpentane	2	-172.09	-170.54	-1.55	-172.09	-171.47	-0.62
2,2-Dimethylbutane	2	-186.10	-182.55	-3.55	-186.10	-184.15	-1.95
2,3-Dimethylbutane	2	-178.28	-179.91	1.63	-178.28	-180.42	2.14
3-Ethylpentane	3	-189.62	-187.82	-1.80	-189.62	-189.84	0.22
2,2,3-Trimethylbutane	5	-204.47	-202.51	-1.97	-204.47	-202.27	-2.20

Comparison of the methyl repulsion correction and Benson's gauche — correction in branched alkane hydrocarbons (gas phase) — Continued

Compound name and formula	Gauche isomers (76BEN)	Benson method (76 BEN) Δ_fH° (kJ/mol)			Methyl repulsion method (This work) Δ_fH° (kJ/mol)		
		expt	calc	resd	expt	calc	resd
3,3-Diethylpentane	8	-232.34	-224.35	-7.99	-232.34	-232.36	0.02
2,2,3,3-Tetramethylbutane	6	-225.60	-231.79	-6.23	-225.60	-219.00	-6.60
2,2,3,3-Tetramethylpentane	8	-236.69	-245.73	9.04	-236.69	-238.99	2.30
2,2,3,4-Tetramethylpentane	6	-236.86	-249.78	12.92	-236.86	-247.50	10.64
2,3,3,4-Tetramethylpentane	6	-236.10	-249.78	13.68	-236.10	-247.50	11.40
2,2,4,4-Tetramethylpentane	4	-241.63	-252.84	11.21	-241.63	-249.63	1.00

Examination of the residuals found above show that the methyl repulsion approach gives better agreement most of the time in contrast to the gauche correction developed by Benson *et al.*, for branched alkane hydrocarbons. Residuals are smaller in 10 out of the above 14 cases when the methyl repulsion approach is used.

Comparison of the methyl repulsion correction and Benson's gauche — correction in branched alkene hydrocarbons (gas phase)

Compound name and formula	Gauche isomers (76BEN)	Benson method (76BEN) Δ_fH° (kJ/mol)			Methyl repulsion method (This work) Δ_fH° (kJ/mol)		
		expt	calc	resd	expt	calc	resd
2-Methylpropene	0	-16.90	-15.90	-1.00	-16.90	-18.58	1.68
2-Methyl-1-butene	0	-35.31	-35.82	0.51	-35.31	-37.20	1.89
2,3-Dimethyl-1-butene	1	-62.38	-62.67	0.04	-62.38	-64.73	2.35
3-Methyl-2-ethyl-1-butene	1	-79.54	-82.59	3.05	-79.54	-83.35	3.81
2,3,3-Trimethyl-1-butene	2	-85.48	-90.04	4.56	-85.48	-92.39	6.91

Calculation of the enthalpies of formation at 298.15 K in the gas phase using either group values developed in this work or those reported by 76BEN yield Δ_fH° 's of about the same magnitudes. For this reason, residuals are also about the same magnitudes from both approaches.

Appendix D. Example Calculations of Total Symmetry Number for Some Hydrocarbon Compounds

The total symmetry number of the molecule, $N(\text{tot})$, is composed of two parts: the internal symmetry number and the external symmetry number. Both are defined below.

(1) The internal symmetry number, $N(\text{int})$, is determined when a terminal rotator has more than one indistinguishable conformation as it makes one complete rotation about an axis colinear with the single bond linking it to the rest of the molecule. If there are n such conformations, then the axis is called an n -fold axis of symmetry and $N(\text{int})$ for that rotor will be equal to n . For example, a methyl group has three such conformations, and therefore, a three-fold axis of symmetry and $N(\text{int})$ equal to 3. The phenyl group has two indistinguishable conformations as it makes one complete rotation about an axis through the bonding carbon atom of the aromatic ring and carbon atom in the para position to it, thus exhibiting two-fold symmetry.

Although nonsymmetrical terminal rotors do not affect $N(\text{tot})$ of a molecule, they need to be considered in determining the symmetry of compound rotors and $N(\text{ext})$ for a molecule.


Several example calculations are provided below within a format which allows for an examination of the internal and external symmetry elements in a molecule followed by a final calculation of the total symmetry number. Using this approach, the symmetries of the terminal rotors are noted. They are then conceptually spun to yield time-averaged symmetry similar to that of a single atom. The time-averaged rotor is called a pseudo-atom and the resulting molecule is called a pseudo-molecule. This process is continued until there are no further terminal rotors and the molecule becomes rigid. The value for $N(\text{int})$ will be the product of the symmetry numbers of all of the rotors.

(2) external symmetry number, $N(\text{ext})$, which is determined when a rigid molecule or a pseudo-molecule, derived from the spinning process, shows additional symmetry. $N(\text{ext})$ is defined as the total number of independent permutations of identical atoms (or pseudo-atoms) in a molecule that can be arrived at by rigid rotations of the entire molecule (or pseudo-molecule).

The value of $N(\text{tot})$ is equal to the product of all $N(\text{int})$ and $N(\text{ext})$.

Example Calculations for Several Hydrocarbon Compounds

methane



Symmetrical terminal rotors: None.


Pseudo-molecule representation: Y.

Internal symmetry: Methane has no internal symmetry.

External symmetry: Methane exhibits tetrahedral symmetry and has four three-fold symmetry axes. Each axis goes through a different C-H bond and includes the rotation of the remaining CH_3 group about this bond. Twelve unique rotational permutations of the methane molecule can be drawn, consequently, $N(\text{ext}) = 12$.

Total symmetry : Methane has $N(\text{tot}) = N(\text{ext}) = 12$.

ethane



Symmetrical terminal rotors: Ethane has two terminal methyl groups each of which have three-fold symmetry.

Pseudo-molecule representation: Y-Y.

Internal symmetry: Ethane has $N(\text{int}) = 3 \times 3 = 9$.

External symmetry: End-to-end symmetry can be visualized in the structural formula or the pseudo-molecule resulting in $N(\text{ext}) = 2$.

Total symmetry: $N(\text{tot}) = N(\text{int}) \times N(\text{ext}) = 3 \times 3 \times 2 = 18$. The value of $N(\text{tot})$ is the same for all of the n -alkanes.

n-butane




Symmetrical terminal rotors: *n* Butane has two methyl groups each of which have three-fold symmetry
 $N(\text{int}) = 3 \times 3 = 9$

Pseudo-molecule representation: Y-CH₃-CH₂-CH₂-Y

Internal symmetry: Spinning of the Y-CH₃ groups in the pseudo molecule does not result in any change in internal symmetry, hence, $N(\text{int}) = 9$

External symmetry: End-to-end symmetry can be visualized in either the structural formula or the pseudo-molecule resulting in $N(\text{ext}) = 2$

Total symmetry: $N(\text{tot}) = N(\text{int}) \times N(\text{ext}) = 3 \times 3 \times 2 = 18$. The value of $n(\text{tot})$ is the same for all of the n -alkanes.

2-Methylpropane


Symmetrical terminal rotors: 2-Methylpropane has three methyl groups each of which have three-fold symmetry. $N(\text{int}) = 3 \times 3 \times 3 = 27$.

Pseudo-molecule representation: $\text{Y}_3\text{-CH}$.

Internal symmetry: The $\text{Y}_3\text{-CH}$ pseudo-molecule itself has three-fold symmetry and contributes an additional $N(\text{int}) = 3$. The final $N(\text{int})$ for the molecule is $N(\text{int}) = 27 \times 3 = 81$.

External symmetry: Spinning of either the structural formula or the pseudo-molecule shows no end-to-end symmetry, hence, $N(\text{ext}) = 1$.

Total symmetry: $N(\text{tot}) = N(\text{int}) \times N(\text{ext}) = 81 \times 1 = 81$.

2,2-Dimethylpropane


Symmetrical terminal rotors: 2,2-Dimethylpropane has four methyl groups each of which have three-fold symmetry. $N(\text{int}) = 3 \times 3 \times 3 \times 3 = 81$.

Pseudo-molecule representation: CY_4 .

Internal symmetry: The CY_4 pseudo-molecule offers no additional elements of internal symmetry, hence, $N(\text{int}) = 81$.

External symmetry: The external symmetry for the CY_4 pseudo-molecule is the same as that for the methane molecule because 12 unique rotational permutations can be drawn for CY_4 , giving $N(\text{ext}) = 12$.

Total symmetry: $N(\text{tot}) = N(\text{int}) \times N(\text{ext}) = 81 \times 12 = 972$.

2-Methylbutane


Symmetrical terminal rotors: 2-Methylbutane has three terminal methyl groups each of which have three-fold symmetry. $N(\text{int}) = 3 \times 3 \times 3 = 27$.

Pseudo-molecule representation: $\text{Y}_2\text{CH}-\text{CH}_2\text{Y}$.

Internal symmetry: Examination of the pseudo-molecule does not result in any additional internal symmetry contributions. $N(\text{int}) = 27$.

External symmetry: Rotation of the pseudo-molecule about the central bond give no contribution to external symmetry. $N(\text{ext}) = 1$.

Total symmetry: $N(\text{int}) \times N(\text{ext}) = 27 \times 1 = 27$.

2,2-Dimethylbutane


Symmetrical terminal rotors: 2,2-Dimethylbutane has four methyl groups each of which have three-fold symmetry. $N(\text{int}) = 3 \times 3 \times 3 \times 3 = 81$.

Pseudo-molecule representation: $\text{Y}_3\text{C}-\text{CH}_2\text{Y}$.

Internal symmetry: Examination of the pseudo-molecule shows that the CY_3 group exhibits an additional amount of three-fold symmetry so that $N(\text{int})$ is now $3 \times 3 \times 3 \times 3 \times 3 = 243$.

External symmetry: Rotation of the pseudo-molecule does not yield any extra elements of external symmetry, hence, $N(\text{ext}) = 1$.

Total symmetry: $N(\text{int}) \times N(\text{ext}) = 243 \times 1 = 243$.

2,2,3,3-Tetramethylbutane


Symmetrical terminal rotors: 2,2,3,3-Tetramethylbutane has six methyl groups each of which have three-fold symmetry, giving $N(\text{int}) = 3 \times 3 \times 3 \times 3 \times 3 \times 3 = 729$.

Pseudo-molecule representation: $\text{Y}_3\text{C}-\text{CY}_3$.

Internal symmetry: Each CY_3 group of the pseudo-molecule exhibits three-fold symmetry enhancing $N(\text{int})$ to $729 \times 3 \times 3 = 6561$.

External symmetry: Rotation of the pseudo-molecule about the central bond shows it to have two-fold symmetry, hence, $N(\text{ext}) = 2$.

Total symmetry: $N(\text{int}) \times N(\text{ext}) = 6561 \times 2 = 13122$.

Ethylene

Symmetrical terminal rotors: None.

Pseudo-molecule representation: $\text{Y}=\text{Y}$.

Internal symmetry: Ethylene does not exhibit internal symmetry because the double bond prevents internal rotation.

External symmetry: Ethylene has two two-fold rotational axes, one horizontally through the two carbon atoms and the other vertically bisecting the two carbon atoms. From this one obtains $N(\text{ext}) = 2 \times 2 = 4$.

Total symmetry: $N(\text{tot}) = N(\text{ext}) = 4$.

Propylene


Symmetrical terminal rotors: Propylene has one methyl group which has three-fold symmetry giving $N(\text{int}) = 3$.

Pseudo-molecule representation: $\text{YCH}=\text{CH}_2$.

Internal symmetry: The pseudo-molecule does not provide any additional elements of internal symmetry, hence, $N(\text{int}) = 3$.

External symmetry: Rotation of the pseudo-molecule does not provide any additional elements of external symmetry, hence, $N(\text{ext}) = 1$.

Total symmetry: $N(\text{tot}) = N(\text{int}) \times N(\text{ext}) = 3 \times 1 = 3$. The latter value for $N(\text{tot})$ is applicable to all 1-olefinic hydrocarbons.

Benzene


Symmetrical terminal rotors: None.

Pseudo-molecule representation: Z.

Internal symmetry: Benzene has no internal symmetry.


External symmetry: Benzene has a six-fold rotational axis perpendicular to the ring which also passes through the center of the ring. Perpendicular to the six-fold axis, there are six two-fold axes through the opposite carbon atoms or bisecting the opposite carbon-carbon bonds, hence, $N(\text{ext}) = 2 \times 6 = 12$.

Total symmetry: $N(\text{tot}) = N(\text{ext}) = 12$.

Toluene

Symmetrical terminal rotors: Toluene has one methyl group which exhibits three-fold symmetry and the phenyl group which shows two-fold symmetry, giving $N(\text{int}) = 3 \times 2 = 6$.


Pseudo-molecule representation: Y-Z or,



Internal symmetry: Further examination of the pseudo-molecule does not give any additional elements of internal symmetry, hence, $N(\text{int}) = 6$.


External symmetry: Rotation of the pseudo-molecule does not give any additional elements of external symmetry, hence, $N(\text{ext}) = 1$.

Total symmetry: $N(\text{tot}) = N(\text{int}) \times N(\text{ext}) = 6 \times 1 = 6$.

t-Butylbenzene

Symmetrical terminal rotors: t-Butylbenzene has three methyl groups each of which have three-fold symmetry. This gives $N(\text{int}) = 3 \times 3 \times 3 = 27$. The phenyl group has two-fold symmetry for which $N(\text{int}) = 2$.

Pseudo-molecule representation: $\text{Y}_3\text{C}-\text{Z}$ or,



Internal symmetry: Examination of the pseudo-molecule shows that the CY_3 group exhibits three-fold symmetry resulting in another $N(\text{int}) = 3$. A final overall $N(\text{int})$ results which is equal to $27 \times 2 \times 3$ or 162.

External symmetry: Rotation of the pseudo-molecule does not provide any additional elements of external symmetry, hence, $N(\text{ext}) = 1$.

Total symmetry: $N(\text{tot}) = N(\text{int}) \times N(\text{ext}) = 162 \times 1 = 162$.

TABLE 1. Additive group contribution values for hydrocarbons at 298.15 K

Groups	Gas phase			Liquid phase			Solid phase		
	$\Delta_f H^\circ$ kJ/mol	C_p° J/mol·K	S°	$\Delta_f H^\circ$ kJ/mol	C_p° J/mol·K	S°	$\Delta_f H^\circ$ kJ/mol	C_p° J/mol·K	S°
Alkane groups									
C-(H)₃(C)	-42.26	25.73	127.32	-47.61	36.48	83.30	-46.74	67.45	56.69
C-(H)₂(C)₂	-20.63	22.89	39.16	-25.73	30.42	32.38	-29.41	21.92	23.01
C-(H)(C)₃	-1.17	20.08	-53.60	-4.77	21.38	-23.89	-5.98		
-CH₃ corr^a	-2.26			-2.18			-2.34		
C-(C)₄	19.20	16.53	-149.49	17.99	10.24	-98.65	12.47	-83.63 ^j	-33.19 ^j
-CH₃ corr^b	-4.56			-4.39			-4.35		
-CH₃ corr^c	-1.80			-1.77			-2.70		
-CH₃ corr^d	-0.64			-0.64			-2.24		
Alkene Groups									
C_d-(H)₂	26.32	21.38	115.52	21.75	28.37	86.19	22.43		
C_d-(H)(C)	36.32	18.74	33.05	31.05	24.60	28.58	25.48		
C_d-(C)₂	44.14	15.10	-50.84	39.16	23.22	-29.83	32.97		
C_d-(H)(C_d)	28.28	18.54	27.74	22.18	31.67	13.30	17.53	35.65	21.75
C_d-(C)(C_d)	36.78	17.57	-61.33	30.42	26.19	-41.92	27.91		
C_d-(C_d)(C_b)							56.07		
[C_d-(H)(C_b)]^e									
C_d-(O)(C_b)	37.95	15.90	-51.97	38.58					
[C_d-(H)(C_b)]^f									
C_d-(C_b)₂	32.88			30.83	25.10		49.90	32.50	
C-(H)₂(C)(C_d)	-20.88	20.63	38.20	-25.73	29.29	31.67	-24.35		
C-(H)(C)₂(C_d)	-1.63	27.49	-50.38	-5.02	30.12	-28.07	-6.49		
-CH₃ corr^a	-2.26			-2.18			-2.34		
C-(C)₃(C_d)	22.13	9.16	-150.23	20.79	28.74	-108.20	12.51		
-CH₃ corr^a	-4.56			-4.39			-4.35		
C-(H)(C)(C_d)₂	-1.00								
C-(H)₂(C_d)₂	-19.79	24.77	42.08	-25.27	40.88	19.32	-21.60		
C-(H)₂(C_d)(C_b)				-24.73					
C-(H)(C)(C_d)(C_b)				-6.90					
cis-correction^g	4.85	-8.03	5.06	5.27			5.73		
Alkyne groups									
C_t-(H)	113.50	22.55	101.96	104.47	39.96	67.57	110.34		
C_t-(C)	115.10	13.2	226.32	107.15	25.59	14.25	101.66		
C_t-(C_d)	121.42	10.71	39.92	114.77					
C_t-(C_b)	120.76	10.17	17.77	119.00			103.28	32.30	
C_t-(C_i)	120.76	14.27	25.94	104.80			103.28		
C-(H)₂(C)(C_i)	-19.70	20.97	42.80	-22.13	30.39	32.36	-29.41		
C-(H)(C)₂(C_i)	-3.16	17.45	-45.69	-2.18			-2.34		
-CH₃ corr^a	-2.26								
C-(C)₃(C_i)				22.83					
-CH₃ corr^b	-4.56			-4.39			-4.35		
C-(H)₂(C_i)₂	-41.14			-39.08					
C-(C)₂(C_i)₂				20.67					

TABLE 1. Additive group contribution values for hydrocarbons at 298.15 K — Continued

Groups	Gas phase			Liquid phase			Solid phase		
	$\Delta_f H^\circ$ kJ/mol	C_p° J/mol·K	S° J/mol·K	$\Delta_f H^\circ$ kJ/mol	C_p° J/mol·K	S° J/mol·K	$\Delta_f H^\circ$ kJ/mol	C_p° J/mol·K	S° J/mol·K
Allenic group									
C_a	142.67	15.86	26.28	134.68	30.04	14.39	131.08		
Aromatic groups									
$C_B-(H)(C_B)_2$	13.81	13.61	48.31	8.16	22.68	28.87	6.53	20.13	22.75
$C_B-(C)(C_B)_2$	23.64	9.75	-35.61	19.16	10.10	-19.50	13.90	-23.26	-5.50
$C_B-(C_d)(C_B)_2$	24.17	14.12	-33.85	19.12	9.44	-9.04	20.27	-20.00	-10.00
$[C_B-(C)(C_B)_2]^b$									
$C_B-(C_B)_3$	21.66	13.12	-36.57	17.21	17.07		17.03	-1.72	-6.00
$C-(H)_2(C)(C_B)$	-21.34	25.61	42.59	-24.81	22.90	47.40	-22.10	49.38	26.90
$C-(H)(C_2)(C_B)$	-4.52	22.45	-48.00	-5.82	17.50	-13.90	-3.50		
$C-(C_B)(C)_3$	18.28	18.28	-147.19	18.70	5.17	-96.10	21.57		
$C-(H)_2(C_B)_2$	-46.43			-26.50	32.91	51.97	-21.44	69.06	22.85
$C-(H)(C)(C_B)_2$				-21.47	11.50	28.12	16.40	43.55	
$C-(H)(C_B)_3$	-6.86						34.48	63.64	-12.62
$C-(C_B)_2(C)$							67.20	39.83	
$C-(C_B)_4$							64.89	58.74	
$C_{BF}-(C_B)(C_B)_2$	20.10			15.83	9.52	-5.54	14.10	2.30	-6.00
$C_{BF}-(C_B)(C_{BF})_2$	16.00			1.50			12.00	5.77	2.00
$C_{BF}-(C_{BF})_3$	3.59			-0.90			1.94	8.00	7.00
ortho corr	1.26	6.40	-2.50	3.26	3.50	0.00	5.00	0.00	0.00
meta corr	-0.63	0.71	0.00	0.00	0.00	0.00	2.00	0.00	0.00

^aUse for each adjacent methyl group.^bUse for each adjacent methyl group when one $C-(C)_4$ is present.^cUse for each adjacent methyl group when there is a $C-(C)_4$ and a $C-(C)_3(H)$ group present in the longest carbon chain.^dUse for each adjacent methyl group when there are two $C-(C)_4$ groups present in the longest carbon chain.^e $C_d-(H)(C_B) = C_d-(H)(C_d)$ ^f $C_d-(H)(C_i) = C_d-(H)(C_d)$ ^gWhen one of the groups is t-butyl, apply the following *cis* correction: $\Delta_f H^\circ(g) = 17.24$; $\Delta_f H^\circ(liq) = 17.48$; $\Delta_f H^\circ(solid) = -17.57$ ^h $C_B-(C_i)(C_B)_2 = C_B-(C_d)(C_B)_2$.ⁱ $C_{BF}-(C_B)_2(C_{BF})$: Do not use ortho, meta corrections for naphthalenesUse: 1. 0 substitution $C_p^\circ(g) = 11.83$; $S^\circ(g) = -19.66$ Use: 2. 1 substitution $C_p^\circ(g) = 14.39$; $S^\circ(g) = -21.50$ Use: 3. 2 substitutions $C_p^\circ(g) = 16.48$; $S^\circ(g) = -23.00$.^jGroup value calculated from experimental data on solid 2,2,3,3-tetramethylbutane (52SCO/DOU)

TABLE 2. Ring strain corrections to be applied to cycloalkanes and related hydrocarbons

Rings	Gas phase			Liquid phase			Solid phase		
	$\Delta_f H^\circ$ kJ/mol	C_p° J/mol·K	S°	$\Delta_f H^\circ$ kJ/mol	C_p° J/mol·K	S°	$\Delta_f H^\circ$ kJ/mol	C_p° J/mol·K	S°
cyclopropane	115.15	-12.73	134.86						
cyclobutane	110.89	-19.34	126.04	106.64	-10.68	51.48			
cyclopentane*	26.75	-31.44	116.22	23.59	-23.32	42.24	34.00		
cyclohexane*	0.39	-31.07	78.18	-2.06	-26.21	10.07	10.30		
cycloheptane	26.34	-37.14	73.97	23.50	-32.19	15.89			
cyclooctane	40.63	-43.17	70.78	38.10	-27.88	2.96			
cyclononane	52.91			50.40					
cyclodecane	43.63			50.61					
cycloundecane	47.56			47.55					
cyclododecane	17.31						46.27		
cyclotridecane	21.84			24.83					
cyclotetradecane	49.58						37.74		
cyclopentadecane	8.03						65.09		
cyclohexadecane	8.41						67.14		
cycloheptadecane	-13.59						69.56		
cyclopropene	224.21								
cyclobutene	125.81	-11.67	126.77						
cyclopentene*	24.31	-26.53	113.76	19.82	-15.82	48.37			
cyclohexene	5.36	-19.50	95.69	2.33	-20.26	29.34			
cycloheptene	21.81			18.26					
cyclooctene	24.65								
1,3-cyclo-pentadiene	24.94			24.79					
1,3-cyclo-hexadiene	18.79			18.18	-26.56	50.18			
1,4-cyclo-hexadiene					-34.22	36.41			
1,3-cyclo-heptadiene	27.54								
1,5-cyclo-octadiene	9.58			9.01	-7.45	23.35			
1,3,5-cyclo-heptatriene	14.90	-18.63	102.26	16.62	-54.00	84.96			
cyclo-octatetraene	69.61	-26.31	116.38	77.07	-68.18	113.89			
spiropentane	248.50	-19.97	286.59	242.58	2.60	162.81			
cis-decalin				-4.02	-54.12	53.75			
trans-decalin				-15.22	-57.63	53.67			
cis-hexahydroindan				16.55	-41.52	86.59			
trans-hexahydroindan				16.45	-46.00	79.98			
2,2-metacyclophane							55.06		
2,2-metaparacyclophane							109.46	-4.02	
2,2-paracyclophane							125.66	-13.18	-1.92
3,3-paracyclophane							65.68	14.90	
adamantane	-6.14						6.28		
bicyclo[2.2.2]octane	27.12			-67.59	-63.45		41.52		
bicyclo[3.3.3]undecane	99.06						124.10		

* Values given are for unsubstituted rings. When the ring is substituted, use the following values:

cyclopentane: C_p° , gas = -27.87, S° , gas = 118.39, liq = 56.65.

cyclohexane: C_p° , gas = -22.82, S° , gas = 83.97, liq = 25.10.

cyclopentene: C_p° , gas = -24.50, S° , gas = 117.11.

TABLE 3 *n*-Alkane hydrocarbons in the gas phase at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)		C_p° (J/mol-K)		S° (J/mol-K)		$N(\text{tot})$			
	expt	calc	resd	expt	calc	resd				
ethane	-83.85	-84.52	0.67	52.63	51.46	1.17	229.49	230.61	-1.12	18
propane	-104.68	-105.15	0.47	73.51	74.35	-0.84	269.91	269.77	0.14	18
butane	-125.65	-125.78	0.13	97.45	97.24	0.21	310.12	308.93	1.19	18
pentane	-146.82	-146.41	-0.41	120.21	120.13	0.08	348.95	348.09	0.86	18
hexane	-167.11	-167.04	-0.07	143.09	143.02	0.07	388.40	387.25	1.15	18
heptane	-187.65	-187.67	0.02	165.98	165.91	0.07	427.90	426.41	1.49	18
octane	-208.57	-208.30	-0.27	188.87	188.80	0.07	466.73	465.57	1.16	18
nonane	-228.24	-228.93	0.69	211.71	211.69	0.02	505.68	504.73	0.95	18
decane	-249.49	-249.56	0.07	234.60	234.58	0.02	544.63	543.89	0.74	18
undecane	-270.91	-270.19	-0.72	257.44	257.47	-0.03	583.58	583.05	0.53	18
dodecane	289.66	290.82	1.16	280.33	280.36	-0.03	622.50	622.21	0.29	18
tridecane				303.21	303.25	-0.04	661.45	661.37	0.08	18
tetradecane				326.06	326.14	-0.08	700.40	700.53	-0.13	18
pentadecane				348.95	349.03	-0.08	739.35	739.69	-0.34	18
hexadecane	-374.76	-373.34	-1.42	371.79	371.92	-0.13	778.31	778.85	-0.54	18
heptadecane				394.68	394.81	-0.13	817.26	818.01	-0.75	18
octadecane				417.56	417.70	-0.14	856.21	857.17	-0.96	18
nonadecane				440.41	440.59	-0.18	895.17	896.33	-1.16	18
eicosane				463.29	463.48	-0.19	934.12	935.49	-1.37	18

TABLE 4. *n*-Alkane hydrocarbons in the liquid phase at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)		C_p° (J/mol-K)		S° (J/mol-K)				
	expt	calc	resd	expt	calc	resd			
propane	-119.75	-120.95	1.20						
butane	-146.65	-146.68	0.03	135.00 ^a	133.80	1.20	231.00	231.36	-0.36
pentane	-173.47	-172.41	-1.06	167.19	164.22	2.97	263.47	263.74	-0.27
hexane	-198.66	-198.14	-0.52	194.97	194.64	0.33	296.06	296.12	-0.06
heptane	-224.22	-223.87	-0.35	224.93	225.06	-0.13	328.57	328.50	0.07
octane	-250.08	-249.60	-0.48	254.14	255.48	-1.34	361.20	360.88	0.32
nonane	-274.68	-275.33	0.65	284.39	285.90	-1.51	393.67	393.26	0.41
decane	-300.87	-301.06	0.19	314.47	316.32	-1.85	425.89	425.64	0.25
undecane	-327.23	-326.79	-0.44	345.05	346.74	-1.69	458.15	458.02	0.13
dodecane	-350.91	-352.52	1.61	375.97	377.16	-1.19	490.66	490.40	0.26
tridecane				406.89	407.58	-0.69	522.87	522.78	0.09
tetradecane				438.44	438.00	0.44	555.43	555.16	0.27
pentadecane				470.07	468.42	1.65	587.52	587.54	0.02
hexadecane	-456.14	-455.44	-0.70	501.45	498.84	2.61	619.65	619.92	0.27
heptadecane				534.34	529.26	5.08	652.24	652.30	0.06

^aObtained from extrapolation of experimental data of 40AST/MES to 298.15 K

TABLE 5. *n*-Alkane hydrocarbons in the solid phase at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
propane	-123.30	-122.89	-0.41						
butane	-153.39	-152.30	-1.09						
pentane	-181.46	-181.71	0.25						
hexane	-211.71	-211.12	-0.59						
heptane	-238.24	-240.53	2.29						
octane	-270.83	-269.94	-0.89						
nonane	-296.44	-299.35	2.91						
decane	-329.57	-328.76	-0.81						
undecane	-356.27	-358.17	1.90						
dodecane	-387.73	-387.58	-0.15						
tridecane	-414.43	-416.99	2.56						
tetradecane	-449.07	-446.40	-2.67						
pentadecane	-473.50	-475.81	2.31						
hexadecane	-509.49	-505.22	-4.27	441.79 ^a	441.78	0.01	434.84 ^a	435.52	-0.68
heptadecane	-532.33	-534.63	2.30						
octadecane	-567.43	-564.04	-3.39	485.64	485.62	0.02	480.20	481.54	-1.34
eicosane				(479.90)	529.46	-49.56	(558.56)	527.56	31.00
tetracosane				(730.94)	617.14	113.80	(651.03)	619.60	31.43
pentacosane				(769.02)	639.06	129.96	(671.11)	642.61	28.50
hexacosane				661.20	660.98	0.22	667.01	665.62	1.39
dotriaccontane	-968.34	-975.78	7.44	(877.38)	792.50	84.88	(851.44)	803.68	47.76
tritriaccontane				(900.82)	814.42	86.40	(877.80)	826.69	51.11

^aObtained from extrapolation of experimental data of 54FIN/GRO to 298.15 K.

TABLE 6. Substituted alkane hydrocarbons (tertiary carbon) in the gas state 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)			N(tot)
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
2-methylpropane	-134.18	-134.73	0.55	96.82	97.27	-0.45	294.64	291.82	2.82	81
2-methylbutane	-153.63	-153.10	-0.53	118.78	120.16	-1.38	343.59	340.12	3.47	27
2-methylpentane	-174.77	-173.73	-1.04	144.18	143.05	1.13	380.53	379.28	1.25	27
2-methylhexane	-194.64	-194.36	-0.28	165.98	165.94	0.04	419.99	418.44	1.55	27
2-methylheptane	-215.35	-214.99	-0.36	188.87	188.83	0.04	455.26	457.60	-2.34	27
2-methyloctane				217.07	211.72	5.35	495.89	496.76	-0.87	27
2-methylnonane				242.09	234.61	7.48	534.46	535.92	-1.46	27
3-methylpentane	-172.09	-171.47	-0.62	143.09	143.05	0.04				
3-methylhexane	-191.33	-192.10	0.77	165.98	165.94	0.04				
3-methylheptane	-212.51	-212.73	0.22	188.87	188.83	0.04	461.58	463.36 ^a	-1.78	27
3-methyloctane				212.59	211.72	0.87				
3-methylnonane				237.61	234.61	3.00				
4-methylheptane	-211.96	-212.73	0.77	188.87	188.83	0.04	453.34	451.83	1.51	54
4-methyloctane				212.59	211.72	0.87				
4-methylnonane				237.61	234.61	3.00				
5-methylnonane				237.61	234.61	3.00				
3-ethylpentane	-189.62	-189.84	0.22	165.98	165.94	0.04	411.50	412.67	-1.17	54
3-ethylhexane	-210.71	-210.47	-0.24	188.87	188.83	0.04	458.19	457.60	0.59	27
3-ethylheptane				208.11	211.72	-3.61				
3-ethyloctane				233.13	234.61	-1.48				

TABLE 6 Substituted alkane hydrocarbons (tertiary carbon) in the gas state 298.15 K — Continued

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)			N (tot)
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
4-ethylheptane				208.11	211.72	-3.61				
4-ethyloctane				233.13	234.61	-1.48				
4-propylheptane				233.13	234.61	-1.48				
4-isopropylheptane				231.00	234.64	-3.64				
2,4-dimethylpentane	-201.71	-201.05	-0.66	165.98	165.97	0.01	396.64	398.94	-2.30	162
2,4-dimethylhexane	-219.24	-219.42	0.18	188.87	188.86	0.01	445.64	449.63 ^a	-3.99	81
2,5-dimethylhexane	-222.51	-221.68	-0.83	188.87	188.86	0.01	439.03	438.10	0.93	162
2,3-dimethylbutane	-178.28	-180.42	2.14	140.54	143.08	-2.54	365.77	359.78	5.99	162
2,3-dimethylpentane	-198.87	-198.79	-0.08	165.98	165.97	0.01	414.05	410.47 ^a	3.58	81
2,3-dimethylhexane	-213.80	-219.42	5.62	188.87	188.86	0.01	443.96	443.86	0.10	81
3,4-dimethylhexane	-212.84	-217.16	4.32	188.87	188.86	0.01	448.32	449.63 ^a	-1.31	81
3-ethyl-2-methylpentane	-211.04	-217.16	6.12	188.87	188.86	0.01	441.12	443.86	-2.74	81
2,3,4-trimethylpentane	-217.32	-226.11	8.79	188.87	188.89	-0.02	428.07	430.13	-2.06	243

^aIncludes correction to entropy for optical isomerism, $R \ln 2$.

TABLE 7 Substituted alkane hydrocarbons (tertiary carbon) in the liquid state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)			
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
2-methylpropane	-153.47	-154.14	0.67							
2-methylbutane	-178.57	-177.69	-0.88							
2-methylpentane	-204.64	-203.42	-1.22	193.72	191.66	2.06	290.58	290.77	-0.19	
2-methylhexane	-229.49	-229.15	-0.34	222.92	222.08	0.84	323.34	323.15	0.19	
2-methylheptane	-255.01	-254.88	-0.13	252.00	252.50	-0.50	356.39	355.53	0.86	
2-methylnonane				313.30	313.34	-0.04	420.07	420.29	0.22	
2-methyldodecane				341.21	343.76	-2.55	453.80	452.67	1.13	
3-methylpentane	-202.38	-201.24	-1.14	190.66	191.66	-1.00				
3-methylhexane	-226.44	-226.97	0.53	214.51	222.08	-7.57	327.31	323.15	4.16	
3-methylheptane	252.34	252.70	0.36	250.20	252.50	-2.30				
3-methylnonane				308.99	313.34	-4.35	427.19	420.29	6.90	
4-methylheptane	-251.63	-252.70	1.07	251.09	252.50	-1.41				
4-methylnonane				317.36	313.34	4.02	425.51	420.29	5.00	
5-methylnonane				314.43	313.34	1.09	423.84	420.29	5.00	
3-ethylpentane	-224.89	-224.79	-0.10	219.58	222.08	2.50	314.55	313.12	6.00	
3-ethylhexane	-250.41	-250.52	0.11							
2,4-dimethylpentane	-234.60	-234.43	-0.17	224.22	219.10	5.11	303.11	300.11	4.00	
2,4-dimethylhexane	-257.02	-257.98	0.96							
2,5-dimethylhexane	-260.37	-260.16	-0.21	249.20	249.52	0.31				
2,3-dimethylbutane	-207.40	-208.70	1.30							
2,3-dimethylpentane	-233.09	-232.25	0.84	215.89	219.10	3.11				
2,3-dimethylhexane	-252.59	-257.98	5.39							
2,7-dimethyloctane				301.61	300.36	1.09				
3,4-dimethylhexane	-251.83	255.80	3.97							
3-ethyl-2-methylpentane	249.58	255.80	6.11							
2,3,4-trimethylpentane	255.01	263.26	8.25	216.13	216.54	0.31	309.13	314.13	1.01	

TABLE 8. Substituted alkane hydrocarbons (tertiary carbon) in the solid state at 198.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
2-methylpropane	-157.99	-153.22	-4.77						
2-methylbutane	-183.72	-180.29	-3.43						
2-methylpentane	-210.92	-209.70	-1.22						
2-methylhexane	-238.66	-239.11	0.45						
2-methylheptane	-266.94	-268.52	1.58						
3-methylpentane	-207.69	-207.36	-0.33						
3-methylhexane	-234.93	-236.77	1.84						
3-methylheptane	-264.01	-266.18	2.17						
3-ethylpentane	-234.43	-234.43	0.00						
4-methylheptane	-262.46	-266.18	3.72						
2,4-dimethylpentane	-241.46	-237.69	-3.77						

TABLE 9. Substituted alkane hydrocarbons (quaternary carbon) in the gas state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)			N(tot)
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
2,2-dimethylpropane	-167.99	-168.08	0.09	121.63	119.45	2.18	306.39	302.59	3.80	972
2,2-dimethylbutane	-186.10	-184.15	-1.95	141.88	142.34	-0.46	358.23	353.28	4.95	243
2,2-dimethylpentane	-205.85	-204.78	-1.07	165.98	165.23	0.75	392.88	392.44	0.44	243
2,2-dimethylhexane	-224.60	-225.41	0.81	188.87	188.12	0.75	431.20	431.60	-0.40	243
3,3-dimethylpentane	-201.17	-200.22	-0.95	165.98	165.23	0.75	399.70	395.81	3.89	162
3,3-dimethylhexane	-219.99	-220.85	0.86	188.87	188.12	0.75	438.06	440.73	-2.67	81
2,2,3-trimethylbutane	-204.47	-202.27	-2.20	164.56	165.26	-0.70	383.60	378.70	4.90	729
2,2,3-trimethylpentane	-219.99	-221.10	1.11	188.87	188.15	0.72	425.18	417.86	7.32	729
2,2,4-trimethylpentane	-224.01	-222.90	-1.11	188.87	188.15	0.72	423.21	417.86	5.35	729
2,3,3-trimethylpentane	-216.27	-221.10	4.83	188.87	188.15	0.72	431.54	427.00	4.54	243
2,2,3,3-tetramethylbutane	-225.60	-219.00	-6.60	192.59	187.44	5.15	389.36	386.10	3.26	13,122
2,2,3,3-tetramethylpentane	-236.69	-238.99	2.30	212.09	210.33	1.76				
2,2,4,4-tetramethylpentane	-241.63	-239.63	-2.00	211.63	210.33	1.30				
2,2,3,4,4-pentamethylpentane				234.43	233.25	1.18				
3-ethyl-3-methylpentane	-214.85	-216.29	1.44	188.87	188.12	0.75	432.96	431.60	1.36	243
3,3-diethylpentane	-232.34	-232.36	0.02	204.18	211.01	-6.83	461.54	459.23	2.31	972

TABLE 10. Substituted alkane hydrocarbons (quaternary carbon) in the liquid state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
2,2-dimethylpropane	-190.33	-190.01	-0.32						
2,2-dimethylbutane	-213.80	-211.35	-2.45	183.18	186.58	-3.40	272.00	266.93	5.07
2,2-dimethylpentane	-238.28	-237.08	-1.20	221.12	217.00	4.12	300.29	299.31	0.98
2,2-dimethylhexane	-261.88	-262.81	0.93						
3,3-dimethylpentane	-234.18	-232.69	-1.49	211.71	217.00	-5.29	293.30	299.31	-6.01
3,3-dimethylhexane	-257.94	-258.42	0.48	246.60	247.42	-0.82			
2,2,3-trimethylbutane	-236.52	-233.68	-2.84	213.51	214.02	-0.51	292.25	293.96	-1.71
2,2,3-trimethylpentane	-256.90	-257.94	0.74						
2,2,4-trimethylpentane	-259.16	-259.41	0.25	238.57	244.44	-5.87	328.03	326.34	1.69
2,3,3-trimethylpentane	-253.51	-257.64	4.13	245.56	244.44	1.14			

TABLE 10 Substituted alkane hydrocarbons (quaternary carbon) in the liquid state at 298.15 K — Continued

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
2,2,3,3-tetramethyl-									
pentane	-278.28	-278.61	0.33						
2,2,4,4-tetramethyl-									
pentane	-279.99	-279.25	-0.74						
3-ethyl-3-methylpentane	-252.84	-254.03	1.19						
3,3-diethylpentane	-275.39	-275.37	-0.02						

TABLE 11. Substituted alkane hydrocarbons (quaternary carbon) in the solid state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
2,2-dimethylpropane									
2,2-dimethylbutane	-196.15	-191.89	-4.26						
2,2-dimethylpentane	-220.08	-216.95	-3.13						
3,3-dimethylpentane	-244.14	-246.36	2.22						
2,2,3-trimethylbutane	-241.25	-242.01	0.76						
3-ethyl-3-methylpentane	-241.08	-240.71	-0.37						
3-ethyl-3-methylpentane	-267.48	-267.07	-0.41						
3,3-diethylpentane	-292.13	-292.13	0.00						
2,2,3,3-tetramethyl-									
butane	-268.95	-268.94	-0.01	237.44	237.44	0.00	273.76	273.76	0.00

TABLE 12. *n*-Alkene hydrocarbons in the gas state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)			$N(\text{tot})$
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
ethylene										
propylene	52.47	52.64	-0.17	43.56	42.76	0.80	219.45	219.51	-0.06	4
1-butene	20.00	20.38	-0.38	63.89	65.85	-1.96	266.94	266.76	0.18	3
1-pentene	0.21	-0.50	0.71	85.65	86.48	-0.83	305.60	304.96	0.64	3
1-hexene	-21.59	-21.13	-0.46	109.58	109.37	0.21	345.81	344.12	1.69	3
1-heptene	-41.67	-41.76	0.09	132.34	132.26	0.08	384.64	383.28	1.36	3
1-octene	-62.30	-62.39	0.09	155.23	155.15	0.08	423.59	422.44	1.15	3
1-nonene	-82.93	-83.02	0.09	178.07	178.04	0.03	462.54	461.60	0.94	3
1-decene	-103.51	-103.65	0.14	200.96	200.93	0.03	501.49	500.76	0.73	3
1-hexadecene	-124.14	-124.28	0.14	223.80	223.82	-0.02	540.45	539.92	0.53	3
cis-2-butene	-247.82	-248.06	0.24	361.04	361.16	-0.12	774.12	774.88	0.76	3
trans-2-butene	-7.07	-7.03	-0.04	78.91	80.91	2.00	300.83	301.77	0.94	18
cis-2-pentene	-11.51	-11.88	0.37	87.82	88.94	1.12	296.48	296.71	0.23	18
trans-2-pentene	-28.49	-27.91	-0.58	101.75	101.54	0.21	346.27	345.73	0.54	9
cis-2-hexene	-32.38	-32.76	0.38	108.45	109.57	1.12	340.41	340.67	0.26	9
trans-2-hexene	-52.34	-48.54	-3.80	125.69	124.43	1.26	386.48	384.89	1.59	9
trans-2-hexene	-53.89	53.39	0.50	132.38	132.46	0.08	380.62	379.83	0.79	9
cis-3-hexene	-47.61	-48.79	1.18	123.64	122.17	1.47	379.61	378.17	1.44	18
trans-3-hexene	-54.43	53.64	0.79	132.84	130.20	2.64	374.84	373.11	1.73	18
cis-2-heptene	-69.16	-69.17	0.01							
trans-2-heptene	-73.55	-74.02	0.47							

TABLE 12. *n*-Alkene hydrocarbons in the gas state at 298.15 K. — Continued

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)			$N(\text{tot})$
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
cis-3-heptene	-68.78	-69.42	0.64							
trans-3-heptene	-73.76	-74.27	0.51							
1,2-butadiene	162.26	163.05	-0.79	80.12	81.71	-1.59	293.01	293.04	-0.03	3
1,2-pentadiene	140.71	142.17	-1.46	105.44	102.34	3.10	333.46	331.24	2.22	3
1,3-butadiene	109.96	109.20	0.76	79.54	79.84	-0.30	278.74	280.76	-2.02	2
cis-1,3-pentadiene	81.25	81.79	-0.54	94.56	94.90	-0.34	324.26	327.30	-3.04	3
trans-1,3-pentadiene	76.07	76.94	-0.87	103.34	102.93	0.41	319.66	322.24	-2.58	3
1,4-pentadiene	105.48	105.49	-0.01	105.02	105.01	0.01	333.46	333.46	0.00	2
1,5-hexadiene	84.06	83.52	0.54							
2,3-pentadiene	133.05	130.79	2.26	101.25	104.80	-3.55	324.68	322.99	1.69	18
allene	190.46	195.31	-4.85	59.25	58.62	0.63	243.93	245.79	-1.86	4

TABLE 13. *n*-Alkene hydrocarbons in the liquid state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)			
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
ethylene		43.50			56.74			172.38		
propylene		5.19			89.45			198.07		
1-butene	-20.38	-20.54	0.16	121.85 ^a	118.74	3.11	227.63 ^b	229.74	-2.11	
1-pentene	-47.03	-46.27	-0.76	155.31	149.16	6.15	262.55	262.12	0.43	
1-hexene	-72.38	-72.00	-0.38	183.30	179.58	3.72	295.18	294.50	0.68	
1-heptene	-98.03	-97.73	-0.30	211.79	210.00	1.79	327.65	326.88	0.77	
1-octene	-123.51	-123.46	-0.05	241.21	240.42	0.79	360.45	359.26	1.19	
1-nonene	-149.03	-149.19	0.16							
1-decene	-174.60	-174.92	0.32	300.33	301.26	-0.93	425.01	424.02	0.99	
1-hexadecene	-328.70	-329.30	0.60							
cis-2-butene	-29.66	-27.85	-1.81	126.15	122.16	3.99	219.91	223.76	-3.85	
trans-2-butene	-33.10	-33.12	0.02	125.90 ^c	122.16	3.74	221.25 ^c	223.76	-2.51	
cis-2-pentene	-54.52	-53.58	-0.94	151.71	151.45	0.26	258.61	255.43	3.18	
trans-2-pentene	-58.83	-58.85	0.02	156.98	151.45	5.53	256.52	255.43	1.09	
cis-2-hexene	-83.89	-79.31	-4.58							
trans-2-hexene	-85.52	-84.58	-0.94							
cis-2-heptene	-105.14	-105.04	-0.10							
trans-2-heptene	-109.54	-110.31	0.77							
cis-2-octene	-135.69	-130.77	-4.92							
trans-2-octene	-135.69	-136.04	0.35							
cis-3-hexene	-78.95	-79.31	0.36							
trans-3-hexene	-86.06	-84.58	-1.48							
cis-3-heptene	-104.31	-105.04	0.73							
trans-3-heptene	-109.33	-110.31	0.98							
1,2-butadiene	138.99	139.87	-0.88							
1,2-pentadiene	150.83	148.78	2.05	244.97	244.13	0.84				
1,3-butadiene	87.91	87.86	0.05	123.65	120.08	3.57	198.99	198.98	0.01	
1,4-pentadiene	80.33	80.33	0.00	146.82	146.82	0.00	248.86	248.86	0.00	
2,3-pentadiene	152.42	152.20	0.22	237.32	238.15	-0.83				
1,5-hexadiene	54.10	54.14	-0.04							

^aObtained from extrapolation of experimental data of 46AST/FIN to 298.15 K.^bCalculated from table XI in 46AST/FIN.^cObtained from extrapolation of experimental data of 45GUT/PIT to 298.15 K.

TABLE 14. *n*-Alkene hydrocarbons in the solid state at 298.15

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
propylene		1.17							
1-butene	-24.23	-23.18	-1.05						
1-pentene	-52.84	-52.59	-0.25						
1-hexene	-82.13	-82.00	-0.13						
1-heptene	-110.58	-111.41	0.83						
1-octene	-138.83	-140.82	1.99						
1-decene	-196.36	-199.64	3.28						
1-hexadecene	-358.90	-376.10	17.20						
cis-2-butene	-36.99	-36.79	-0.20						
trans-2-butene	-42.84	-42.52	-0.32						
cis-2-pentene	-61.63	-61.14	-0.49						
trans-2-pentene	-67.20	-66.87	-0.33						
2-hexene ^a	-97.78	-96.28	-1.50						
2-heptene ^a	-124.68	-125.69	1.01						
2-octene ^a	-153.64	-155.10	1.46						
1,2-butadiene	132.05	132.25	-0.20						
1,3-butadiene	79.91	79.92	-0.01						
1,4-pentadiene	74.22	74.22	0.00						

^aAssumed to be the *trans* isomer.

TABLE 15. Substituted alkene hydrocarbons in the gas state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)			N(tot)
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
2-methylpropene	-16.90	-18.58	1.68	89.12	87.94	1.18	293.59	295.29	-1.70	18
2-methyl-1-butene	-35.31	-37.20	1.89	111.63	108.57	3.06	339.53	339.25	0.28	9
2-methyl-1-pentene	-59.37	-57.83	-1.54	135.60	131.46	4.14	382.17	378.41	3.76	9
2-methyl-2-butene				105.02	111.03	-6.01	338.57	345.90	-7.33	9
2-methyl-2-pentene				126.61	131.66	-5.05	378.44	384.10	-5.66	9
2-ethyl-1-butene	-56.02	-55.82	-0.20	133.55	129.20	4.35	376.60	371.69	4.91	18
3-methyl-1-butene	-27.45	-28.03	0.58	118.62	119.07	-0.45	333.46	334.56	-1.10	9
3-methyl-1-pentene	-49.50	-48.66	-0.84	142.42	141.96	0.46	376.81	373.72	3.09	9
cis-3-methyl-2-pentene	-62.30	-64.61	2.31	126.61	123.63	2.98	378.44	380.03	1.59	11
trans-3-methyl-2-pentene	-63.14	-69.46	6.32	126.61	131.66	-5.05	381.83	374.97	6.86	11
3-methyl-cis-3-hexene	-79.41	-85.49	6.08							
3-methyl-trans-3-hexene	-76.82	-90.34	13.52							
4-methyl-1-pentene				126.48	132.29	-5.81	367.73	369.54	1.81	
cis-4-methyl-2-pentene	-57.49	-55.44	-2.05	133.55	134.13	-0.58	373.34	375.34	1.00	
trans-4-methyl-2-pentene	-61.50	-60.29	-1.21	141.42	142.16	-0.74	368.28	370.28	1.00	
<i>cis</i> -2,2-dimethyl-										
3-hexene	-89.29	-91.59	2.30							
<i>trans</i> -2,2-dimethyl-										
3-hexene	-107.65	-108.83	1.18							
2,3-dimethyl-1-butene	-62.38	-64.73	2.35	143.47	141.16	3.31	367.64	369.57	1.00	
2,3-dimethyl-2-butene				123.60	133.12	9.53	367.64	370.57	1.00	
2,4-dimethyl-1-pentene	-83.81	-85.15	1.34							
2,4-dimethyl-2-pentene	-88.70	-99.25	10.55							
3,3-dimethyl-1-butene	-60.67	-55.69	-4.98	126.48	126.48	0.00	374.60	375.60	1.00	
<i>cis</i> -4,4-dimethyl-										
2-pentene	-72.63	-70.71	1.92							
<i>trans</i> -4,4-dimethyl-										
2-pentene	-88.78	87.95	0.83							
2,3,3-trimethyl-										
1-butene	85.48	92.39	6.91							

TABLE 15. Substituted alkene hydrocarbons in the gas state at 298.15 K — Continued

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)			N(tot)
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
2,4,4-trimethyl-1-pentene	-110.37	-116.20	5.83							
2,4,4-trimethyl-2-pentene	-104.89	-126.91	22.02							
2-methyl-3-ethyl-1-pentene	-100.29	-101.47	1.18							
3-methyl-2-ethyl-1-butene	-79.54	-83.35	3.81							
2-methyl-1,3-butadiene	75.44	73.18	2.26	104.60	104.60	0.00	315.64	315.64	0.00	3
3-methyl-1,2-butadiene				105.44	103.80	1.64	319.66	321.57	-1.91	18

TABLE 16. Substituted alkene hydrocarbons in the liquid state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)			
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
2-methylpropene		-34.31			124.55					
2-methyl-1-butene	-61.13	-60.04	1.09	157.19	153.84	3.35	253.97	254.63	-0.66	
2-methyl-1-pentene	-89.96	-87.95	-2.01							
2-methyl-2-butene	-68.58	-76.98	8.40	152.80	157.26	-4.46	251.04	248.65	2.39	
2-ethyl-1-butene	-87.11	-85.77	-1.34							
3-methyl-1-butene	-51.38	-51.80	0.42	156.06	156.05	0.01	253.30	253.30	0.00	
3-methyl-1-pentene	-78.16	-77.53	-0.63							
cis-3-methyl-2-pentene	-94.47	-95.26	0.79							
trans-3-methyl-2-pentene	-94.56	-100.53	5.97							
cis-4-methyl-2-pentene	-87.03	-84.84	-2.19							
trans-4-methyl-2-pentene	-91.55	-90.11	-1.44							
3-methyl-cis-3-hexene	-115.94	-120.99	5.05							
3-methyl-trans-3-hexene	-112.72	-126.26	13.54							
cis-2,2-dimethyl-3-hexene	-126.44	-128.97	2.53							
trans-2,2-dimethyl-3-hexene	-144.93	-146.45	1.52							
2,3-dimethyl-1-butene	-92.63	93.48	0.85							
2,3-dimethyl-2-butene	-101.46	-120.84	19.38	174.68	192.36	-17.68	270.20	273.54	-3.34	
2,4-dimethyl-1-pentene	-116.98	-118.96	1.98							
2,4-dimethyl-2-pentene	-123.09	-133.97	10.88							
3,3-dimethyl-1-butene	-87.40	-82.41	-4.99	191.17	191.15	0.02	256.50	256.47	0.03	
cis-4,4-dimethyl-2-pentene	-105.31	-103.24	-2.07							
trans-4,4-dimethyl-2-pentene	-121.71	-120.72	-0.99							
cis-2,5-dimethyl-3-hexene	-151.04	-141.83	-9.21							
trans-2,5-dimethyl-3-hexene	-159.24	-147.10	-12.14							
2,3,3-trimethyl-1-butene	-117.70	-124.09	6.39							
2,4,4-trimethyl-1-pentene	-146.15	-152.62	6.47	240.20	237.04	3.16	311.71	322.58	-10.87	
2,4,4-trimethyl-2-pentene	-142.42	-164.58	22.16							
2-methyl-3-ethyl-1-pentene	-136.36	-140.58	4.22							
3-methyl-2-ethyl-1-butene	-114.06	-117.03	2.97							

TABLE 16. Substituted alkene hydrocarbons in the liquid state at 298.15 K — Continued

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
2-methyl-1,3-butadiene	48.49	46.31	2.18	151.08	151.08	0.00	228.28	227.06	1.22
3-methyl-1,2-butadiene	101.17	96.01	5.16	152.42	154.59	-2.17	231.79	237.35	-5.56
2,3-dimethyl-1,3-butadiene	14.14	4.76	9.38						

TABLE 17. Substituted alkene hydrocarbons in the solid state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
2-methylpropene									
2-methyl-1-butene	-69.04	-64.77	-4.27						
2-methyl-2-butene	-76.44	-86.45	10.01						
3-methyl-1-butene	-56.74	-56.74	0.00						
2,3-dimethyl-2-butene	-111.13	-130.38	19.25						
3,3-dimethyl-1-butene	-92.84	-92.85	0.01						
2,4,4-trimethyl-1-pentene	-154.89	-158.83	3.94						
2,4,4-trimethyl-2-pentene	-149.16	-180.47	31.31						
2-methyl-1,3-butadiene	43.56	41.22	2.34						
3-methyl-1,2-butadiene	93.22	88.32	4.90						

TABLE 18. Alkyne hydrocarbons in the gas state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)			$N(\text{tot})$
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
acetylene	228.24	227.00	1.24	43.93	45.10	-1.17	200.83	198.16	2.67	2
propyne	184.93	186.34	-1.41	60.67	61.50	-0.83	248.11	246.47	1.64	3
1-hutyne	165.23	166.64	-1.41	81.42	82.47	-1.05	290.83	289.27	1.56	3
1-pentyne				106.69	105.36	1.33	329.78	328.43	1.35	3
1-hexyne				128.24	128.25	-0.01	368.74	367.59	1.15	3
1-heptyne				151.08	151.14	-0.06	407.69	406.75	0.94	3
1-octyne				173.97	174.03	-0.06	446.64	445.91	0.73	3
1-nonyne				196.82	196.92	-0.10	485.60	485.07	0.53	3
1-decyne				219.70	219.81	-0.11	524.51	524.23	0.28	3
1-hexadecyne				356.94	357.15	-0.21	758.22	759.19	-0.97	3
2-butyne	145.69	145.68	0.01	77.95	77.90	0.05	283.30	283.25	0.05	18
2-pentyne	128.87	125.98	2.89	98.70	98.87	-0.17	331.79	331.81	-0.02	9
3-methyl-1-butyne	136.40	136.40	0.00	104.68	104.68	0.00	318.95	318.96	-0.01	9
1-buten-3-yne				73.18	73.18	0.00	279.37	279.38	-0.01	2
cis-3-penten-1-yne	253.55	262.11	-8.56							
trans-3-penten-1-yne	254.89	257.26	-2.37							
cis-3-decen-1-yne	151.67	158.71	-7.04							
trans-3-decen-1-yne	154.60	153.86	0.74							
butadiyne				73.64	73.64	0.00	250.04	250.04	0.00	2
1,5-hexadiyne	416.06	416.06	0.00							

TABLE 19. Alkyne hydrocarbons in the liquid state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
1-butyne	141.88	141.88	0.00	132.42	132.42	0.00	197.48	197.48	0.00
2-butyne	119.08	119.08	0.00	124.14	124.14	0.00	195.10	195.10	0.00
cis-3-penten-1-yne	226.48	230.13	-3.65						
trans-3-penten-1-yne	228.15	224.86	3.29						
1-octene-3-yne	140.71	144.65	-3.94						
cis-3-decen-1-yne	99.20	101.48	-2.28						
trans-3-decen-1-yne	100.46	96.21	4.25						
1,5-hexadiyne	384.17	384.16	0.01						
1,7-octadiyne	334.43	327.52	6.91						
3,9-dodecadiyne	197.82	193.40	4.42						
5,7-dodecadiyne	181.50	181.50	0.00						
3,3-dimethyl-1-butyne	78.45	78.45	0.00						
3,3-dimethylpenta-1,4-diyne	348.69	348.69	0.00						
3,3,6,6-tetramethyl-octa-1,7-diyne	211.08	209.44	1.64						

TABLE 20. Alkyne hydrocarbons in the solid state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
1-butyne	135.85	135.85	0.00						
2-butyne	109.85	109.84	0.01						

TABLE 21. Aromatic hydrocarbons in the gas state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)			$N(\text{tot})$
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
benzene	82.89	82.86	0.03	81.67	81.66	0.01	269.20	269.20	0.00	12
toluene	50.17	50.43	-0.26	103.64	103.53	0.11	320.66	318.36	2.30	6
1,2-dimethylbenzene (o-xylene)	19.08	19.26	-0.18	133.26	131.80	1.46	352.75	350.13	2.62	18
1,3-dimethylbenzene (m-xylene)	17.32	17.37	-0.05	127.57	126.11	1.46	357.69	352.63	5.06	18
1,4-dimethylbenzene (p-xylene)	18.03	18.00	0.03	126.86	125.40	1.46	352.42	352.63	-0.21	18
1,2,3-trimethylbenzene	-9.46	-12.54	3.08	154.18	160.78	-6.60	384.84	381.89	2.95	54
1,2,4-trimethylbenzene	-13.85	-13.80	-0.05	154.01	154.38	-0.37	395.76	390.16	5.60	27
1,3,5-trimethylbenzene	-15.94	-16.32	0.38	150.25	149.40	0.85	385.30	377.76	7.54	162
1,2,3,4-tetramethylbenzene				189.58	189.76	-0.18	416.52	413.66	2.86	162
1,2,3,5-tetramethylbenzene				185.73	183.36	2.37	422.54	416.16	6.38	162
1,2,4,5-tetramethylbenzene				186.52	183.36	3.16	418.53	410.40	8.13	324
pentamethylbenzene				216.48	219.45	-2.97	443.88	445.47	-1.54	486
hexamethylbenzene	-86.82	-107.31	20.49	248.61	254.83	-6.22	452.37	459.79	-7.42	8,748
ethylbenzene	29.92	29.09	0.83	128.41	129.14	-0.73	360.45	360.95	-0.50	6
propylbenzene	7.91	8.46	-0.55	152.34	152.03	0.31	400.66	400.11	0.55	6

TABLE 21. Aromatic hydrocarbons in the gas state at 298.15 K — Continued

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)			$N(\text{tot})$	
	expt	calc	resd	expt	calc	resd	expt	calc	resd		
butylbenzene	-13.05	-12.17	-0.88	175.10	174.92	0.18	439.49	439.27	0.22	6	
pentylbenzene				197.99	197.81	0.18	478.94	478.43	0.51	6	
hexylbenzene				220.87	220.70	0.17	517.90	517.59	0.31	6	
heptylbenzene				243.72	243.59	0.13	556.85	556.75	0.10	6	
octylbenzene				266.60	266.48	0.12	595.80	595.91	-0.11	6	
nonylbenzene				289.45	289.37	0.08	634.75	635.07	-0.32	6	
decylbenzene				312.34	312.26	0.08	673.71	674.23	-0.52	6	
undecylbenzene				335.22	335.15	0.07	712.62	713.39	-0.77	6	
dodecylbenzene				358.07	358.04	0.03	751.57	752.55	-0.98	6	
1-methyl-2-ethylbenzene	1.30	-2.08	3.38								
1-methyl-3-ethylbenzene	-1.80	-3.97	2.17								
1-methyl-4-ethylbenzene	-3.18	-3.34	0.16	151.54	151.01	0.53	398.90	395.22	3.68	18	
1,4-diethylbenzene				176.15	176.62	-0.47	434.01	437.81	-3.80	18	
1,2,3-triethylbenzene				228.11	237.61	-9.50	507.23	509.66	-2.43	54	
1,2,4-triethylbenzene				227.94	231.21	-3.27	518.15	517.93	0.22	27	
1,3,5-triethylbenzene				224.18	226.23	-2.05	507.69	505.53	-2.16	162	
pentaethylbenzene				339.70	347.50	-7.80	647.89	658.37	-10.48	486	
hexaethylbenzene				396.48	408.49	-12.01	697.14	715.33	-18.19	8,748	
isopropylbenzene	4.02	-0.87	4.89	151.71	151.71	0.00	388.57	388.55	0.02	18	
sec-butylbenzene	-17.36	-19.24	1.88								
isobutylbenzene	-21.51	-18.86	-2.65								
t-butylbenzene	-22.59	-22.59	0.00								
styrene	147.82	147.82	0.00	122.09	122.09	0.00	345.10	345.20	-0.10	2	
o-methylstyrene				145.18	150.36	-5.18	383.67	382.73	0.94	3	
m-methylstyrene				145.18	144.67	0.51	389.53	385.23	4.30	3	
p-methylstyrene				145.18	143.96	1.22	383.67	379.46	4.21	6	
1-isopropenylbenzene											
(α -methylstyrene)	112.97	112.97	0.00	145.18	145.18	0.00	383.67	383.67	0.00	6	
cis-1-propenylbenzene					145.18	137.15	8.03	383.67	385.97	-2.30	6
(cis- β -methylstyrene)											
trans-1-propenylbenzene					146.02	145.18	0.84	380.33	380.91	-0.58	6
(trans- β -methylstyrene)					114.89	114.89	0.00	321.67	321.67	0.00	2
ethynylbenzene	327.65	327.48	0.17								
1,1-diphenylethylene	245.64	245.64	0.00								
cis-stilbene	252.34	247.85	4.49								
trans-stilbene	236.14	243.00	-6.86								
diphenylmethane	138.95	138.95	0.00								
1,2-diphenylethane	142.93	142.70	0.23								
biphenyl	181.42	181.42	0.00	162.34	162.34	0.00	392.67	392.67	0.00	8	
naphthalene	150.29	150.68	-0.39	132.55	132.54	0.01	335.64	335.63	0.01	4	
1-methylnaphthalene	113.51	118.25	-4.74	159.54	159.53	0.01	377.44	377.75	-0.31	3	
2-methylnaphthalene	110.54	118.25	-7.71	159.79	159.53	0.26	380.03	377.75	2.28	3	
1-ethylnaphthalene					184.18	185.14	-0.96	418.15	420.34	-2.19	3
2-ethylnaphthalene					184.43	185.14	-0.71	420.74	420.34	0.40	3
1-propylnaphthalene					208.11	208.03	0.08	458.36	459.50	-1.14	3
2-propylnaphthalene					208.36	208.03	0.33	460.99	459.50	1.49	3
1-butylnaphthalene					230.87	230.92	-0.05	497.18	498.66	-1.48	3
2-butylnaphthalene					231.12	230.92	0.20	499.82	498.66	1.16	3
1-pentylnaphthalene					253.76	253.81	-0.05	536.64	537.82	-1.18	3
2-pentylnaphthalene					254.01	253.81	0.20	539.28	537.82	1.46	3
1,2-dimethylnaphthalene					184.85	185.58	-0.73	406.81	409.01	-2.20	9
1,3-dimethylnaphthalene					185.10	185.58	-0.48	409.45	409.01	0.44	9
1,4-dimethylnaphthalene					184.85	185.58	-0.73	401.08	403.25	-2.17	18

TABLE 21. Aromatic hydrocarbons in the gas state at 298.15 K — Continued

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)			$N(\text{tot})$
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
1,5-dimethylnaphthalene				184.85	185.58	-0.73	401.08	409.01	-7.93	9
1,6-dimethylnaphthalene				185.10	185.58	-0.48	409.45	409.01	0.44	9
1,7-dimethylnaphthalene				185.10	185.58	-0.48	409.45	409.01	0.44	9
1,8-dimethylnaphthalene	108.66	85.82	22.84	185.81	185.58	0.23	410.95	403.25	7.70	18
2,3-dimethylnaphthalene										
2,6-dimethylnaphthalene				187.07	185.58	1.49	408.69	409.01	-0.32	9
2,7-dimethylnaphthalene				187.07	185.58	1.49	408.69	403.25	5.44	18
2-ethyl-3-methyl-naphthalene				210.46	211.19	-0.73	457.44	451.60	5.84	9
2-ethyl-6-methyl-naphthalene				211.71	211.19	0.52	455.18	451.60	3.58	9
2-ethyl-7-methyl-naphthalene				211.71	211.19	0.52	455.18	451.60	3.58	9
anthracene	230.91	218.50	12.41							
naphthacene	291.42	286.32	5.10							
phenanthrene	207.48	210.30	-2.82							
triphenylene	274.18	269.92	4.26							
chrysene	269.78	269.92	-0.14							
pyrene	225.68	225.68	0.00							
1,2-benzanthracene	294.14	278.12	16.02							

TABLE 22. Aromatic hydrocarbons in the liquid state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)			
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
benzene	48.95	48.96	-0.01	136.06	136.08	-0.02	173.26	173.22	0.04	
toluene	12.38	12.35	0.03	157.23	159.98	-2.75	220.96	208.15	12.81	
1,2-dimethylbenzene (o-xylene)	-24.35	-21.00	-3.35	187.82	187.38	0.44	246.02	243.08	2.94	
1,3-dimethylbenzene (m-xylene)	-25.36	-24.26	-1.10	183.18	183.88	-0.70	253.80	243.08	10.72	
1,4-dimethylbenzene (p-xylene)	-24.35	-24.26	-0.09	183.76	183.88	-0.12	243.51	243.08	0.43	
1,2,3-trimethylbenzene	-58.53	-54.35	-4.18	216.44	214.78	1.66	267.94	278.01	-10.07	
1,2,4-trimethylbenzene	-61.80	-57.61	-4.19	214.97	211.28	3.69	283.38	278.01	5.37	
1,3,5-trimethylbenzene	-63.43	-60.87	-2.56	209.53	207.78	1.75	273.55	278.01	-4.46	
1,2,3,4-tetramethylbenzene	-90.21	-87.70	-2.51	235.98	242.18	-6.20	290.79	312.94	-22.15	
1,2,3,5-tetramethylbenzene	-96.36	-90.96	-5.40	240.16	238.68	1.48	310.03	312.94	-2.91	
1,2,4,5-tetramethylbenzene	-98.99	-90.96	-8.03							
pentamethylbenzene	-122.97	-121.05	-1.92							
hexamethylbenzene	-139.14	-151.14	12.00							
ethylbenzene	-12.34	-12.46	0.12	185.81	182.88	2.93	255.01	255.55	-0.54	
propylbenzene	-38.33	-38.19	-0.14	214.72	213.30	1.42	287.78	287.93	-0.15	
butylbenzene	-63.18	-63.92	0.74	243.34	243.72	-0.38	321.21	320.31	0.90	
1-methyl-2-ethylbenzene	-46.40	-45.81	-0.59							
1-methyl-3-ethylbenzene	-48.70	-49.07	0.37							
1-methyl-4-ethylbenzene	-49.79	-49.07	-0.72							
1-methyl-2-propylbenzene	-72.47	-71.54	-0.93							

TABLE 22. Aromatic hydrocarbons in the liquid state at 298.15 K — Continued

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
1-methyl-3-propylbenzene	-76.23	-74.80	-1.43						
1-methyl-4-propylbenzene	-75.06	-74.80	-0.26						
1-methyl-									
2-isopropylbenzene	-73.30	-78.79	5.49						
1-methyl-									
3-isopropylbenzene	-78.62	-82.05	3.43						
1-methyl-									
4-isopropylbenzene	-78.03	-82.05	4.02	236.40	237.86	-1.46	306.69	312.48	-5.79
3-ethyl-1,2-dimethylbenzene	-80.50	-79.16	-1.34						
4-ethyl-1,2-dimethylbenzene	-86.02	-82.42	-3.60						
2-ethyl-1,3-dimethylbenzene	-80.12	-79.16	-0.96						
4-ethyl-1,3-dimethylbenzene	-84.10	-82.42	-1.68						
5-ethyl-1,3-dimethylbenzene	-87.78	-85.68	-2.10						
2-ethyl-1,4-dimethylbenzene	-84.81	-82.42	-2.39						
1,2-diethylbenzene	-68.49	-70.62	2.13						
1,3-diethylbenzene	-73.51	-73.88	0.37						
1,4-diethylbenzene	-72.84	-73.88	1.04						
isopropylbenzene	-41.13	-45.44	4.31	215.40	213.96	1.44	277.57	277.55	0.02
(1-methylpropyl)benzene	-66.40	-68.99	2.59						
(2-methylpropyl)benzene	-69.79	-69.20	-0.59						
t-butylbenzene	-70.71	-70.71	0.00	238.11	238.11	0.00	278.65	278.65	0.00
styrene	103.85	103.85	0.00	182.88	182.90	-0.02	237.57	234.80	2.77
isopropenylbenzene (α -methylstyrene)	70.46	70.46	0.00						
2-propenylbenzene (allylbenzene)	88.03	88.03	0.00						
1-methyl-2-propenylbenzene	56.07	56.07	0.00						
ethynylbenzene	283.47	283.39	0.08						
diphenylmethane	89.66	93.42	-3.76	279.91	279.91	0.00	301.67	301.67	0.00
4-methyldiphenylmethane	61.55	56.81	4.74						
2,5-dimethyl-diphenylmethane	24.69	23.46	1.23						
1,1-diphenylethane	48.66	48.66	0.00	294.97	294.98	-0.01			
1,1-diphenyldodecane	593.71	599.18	-5.47	684.92	684.92	0.00			
cis-stilbene	183.30	169.47	13.83						
1,1-diphenylethylene	172.42	172.42	0.00	299.16	299.15	0.01			
biphenyl	116.02	116.02	0.00						
2-methylbiphenyl	108.16	82.67	25.49						
3-methylbiphenyl	85.56	79.41	6.15						
isopropylbiphenyl				338.49	342.32	1.81			
naphthalene	96.94	96.94	0.00						
1-methylnaphthalene	56.19	60.33	-4.14	224.39	224.31	-0.01			
2-methylnaphthalene	62.58	60.33	2.25						
1,8-dimethylnaphthalene	41.76	23.72	18.04						
anthracene	158.57	144.92	13.65						
phenanthrene	132.87	136.26	3.49						
triphenylene	176.52	175.88	0.94						
pyrene	143.13	143.13	0.01						

TABLE 23. Aromatic hydrocarbons in the solid state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
benzene	39.08	39.18	-0.10	120.79	120.78	0.01	136.48	136.50	-0.02
toluene	5.73	-0.19	5.92						
1,2-dimethylbenzene (o-xylene)	-37.95	-34.56	-3.39						
1,3-dimethylbenzene (m-xylene)	-36.90	-37.56	0.66						
1,4-dimethylbenzene (p-xylene)	-41.46	-39.56	-1.90						
1,2,3-trimethylbenzene	-68.70	-66.93	-1.77						
1,2,4-trimethylbenzene	-74.98	-71.93	-3.05						
1,3,5-trimethylbenzene	-72.93	-72.93	0.00						
1,2,3,4-tetramethylbenzene	-101.42	-99.30	-2.12						
1,2,3,5-tetramethylbenzene	-109.29	-104.30	-4.99						
1,2,4,5-tetramethylbenzene	-119.87	-104.30	-15.57	220.08	217.02	3.06	245.60	250.26	-4.66
pentamethylbenzene	-133.64	-129.67	-3.97	270.29	241.08	29.21	294.14	278.70	15.44
hexamethylbenzene	-161.54	-157.04	-4.50	245.64	265.14	-19.50	306.31	307.14	-0.83
ethylbenzene	-21.51	-22.29	0.78						
propylbenzene	-56.11	-51.70	-4.41						
butylbenzene	-74.39	-81.11	6.72						
1-methyl-4-isopropylbenzene	-87.70	-94.48	6.78						
isopropylbenzene	-48.45	-55.11	6.66						
t-butylbenzene	-79.12	-79.12	0.00						
styrene	92.88	92.88	0.00						
diphenylmethane	71.46	71.66	-0.20	223.84	223.84	0.00	239.32	239.35	-0.03
trans-stilbene	136.94	140.90	-3.96	232.63	232.60	0.03	251.04	251.00	0.04
1,2-diphenylethane	51.55	48.90	2.65	253.55	253.54	0.01	270.29	270.30	-0.01
triphenylmethane				295.81	295.81	0.00	312.13	312.13	0.00
tetraphenylmethane				368.19	368.30	-0.11			
1,1,2-triphenylethane				319.66	325.10	-5.44			
1,1,1-triphenylethane				316.73	339.45	-22.72			
1,1,1,2-tetraphenylethane				395.39	398.77	-3.38			
1,1,2,2-tetraphenylethane				396.64	396.66	-0.02			
pentaphenylethane				473.63	470.33	3.30			
triphenylethylenes				309.20	310.10	-0.90			
diphenylacetylene	312.40	312.00	0.40	225.90	225.90	0.00			
biphenyl	99.37	99.36	0.01	197.90	197.86	0.04	205.85	215.50	-9.65
4-methylbiphenyl	55.44	59.99	-4.55						
4,4'-dimethylbiphenyl	14.14	20.62	-6.48						
o-terphenyl				274.34	274.94	-0.60	298.82	294.50	4.32
1,3,5-triphenylbenzene				358.32	352.02	6.30	367.36	373.50	-6.14
naphthalene	77.90	80.44	-2.54	165.69	165.64	0.05	167.40	170.00	-2.60
1-methylnaphthalene	44.27	41.07	3.20						
2-methylnaphthalene	44.85	41.07	3.78	195.98	189.70	6.28	219.99	198.44	21.55
1,8-dimethylnaphthalene	25.94	1.70	24.24	242.80	213.76	29.04	224.72	226.88	-2.16
2,3-dimethylnaphthalene	-0.23	1.70	-1.93						
2,6-dimethylnaphthalene	-5.69	1.70	-7.39	203.55	213.76	-10.21	227.86	226.88	0.98
tetraphenylethylenes	311.50	311.48	0.02	387.60	387.60	0.00			
2,7-dimethylnaphthalene	-5.40	1.70	7.10	204.39	213.76	-9.37	228.57	226.88	1.69
anthracene	129.20	121.70	7.50	210.50	210.50	0.00	207.15	203.50	3.65
naphthacene	158.78	162.96	-4.18	236.56	255.36	-18.80	215.39	237.00	-21.61
phenanthrene	116.20	117.50	-1.30	220.62	217.44	3.18	215.06	219.50	-4.44

TABLE 23. Aromatic hydrocarbons in the solid state at 298.15 K — Continued

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
triphenylene	151.80	154.56	-2.76	259.20	269.24	-10.04	254.68	269.00	-14.32
fluoranthene	230.25	234.68	-4.43	230.58	236.50	-5.92			
pyrene	125.48	125.58	-0.10	227.65	226.50	1.15	224.89	217.50	7.39
perylene				274.93	285.24	-10.31	264.55	283.00	-18.45
coronene				313.76	303.36	10.40	280.87	279.00	1.87

TABLE 24. Cycloalkane and related hydrocarbons in the gas state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol·K)			S° (J/mol·K)			N(tot)
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
cyclopropane	53.26	53.26	0.00	55.94	55.94	0.00	237.44	237.44	0.00	6
cyclobutane	28.37	28.37	0.00	72.22	72.22	0.00	265.39	265.39	0.00	8
cyclopentane	-76.40	-76.40	0.00	83.01	83.01	0.00	292.88	292.88	0.00	10
cyclohexane	-123.39	-123.39	0.00	106.27	106.27	0.00	298.24	298.24	0.00	6
cycloheptane	-118.07	-118.07	0.00	123.09	123.09	0.00	342.33	342.33	0.00	2
cyclooctane	-124.39	-124.39	0.00	139.95	139.95	0.00	366.77	366.77	0.00	8
cyclononane	-132.76	-132.76	0.00							
cyclodecane	-162.67	-162.67	0.00							
cycloundecane	-179.37	-179.37	0.00							
cyclododecane	-230.25	-230.25	0.00							
cyclotridecane	-246.35	-246.35	0.00							
cyclotetradecane	-239.24	-239.24	0.00							
cyclopentadecane	-301.42	-301.42	0.00							
cyclohexadecane	-321.67	-321.67	0.00							
cycloheptadecane	-364.30	-364.30	0.00							
cyclopropene	277.06	277.06	0.00							
cyclobutene	156.69	156.69	0.00	67.07	67.07	0.00	263.51	263.51	0.00	2
cyclopentene	34.56	34.56	0.00	75.10	75.10	0.00	289.66	289.66	0.00	2
cyclohexene	-5.02	-5.02	0.00	105.02	105.02	0.00	310.75	310.75	0.00	2
cycloheptene	-9.20	-9.20	0.00							
cyclooctene	-26.99	-26.99	0.00							
1,3-cyclopentadiene	134.35	134.35	0.00							
1,3-cyclohexadiene	106.23	106.23	0.00							
1,3-cycloheptadiene	94.35	94.35	0.00							
1,5-cyclooctadiene	71.34	71.34	0.00							
1,3,5-cyclo-										
heptatriene	180.87	180.87	0.00	117.78	117.78	0.00	315.64	315.64	0.00	2
cyclooctatetraene	295.85	295.85	0.00	122.01	122.01	0.00	326.77	326.77	0.00	4
spiropentane	185.18	185.18	0.00	88.12	88.12	0.00	282.21	282.21	0.00	4
methylene cyclobutane	121.55	119.46	2.09							
ethylcyclobutane	-26.32	-15.06	11.26							
methylcyclopentane	-106.20	-101.46	-4.74	109.79	109.50	0.29	339.91	339.62	0.29	3
methylene cyclopentane	12.01	14.19	-2.18							
1,1-dimethyl-										
cyclopentane	-138.24	-130.21	-8.03	133.30	131.68	1.62	359.28	356.15	3.13	18
trans-1,2-dimethyl-										
cyclopentane	-136.65	-126.52	-10.13	134.47	132.42	2.05	366.81	359.28	7.53	18
trans-1,3-dimethyl-										
cyclopentane	-133.55	-126.52	-7.03	134.47	132.42	2.05	366.81	359.28	7.53	18

TABLE 24. Cycloalkane and related hydrocarbons in the gas state at 298.15 K — Continued

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)			$N(\text{tot})$
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
ethylcyclopentane	-126.94	-119.83	-7.11	131.75	132.39	-0.64	378.32	378.78	-0.46	3
propylcyclopentane	-147.74	-140.46	-7.28	154.64	155.28	-0.64	417.27	417.94	-0.67	3
butylcyclopentane				177.49	178.17	-0.68	456.22	457.10	-0.88	3
pentylcyclopentane				200.37	201.06	-0.69	495.18	496.26	-1.08	3
hexylcyclopentane				223.22	223.95	-0.73	534.13	535.42	-1.29	3
heptylcyclopentane				246.10	246.84	-0.74	573.04	574.58	-1.54	3
octylcyclopentane				268.99	269.73	-0.74	611.99	613.74	-1.75	3
nonylcyclopentane				291.83	292.62	-0.79	650.95	652.90	-1.95	3
decylcyclopentane	-301.42	-284.87	-16.55	314.72	315.51	-0.79	689.90	692.06	-2.16	3
1-methylcyclopentene	-3.81	0.12	-3.93	100.83	99.22	1.61	326.35	333.07	-6.72	3
3-methylcyclopentene	7.36	9.29	-1.93	100.00	109.72	-9.72	330.54	328.38	2.16	3
4-methylcyclopentene	14.64	9.50	5.14	100.00	100.05	-0.05	328.86	324.20	4.66	3
1-ethylcyclopentene	-19.75	-20.76	1.01							
methylcyclohexane	154.72	-148.47	-6.27	135.02	137.44	-2.42	343.34	344.36	-1.02	3
methylene cyclohexane	-25.23	-32.80	7.57							
1,1-dimethylcyclohexane	-180.87	-177.20	-3.67	154.39	159.62	-5.23	365.01	366.65	-1.64	9
trans-1,2-dimethyl-										
cyclohexane	-179.87	-173.51	-6.36	158.99	160.36	-1.37	370.91	369.78	1.13	9
trans-1,3-dimethyl-										
cyclohexane	-176.48	-173.51	-2.97	157.32	160.36	-3.04	376.23	369.78	6.45	9
trans-1,4-dimethyl-										
cyclohexane	-184.51	-173.51	-11.00	157.74	160.36	-2.62				
ethylcyclohexane	-171.88	-166.82	-5.06	158.82	160.33	-1.51	382.58	383.52	-0.94	3
propylcyclohexane				184.22	183.22	1.00	419.53	422.68	-3.15	3
butylcyclohexane	-213.30	-208.08	-5.22	207.11	206.11	1.00	458.48	461.84	-3.36	3
pentylcyclohexane				229.95	229.00	0.95	497.44	501.00	-3.56	3
dodecylcyclohexane	-378.70	-373.12	-5.58							
1-methylcyclohexene	-43.26	-41.72	-1.54							
1-ethylcyclohexene	-63.43	-60.34	-3.09							
bicyclo[2.2.2]octane	-99.00	-99.00	0.00							
adamantane	-134.60	-134.60	0.00							
bicyclo[3.3.3]undecane	-88.95	-88.95	0.00							

TABLE 25. Cycloalkane and related hydrocarbons in the liquid state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)			
	expt	calc	resd	expt	calc	resd	expt	calc	resd	
cyclobutane	3.72	3.72	0.00	111.00 ^a	111.00	0.00	181.00 ^a	181.00	0.00	
cyclopentane	-105.06	-105.06	0.00	128.78	128.78	0.00	204.14	204.14	0.00	
cyclohexane	-156.44	-156.44	0.00	156.31	156.31	0.00	204.35	204.35	0.00	
cycloheptane	-156.61	-156.61	0.00	180.75	180.75	0.00	242.55	242.55	0.00	
cyclooctane	-167.74	-167.74	0.00	215.48	215.48	0.00	262.00	262.00	0.00	
cyclononane	-181.17	-181.17	0.00							
cyclodecane	-206.69	-206.69	0.00							
cycloundecane	-235.48	-235.48	0.00							
cyclododecane	-306.65	-306.65	0.00							
cyclotridecane	-309.66	-309.66	0.00							
cyclopentene	4.73	4.73	0.00	122.38	122.38	0.00	201.25	201.25	0.00	
cyclohexene	-38.49	-38.49	0.00	148.36	148.36	0.00	214.60	214.60	0.00	
cyclooctene	-74.02	-74.02	0.00							
1,3-cyclopentadiene	105.98	105.98	0.00							
1,3-cyclohexadiene	73.18	73.18	0.00	144.56	144.56	0.00	197.28	197.28	0.00	

TABLE 25. Cycloalkane and related hydrocarbons in the liquid state at 298.15 K — Continued

Name	ΔH° (kJ/mol)			C_p° (J/mol-K)			S° (J/mol-K)		
	expt	calc	resd	expt	calc	resd	expt	calc	resd
1,4-cyclohexadiene				145.94	145.94	0.00	189.37	189.37	0.00
1,5-cyclooctadiene	30.29	30.29	0.00	208.11	208.11	0.00	264.35	264.35	0.00
1,3,5-cycloheptatriene	142.17	142.17	0.00	162.76	162.76	0.00	214.64	214.64	0.00
cyclooctatetraene	254.51	254.51	0.00	185.18	185.18	0.00	220.29	220.29	0.00
spiropentane	157.65	157.65	0.00	134.52	134.52	0.00	193.68	193.68	0.00
methylene cyclobutane	93.85	90.36	3.49	131.13	132.17	-1.04	210.20	204.98	5.22
methylcyclobutane	-44.48	-25.11	-19.37						
ethylcyclobutane	-58.95	-48.66	-10.29						
methylcyclopentane				158.70	155.25	3.45	247.78	245.58	2.20
methylene cyclopentane	-20.08	-18.42	-1.66						
1,1-dimethylcyclopentane	-172.05	-165.34	-6.71	187.36	180.59	6.77	265.01	254.12	10.89
trans-1,2-dimethylcyclopentane	-171.21	-162.72	-8.49	187.40	183.66	3.74	269.90	272.61	-2.71
trans-1,3-dimethylcyclopentane	-168.07	-162.72	-5.35						
ethylcyclopentane	-163.43	-157.44	-5.99	185.31	186.64	-1.33	279.91	277.96	1.95
ethylidenecyclopentane				181.17	182.66	-1.49			
ethenylcyclopentane	-34.81	-31.55	-3.26						
allylcyclopentane				202.92	202.00	.92			
propylcyclopentane	-188.82	-183.17	-5.65	216.27	217.06	-0.79	310.83	310.34	0.49
butylcyclopentane				245.35	247.48	-2.13	343.84	342.72	1.12
decylcyclopentane	-367.31	-363.28	-4.03						
11-cyclopentylheneicosane	-648.60	-647.23	-1.37						
bicyclopentyl				238.91	239.48	-0.57			
1-methylcyclopentene	-36.44	-34.77	-1.67	153.10	157.48	-4.38			
3-methylcyclopentene	-17.57	-24.35	6.78						
1-ethylcyclopentene	-58.28	-60.50	2.22	188.28	186.77	1.51			
methylcyclohexane	-190.08	-185.27	-4.81	184.51	183.75	0.76	247.90	246.41	1.49
methylene cyclohexane	-61.30	-69.80	8.50	177.40	175.22	2.18			
1,1-dimethylcyclohexane	-218.74	-216.72	-2.02	209.24	209.09	0.15	267.23	254.95	12.28
1,2-dimethylcyclohexane	-218.24	-214.10	-4.14	212.76	211.19	1.57	273.22	273.44	-0.22
1,3-dimethylcyclohexane	-215.69	-214.10	-1.59	212.84	211.19	1.65	276.27	273.44	2.83
1,4-dimethylcyclohexane	-222.38	-214.10	-8.28	210.25	211.19	-0.94	268.03	273.44	-5.41
ethylcyclohexane	-212.13	-208.82	-3.31	211.79	214.17	-2.38	280.91	278.79	2.12
allylcyclohexane				233.47	229.53	3.94			
ethylidenecyclohexane				203.76	207.93	-4.17			
propylcyclohexane				242.04	244.59	-2.55	311.88	311.17	0.71
butylcyclohexane	-263.09	-260.28	-2.81	271.04	275.01	-3.97	344.97	343.55	1.42
dodecylcyclohexane	-467.56	-466.12	-1.44						
3-cyclohexyleicosane	-666.18	-672.88	6.70						
9-cyclohexyleicosane	-674.13	-672.88	-1.25						
11-cyclohexylheneicosane	-689.44	-698.61	9.17						
13-cyclohexylpentacosane	-792.66	-801.53	8.87						
1-methylcyclohexene	-81.17	-80.17	-1.00						
1-ethylcyclohexene	-106.69	-103.72	-2.97						
cis-hexahydroindan	-173.10	-173.10	0.00	214.18	214.18	0.00	265.47	265.47	0.00
trans-hexahydroindan	-173.20	-173.20	0.00	209.70	209.70	0.00	258.86	258.86	0.00
cis-decalin	-219.40	-219.40	0.00	232.00	232.00	0.00	265.01	265.01	0.00
trans-decalin	-230.60	-230.60	0.00	228.49	228.49	0.00	264.93	264.93	0.00

*Obtained from extrapolation of experimental data of 53RAT/GWI to 298.15 K.

TABLE 26. Cycloalkane and related hydrocarbons in the solid state at 298.15 K

Name	$\Delta_f H^\circ$ (kJ/mol)		expt	calc	C_p °(J/mol-K)		expt	S° (J/mol-K)		resd
	expt	calc			resd	calc		calc	resd	
cyclopentane	-113.05	-113.05			0.00					
cyclohexane	-166.16	-166.16			0.00					
cyclododecane	-306.65	-306.65			0.00					
cyclotetradecane	-374.00	-374.00			0.00					
cyclopentadecane	-376.06	-376.06			0.00					
cyclohexadecane	-403.42	-403.42			0.00					
cycloheptadecane	-430.41	-430.41			0.00					
decylcyclopentane	-401.00	-401.05			-0.05					
dodecylcyclohexane	-514.00	-512.98			-1.02					
2,2-metacyclophane	78.50	78.50			0.00					
2,2-metaparacyclophane	130.90	130.90			0.00	261.50	261.50	0.00		
2,2-paracyclophane	145.10	145.10			0.00	252.34	252.34	0.00	265.68	265.68
3,3-paracyclophane	26.30	26.30			0.00	324.26	324.26	0.00		0.00
bicyclo[2.2.2]octane	-146.90	-146.90			0.00					
adamantane	194.10	-194.10			0.00					