

¹³C Chemical Shieldings in Solids

T. M. Duncan

AT&T Bell Laboratories, Murray Hill, New Jersey 07974

Received June 20, 1986; revised manuscript received October 3, 1986

Analogous to the importance of ¹³C isotropic shieldings for chemical analysis of liquids with nuclear magnetic resonance spectroscopy, ¹³C chemical shielding anisotropies are proving to be valuable in the characterization of solids. Specifically, molecular geometry is revealed by the full shielding anisotropy and molecular motion may be characterized by changes in the powder pattern. In particular, the principal components of the shielding reveal differences in bonding geometry which may not be correlated to monotonic changes in the isotropic shift. This report is a comprehensive, critical compilation of ¹³C chemical shieldings in solids, organized by carbon functionality. From these data, representative shieldings of common carbon functionalities are calculated.

Key words: ¹³C chemical shielding; chemical shift anisotropy; critical review; nuclear magnetic resonance; solid state.

Contents

1. Introduction	125	7. Carbon–nitrogen functionalities	142
2. Criteria for Selection of Data	126	8. Aromatic carbons	143
3. Description of Tables	126	9. Carbon monoxide and inorganic carbonyls	146
3.1. Data Tables	126	10. Miscellaneous compounds	147
3.2. Summary Table	127	11. Typical chemical shieldings of carbon functiona-	
4. Text References	128	lities	149
5. Table References	150		

List of Tables

1. Paraffinic carbons	129
2. Olefinic carbons	134
3. Acetylenic carbons	135
4. Alcohols and ethers	136
5. Aldehydes and ketones	138
6. Carboxylic acids and esters	139

List of Figures

1. Typical chemical shielding spectra of <i>sp</i> ³ -hybridized carbon functionalities	127
2. Typical chemical shielding spectra of <i>sp</i> ² -hybridized carbon functionalities	127
3. Typical chemical shielding spectra of <i>sp</i> -hybridized and aromatic carbon functionalities	128
4. Chemical structures of selected compounds	128

1. Introduction

The chemical shielding at a nucleus in a diamagnetic material is the magnetic field caused by electronic currents induced by an applied magnetic field.^{1,2} The applied field causes two effects: (1) induced precession of electron orbits, quenched by electrostatic charges of the molecule (diamagnetic contribution) and (2) mixing of the electronic ground state with excited states, which polarizes the electron distribution and thus alters the current path (paramagnetic contribution). The diamagnetic effect decreases the net magnet-

ic field at a nucleus, whereas the paramagnetic effect increases the field. Both effects, and therefore the net chemical shielding, are proportional to the applied magnetic field. Consequently, chemical shieldings are reported as fractional shifts of the applied magnetic field, relative to the shift of an arbitrary liquid standard. For ¹³C, chemical shifts are generally ~0.001%–0.02% (10–200 ppm) of the applied field and the standard shift reference is tetramethylsilane (TMS).

The orientational anisotropy of the chemical shielding may be represented by a second-rank, dimensionless Cartesian tensor, σ .^{3–5} Each chemically and crystallographically distinct site has a distinct chemical shielding tensor. The trace of σ yields the isotropic shift, σ_{iso} , which is the observed shift of molecular sites undergoing rapid isotropic reorientation, such as liquids. The chemical shift of a fixed

©1987 by the U. S. Secretary of Commerce on behalf of the United States. This copyright is assigned to the American Institute of Physics and the American Chemical Society.

Reprints available from ACS; see Reprints List at back of issue.




FIG. 1. Typical ¹³C chemical shielding powder patterns of *sp*³-hybridized carbon functionalities, calculated from the data in Tables 1–10. All spectra are convoluted with Lorentzian broadening of halfwidth 2 ppm. A dangling bond in the chemical structures implies a bond to a carbon atom.

monoxide and inorganic carbonyls, and miscellaneous. Each entry contains the compound name (proper and common), the structural chemical formula, the temperature of the measurement, the three chemical shielding principal components, the isotropic shift, and the literature reference. All studies meeting the above criteria were included, thus some species have multiple entries. The shieldings are reported relative to TMS, on the δ scale, such that downfield shifts are positive. Studies with frequency references other than TMS were converted using the isotropic shifts given in Stothers's monograph.¹⁰ The most common alternate references are benzene (128.7 ppm relative to TMS on the δ scale) and CS₂ (192.8 ppm). When no shift reference is given, the isotropic shift of the solid is taken as that of the liquid, given in Stothers's monograph; these entries are indicated by a footnote in the tables. The convention for the relative assignment of the Cartesian components σ_{xx} , σ_{yy} , and σ_{zz} varies with investigator. The principal components are tabulated here as σ_{11} , σ_{22} , and σ_{33} , such that $\sigma_{11} < \sigma_{22} < \sigma_{33}$, with no inference to the Cartesian axes. In some cases the principal components were calculated from reports of the anisotropy and asymmetry, definitions of which also vary. Some define the anisotropy as the total width of the powder pattern ($\sigma_{11} - \sigma_{33}$), whereas others define it as the difference between the isotropic shift and the extreme shielding component ($\sigma_{11} - \sigma_{\text{iso}}$, or $\sigma_{\text{iso}} - \sigma_{33}$). However, in each of these cases the principal




FIG. 2. Typical ¹³C chemical shielding powder patterns of *sp*²-hybridized carbon functionalities, calculated from the data in Tables 1–10. All spectra are convoluted with Lorentzian broadening of halfwidth 2 ppm. A dangling bond in the chemical structures implies a bond to a carbon atom.

components were obtained without ambiguity.

The quality of the experimental data vary and confidence limits (when reported) range from ± 20 to ± 0.2 ppm; confidence limits are not tabulated here. The tabulated data are not corrected for second-order dipolar perturbations, which are typically < 1 ppm upfield.¹¹ It is important to note the temperature at which each spectrum was obtained because some species (especially methyl and aromatic groups) may possess some residual motion and the spectrum will not reflect the full anisotropy. Residual motional averaging is often indicated by the degeneracy of two principal shielding components for a site that does not have C_{3v} (or higher) symmetry. In summary, readers are advised to consult the original references to assess individual spectra.

3.2. Summary Table

Table 11 contains the typical shielding components for 23 carbon functionalities, obtained from averages of the data in Tables 1–10. Only the spectra of compounds free of conformational restraints were used in the averages (e.g., no data from cyclic compounds with other than six atoms). In some cases, these typical spectra are based on small data bases and are expected to change as more spectra are reported and as accuracy improves. Powder patterns of the 23

TABLE 1. Paraffinic carbons—Continued

Compound	Formula	T (K)	Chemical shielding parameters ^(a)				Reference
			σ_{11}	σ_{22}	σ_{33}	σ_{iso}	
<i>Methyl carbons, continued</i>							
trimethyl phosphine	P(CH ₃) ₃	(lc)	16	16	16	16	Albrand <i>et al.</i> (1977)
trimethyl phosphorus oxide	O=P(CH ₃) ₃	(lc)	25	25	-9	14	Albrand <i>et al.</i> (1977)
trimethyl phosphorus sulfide	S=P(CH ₃) ₃	(lc)	24	24	16	21	Albrand <i>et al.</i> (1977)
trimethyl phosphorus selenide	Se=P(CH ₃) ₃	(lc)	27	27	9	21	Albrand <i>et al.</i> (1977)
methyl fluoride	CH ₃ F	15 (lc)	105 104	105 104	15 17	75 75	Zilm and Grant (1981) Jokisaari <i>et al.</i> (1986)
methyl iodide	CH ₃ I	(lc)	32	32	-1	21	Yannoni and Whipple (1967)
<i>Methylene carbons</i>							
<i>n</i> -eicosane	CH ₃ C [*] H ₂ (CH ₂) ₁₀ C [*] H ₂ CH ₃ CH ₃ CH ₂ (C [*] H ₂) ₁₆ CH ₂ CH ₃	RT RT	37 50	30 38	15 17	27 35	VanderHart (1976)
polyethylene	(-CH ₂ -) _x	RT RT	52 50	39 37	14 13	35 33	Opella and Waugh (1977) Urbina and Waugh (1974)
polymethyl-methacrylate	(-C [*] H ₂ C(CH ₃)(C(O)(OCH ₃))-) _n	RT	79	48	29	52	Edzes (1983)
poly(vinylidene fluoride)	(-C [*] H ₂ CF ₂ -) _n	297	56	49	30	45	Fleming <i>et al.</i> (1980)
L-asparagine monohydrate	NH ₂ C(O)C [*] H ₂ CH(NH ₃)COOH	RT	49	43	16	36	Naito and McDowell (1984)
malonic acid	HOOCC [*] H ₂ COOH	RT	62	50	18	43	Tegenfeldt <i>et al.</i> (1980)
ammonium hydrogen malonate	(OOCC [*] H ₂ COO)(H)(NH ₄)	RT	63	52	19	45	Chang <i>et al.</i> (1975)
dihydro muconic acid	HOOCC [*] H ₂ CH=CHC [*] H ₂ COOH	RT	40	38	7	28	Wolff <i>et al.</i> (1977)
hexaethyl benzene	C ₆ (C [*] H ₂ CH ₃) ₆	RT	27	25	7	20	Pausak <i>et al.</i> (1974)
malononitrile	N≡CC [*] H ₂ C≡N	20	30	11	-11	10	Facelli <i>et al.</i> (1985)
dimedone	 CH ₃ C [*] H ₂ - C(O) CH ₃ CH ₃ C [*] H ₂ - C(OH)	RT	57 61	48 55	28 33	44 49	Takegoshi <i>et al.</i> (1985)

TABLE 1. Paraffinic carbons—Continued

Compound	Formula	T (K)	Chemical shielding parameters ^(a)				Reference
			σ_{11}	σ_{22}	σ_{33}	σ_{iso}	
<i>Methylene carbons, continued</i>							
2-isopropyl, 5-methyl-cyclohexanol (1-menthol) ^(b)		RT	33 48 72	27 38 52	21 30 24	27 39 50	Maciel et al. (1985)
<i>tert</i> -butyl-cyclohexane		20	57	25	9	30	Facelli et al. (1985)
1,4-cyclohexadiene		20	44	21	10	25	Facelli et al. (1985)
1,3,5-cycloheptatriene (tropilidene)		20	55	25	16	32	Facelli et al. (1985)
cyclopropane		20	22	2	-36	-4	Zilm et al. (1981)
spiropentane (bowtie-ane)		20	37	16	-23	10	Facelli et al. (1985)
bicyclo[2.2.1]hepta-2,5-diene (norbornadiene)		20	97	69	63	76	Facelli et al. (1985)
bicyclo[1.1.1]pentane		20	56	54	38	49	Facelli et al. (1985)
bicyclo[1.1.1]pentanone		20	87	47	1	45	Facelli et al. (1985)
cyclopropene		20	40	29	-59	3	Zilm et al. (1980)
cyclobutane		20	39	23	14	25	Facelli et al. (1985)
trimethylene oxide (oxetane)		20	41	22	4	22	Facelli et al. (1985)
cyclobutene		30	43	33	23	33	Zilm et al. (1980)

TABLE 4. Alcohols and ethers—Continued

Compound	Formula	T (K)	Chemical shielding parameters ^(a)				Reference
			σ_{11}	σ_{22}	σ_{33}	σ_{iso}	
<i>Ethers, cont'd</i>							
polyoxymethylene	$(-\text{OCH}_2-)_n$	RT	114	87	64	88	Veeman (1984)
1,2-epoxy ethane (ethylene oxide)		20	93	19	19	44	Facelli et al. (1985)
trimethylene oxide (oxetane)		20	104	93	15	71	Facelli et al. (1985)
1,4-epoxybutane (tetrahydrofuran)		20	110	75	21	68	Facelli et al. (1985)
1,4-dioxane		20	86	81	37	68	Facelli et al. (1985)
trimethoxymethane	$\text{C}^*\text{H}(\text{OCH}_3)_3$	20	123	116	99	113	Facelli et al. (1986)
dimethyl dimethoxy silane	$\text{Si}(\text{CH}_3)_2(\text{OC}^*\text{H}_3)_2$	87	74	69	-3	47	Waugh et al. (1973)
tetramethoxy silane	$\text{Si}(\text{OCH}_3)_4$	87	72	66	2	47	Waugh et al. (1973)

(a) relative to TMS (tetramethylsilane), on the δ scale, such that $\text{C}_6\text{H}_{6(0)} = 128.7$ ppm and $\text{CS}_2 = 192.8$ ppm (i.e., upfield is negative).

(b) peak assignments not given in original reference; assignments proposed here are based on isotropic shifts given in Stothers (1972).

(c) oriented in a liquid crystal, only anisotropy given; principal components calculated by assuming isotropic shift is that of neat liquid given in Stothers (1972).

TABLE 5. Aldehydes and ketones

Compound	Formula	T (K)	Chemical shielding parameters ^(a)				Reference
			σ_{11}	σ_{22}	σ_{33}	σ_{iso}	
<i>Aldehydes</i>							
acetaldehyde	$\text{CH}_3\text{C}^*\text{HO}$	87 15	276 285	234 231	87 84	199 200	Pines <i>et al.</i> (1972) Zilm and Grant (1981)
1,1,2,2-tetra-acetylethane	$\text{CH}_3\text{C}(\text{OH})=\text{CC}^*(\text{O})\text{CH}_3$ $\text{CH}_3\text{C}(\text{OH})=\text{CC}^*(\text{O})\text{CH}_3$	RT RT	278	240	85	201	Takegoshi and McDowell (1986)
3,4-dimethoxy-benzaldehyde (veratraldehyde)	$\text{HC}^*(\text{O})-\text{C}_6\text{H}_3-(\text{OCH}_3)_2$ (see Fig. 4)	RT	264	211	101	192	Terao <i>et al.</i> (1984b)
<i>Ketones</i>							
dimethyl ketone (acetone)	$\text{CH}_3\text{C}^*(\text{O})\text{CH}_3$	87	279	265	79	208	Pines <i>et al.</i> (1972)
methylphenyl ketone (acetophenone)	$(\text{C}_6\text{H}_5)\text{C}^*(\text{O})\text{CH}_3$	100	279	231	84	198	Van Dongen Torman <i>et al.</i> (1978)
diphenyl ketone (benzophenone)	$\text{C}_6\text{H}_5\text{C}^*(\text{O})\text{C}_6\text{H}_5$	RT	272	229	99	200	Kempf <i>et al.</i> (1972)
squaric acid (3,4-dihydroxy-3-cyclobutene-1,2-dione)	$(\text{HO})\text{C}=\text{C}(\text{OH})$ $(\text{O})\text{C}^*-\text{C}^*(\text{O})$	RT	241 236	230 235	84 84	185 182	Mehring and Becker (1981)
dimedone		RT	284	255	79	206	Takegoshi <i>et al.</i> (1985)

(a) relative to TMS (tetramethylsilane), on the δ scale, such that $\text{C}_6\text{H}_{6(0)} = 128.7$ ppm and $\text{CS}_2 = 192.8$ ppm (*i.e.*, upfield is negative).

TABLE 6. Carboxylic acids and esters—Continued

Compound	Formula	T (K)	Chemical shielding parameters ^(a)				Reference
			σ_{11}	σ_{22}	σ_{33}	σ_{iso}	
<i>Esters, cont'd</i>							
silver trifluoroacetate	CF ₃ C [*] OOAg	296	245	129	129	168	Waugh <i>et al.</i> (1973)
silver benzoate	(C ₆ H ₅)C [*] OOAg	RT	247	176	93	172	Kempf <i>et al.</i> (1974)
ammonium hydrogen malonate	(OOC [*] CH ₂ C [*] OO)(H)(NH ₄)	RT	242	175	102	173	Chang <i>et al.</i> (1975)
potassium hydrogen malonate	(OOC [*] CH ₂ C [*] OO)(H)(K)	RT	253	173	97	174	Schroter <i>et al.</i> (1983)
potassium hydrogen succinate	(OOC [*] CH ₂ CH ₂ C [*] OO)(H)(K)	RT	263 ^(b)	179 ^(b)	105 ^(b)	182	Schroter <i>et al.</i> (1983)
ammonium d-tartrate	(-CH(OH)C [*] OONH ₄) ₂	RT	239	190	107	179	Pines <i>et al.</i> (1974)
ammonium hydrogen maleate (cis-butenedioic acid)	(OOC [*] CH=CHC [*] OO)(H)(NH ₄)	RT	240 ^(c)	172 ^(c)	104 ^(c)	172	Schroter <i>et al.</i> (1983)
ammonium hydrogen fumarate (trans-butenedioic acid)	(OOC [*] CH=CHC [*] OO)(H)(NH ₄)	RT	249 ^(b)	172 ^(b)	101 ^(b)	174	Schroter <i>et al.</i> (1983)
ammonium hydrogen glutarate (pentanedioic acid)	(OOC [*] CH=CHC [*] OO)(H)(NH ₄)	RT	251 ^(c)	179 ^(c)	110 ^(c)	180	Schroter <i>et al.</i> (1983)
dimethyl maleic anhydride	(CH ₃) ₂ C=C(CH ₃) (O)C [*] -O-C [*] (O)	RT	258 263	126 128	120 122	168 171	Igner and Fiat (1982)

(a) relative to TMS (tetramethylsilane), on the δ scale, such that C₆H₆₍₀₎ = 128.7 ppm and CS₂ = 192.8 ppm (i.e., upfield is negative).

(b) determined from spinning sidebands of a rotating sample.

(c) determined from narrowed powder pattern obtained spinning 7° off magic angle.

TABLE 8. Aromatic carbons—Continued

Compound	Formula	T (K)	Chemical shielding parameters ^(a)				Reference
			σ_{11}	σ_{22}	σ_{33}	σ_{iso}	
<i>Heterocyclic aromatics</i>							
azine (pyridine)	C ₅ H ₅ N	(lc)	241 205 233	173 158 170	36 8 4	150 124 136	Parhami and Fung (1985)
1,2-diazine (pyridazine)	C ₄ H ₄ N ₂	(lc)	293 190	130 186	31 3	152 126	Parhami and Fung (1985)
1,3-diazine (pyrimidine)	C ₄ H ₄ N ₂	(lc)	246 239 221	176 201 138	55 30 4	159 157 121	Parhami and Fung (1985)
1,4-diazine (pyrazine)	C ₄ H ₄ N ₂	(lc)	223	173	39	145	Parhami and Fung (1985)
<i>Miscellaneous</i>							
cyclopentadienide anion	C ₅ H ₅ ⁻	20	182	114	21	106	Strub <i>et al.</i> (1983)
tropylium cation	C ₇ H ₇ ⁺	20	280	168	22	167	Strub <i>et al.</i> (1983)

(a) relative to TMS (tetramethylsilane), on the δ scale, such that C₆H₆₍₀₎ = 128.7 ppm and CS₂ = 192.8 ppm (i.e., upfield is negative).

(b) peak assignments not given in original reference; assignments proposed here are based on isotropic shifts given in Stothers (1972).

(c) non-substituted ring carbon.

TABLE 10. Miscellaneous compounds

Compound	Formula	T (K)	Chemical shielding parameters ^(a)				Reference
			σ_{11}	σ_{22}	σ_{33}	σ_{iso}	
<i>Carbon-oxygen</i>							
carbon dioxide	CO ₂	20	245	245	-90	133	Beeler <i>et al.</i> (1984)
carbonyl sulfide	SCO	20	275	275	-90	154	Beeler <i>et al.</i> (1984)
calcium carbonate	CaCO ₃	RT	194	194	119	169	Lauterbur (1958)
		RT	195	195	119	169	Pines <i>et al.</i> (1971)
dimethyl carbonate	(CH ₃ O) ₂ C [*] O	87	231	110	110	150	Pines <i>et al.</i> (1972)
<i>Carbon-sulfur</i>							
carbon disulfide	CS ₂	73	333	333	-92	193	Pines <i>et al.</i> (1971)
		20	332	332	-90	191	Zilm <i>et al.</i> (1978)
thioacetic acid	CH ₃ C [*] OSH	87	264	230	101	202	Waugh <i>et al.</i> (1973)
dimethyl sulfoxide	(C [*] H ₃) ₂ SO	226	66	44	18	43	Pines <i>et al.</i> (1972)
dimethyl sulfone	(C [*] H ₃) ₂ SO ₂	295	63	63	7	44	Solum <i>et al.</i> (1983)
dimethyl disulfide	C [*] H ₃ S—SC [*] H ₃	87	48	18	6	24	Pines <i>et al.</i> (1972)
ethylene sulfide	$\begin{matrix} \text{CH}_2 & \text{CH}_2 \\ & \backslash \\ & \text{S} \end{matrix}$	20	54	11	7	24	Facelli <i>et al.</i> (1985)
<i>Carbon-fluorine</i>							
trifluoro silver acetate	C [*] F ₃ COOAg	296	129	129	110	123	Waugh <i>et al.</i> (1973)
trifluoro acetic anhydride	(C [*] F ₃ CO) ₂ O	109	119	119	102	113	Waugh <i>et al.</i> (1973)
poly(vinylidene fluoride)	(—CH ₂ C [*] F ₂ —) _n	297	131	120	111	121	Fleming <i>et al.</i> (1980)
<i>Inorganic carbides</i>							
boron carbide	B ₁₂ (CC [*] C)	RT	190	170	-105	85	Duncan (1984)
tungsten carbide	WC	RT	348	286	286	307	Duncan (1986)
molybdenum carbide	Mo ₂ C	RT	307	274	244	275	Duncan (1986)
—	CsZr ₆ I ₁₄ C	RT	54	54	15	41	Fry <i>et al.</i> (1986)

TABLE 10. Miscellaneous compounds—Continued

Compound	Formula	T (K)	Chemical shielding parameters ^(a)				Reference
			σ_{11}	σ_{22}	σ_{33}	σ_{iso}	
<i>Metal-sandwich compounds</i>							
ferrocene	$(C_5H_5)_2Fe$	—	95	95	19	70	Wemmer and Pines (1981)
bis(cyclopentadienyl) ruthenium	$(C_5H_5)_2Ru$	—	100	100	21	74	Wemmer and Pines (1981)
bis(cyclopentadienyl) magnesium	$(C_5H_5)_2Mg$	—	153	153	21	109	Wemmer and Pines (1981)
bis(cyclopentadienyl) titanium dichloride	$(C_5^*H_5)_2TiCl_2$	—	168	168	13	116	Wemmer and Pines (1981)
permethyl ferrocene	$(C_5^*(CH_3)_5)_2Fe$	— 88	128 102 128	87 102 116	24 35 24	45 62 89	Wemmer and Pines (1981) Wemmer <i>et al.</i> (1981)
bis(pentamethyl cyclopentadienyl) cobalt chloride	$(C_5^*(CH_3)_5)_2CoCl$	—	180 172	124 111	39 39	114 107	Wemmer and Pines (1981)
bis(benzene) chromium	$(C_6H_6)_2Cr$	—	115	115	1	77	Wemmer and Pines (1981)
<i>Others</i>							
graphite	C _n	RT RT	178 251	178 20	0 20	119 97	Resing <i>et al.</i> (1982) Duncan <i>et al.</i> (1985)

(a) relative to TMS (tetramethylsilane), on the δ scale, such that $C_6H_{6(0)} = 128.7$ ppm and $CS_2 = 192.8$ ppm (i.e., upfield is negative).

TABLE 11. Typical chemical shieldings of carbon functionalities

Compound	Formula ^(a)	Chemical shielding parameters ^(b)						Number of Reference Spectra
		σ_{11}	σ_{22}	σ_{33}	σ_{iso}			
methyl	-CH ₃	38 (10)	29 (10)	3 (5)	23 (6)			38
methylene	-CH ₂ -	54 (13)	41 (10)	16 (6)	37 (8)			9
methylidyne	-CH-	40 (15)	32 (10)	28 (10)	33 (11)			5
alcohol	-CH ₂ OH	80	70	25	58			2
	-CH-OH	90 (4)	83 (5)	42 (8)	72 (4)			6
ether	-O-CH ₃	83 (7)	71 (4)	12 (3)	55 (3)			10
	-O-CH ₂ -	90 (9)	83 (3)	27 (7)	67 (3)			3
amine	-CH-NH ₃	68 (2)	54 (7)	36 (11)	52 (6)			6
olefinic		223 (8)	127 (14)	34 (11)	128 (6)			9
aldehyde	-C(O)H	276 (9)	229 (13)	90 (8)	198 (4)			4
ketone	-C(O)-	279 (5)	245 (18)	85 (10)	203 (5)			4
acid	-C(O)OH	247 (10)	171 (18)	106 (4)	175 (6)			24
ester	-C(O)-O-	258 (13)	132 (18)	107 (14)	166 (10)			9
salt	-C(O)-O-M	239 (13)	188 (19)	105 (8)	176 (6)			24
amide	-C(O)-NH ₂	243 (7)	179 (6)	94 (9)	172 (3)			14
acetylenic	-C≡C-	151 (9)	151 (9)	-79 (17)	75 (8)			8
CO	C≡O	316	316	-90	181			1
carbonyl (t)	M-C≡O	327 (22)	320 (21)	-64 (28)	195 (15)			21
carbonyl (br)		311 (18)	292 (18)	125 (37)	242 (21)			4
nitrile	-C≡N	227 (4)	208 (19)	-89 (6)	115 (5)			3

