# **REFERENCE MATERIALS FOR NUCLEAR FORENSICS**

J.L. Mann (1), M. Tyra (1), R.D. Vocke (2), R. Essex (3), J. Morrison (4), J. Leggitt (5), S. Jerome (6) (1) Radiation Physics Division, NIST, Gaithersburg, MD 20899; (2) Chemical Sciences Division, NIST, Gaithersburg, MD 20899; (3) New Brunswick Laboratory, U.S. Department of Energy, Argonne, IL 60439; (4) Domestic Nuclear Detection Office, Department Homeland of Security, Washington, DC 20528; (5) Laboratory Division Federal Bureau of Investigation, Quantico, VA 22135; (6) National Physical Laboratory, Teddington, Middlesex, TW11 OLW



For further information please contact: Jacqueline L. Mann, NIST, 100 Bureau Drive, Stop 8462, Gaithersburg, MD 20899, USA (phone: 301-975-4472, Email: jmann@nist.gov)

### Background

The discipline of nuclear forensics was developed because of an increasing concern nuclear or radiological material will be used in a terrorist attack. Nuclear forensics is the thorough collection, analysis and evaluation of radiological and nuclear material in pre-detonation and post-detonation scenarios resulting from a nuclear detonation. Conclusions drawn from this collected data coupled with law enforcement and intelligence information, may support nuclear attribution – the identification of those responsible for planned and actual attacks. The development of reference materials (RMs) for the improvement of traceability, accuracy, and precision of nuclear forensic measurements is currently being pursued by NIST as well as the USDOE New Brunswick Laboratory (NBL) and partner organizations (DHS DNDO NTNFC, FBI, DOD, DOE National Laboratories, and international partners), with support from DHS National Technical Nuclear Forensic Center and the FBI. Using a proper RM standard to constrain measurement accuracy and uncertainty bolsters the veracity of measurement results and provides a strong base to stand up to legal scrutiny.







Projects

## **Isotope Dilution Mass Spectrometry Tracer Reference Material**



#### Purpose For accurate <sup>230</sup>Th assay for <sup>234</sup>U-<sup>230</sup>Th age-dating.

#### **Project Description**

- Production of ~160 <sup>229</sup>Th IDMS tracer units (prepared from existing NIST Stock <sup>229</sup>Th Solution).
- Production of ~ 80 <sup>232</sup>Th calibration units (prepared from existing NIST stock <sup>232</sup>Th Solution).
- Primary: Certification of <sup>229</sup>Th tracer material for assay & isotopic composition.
- Secondary: Higher accuracy determination of <sup>229</sup>Th halflife.

#### **Current Efforts**

• Unit production completed (NIST).

<sup>134</sup>Ba **Isotope Dilution Mass Spectrometry Tracer Reference Material** 



#### Purpose Tracer for accurate <sup>137</sup>Ba assay for <sup>137</sup>Cs-<sup>137</sup>Ba age-dating.

#### **Project Description**

- Production of enriched <sup>134</sup>Ba (>99 %) isotopic tracer.
- Units of tracer to be ampouled in high purity quartz.
- Certification of <sup>134</sup>Ba tracer material for assay & isotopic composition.

#### **Current Efforts**

Completing production of <sup>134</sup>Ba material. Production of tracer material sample units. Evaluation of test samples for isotopics complete (Table 1).

### Surrogate Post-Detonation Urban Debris **Standard Reference Material**

#### Purpose

"What will happen when the explosions come—when a part of New York or Cairo or Adelaide has been hollowed out by a device in the kiloton range? Since even a so called fizzle yield could kill a number

of thousands of people, how many nuclear detonations can the world tolerate?"

—John McPhee, The Curve of Binding Energy (Farrar, Straus and Giroux, New York, 1974).

To test post-detonation nuclear forensic measurement capability by providing a surrogate postdetonation material to enable:

- method development and validation,
- testing and readiness,
- and enable labs to legally and scientifically demonstrate the efficacy, accuracy, and precision of analysis methods and establish traceability.

What Is It? Surrogate debris urban material with fresh fission products that mimics "rubble" of a blown up city after an Improvised Nuclear Device



- Samples prepared (NIST & NBL).
- Sample assay and isotopic analysis complete (LANL, LLNL, CEA).
- Preliminary data evaluation complete (NBL & NIST) (Figure 1).



**Figure 1.** The <sup>229</sup>Th Tracer solution concentration data from the 3 labs. It is clear that the measured concentration is consistent with the concentration determined from <sup>229</sup>Th activity (green circle) and IDMS concentration determinations using the SRM 4342A (<sup>230</sup>Th) solution as a tracer (red circles). A systematic bias, however, appears to exist between the assay values determined using the <sup>232</sup>Th calibrant versus the <sup>229</sup>Th activity determination and the <sup>230</sup>Th calibrant. Error bars are expanded uncertainties (k=2).

	NIST	INL
	lso %	lso%
<sup>130</sup> Ba	1.2005	<0.05
<sup>132</sup> Ba	0.0421	<0.08
<sup>134</sup> Ba	90.287	93.6
<sup>135</sup> Ba	1.5527	1.87
<sup>136</sup> Ba	0.2875	0.591
<sup>137</sup> Ba	5.9751	0.625
<sup>138</sup> Ba	0.6552	3.45

Table 1. The isotopic abundance results from NIST and Idaho National Laboratory (INL) for the Ba isotopes are listed. One concern with producing highly-enriched materials is natural Ba contamination, in this case contamination from natural Ba which is dominated by <sup>138</sup>Ba. INL requested NIST assess whether natural Ba had contaminated the purified material. In this case we did not find an issue with natural contamination but we did find contamination at the <sup>137</sup>Ba isotope.

#### detonation.

- vitrified mixture of cement/concrete/steel material (CaAlFeSi) doped with HEU  $(^{235}\text{U}-22\%)$  and major, minor & trace elements
- Particle size 150-300 microns,100 units of 25g each



#### How Would It Work?

- Known starting material (known chemical) concentrations and isotope composition)
- Expose to known neutron energy & fluence (reactor)
- Predict "known" activation and fission product yields

<b>Certification Stages</b>	Lab	Analysis Method	Funding Source	Status
Production	NPL		FBI	Complete
Homogeneity Characterization	DoD, FBI, NIST	INAA, µXRF	NIST	ln progress

#### **Future Efforts (Verification Analyses)**

Verification Analysis.

- Production of Th Metal IDMS Solutions.
- IDMS sample Production and Analysis.
- Data evaluation and production report.
- Completion of Certification Report.

#### **Future Efforts**

- Analysis of additional test samples to assess potential contamination.
- Production of purified tracer material sample units.
- Assay & isotopic characterization (true) of units.
- Data evaluation, report, CRM certificate.

Dissolution Procedures	FBI, NIST	Microwave digestion, Parr bomb	FBI, NIST	In Progress
Elemental/Actinide Assay, U Isotopics	NIST	INAA, HR- ICP-MS	NIST (?), Other	Scoping
Modelling	NIST		NIST	Scoping
Certification & Release	NIST		NIST	No Activity

### **SUMMARY**

The projects highlighted here are some of the first reference materials to be developed for the field of nuclear forensics. Demonstrating the validity of nuclear forensic data is critical as this data could be an important component of the attribution process resulting in prosecution in a court of law or development of actionable intelligence. Thus, using a proper reference material to constrain measurement accuracy and uncertainty bolsters the veracity of measurement results and provides a strong base to stand up to legal scrutiny.