RESOURCE LETTER

Resource Letters are guides for college and university physicists, astronomers, and other scientists to literature, websites, and other teaching aids. Each Resource Letter focuses on a particular topic and is intended to help teachers improve course content in a specific field of physics or to introduce nonspecialists to this field. The Resource Letters Editorial Board meets at the AAPT Winter Meeting to choose topics for which Resource Letters will be commissioned during the ensuing year. Items in the Resource Letter below are labeled with the letter E to indicate elementary level or material of general interest to persons seeking to become informed in the field, the letter I to indicate intermediate level or somewhat specialized material, or the letter A to indicate advanced or specialized material. No Resource Letter is meant to be exhaustive and complete; in time there may be more than one Resource Letter on a given subject. A complete list by field of all Resource Letters published to date is at the website www.kzoo.edu/ajp/letters.html. Suggestions for future Resource Letters, including those of high pedagogical value, are welcome and should be sent to Professor Roger H. Stuewer, Editor, AAPT Resource Letters, School of Physics and Astronomy, University of Minnesota, 116 Church Street SE, Minneapolis, MN 55455; e-mail: rstuewer@physics.umn.edu

Resource Letter FC-1: The physics of fundamental constants

Peter J. Mohr[a] and David B. Newell[b]
National Institute of Standards and Technology, Gaithersburg, Maryland 20899

(Received 7 October 2009; accepted 7 December 2009)

This Resource Letter provides a guide to the literature on the physics of fundamental constants and their values as determined within the International System of Units (SI). Journal articles, books, and websites that provide relevant information are surveyed. Literature on redefining the SI in terms of exact values of fundamental constants is also included.
[DOI: 10.1119/1.3279700]

I. INTRODUCTION

This Resource Letter is a guide to the literature describing both experimental and theoretical works on the determination of values of certain fundamental physical constants.

In this Resource Letter, fundamental physical constants are taken to be the experimentally determined parameters in the equations that describe the basic laws of physics as they are currently understood. The numerical values of these constants are needed to make quantitative predictions by theory for comparison to the results of measurements. In fact, numerical values of the constants are periodically determined as being those that give the best agreement between theoretical predictions and experimental measurements.

Some of these constants have dimensions and others do not. For example, the speed of light, a fundamental constant associated with special relativity and electromagnetism, has the dimension of velocity, or distance divided by time. In order to give a unique meaning to the value of such a constant, it is necessary to specify the system of units in which it is expressed since the numerical value will depend on the unit definitions. For example, the speed of light is about 3×10^8 meters/second, which is the same as about 6.7×10^8 miles/hour. For such constants with dimensions, the International System of Units (SI) is widely accepted in science and technology and is presently the standard agreed to through a treaty among 54 nations. On the other hand, dimensionless constants, such as the fine-structure constant, the constant that characterizes the strength of electromagnetic interactions, are independent of the unit system.

Many of the constants considered here are of use in practical metrological applications, while others are included because they have traditionally been evaluated in the periodic least-squares adjustments of the constants. In general, these constants have values that have been accurately determined, and they have a well-defined place in a fundamental theory such as quantum mechanics or relativity. This includes universal constants that apply to broad physical laws as well as properties of particular particles such as their mass. In view of the critical role of the system of units for expressing the values of constants, literature on changes that may be made to the SI definitions in the near future is also surveyed. Not included here is the broad topic of possible time variation of the constants.

Besides the references given in this Resource Letter, relevant papers may be located on the searchable database of the NIST Fundamental Constants Data Center at http://physics.nist.gov/constantsbib.

II. BASIC RESOURCES

A. Journals

Most of the physics leading to values of the fundamental constants is reported in the following journals. The listing is in decreasing order of the number of times articles in the journal are cited in the latest adjustment of the values of the constants.

Physical Review Letters
Physical Review A
Metrologia
IEEE Transactions on Instrumentation and Measurement
Physical Review D
Physics Letters B
Canadian Journal of Physics
Zhurnal Eksperimental' noi i Teoreticheskoi Fiziki [Journal of Experimental and Theoretical Physics (JETP)]

[a]Electronic mail: mohr@nist.gov
[b]Electronic mail: dnewell@nist.gov

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

Philosophical Transactions of the Royal Society of London
Physics Letters A
International Journal of Mass Spectrometry
Reviews of Modern Physics
Measurement Science and Technology
Reports on Progress in Physics
Physical Review
Nuclear Physics B (Proceedings Supplements)
Modern Physics Letters A
Journal of Physics G
Journal of Physics B
Izmeritel'naya Tekhnika [Measurement Techniques]
Annals of Physics (N.Y.)

B. Conference proceedings

A number of conference series are focused on work that is often relevant to the physics and values of fundamental constants. Published proceedings of these conferences are listed here. Other relevant conference proceedings that are not part of a series are also given.

1. Conference on Precision Electromagnetic Measurements (CPEM)

This series is focused on measurements and theory that are of high relevance to knowledge of the fundamental constants.

1. **2008 Conference on Precision Electromagnetic Measurements (CPEM)**, edited by T. E. Lipe and Y.-H. Tang, IEEE Trans. Instrum. Meas. **58**(4) 748–1267 (2009). (A)
2. **Special issue on CPEM 2006**, edited by G. M. Reedtz, IEEE Trans. Instrum. Meas. **56**(2) 209–676 (2007). (A)
3. **Special issue on CPEM 2004**, edited by G. Jones, IEEE Trans. Instrum. Meas. **54**(2) 473–936 (2005). (A)
4. **Special issue on CPEM 2002**, edited by U. Feller, IEEE Trans. Instrum. Meas. **52**(2) 221–651 (2003). (A)
5. **Special issue on Selected Papers CPEM'2000**, edited by B. W. Ricketts, IEEE Trans. Instrum. Meas. **50**(2) 173–655 (2001). (A)
6. **Special issue on Selected Papers CPEM'98**, edited by B. A. Bell, IEEE Trans. Instrum. Meas. **48**(2) 145–671 (1999). (A)
7. **Special issue on Selected Papers CPEM'96**, edited by R. J. Cook, IEEE Trans. Instrum. Meas. **46**(2) 89–646 (1997). (A)
8. **Special issue on Selected Papers CPEM/94**, edited by M. Young, IEEE Trans. Instrum. Meas. **44**(2) 77–622 (1995). (A)
9. Editor J. McA Steele, IEEE Trans. Instrum. Meas. **42**(2) i–679 (1993). (A)
10. **Special issue on Selected Papers CPEM '90**, edited by N. B. Belecki, IEEE Trans. Instrum. Meas. **40**(2) 65–535 (1991). (A)

2. Enrico Fermi Schools on Metrology and Fundamental Constants

This series is of general interest for fundamental constants.

11. **Metrology and Fundamental Constants**, edited by T. Hänsch, S. Leschiutta, and A. J. Wallard (IOS, Amsterdam, 2007). (A)
12. **Recent Advances in Metrology and Fundamental Constants**, edited by T. J. Quinn, S. Leschiutta, and P. Tavella (IOS, Amsterdam, 2001). (A)
13. **Metrology at the Frontiers of Physics and Technology**, edited by L. Crovini and T. J. Quinn (North-Holland, Amsterdam, 1992). (A)
14. **Metrology and Fundamental Constants**, edited by A. F. Milone, P. Giacomo, and S. Leschiutta (North-Holland, Amsterdam, 1980). (A)

3. International Conference on Atomic Physics (ICAP)

These conferences cover a broader range of topics but include work relevant to the constants.

15. **Proceedings of the XXI International Conference on Atomic Physics, Pushing the Frontiers of Atomic Physics**, edited by R. Côté, P. L. Gould, M. Rozman, and W. W. Smith (World Scientific, Singapore, 2009). (A)
16. **Atomic Physics 20: XX International Conference on Atomic Physics**, edited by C. Roos, H. Häffner, and R. Blatt, AIP Conference Proceedings 869 (AIP, Melville, NY, 2006). (A)
17. **Atomic Physics 19: XIX International Conference on Atomic Physics, ICAP 2004**, edited by L. G. Marcassa, V. S. Bagnato, and K. Helmerson, AIP Conference Proceedings 770 (AIP, Melville, NY, 2005). (A)
18. **Proceedings of the XVIII International Conference on Atomic Physics: The Expanding Frontier of Atomic Physics**, edited by H. R. Sadeghpour, E. J. Heller, and D. E. Pritchard (World Scientific, Singapore, 2003). (A)
19. **Atomic Physics 17: XVII International Conference, ICAP 2000**, edited by P. De Natale, M. Inguscio, and E. Arimondo, AIP Conference Proceedings 551 (AIP, Melville, NY, 2000). (A)

4. Precision Physics of Simple Atomic Systems

As the titles indicate, these conferences are focused on precision work on atomic structure.

20. **Precision Physics of Simple Atoms and Molecules**, edited by S. G. Karshenboim, Lecuture Notes in Physics 745 (Springer, Berlin, 2008). (A)
21. **Precision Physics of Simple Atomic Systems**, edited by S. G. Karshenboim and V. B. Smirnov, Lecture Notes in Physics 627 (Springer, Berlin, 2003. (A)
22. **The Hydrogen Atom: Precision Physics of Simple Atomic Systems**, edited by S. G. Karshenboim, F. S. Pavone, F. Bassani, M. Inguscio, and T. W. Hänsch, Lecture Notes in Physics 570 (Springer, Berlin, 2001). (A)

5. Other conferences

23. **The Fundamental Constants of Physics, Precision Measurements and the Base Units of the SI**, edited by T. Quinn and K. Burnett, Philos. Trans. R. Soc. London, Ser. A **363**(1834), 2097–2327 (2005). (A)

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

24. **Gravitational Measurements, Fundamental Metrology and Constants**, edited by V. De Sabbata and V. N. Melnikov, NATO ASI Series C, Vol. 230 (Kluwer Academic, Dordrecht, 1988). (A)
25. **Precision Measurement and Fundamental Constants II**, edited by B. N. Taylor and W. D. Phillips, NBS Special Publication 617 (U.S. Government Printing Office, Washington, DC, 1984). (A)
26. **Quantum Metrology and Fundamental Physical Constants**, edited by P. H. Cutler and A. A. Lucas, NATO ASI Series B Vol. 98 (Plenum, New York, 1983). (A)

C. Books and monographs

27. H. Fritzsch, **The Fundamental Constants: A Mystery of Physics** (World Scientific, Singapore, 2009). (I)
28. **Quantum Metrology and Fundamental Constants**, edited by F. Piquemal and B. Jeckelmann, Eur. Phys. J. Special Topics **172**, 1–408 (2009). (A)
29. **Atomic Clocks and Fundamental Constants**, edited by S. G. Karshenboim and E. Peik, Eur. Phys. J. Special Topics **163**, 1–332 (2008). (A)
30. J.-P. Uzan and B. Leclercq, **The Natural Laws of the Universe: Understanding Fundamental Constants** (Springer, Berlin, 2008). (I)
31. **Astrophysics, Clocks and Fundamental Constants**, edited by S. G. Karshenboim and E. Peik, Lecture Notes in Physics 648 (Springer, Berlin, 2004). (A)
32. J. D. Barrow, **The Constants of Nature** (Random House, New York, 2003). (I)
33. H. Bachmair *et al.*, **Fundamental Constants in Physics and Chemistry (Numerical Data and Functional Relationships in Science and Technology)** (Springer, Berlin, 1992). (A)
34. B. W. Petley, **The Fundamental Physical Constants and the Frontier of Measurement** (Adam Hilger, Bristol, 1985). (A)
35. B. N. Taylor, W. H. Parker, and D. N. Langenberg, **The Fundamental Constants and Quantum Electrodynamics** (Academic, New York, 1969). (A)

D. Electronic archives and websites

Papers on fundamental physical constants may be found in various electronic archives. The more frequently used ones are the following:

36. <http://arxiv.org/archive/physics.gen-ph/GeneralPhysics> (A)
37. <http://arxiv.org/archive/physics.atom-ph/AtomicPhysics> (A)
38. <http://arxiv.org/archive/astro-ph/Astrophysics> (A)
39. <http://arxiv.org/archive/hep-th/HighEnergyPhysics-Theory>. (A)
40. <http://arxiv.org/archive/hep-ph/HighEnergyPhysics-Phenomenology>. (A)
41. <http://arxiv.org/archive/gr-qc/GeneralRelativityandQuantumCosmology> (A)

Relevant websites are the following:

42. ⟨http://www.aps.org/units/gpmfc/⟩, The website for the American Physical Society's Topical Group on Precision Measurement and Fundamental Constants. (E)
43. ⟨http://www.codata.org/taskgroups/TGfundconst/⟩, A description of the CODATA Task Group on Fundamental Physical constants within the home page of the Committee on Data for Science and Technology (CODATA). (E)
44. ⟨http://www.bipm.org/extra/codata/⟩, The home page of the CODATA Task Group on Fundamental Physical constants within the home page of the International Bureau of Weights and Measures (BIPM). (E)
45. ⟨http://physics.nist.gov/cuu/⟩, The National Institute of Standards and Technology Reference on Constants, Units, and Uncertainty. (E)
46. ⟨http://amdc.in2p3.fr/web/amdcw_en.html⟩, The home page for the Atomic Mass Data Center which provides the 2003 Atomic Mass Evaluation. (I)
47. ⟨http://www.iupap.org/commissions/c2/⟩, The website of the International Union of Pure and Applied Physics (IUPAP) Commission on Symbols, Units, Nomenclature, Atomic Masses and Fundamental Constants (SUNAMCO). (E)

III. SURVEYS OF CONSTANTS AND UNITS

A. Overviews of the constants

48. "Adjusted recommended values of the fundamental physical constants," S. G. Karshenboim, Eur. Phys. J. Spec. Top. **172**, 385–397 (2009). (I)
49. "Universal constants, standard models and fundamental metrology," G. Cohen-Tannoudji, Eur. Phys. J. Spec. Top. **172**, 5–24 (2009). (I)
50. "New recommended values of the fundamental physical constants (CODATA 2006)," S. G. Karshenboim, Phys. Usp. **51**(10), 1019–1026 (2008). (I)
51. "The fundamental physical constants," P. J. Mohr, B. N. Taylor, and D. B. Newell, Phys. Today **60**(7), 52–55 (2007). (I)
52. "Fundamental physical constants: Looking from different angles," S. G. Karshenboim, Can. J. Phys. **83**(8), 767–811 (2005). (I)
53. "Adjusting the Values of the Fundamental Constants," P. J. Mohr and B. N. Taylor, Phys. Today **54**(3), 29–34 (2001). (I)

B. Evaluations of fundamental constants

Starting with the pioneering work of Raymond Birge at the University of California, Berkeley, in 1929, various evaluations of the fundamental constants, based on the data available at the time, have been made. The Committee on Data for Science and Technology (CODATA) Task Group on Fundamental Constants was founded in 1969, and since then, they have recommended values for the constants that are internationally recognized and used in research and publications.

54. "CODATA recommended values of the fundamental physical constants: 2006," P. J. Mohr, B. N. Taylor, and

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

D. B. Newell, J. Phys. Chem. Ref. Data **37**(3), 1187–1284 (2008). (A)
55. "CODATA recommended values of the fundamental physical constants: 2006," P. J. Mohr, B. N. Taylor, and D. B. Newell, Rev. Mod. Phys. **80**(2), 633–730 (2008). (A)
56. "CODATA recommended values of the fundamental physical constants: 2002," P. J. Mohr and B. N. Taylor, Rev. Mod. Phys. **77**(1), 1–107 (2005). (A)
57. "CODATA recommended values of the fundamental physical constants: 1998," P. J. Mohr and B. N. Taylor, Rev. Mod. Phys. **72**(2), 351–495 (2000). (A)
58. "CODATA recommended values of the fundamental physical constants: 1998," P. J. Mohr and B. N. Taylor, J. Phys. Chem. Ref. Data **28**(6), 1713–1852 (1999). (A)
59. "Recommended values of the fundamental physical constants: A status report," B. N. Taylor and E. R. Cohen, J. Res. Natl. Inst. Stand. Technol. **95**(5), 497–523 (1990). (A)
60. "The 1986 adjustment of the fundamental physical constants," E. R. Cohen and B. N. Taylor, Rev. Mod. Phys. **59**(4), 1121–1148 (1987). (A)
61. "The 1973 least-squares adjustment of the fundamental constants," E. R. Cohen and B. N. Taylor, J. Phys. Chem. Ref. Data **2**(4), 663–734 (1973). (A)
62. "Determination of e/h using macroscopic quantum phase coherence in superconductors: Implications for quantum electrodynamics and the fundamental physical constants," B. N. Taylor, W. H. Parker, and D. N. Langenberg, Rev. Mod. Phys. **41**(3), 375–496 (1969). (A)
63. "Our knowledge of the fundamental constants of physics and chemistry in 1965," E. R. Cohen and J. W. M. DuMond, Rev. Mod. Phys. **37**(4), 537–594 (1965). (A)
64. "Status of knowledge of the fundamental constants of physics and chemistry as of January 1959," J. W. M. DuMond, Ann. Phys. (N.Y.) **7**(4), 365–403 (1959). (A)
65. "Résumé of atomic constants," J. A. Bearden and J. S. Thomsen, Am. J. Phys. **27**(8), 569–576 (1959). (A)
66. "A survey of the systematic evaluation of the universal physical constants," R. T. Birge, Nuovo Cimento, Suppl. **6**(1), 39–67 (1957). (A)
67. "A survey of atomic constants," J. A. Bearden and J. S. Thomsen, Nuovo Cimento, Suppl. **5**(2), 267–360 (1957). (A)
68. "Present status of the atomic constants," J. A. Bearden, M. D. Earle, J. M. Minkowski, and J. S. Thomsen, Phys. Rev. **93**(3), 629–630 (1954). (A)
69. "Least-squares adjustment of the atomic constants, 1952," J. W. M. DuMond and E. R. Cohen, Rev. Mod. Phys. **25**(3), 691–708 (1953). (A)
70. "A Re-evaluation of the fundamental atomic constants," J. A. Bearden and H. M. Watts, Phys. Rev. **81**(1), 73–81 (1951). (A)
71. "Our knowledge of the atomic constants, F, N, m, and h in 1947, and of other constants derivable therefrom," J. W. M. DuMond and E. R. Cohen, Rev. Mod. Phys. **20**(1), 82–108 (1948); Erratum **21**(4), 651–652 (1949).
72. "Probable values of the general physical constants," R. T. Birge, Rev. Mod. Phys. **1**(1), 1–73 (1929). (A)

C. The International System of Units (SI)

The following are various resources describing the International System of Units and related concepts:

73. **International System of Units (SI)**, 8th ed. (Bureau International des Poids et Mesures, Sèvres, France, 2006), ⟨http://www.bipm.org/en/si/si_brochure/⟩. (I)
74. B. N. Taylor and A. Thompson, **The International System of Units (SI)**, 2008 ed., NIST Spec. Pub. 330 (U.S. Government Printing Office, Washington, DC, 2008) . (I)
75. A. Thompson and B. N. Taylor, **Guide for the Use of the International System of Units (SI)**, 2008 ed., NIST Spec. Publ. 811 (U.S. Government Printing Office, Washington, DC, 2008). (I)
76. E. R. Cohen, T. Cvitaš, J. G. Frey, B. Holmström, K. Kuchitsu, R. Marquardt, I. Mills, F. Pavese, M. Quack, J. Stohner, H. L. Strauss, M. Takami, and A. J. Thor, **Quantities, Units and Symbols in Physical Chemistry, IUPAC (International Union of Pure and Applied Chemistry)**, 3rd ed. (Royal Society of Chemistry, Cambridge, 2007). (I)
77. E. R. Cohen and P. Giacomo, **Symbols, Units, Nomenclature and Fundamental Constants in Physics, IUPAP-25 (IUPAP-SUNAMCO 87-1)** (International Union of Pure and Applied Physics, 1987); also published in Physica (Utrecht) **146A**(1–2), 1–68 (1987). (I)
See also Ref. 469.

IV. EXACT CONSTANTS

A number of fundamental constants have exact values either by the definition of the constant itself or as a consequence of the definitions of the SI units. Because of the possible dependence on SI unit definitions, which are modified from time to time, the set of constants that are exact may also change.

A. Speed of light c

The primary example of an exact constant is the speed of light $c = 299\,792\,458$ m/s. Before 1983, the speed of light was a measured quantity based on the meter and the second, which were defined independently. In 1983, the SI meter was redefined to be the distance light travels in a certain interval of time. From then on, the speed of light has been exactly one meter divided by that specified time interval. This definition takes into account the result of special relativity, that the speed of light in vacuum is the same in all inertial frames. This redefinition represented a new concept in units in which a general property of nature, namely, the invariability of the speed of light, could be used as the basis for measuring and calibrating any other velocity or, in combination with the definition of the second, as the basis for defining the meter.

B. Hyperfine frequency in cesium ν_{Cs}

Another constant that is exact as the result of a definition of a unit is the ground-state hyperfine frequency in cesium $\nu_{Cs} = 9\,192\,631\,770$ Hz. This constant is qualitatively different from the speed of light in that it refers to a particular property of a particular atom. As such, it is not a universal constant like c. In the SI, the second is defined to be the period of time for a certain number of cycles of this transition frequency to elapse. In view of this definition, the hyperfine frequency is an exact number. Other frequencies are determined by comparing them to ν_{Cs}.

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

C. Magnetic constant μ_0 and electric constant ϵ_0

The magnetic constant, or the permeability of free space, is fixed by the SI definition of the ampere to be $\mu_0 = 4\pi \times 10^{-7}$ N A^{-2}. From electromagnetic theory in SI units, the electric constant, or permittivity of free space, is $\epsilon_0 = 1/\mu_0 c^2$ so that ϵ_0 is also exact.

V. DIRECTLY MEASURED CONSTANTS

As mentioned in the introduction, the values of the fundamental constants are obtained by comparing experimental results to the predictions of theory in which the constants appear as parameters. The best values of the constants are taken to be those for which the theoretical predictions best match the experimental results as determined by the method of least squares. That work and its methodology are described in the papers cited in Sec. III B. In this section, references for the more recent theoretical and experimental work relevant to the determination of the values of particular constants are given and organized according to the constant to which the research is relevant. The focus is on recent papers and some older papers that are of particular significance.

A. Fine-structure constant α

The most accurate determination of the fine-structure constant α is from the combination of measurements of the anomalous magnetic moment of the electron, a_e, and quantum electrodynamics (QED) theory. Also given are determinations of α from atomic recoil, neutron diffraction, and electromagnetic measurements.

1. Electron anomalous magnetic moment

The QED part of the theoretical expression for the anomalous magnetic moment of the electron a_e is

$$a_e(\mathrm{QED}) = C_e^{(2)}\left(\frac{\alpha}{\pi}\right) + C_e^{(4)}\left(\frac{\alpha}{\pi}\right)^2 + C_e^{(6)}\left(\frac{\alpha}{\pi}\right)^3 + C_e^{(8)}\left(\frac{\alpha}{\pi}\right)^4 + \ldots .$$

There are additional contributions from weak and strong interaction effects, but these are relatively small. A value for α may be obtained by equating the theoretical expression to the measured a_e.

The first coefficient $C_e^{(2)}$ was calculated about 60 years ago by Schwinger, and subsequent work has led to values for the other coefficients. Work is still underway refining the value of $C_e^{(8)}$, and calculations have only begun on the next term in the series.

78. "Renormalization of QED and its experimental test over 60 years," T. Kinoshita, Prog. Theor. Phys. Suppl. **167**, 62–75 (2007). (I)

79. "Automated calculation scheme for α^n contributions of QED to lepton $g-2$: New treatment of infrared divergence for diagrams without lepton loops," T. Aoyama, M. Hayakawa, T. Kinoshita, and M. Nio, Nucl. Phys. B **796**(1–2), 184–210 (2008). (A)

80. "Revised value of the eighth-order QED contribution to the anomalous magnetic moment of the electron," T. Aoyama, M. Hayakawa, T. Kinoshita, and M. Nio, Phys. Rev. D **77**, 053012 (2008), 24 pp. (A)

81. "Tenth-order lepton anomalous magnetic moment: Second-order vertex containing two vacuum polarization subdiagrams, one within the other," T. Aoyama, M. Hayakawa, T. Kinoshita, and M. Nio, Phys. Rev. D **78**, 113006 (2008), 7 pp. (A)

82. "Eighth-order vacuum-polarization function formed by two light-by-light-scattering diagrams and its contribution to the tenth-order electron $g-2$," T. Aoyama, M. Hayakawa, T. Kinoshita, M. Nio, and N. Watanabe, Phys. Rev. D **78**, 053005 (2008), 14 pp. (A)

83. "Revised value of the eighth-order contribution to the electron $g-2$," T. Aoyama, M. Hayakawa, T. Kinoshita, and M. Nio, Phys. Rev. Lett. **99**, 110406 (2007), 4 pp. (A)

84. "Precise mass-dependent QED contributions to leptonic $g-2$ at order α^2 and α^3," M. Passera, Phys. Rev. D **75**, 013002 (2007), 6 pp. (A)

85. "Improved α^4 term of the electron anomalous magnetic moment," T. Kinoshita and M. Nio, Phys. Rev. D **73**, 013003 (2006), 28 pp. (A)

86. "New results on the hadronic contributions to $\alpha(M_Z^2)$ and to $(g-2)_\mu$," M. Davier and A. Höcker, Phys. Lett. B **435**(3–4), 427–440 (1998). (A)

87. "Higher-order hadronic contributions to the anomalous magnetic moment of leptons," B. Krause, Phys. Lett. B **390**(1–4), 392–400 (1997). (A)

88. "The analytical value of the electron $(g-2)$ at order α^3 in QED," S. Laporta and E. Remiddi, Phys. Lett. B **379**(1–4), 283–291 (1996). (A)

89. "The analytical value of the corner-ladder graphs contribution to the electron $(g-2)$ in QED," S. Laporta, Phys. Lett. B **343**(1–4), 421–426 (1995). (A)

90. "Progress in the analytical evaluation of the electron $(g-2)$ in QED: The scalar part of the triple-cross graphs," S. Laporta and E. Remiddi, Phys. Lett. B **356**(2–3), 390–397 (1995). (A)

91. "Analytical value of some sixth-order graphs to the electron $g-2$ in QED," S. Laporta, Phys. Rev. D **47**(10), 4793–4795 (1993). (A)

92. "The Analytical Contribution of the sixth-order graphs with vacuum polarization insertions to the muon $(g-2)$ in QED," S. Laporta, Nuovo Cimento **106**(5), 675–683 (1993). (A)

93. "Precise mass-ratio dependence of fourth-order lepton anomalous magnetic moments: Effect of a new measurement of m_τ," G. Li, R. Mendel, and M. A. Samuel, Phys. Rev. D **47**(4), 1723–1725 (1993). (A)

94. "The analytic value of the light-light vertex graph contributions to the electron $g-2$ in QED," S. Laporta and E. Remiddi, Phys. Lett. B **265**, 182–184 (1991). (A)

95. "Improved analytic theory of the muon anomalous magnetic moment," M. A. Samuel and G. Li, Phys. Rev. D **44**(12), 3935–3942 (1991); Erratum **48**(4), 1879–1881 (1993). (A)

96. R. Z. Roskies, E. Remiddi, and M. J. Levine, "Analytic evaluation of sixth-order contributions to the electron's g factor," in **Quantum Electrodynamics**, edited by T. Kinoshita (World Scientific, Singapore, 1990), Chap. 6, pp. 162–217. (A)

97. T. Kinoshita, "Theory of the anomalous magnetic moment of the electron—numerical approach," in **Quantum Electrodynamics**, edited by T. Kinoshita (World

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

Scientific, Singapore, 1990), Chap. 7, pp. 218–321. (A)
98. "Fourth-order magnetic moment of the electron," A. Petermann, Nucl. Phys. **5**(4), 677–683 (1958). (A)
99. "The magnetic moment of the electron," C. M. Sommerfield, Ann. Phys. (N.Y.) **5**(1), 26–57 (1958). (A)
100. "Fourth-order magnetic moment of the electron," A. Petermann, Helv. Phys. Acta **30**(5), 407–408 (1957). (A)
101. "Magnetic dipole moment of the electron," C. M. Sommerfield, Phys. Rev. **107**(1), 328–329 (1957). (A)
102. "Quantum electrodynamics III. The electromagnetic properties of the electron—radiative corrections to scattering," J. Schwinger, Phys. Rev. **76**(6), 790–817 (1949). (A)
103. "On quantum-electrodynamics and the magnetic moment of the electron," J. Schwinger, Phys. Rev. **73**(4), 416–417 (1948). (A)

Accurate measurements of the anomalous magnetic moment of the electron are made by confining electrons in a Penning trap.

104. "The anomalous magnetic moment of the electron," M. Vogel, Contemp. Phys. **50**(3), 437–452 (2009). (I)
105. "Why is sideband mass spectrometry possible with ions in a penning trap?," G. Gabrielse, Phys. Rev. Lett. **102**, 172501 (2009), 4 pp. (A)
106. "The true cyclotron frequency for particles and ions in a Penning trap," G. Gabrielse, Int. J. Mass Spectrom. **279**(2–3), 107–112 (2009). (A)
107. "New measurement of the election magnetic moment and the fine structure constant," D. Hanneke, S. Fogwell, and G. Gabrielse, Phys. Rev. Lett. **100**, 120801 (2008), 4 pp. (A)
108. "New determination of the fine structure constant from the electron g value and QED," G. Gabrielse, D. Hanneke, T. Kinoshita, M. Nio, and B. Odom, Phys. Rev. Lett. **97** 030802 (2006), 4 pp.; Erratum **99**, 039902 (2007), 2 pp. (A)
109. "New measurement of the electron magnetic moment using a one-electron quantum cyclotron," B. Odom, D. Hanneke, B. D'Urso, and G. Gabrielse, Phys. Rev. Lett. **97**, 030801 (2006), 4 pp. (A)
110. "Nobel lecture: Experiments with an isolated subatomic particle at rest," H. Dehmelt, Rev. Mod. Phys. **62**(3), 525–530 (1990). (I)
111. "New high-precision comparison of electron and positron g factors," R. S. Van Dyck, Jr., P. B. Schwinberg, and H. G. Dehmelt, Phys. Rev. Lett. **59**(1), 26–29 (1987). (A)

2. Quotient of Planck constant and particle mass experiments

The quotient of the Planck constant h to the mass of a cesium or rubidium atom has been obtained by measuring the frequency shift due to recoil of the atoms when photons are absorbed or emitted. The quotient of the Planck constant to the mass of the neutron has been obtained by Bragg reflection of neutrons from a crystal, which gives the neutron de Broglie wavelength. All of these measurements lead to a value of the fine-structure constant from the definition of the Rydberg constant, given in Sec. V C, by writing

$$\alpha^2 = \frac{2R_\infty}{c}\frac{m_x}{m_e}\frac{h}{m_x},$$

where m_x is the mass of the particle in the measurement and m_e is the mass of the electron. The Rydberg constant and the mass ratios are very accurately known, so the quotient gives a value for α limited mainly by the accuracy of the measurement of h/m_x.

112. "Combination of Bloch oscillations with a Ramsey–Bordé interferometer: New determination of the fine structure constant," M. Cadoret, E. de Mirandes, P. Cladé, S. Guellati-Khélifa, C. Schwob, F. Nez, L. Julien, and R. Biraben, Phys. Rev. Lett. **101**, 230801 (2008), 4 pp. (A)
113. "Precise measurement of h/m_{Rb} using Bloch oscillations in a vertical optical lattice: Determination of the fine-structure constant," P. Cladé, E. de Mirandes, M. Cadoret, S. Guellati-Khélifa, C. Schwob, F. Nez, L. Julien, and F. Biraben, Phys. Rev. A **74**, 052109 (2006), 18 pp. (A)
114. "Determination of the fine structure constant based on Bloch oscillations of ultracold atoms in a vertical optical lattice," P. Cladé, E. de Mirandes, M. Cadoret, S. Guellati-Khélifa, C. Schwob, F. Nez, L. Julien, and F. Biraben, Phys. Rev. Lett. **96**, 033001 (2006), 4 pp. (A)
115. "A new photon recoil experiment: towards a determination of the fine structure constant," H. Müller, S.-W. Chiow, Q. Long, C. Vo, and S. Chu, Appl. Phys. B: Lasers Opt. **84**, 633–642 (2006). (A)
116. "A preliminary measurement of the fine structure constant based on atom interferometry," A. Wicht, J. M. Hensley, E. Sarajlic, and S. Chu, Phys. Scr. **T102**, 82–88 (2002). (A)
117. "Re-evaluation of a precise measurement of h/m_n," E. Krüger, W. Nistler, and W. Weirauch, Metrologia **36**(2), 147–148 (1999). (A)

3. Quantum Hall effect experiments and the Thompson–Lampard theorem

When calibrated with an independent value for the ohm, the quantum Hall effect provides a condensed-matter determination of α. In terms of the von Klitzing constant R_K, α is given by

$$\alpha = \frac{\mu_0 c}{2R_K},$$

(see also Sec. VI). An accurate SI value of the ohm can be determined by means of a cross capacitor and a theorem in electrostatics developed by Thompson and Lampard that allows the capacitance per unit length to be calculated.

118. "Determination of the unit of resistance and the von Klitzing constant R_K based on a calculable capacitor," H. Bachmair, Eur. Phys. J. Spec. Top. **172**, 257–266 (2009). (I)
119. "A curious and useful theorem in two-dimensional

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

electrostatics," J. D. Jackson, Am. J. Phys. **67**(2), 107–115 (1999). (I)
120. "A primer on electrical units in the Système International," N. M. Zimmerman, Am. J. Phys. **66**(4), 324–331 (1998). (I)
121. "Determination of the von Klitzing constant R_K in terms of the BNM calculable capacitor—Fifteen years of investigations," G. Trapon, O. Thévenot, J. C. Lacueille, and W. Poirier, Metrologia **40**(4), 159–171 (2003). (A)
122. "Determination of the von Klitzing constant and the fine-structure constant through a comparison of the quantized Hall resistance and the ohm derived from the NIST calculable capacitor," A. Jeffery, R. E. Elmquist, J. Q. Shields, L. H. Lee, M. E. Cage, S. H. Shields, and R. F. Dziuba, Metrologia **35**(2), 83–96 (1998). (A)
123. "NIST comparison of the quantized Hall resistance and the realization of the SI OHM through the calculable capacitor," A.-M. Jeffery, R. E. Elmquist, L. H. Lee, J. Q. Shields, and R. F. Dziuba, IEEE Trans. Instrum. Meas. **46**(2), 264–268 (1997). (A)
124. "A new determination of the quantized Hall resistance in terms of the NML calculable cross capacitor," G. W. Small, B. W. Ricketts, P. C. Coogan, B. J. Pritchard, and M. M. R. Sovierzoski, Metrologia **34**(3), 241–243 (1997). (A)
125. "A determination of the SI value of QHR by the cross-capacitor in NIM," Z. Zhang, X. Wang, D. Wang, X. Li, Q. He, and Y. Ruan, Acta Metrologia Sinica **16**(1), 1–5 (1995). (A)
126. "SI value of quantized Hall resistance based on ETL's calculable capacitor," K. Shida, T. Wada, H. Nishinaka, K. Segawa, and T. Igarashi, IEEE Trans. Instrum. Meas. **38**(2), 252–255 (1989). (A)
127. "The relationship among the SI ohm, the ohm at NPL, and the quantized Hall resistance," A. Hartland, R. G. Jones, B. P. Kibble, and D. J. Legg, IEEE Trans. Instrum. Meas. **IM-36**(2), 208–213 (1987). (A)
128. "Absolute determination of the farad and the ohm, and measurement of the quantized Hall resistance $R_H(2)$ at LCIE," F. Delahaye, A. Fau, D. Dominguez, and M. Bellon, IEEE Trans. Instrum. Meas. **IM-36**(2), 205–207 (1987). (A)
129. "The cylindrical cross-capacitor as a calculable standard," A. M. Thompson, Proc. Inst. Electr. Eng. **106B**, 307–310 (1959). (A)
130. "A new theorem in electrostatics with applications to calculable standards of capacitance," D. G. Lampard, Proc. Inst. Electr. Eng. **104C**, 271–280 (1957). (A)
131. "A new theorem in electrostatics and its application to calculable standards of capacitance," A. M. Thompson and D. G. Lampard, Nature (London) **177**, 888–890 (1956). (A)

4. Low-field gyromagnetic ratio experiments

The gyromagnetic ratio γ' of a bound particle with a magnetic moment μ can be determined using a low magnetic field generated by a precision single-layer solenoid carrying an electric current, where the dimensions of the solenoid and the current are accurately measured. The results can be related to the fine-structure constant and other constants by

$$\alpha^3 = \frac{4\mu_0 R_\infty}{g_e K_J R_K}\frac{\mu_e}{\mu}\gamma',$$

where R_∞ is the Rydberg constant, μ_e is the magnetic moment of the electron, g_e is the g-factor of the electron, and K_J is the Josephson constant.

132. "The proton gyromagnetic ratio by low field method measured at KRISS," P. G. Park, C. H. Choi, C. S. Kim, V. Ya. Shifrin, and V. N. Khorev, J. Korean Phys. Soc. **34**(4), 327–332 (1999). (A)
133. "Determination of the tesla-to-ampere ratio for the KRISS/VNIIM γ'_p-experiment," V. Y. Shifrin, P. G. Park, V. N. Khorev, C. H. Choi, and S. Lee, IEEE Trans. Instrum. Meas. **48**(2), 196–199 (1999). (A)
134. "A new low-field determination of the proton gyromagnetic ratio in water," V. Ya. Shifrin, P. G. Park, V. N. Khorev, C. H. Choi, and C. S. Kim, IEEE Trans. Instrum. Meas. **47**(3), 638–643 (1998). (A)
135. "Low-field method for measuring proton gyromagnetic ratio," C. G. Kim, B. C. Woo, P. G. Park, K. S. Ryu, and C. S. Kim, IEEE Trans. Instrum. Meas. **44**(2), 484–487 (1995). (A)
136. "A recent determination for the SI values of γ'_p and $2e/h$ at NIM," R. Liu, H. Liu, T. Jin, Z. Lu, X. Du, S. Xue, J. Kong, B. Yu, X. Zhou, T. Liu, and W. Zhang, Acta Metrologia Sinica **16**(3), 161–168 (1995). (A)
137. "A low field determination of the proton gyromagnetic ratio in water," E. R. Williams, G. R. Jones, Jr., S. Ye, R. Liu, H. Sasaki, P. T Olsen, W. D. Phillips, and H. P. Layer, IEEE Trans. Instrum. Meas. **38**(2), 233–237 (1989). (A)

B. Newtonian constant of gravitation *G*

The Newtonian constant of gravitation G is the proportionality constant in the universal law of gravitational attraction force F between two masses m_1 and m_2 separated by a distance r

$$F = G\frac{m_1 m_2}{r^2}.$$

Mainly due to the intrinsically weak nature of gravity, the Newtonian constant continues to be a relatively poorly determined physical constant.

138. "Newton's constant and the twenty-first century laboratory," C. C. Speake, Philos. Trans. R. Soc. London, Ser. A **363**(1834), 2265–2287 (2005). (I)
139. "Precision gravimetry with atomic sensors," M. de Angelis, A. Bertoldi, L. Cacciapuoti, A. Giorgini, G. Lamporesi, M. Prevedelli, G. Saccorotti, F. Sorrentino, and G. M. Tino, Meas. Sci. Technol. **20**, 022001 (2009), 16 pp. (I)
140. "Determination of the Newtonian gravitational constant G with time-of-swing method," J. Luo, Q. Liu, L.-C. Tu, C.-G. Shao, L.-X. Liu, S.-Q. Yang, Q. Li, and Y.-T. Zhang, Phys. Rev. Lett. **102**, 240801 (2009), 4 pp. (A)
141. "Determination of the Newtonian gravitational constant using atom interferometry," G. Lamporesi, A. Bertoldi, L. Cacciapuoti, M. Prevedelli, and G. M. Tino, Phys. Rev. Lett. **100**, 050801 (2008), 4 pp. (A)

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

142. "Atom interferometer measurement of the Newtonian constant of gravity," J. B. Fixler, G. T. Foster, J. M. McGuirk, and M. A. Kasevich, Science **315**(5808), 74–77 (2007). (A)
143. "Measurement of Newton's gravitational constant," St. Schlamminger, E. Holzschuh, W. Kündig, F. Nolting, R. E. Pixley, J. Schurr, and U. Straumann, Phys. Rev. D **74**, 082001 (2006), 25 pages. (A)
144. "Correction of source mass effects in the HUST-99 measurement of G," Z.-K. Hu, J.-Q. Guo, and J. Luo, Phys. Rev. D **71**, 127505 (2005), 3 pp. (A)
145. "New measurement of G using the measurement standards laboratory torsion balance," T. R. Armstrong and M. P. Fitzgerald, Phys. Rev. Lett. **91**, 201101 (2003), 4 pp. (A)
146. "A new determination of G using two methods," T. J. Quinn, C. C. Speake, S. J. Richman, R. S. Davis, and A. Picard, Phys. Rev. Lett. **87**, 111101 (2001), 4 pp. (A)
147. "Measurement of Newton's constant using a torsion balance with angular acceleration feedback," J. H. Gundlach and S. M. Merkowitz, Phys. Rev. Lett. **85**(14), 2869–2872 (2000). (A)
148. "A cryogenic torsion pendulum: Progress report," M. K. Bantel and R. D. Newman, Class. Quantum Grav. **17**, 2313–2318 (2000). (A)
149. "Absolute measurement of the Newtonian force and a determination of G," U. Kleinevoß, H. Meyer, A. Schumacher, and S. Hartmann, Meas. Sci. Technol. **10**(6), 492–494 (1999). (A)
150. "A free-fall determination of the Newtonian Constant of gravity," J. P. Schwarz, D. S. Robertson, T. M. Niebauer, and J. E. Faller, Science **282**(5397), 2230–2234 (1998). (A)
151. "Preliminary results of a determination of the Newtonian constant of gravitation: A test of the Kuroda hypothesis," C. H. Bagley and G. G. Luther, Phys. Rev. Lett. **78**(16), 3047–3050 (1997). (A)
152. "Measurement of the gravitational constant with a torsion balance," O. V. Karagioz and V. P. Izmailov, Meas. Tech. **39**(10), 979–987 (1996). (A)

C. Rydberg constant R_∞

The Rydberg constant is obtained by comparing theory and experiment for transition frequencies in hydrogen and deuterium atoms. Many of the recent experiments are based on use of a laser frequency comb to make precise frequency measurements. The quantity actually measured is cR_∞, which has the dimension of frequency. The Rydberg constant is related to other constants by

$$R_\infty = \frac{\alpha^2 m_e c}{2h}.$$

1. Theory

The frequency is expressed as cR_∞ times a factor that depends on the detailed theory, including effects from the finite size of the nucleus and quantum electrodynamics (QED). The papers cited in this section describe the various calculations that contribute to that factor.

153. M. I. Eides, H. Grotch, and V. A. Shelyuto, **Theory of Light Hydrogenic Bound States**, Springer Tracts in Modern Physics, Vol. 222 (Springer, Berlin, 2007). (I)
154. "Precision physics of simple atoms: QED tests, nuclear structure and fundamental constants," S. G. Karshenboim, Phys. Rep. **422**(1–2), 1–63 (2005). (I)
155. "Precise calculation of transition frequencies of hydrogen and deuterium based on a least-squares analysis," U. D. Jentschura, S. Kotochigova, E.-O. Le Bigot, P. J. Mohr, and B. N. Taylor, Phys. Rev. Lett. **95**, 163003 (2005), 4 pp. (A)
156. "Theory of light hydrogenlike atoms," M. I. Eides, H. Grotch, and V. A. Shelyuto, Phys. Rep. **342**(2–3), 63–261 (2001). (I)
157. J. R. Sapirstein and D. R. Yennie, "Theory of hydrogenic bound states," in **Quantum Electrodynamics**, edited by T. Kinoshita (World Scientific, Singapore, 1990), Chap. 12, pp. 560–672. (I)
158. "Recent developments in the comparison between theory and experiments in quantum electrodynamics," B. E. Lautrup, A. Peterman, and E. de Rafael, Phys. Rep. **3**(4), 193–259 (1972). (I)

Dirac energy with mass corrections:

159. "Reduction of relativistic two-particle wave equations to approximate forms. III," W. A. Barker and F. N. Glover, Phys. Rev. **99**(1), 317–324 (1955). (A)

Relativistic recoil corrections:

160. "Higher-order recoil corrections to energy levels of two-body systems," K. Pachucki and S. G. Karshenboim, Phys. Rev. A **60**(4), 2792–2798 (1999). (A)
161. "$O(m\alpha^7 \ln^2 \alpha)$ corrections to positronium energy levels," K. Melnikov and A. Yelkhovsky, Phys. Lett. B **458**(1), 143–151 (1999). (A)
162. "Relativistic recoil correction to hydrogen energy levels," V. M. Shabaev, A. N. Artemyev, T. Beier, and G. Soff, J. Phys. B **31**(8), L337–L339 (1998). (A)
163. "Recoil corrections of order $(Z\alpha)^6(m/M)m$ to the hydrogen energy levels recalculated," M. I. Eides and H. Grotch, Phys. Rev. A **55**(5), 3351–3360 (1997). (A)
164. "Order $(Z\alpha)^4(m/M)R_\infty$ correction to the hydrogen energy levels," A. S. Elkhovskiĭ, JETP **83**(2), 230–235 (1996). (A)
165. "Higher-order binding corrections to the Lamb shift of 2P states," U. Jentschura and K. Pachucki, Phys. Rev. A **54**(3), 1853–1861 (1996). (A)
166. "Pure recoil corrections to hydrogen energy levels," K. Pachucki and H. Grotch, Phys. Rev. A **51**(3), 1854–1862 (1995). (A)
167. "Order $\alpha^4(m/M)R_\infty$ corrections to hydrogen P levels," É. A. Golosov, A. S. Elkhovskiĭ, A. I. Mil'shteĭn, and I. B. Khriplovich, JETP **80**(2), 208–211 (1995). (A)
168. "Energy levels of one-electron atoms," G. W. Erickson, J. Phys. Chem. Ref. Data **6**(3), 831–869 (1977). (A)

Nuclear polarization corrections:

169. "Comment on 'Proton polarization shifts in electronic and muonic hydrogen'," I. B. Khriplovich and R. A.

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

Sen'kov, Phys. Lett. B **481**(2–4), 447–449 (2000). (A)
170. "Higher-order nuclear-polarizability corrections in atomic hydrogen," J. L. Friar and G. L. Payne, Phys. Rev. C **56**(2), 619–630 (1997). (A)

Self-energy radiative correction:

171. "Precise theory of levels of hydrogen and deuterium: The one-photon self energy correction," U. J. Jentschura, S. Kotochigova, E.-O. Le Bigot, and P. J. Mohr, NIST Tech. Note 1469 (2005), 3 pp., ⟨http://physics.nist.gov/hdel⟩. (A)
172. "Nonrelativistic QED approach to the Lamb shift," U. D. Jentschura, A. Czarnecki, and K. Pachucki, Phys. Rev. A **72**, 062102 (2005), 20 pp. (A)
173. "Self-energy values for P states in hydrogen and low-Z hydrogenlike ions," U. D. Jentschura and P. J. Mohr, Phys. Rev. A **72**, 014103 (2005), 3 pp. (A)
174. "Electron self-energy for higher excited S levels," U. D. Jentschura and P. J. Mohr, Phys. Rev. A **69**, 064103 (2004), 2 pp. (A)
175. "Electron self-energy for K and L shells at low nuclear charge," U. D. Jentschura, P. J. Mohr, and G. Soff, Phys. Rev. A **63**, 042512 (2001), 19 pp. (A)
176. "Calculation of the electron self-energy for low nuclear charge," U. D. Jentschura, P. J. Mohr, and G. Soff, Phys. Rev. Lett. **82**(1), 53–56 (1999). (A)
177. "Higher-order binding corrections to the Lamb shift," K. Pachucki, Ann. Phys. (N.Y.) **226**(1), 1–87 (1993). (A)
178. "Bethe logarithms for hydrogen up to $n=20$, and approximations for two-electron atoms," G. W. F. Drake and R. A. Swainson, Phys. Rev. A **41**(3), 1243–1246 (1990). (A)
179. "Radiative level shifts, I. Formulation and lowest order Lamb shift," G. W. Erickson and D. R. Yennie, Ann. Phys. (N.Y.) **35**(2), 271–313 (1965). (A)

Vacuum polarization correction:

180. "Precise energies of highly excited hydrogen and deuterium," S. Kotochigova, P. J. Mohr, and B. N. Taylor, Can. J. Phys. **80**(11), 1373–1382 (2002). (A)
181. "Hadronic vacuum polarization and the Lamb shift," J. L. Friar, J. Martorell, and D. W. L. Sprung, Phys. Rev. A **59**(5), 4061–4063 (1999). (A)
182. "Muonic vacuum polarization contribution to the energy levels of atomic hydrogen," S. G. Karshenboim, J. Phys. B **28**(4), L77–L79 (1995). (A)
183. "Energy levels of hydrogen-like atoms predicted by quantum electrodynamics, $10 \leq Z \leq 40$," P. J. Mohr, At. Data. Nucl. Data Tables **29**(3), 453–466 (1983). (A)
184. "Self-energy of the $n=2$ states in a strong Coulomb field," P. J. Mohr, Phys. Rev. A **26**(5), 2338–2354 (1982).
185. P. J. Mohr, "Lamb shift in hydrogen-like ions," in **Beam-Foil Spectroscopy**, edited by I. A. Sellin and D. J. Pegg, Vol. 1 (Plenum, New York, 1975), pp. 89–96. (A)
186. "Vacuum polarization on a strong Coulomb field," E. H. Wichmann and N. M. Kroll, Phys. Rev. **101**(2), 843–859 (1956). (A)

Two-photon radiative corrections:

187. "Two-loop QED corrections in few-electron ions," V. A. Yerokhin, P. Indelicato, and V. M. Shabaev, Can. J. Phys. **85**(5), 521–529 (2007). (A)
188. "Two-loop self-energy correction in a strong Coulomb nuclear field," V. A. Yerokhin, P. Indelicato, and V. M. Shabaev, JETP **101**(2), 280–293 (2005). (A)
189. "Two-loop self-energy contribution to the Lamb shift in H-like ions," V. A. Yerokhin, P. Indelicato, and V. M. Shabaev, Phys. Rev. A **71**, 040101 (2005), 4 pp. (A)
190. "Calculation of the one- and two-loop Lamb shift for arbitrary excited hydrogenic states," A. Czarnecki, U. D. Jentschura, and K. Pachucki, Phys. Rev. Lett. **95**, 180404 (2005), 4 pp. (A)
191. "Radiative energy shifts induced by local potentials," U. D. Jentschura, J. Phys. A **36**(15), L229–L236 (2003). (A)
192. "Two-loop Bethe-logarithm correction in hydrogenlike atoms," K. Pachucki and U. D. Jentschura, Phys. Rev. Lett. **91**, 113005 (2003), 4 pp. (A)
193. "Two-loop self-energy correction in high-Z hydrogenlike ions," V. A. Yerokhin, P. Indelicato, and V. M. Shabaev, Phys. Rev. Lett. **91**, 073001 (2003), 4 pp. (A)
194. "Double-logarithmic two-loop self-energy corrections to the Lamb shift," U. D. Jentschura and I. Nándori, Phys. Rev. A **66**, 022114 (2002), 9 pp. (A)
195. "Logarithmic two-loop corrections to the Lamb shift in hydrogen," K. Pachucki, Phys. Rev. A **63**, 042503 (2001), 8 pp. (A)
196. "Loop-after-loop contribution to the second-order Lamb shift in hydrogenlike low-Z atoms," V. A. Yerokhin, Phys. Rev. A **62**, 012508 (2000), 6 pp. (A)
197. "Logarithms of α in QED bound states from the renormalization group," A. V. Manohar and I. W. Stewart, Phys. Rev. Lett. **85**(11), 2248–2251 (2000). (A)
198. "Analytic contribution of order $\alpha^2(Z\alpha)^5 m$ to the Lamb shift," M. I. Eides, H. Grotch, and V. A. Shelyuto, Phys. Rev. A **55**(3), 2447–2449 (1997). (A)
199. "Two-loop logarithmic corrections in the hydrogen Lamb shift," S. G. Karshenboim, J. Phys. B **29**(2), L29–L31 (1996). (A)
200. "New corrections to hyperfine splitting and Lamb shift and the value of the Rydberg constant," M. I. Eides and V. A. Shelyuto, JETP Lett. **61**(6), 478–484 (1995). (A)
201. "Corrections of order $\alpha^2(Z\alpha)^5$ to the hyperfine splitting and the Lamb shift," M. I. Eides and V. A. Shelyuto, Phys. Rev. A **52**(2), 954–961 (1995). (A)
202. "The Lamb shift in the hydrogen atom," S. G. Karshenboĭm, JETP **79**(2), 230–235 (1994). (A)
203. "Complete two-loop binding correction to the Lamb shift," K. Pachucki, Phys. Rev. Lett. **72**(20), 3154–3157 (1994). (A)
204. "Contributions to the binding, two-loop correction to the Lamb shift," K. Pachucki, Phys. Rev. A **48**(4), 2609–2614 (1993). (A)

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

205. "New logarithmic contributions in muonium and positronium," S. G. Karshenboĭm, JETP **76**(4), 541–546 (1993). (A)

Three-photon radiative corrections:

206. "Three-loop radiative corrections to Lamb shift and hyperfine splitting," M. I. Eides and V. A. Shelyuto, Can. J. Phys. **85**(5), 509–519 (2007). (A)
207. "Three-loop slope of the Dirac form factor and the 1S Lamb shift in hydrogen," K. Melnikov and T. van Ritbergen, Phys. Rev. Lett. **84**(8), 1673–1676 (2000). (A)
208. P. A. Baikov and D. J. Broadhurst, "Three-loop QED vacuum polarization and the four-loop muon anomalous magnetic moment," in **New Computing Techniques in Physics Research IV. International Workshop on Software Engineering, Artificial Intelligence, and Expert Systems for High Energy and Nuclear Physics**, edited by B. Denby and D. Perret-Gallix (World Scientific, Singapore, 1995), pp. 167–172. (A)
209. "Corrections of order $\alpha^3(Z\alpha)^4$ and $\alpha^2(Z\alpha)^6$ to the Lamb shift," M. I. Eides and H. Grotch, Phys. Rev. A **52**(4), 3360–3361 (1995). (A)

See also Ref. 88.

Finite-nuclear-size correction:

210. "Higher-order nuclear-size corrections in atomic hydrogen," J. L. Friar and G. L. Payne, Phys. Rev. A **56**(6), 5173–5175 (1997). (A)
211. "The Lamb shift of excited S-levels in hydrogen and deuterium atoms," S. G. Karshenboim, Z. Phys. D: At., Mol. Clusters **39**(2), 109–113 (1997). (A)
212. "Radiative correction to the electron charge density in the hydrogen atom," K. Pachucki, Phys. Rev. A **48**(1), 120–128 (1993). (A)
213. "The nuclear size correction to the energy levels of the light hydrogen-like and muonic atoms," L. A. Borisoglebsky and E. E. Trofimenko, Phys. Lett. B **81**(2), 175–177 (1979). (A)
214. "Nuclear finite-size effects in light muonic atoms," J. L. Friar, Ann. Phys. (N.Y.) **122**(1), 151–196 (1979). (A)

Nuclear-size correction to the self-energy and vacuum polarization:

215. "Finite nuclear size and Lamb shift of p-wave atomic states," A. I. Milstein, O. P. Sushkov, and I. S. Terekhov, Phys. Rev. A **67**, 062111 (2003), 5 pp. (A)
216. "Calculation of radiative corrections to the effect of parity nonconservation in heavy atoms," A. I. Milstein, O. P. Sushkov, and I. S. Terekhov, Phys. Rev. A **67**, 062103 (2003), 12 pp. (A)
217. "Radiative corrections and parity nonconservation in heavy atoms," A. I. Milstein, O. P. Sushkov, and I. S. Terekhov, Phys. Rev. Lett. **89**, 283003 (2002), 4 pp. (A)
218. "Radiative correction to the nuclear-size effect and hydrogen-deuterium isotopic shift," M. I. Eides and H. Grotch, Phys. Rev. A **56**(4), R2507–R2509 (1997). (A)
219. "Finite-nuclear-size corrections to the Uehling potential," D. J. Hylton, Phys. Rev. A **32**(3), 1303–1309 (1985). (A)
220. "Approximate normalized wave functions for the Finite-Size Coulomb problem," J. L. Friar, Z. Phys. A **292**(1), 1–6 (1979); Erratum **303**(1), 84 (1981). (A)

Radiative-recoil corrections:

221. "Expansion of bound-state energies in powers of m/M," A. Czarnecki and K. Melnikov, Phys. Rev. Lett. **87**, 013001 (2001), 4 pp. (A)
222. "Radiative-recoil corrections of order $\alpha(Z\alpha)^5(m/M)m$ to the Lamb shift revisited," M. I. Eides, H. Grotch, and V. A. Shelyuto, Phys. Rev. A **63**, 052509 (2001), 8 pp. (A)
223. "Corrections of order α^6 to S levels of two-body systems," M. I. Eides and H. Grotch, Phys. Rev. A **52**(2), 1757–1760 (1995). (A)
224. "Radiative recoil correction to the Lamb shift," K. Pachucki, Phys. Rev. A **52**(2), 1079–1085 (1995). (A)

See also Refs. 161 and 160.

Nucleus self-energy correction:

See Refs. 156 and 224.

2. Experiments

The Rydberg constant is determined from precision frequency measurements on hydrogen and deuterium atoms. Frequencies are the physical quantities that can be measured with the highest precision. The quest to improve the precision of frequency measurements was motivated largely by efforts to improve the accuracy of the Rydberg constant. This effort eventually led to the development of the frequency comb, which resulted in a recent Nobel Prize in physics.

225. "Femtosecond optical frequency combs," Th. Udem, R. Holzwarth, and Th. Hänsch, Eur. Phys. J. Spec. Top. **172**, 69–79 (2009). (I)
226. "Spectroscopy of atomic hydrogen," F. Biraben, Eur. Phys. J. Spec. Top. **172**, 109–119 (2009). (I)
227. "Nobel lecture: Passion for precision," T. W. Hänsch, Rev. Mod. Phys. **78**(4), 1297–1309 (2006). (I)
228. "Nobel lecture: Defining and measuring optical frequencies," J. L. Hall, Rev. Mod. Phys. **78**(4), 1279–1295 (2006). (I)
229. "New limits on the drift of fundamental constants from laboratory measurements," M. Fischer, N. Kolachevsky, M. Zimmermann, R. Holzwarth, Th. Udem, T. W. Hänsch, M. Abgrall, J. Grünert, I. Maksimovic, S. Bize, H. Marion, F. P. Dos Santos, P. Lemonde, G. Santarelli, P. Laurent, A. Clairon, C. Salomon, M. Haas, U. D. Jentschura, and C. H. Keitel, Phys. Rev. Lett. **92**, 230802, (2004), 4 pp. (A)
230. "Optical frequency measurement of the 2S−12D transitions in hydrogen and deuterium: Rydberg constant

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

and Lamb shift determinations," C. Schwob, L. Jozefowski, B. de Beauvoir, L. Hilico, F. Nez, L. Julien, F. Biraben, O. Acef, and A. Clairon, Phys. Rev. Lett. **82**(25), 4960–4963 (1999); Erratum **86**(18), 4193 (2001). (A)

231. "Hydrogen-deuterium 1S−2S isotope shift and the structure of the deuteron," A. Huber, Th. Udem, B. Gross, J. Reichert, M. Kourogi, K. Pachucki, M. Weitz, and T. W. Hänsch, Phys. Rev. Lett. **80**(3), 468–471 (1998). (A)

232. "Absolute frequency measurement of the 2S−8S/D transitions in hydrogen and deuterium: New determination of the Rydberg constant," B. de Beauvoir, F. Nez, L. Julien, B. Cagnac, F. Biraben, D. Touahri, L. Hilico, O. Acef, A. Clairon, and J. J. Zondy, Phys. Rev. Lett. **78**(3), 440–443 (1997). (A)

233. "High resolution spectroscopy of the hydrogen atom: Determination of the 1S Lamb shift," S. Bourzeix, B. de Beauvoir, F. Nez, M. D. Plimmer, F. de Tomasi, L. Julien, F. Biraben, and D. N. Stacey, Phys. Rev. Lett. **76**(3), 384–387 (1996). (A)

234. "Precision measurement of the 1S ground-state Lamb shift in atomic hydrogen and deuterium by frequency comparison," M. Weitz, A. Huber, F. Schmidt-Kaler, D. Leibfried, W. Vassen, C. Zimmermann, K. Pachucki, T. W. Hänsch, L. Julien, and F. Biraben, Phys. Rev. A **52**(4), 2664–2681 (1995). (A)

235. "Precise optical measurement of Lamb shifts in atomic hydrogen," D. J. Berkeland, E. A. Hinds, and M. G. Boshier, Phys. Rev. Lett. **75**(13), 2470–2473 (1995). (A)

236. "Separated oscillatory field measurement of hydrogen $2S_{1/2}-2P_{3/2}$ fine structure interval," E. W. Hagley and F. M. Pipkin, Phys. Rev. Lett. **72**(8), 1172–1175 (1994). (A)

237. "Separated oscillatory field measurement of the Lamb shift in H, $n=2$," S. R. Lundeen and F. M. Pipkin, Metrologia **22**(1), 9–54 (1986). (A)

238. "A precision determination of the Lamb shift in hydrogen," G. Newton, D. A. Andrews, and P. J. Unsworth, Philos. Trans. R. Soc. London, Ser. A **290**(1373), 373–404 (1979). (A)

239. "Status of the muonic hydrogen Lamb-shift experiment," T. Nebel, F. D. Amaro, A. Antognini, F. Biraben, J. M. R. Cardoso, C. A. N. Conde, A. Dax, S. Dhawan, L. M. P. Fernandes, A. Giesen, T. W. Hänsch, P. Indelicato, L. Julien, P. E. Knowles, F. Kottmann, E.-O. Le Bigot, Y.-W. Liu, J. A. M. Lopes, L. Ludhova, C. M. B. Monteiro, F. Mulhauser, F. Nez, R. Pohl, P. Rabinowitz, J. M. F. dos Santos, L. A. Schaller, K. Schuhmann, C. Schwob, D. Taqqu, and J. F. C. A. Veloso, Can. J. Phys. **85**(5), 469–478 (2007). (A)

240. "Precise proton radii from electron scattering," I. Sick, Can. J. Phys. **85**(5), 409–418 (2007). (A)

241. "On the rms-radius of the proton," I. Sick, Phys. Lett. B **576**(1–2), 62–67 (2003). (A)

242. "On the rms radius of the deuteron," I. Sick and D. Trautmann, Nucl. Phys. **A637**(4), 559–575 (1998). (A)

D. Planck constant *h*

The Planck constant h appears in various roles in quantum mechanics. It is the conversion factor between frequency ν and energy E of electromagnetic radiation

$$E = h\nu,$$

and it appears in the von Klitzing and Josephson relations associated with electrical phenomena in condensed-matter physics. The reduced Planck constant $\hbar = h/2\pi$ is two times the quantum unit of angular momentum, is a proportionality factor for action, and appears in the Heisenberg uncertainty relation.

The most accurate measurements of the Planck constant are made with the watt-balance experiment. Measurements on crystalline silicon, discussed in a subsequent section on the Avogadro constant, are also capable of providing information on the Planck constant through a theoretical relation.

243. "Determination of the Planck constant by means of a watt balance," A. Eichenberger, G. Genevès, and P. Gournay, Eur. Phys. J. Spec. Top. **172**, 363–383 (2009). (I)

244. "Toward the redefinition of the kilogram: Measurements of Planck's constant using watt balances," I. A. Robinson, IEEE Trans. Instrum. Meas. **58**(4), 942–948 (2009). (I)

245. "Planck, units, and modern metrology," J. L. Flowers and B. W. Petley, Ann. Phys. (Berlin) **17**(2–3), 101–114 (2008). (I)

246. "Tracing Planck's constant to the kilogram by electromechanical methods," A. Eichenberger, B. Jeckelmann, and P. Richard, Metrologia **40**(6), 356–365 (2003). (I)

247. "Progress on the BIPM watt balance," A. Picard, H. Fang, A. Kiss, E. de Mirandés, M. Stock, and C. Urano, IEEE Trans. Instrum. Meas. **58**(4), 924–929 (2009). (A)

248. "Toward a final result from the NPL Mark II watt balance," I. A. Robinson, IEEE Trans. Instrum. Meas. **58**(4), 936–941 (2009). (A)

249. "An initial measurement of Planck's constant using the NPL Mark II watt balance," I. A. Robinson, and B. P. Kibble, Metrologia **44**(6), 427–440 (2007). (A)

250. "Uncertainty improvements of the NIST electronic kilogram," R. L. Steiner, E. R. Williams, R. Liu, and D. B. Newell, IEEE Trans. Instrum. Meas. **56**(2), 592–596 (2007). (A)

251. "The watt balance: Towards a new definition of the mass unit?," G. Genevès, P. Gournay, F. Villar, P. Pinot, P. Juncar, M. Lecollinet, L. Chassagne, A. Clairon, A. Landragin, D. Holleville, F. Pereira Dos Santos, J. David, M. Besbes, F. Alves, S. Topçu, D. Haddad, A. Gosset, Z. Silvestri, P.-A. Meury, T. Madec, and S. Macé, Revue française de métrologie **2007-1**(9), 3–34 (2007). (A)

252. "The BIPM watt balance," A. Picard, M. Stock, H. Fang, T. J. Witt, and D. Reymann, IEEE Trans. Instrum. Meas. **56**(2), 538–542 (2007). (A)

253. "The BNM watt balance project," G. Genevès, P. Gournay, A. Gosset, M. Lecollinet, F. Villar, P. Pinot, P. Juncar, A. Clairon, A. Landragin, D. Holleville, F. Pereira Dos Santos, J. David, M. Besbes, F. Alves, L. Chassagne, and S. Topçu, IEEE Trans. Instrum. Meas. **54**(2), 850–853 (2005). (A)

254. "Details of the 1998 watt balance experiment determining the Planck constant," R. Steiner, D. Newell, and E. Williams, J. Res. Natl. Inst. Stand. Technol. **110**(1), 1–26 (2005). (A)

255. "Status of the METAS watt balance experiment," W.

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

Beer, A. L. Eichenberger, B. Jeanneret, B. Jeckelmann, A. R. Pourzand, P. Richard, and J. P. Schwarz, IEEE Trans. Instrum. Meas. **52**(2), 626–630 (2003). (A)

256. "Accurate measurement of the Planck constant," E. R. Williams, R. L. Steiner, D. B. Newell, and P. T. Olsen, Phys. Rev. Lett. **81**(12), 2404–2407 (1998). (A)

257. "A realization of the SI watt by the NPL moving-coil balance," B. P. Kibble, I. A. Robinson, and J. H. Belliss, Metrologia **27**(4), 173–192 (1990). (A)

E. Avogadro constant N_A

The Avogadro constant gives the number of entities in a mole. According to the definition of the mole in the SI, this number is the same as the number of carbon-12 atoms in 12 g of carbon. The main effort to directly determine this constant has been focused on determining the lattice spacing, mass, volume, and isotopic composition of spherical crystals of silicon.

258. "Determination of the Avogadro constant—A contribution to the new definition of the mass unit kilogram," P. Becker, Eur. Phys. J. Spec. Top. **172**, 343–352 (2009). (I)

259. "Towards an atomic realization of the kilogram: The measurements of N_A and $N_A h$," P. Becker, Eur. Phys. J. Spec. Top. **163**, 127–139 (2008). (I)

260. "High-purity single-crystal monoisotopic silicon-28 for precise determination of Avogadro's number," G. G. Devyatykh, A. D. Bulanov, A. V. Gusev, I. D. Kovalev, V. A. Krylov, A. M. Potapov, P. G. Sennikov, S. A. Adamchik, V. A. Gavva, A. P. Kotkov, M. F. Churbanov, E. M. Dianov, A. K. Kaliteevskii, O. N. Godisov, H.-J. Pohl, P. Becker, H. Riemann, and N. V. Abrosimov, Dokl. Chem. **421**(1), 157–160 (2008). (A)

261. "Accumulation of 38 mg of bismuth in a cylindrical collector from a 2.5 mA ion beam," C. Schlegel, F. Scholz, M. Gläser, M. Mecke, and G. Bethke, Metrologia **44**(1), 24–28 (2007). (A)

262. "Proposal for new experimental schemes to determine the Avogadro constant," F. Biraben, M. Cadoret, P. Cladé, G. Genevès, P. Gournay, S. Guellati-Khélifa, L. Julien, P. Juncar, E. de Mirandes, and F. Nez, Metrologia **43**(6), L47-L50 (2006). (A)

263. "Re-assessment of silicon isotope reference materials using high-resolution multi-collector ICP-MS," B. C. Reynolds, R. B. Georg, F. Oberli, U. Wiechert, and A. N. Halliday, J. Anal. At. Spectrom. **21**(3), 266–269 (2006). (A)

264. "Large-scale production of highly enriched ^{28}Si for the precise determination of the Avogadro constant," P. Becker, D. Schiel, H.-J. Pohl, A. K. Kaliteevski, O. N. Godisov, M. F. Churbanov, G. G. Devyatykh, A. V. Gusev, A. D. Bulanov, S. A. Adamchik, V. A. Gavva, I. D. Kovalev, N. V. Abrosimov, B. Hallmann-Seiffert, H. Riemann, S. Valkiers, P. Taylor, P. De Bièvre, and E. M. Dianov, Meas. Sci. Technol. **17**(7), 1854–1860 (2006). (A)

265. "Present state of the Avogadro constant determination from silicon crystals with natural isotopic compositions," K. Fujii, A. Waseda, N. Kuramoto, S. Mizushima, P. Becker, H. Bettin, A. Nicolaus, U. Kuetgens, S. Valkiers, P. Taylor, P. De Bièvre, G. Mana, E. Massa, R. Matyi, E. G. Kessler, Jr., and M. Hanke, IEEE Trans. Instrum. Meas. **54**(2), 854–859 (2005). (A)

266. "Silicon isotope abundance ratios and atomic weights of NBS-28 and other reference materials," T. Ding, D. Wan, R. Bai, Z. Zhang, Y. Shen, and R. Meng, Geochim. Cosmochim. Acta **69**(23), 5487–5494 (2005). (A)

267. "Determination of the Avogadro constant via the silicon route," P. Becker, H. Bettin, H.-U. Danzebrink, M. Gläser, U. Kuetgens, A. Nicolaus, D. Schiel, P. De Bièvre, S. Valkiers, and P. Taylor, Metrologia **40**(5), 271–287 (2003). (A)

268. "The Avogadro constant," G. Mana in **Proceedings of the International School of Physics "Enrico Fermi," Course CXLVI, Recent Advances in Metrology and Fundamental Constants**, edited by T. J. Quinn, S. Leschiutta, and P. Tavella (IOS, Amsterdam, 2001), pp. 177–193. (A)

269. "History and progress in the accurate determination of the Avogadro constant," P. Becker, Rep. Prog. Phys. **64**(12), 1945–2008 (2001). (A)

270. "Prompt (n,γ) mass measurements for the AVOGADRO Project," A. Paul, S. Röttger, A. Zimbal, and U. Keyser, Hyperfine Interact. **132**(1–4), 189–194 (2001). (A)

271. "Determination of the Avogadro constant by accurate measurement of the molar volume of a silicon crystal," K. Fujii, M. Tanaka, Y. Nezu, K. Nakayama, H. Fujimoto, P. De Bièvre, and S. Valkiers, Metrologia **36**(5), 455–464 (1999). (A)

272. "Just how perfect can a perfect crystal be?," R. D. Deslattes, E. G. Kessler, Jr., S. Owens, D. Black, and A. Henins, J. Phys. D **32**, A3-A7 (1999). (A)

273. "The determination of the Avogadro constant—Not simply a metrological problem," P. Becker, IEEE Trans. Instrum. Meas. **48**(2), 225–229 (1999). (A)

274. "Conclusions (Regarding the Avogadro constant)," T. J. Quinn, Metrologia **31**(3), 275–276 (1994). (A)

275. "Determining the Avogadro constant from electrical measurements," B. N. Taylor, Metrologia **31**(3), 181–194 (1994). (A)

276. "A determination of the Avogadro constant," P. Seyfried, P. Becker, A. Kozdon, F. Lüdicke, F. Spieweck, J. Stümpel, H. Wagenbreth, D. Windisch, P. De Bièvre, H. H. Ku, G. Lenaers, T. J. Murphy, H. S. Peiser, and S. Valkiers, Z. Phys. B: Condens. Matter **87**(3), 289–298 (1992). (A)

277. "Determination of the Avogadro constant," R. D. Deslattes, A. Henins, H. A. Bowman, R. M. Schoonover, C. L. Carroll, I. L. Barnes, L. A. Machlan, L. J. Moore, and W. R. Shields, Phys. Rev. Lett. **33**(8), 463–466 (1974). (A)

F. Molar gas constant R

The molar gas constant is the macroscopic link between energy and thermodynamic temperature expressed in units of J K^{-1} mol^{-1}. To date, the most accurate value of R has been determined by measuring the speed of sound in argon in a spherical acoustic resonator.

278. A. R. Colclough, "Methods for the determination of the gas constant," in **Precision Measurement and Fundamental Constants II**, edited by B. N. Taylor and W. D.

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

Phillips, NBS Special Publication 617 (U.S. Government Printing Office, Washington, DC, 1984), pp. 263–275. (I)
279. "Progress towards an acoustic/microwave determination of the Boltzmann constant at LNE-INM/CNAM," L. Pitre, C. Guianvarc'h, F. Sparasci, A. Richard, and D. Truong, Int. J. Thermophys. **29**(5), 1730–1739 (2008). (A)
280. "Measurement of the universal gas constant R with a spherical resonator," M.-G. He and Z.-G. Liu, Acta Phys. Sin.-Ch. Ed. **51**(5), 1004–1010 (2002). (A)
281. "Measurement of the universal gas constant R using a spherical acoustic resonator," M. R. Moldover, J. P. M. Trusler, T. J. Edwards, J. B. Mehl, and R. S. Davis, Phys. Rev. Lett. **60**(4), 249–252 (1988). (A)
282. "An acoustic redetermination of the gas constant," A. R. Colclough, T. J. Quinn, and T. R. D. Chandler, Proc. R. Soc. London, Ser. A **368**(1732), 125–139 (1979). (A)
283. "A new determination of the gas constant by an acoustical method," T. J. Quinn, A. R. Colclough, and T. R. D. Chandler, Philos. Trans. R. Soc. London, Ser. A **283**(1314), 367–420 (1976). (A)

G. Boltzmann constant k

The Boltzmann constant is the microscopic link between energy and thermodynamic temperature expressed in units of J K^{-1}. In practical determinations of temperature, it is the product kT that is proportional to the energy associated with the temperature that is measured. Although there are direct measurements of k, the most accurate determination is through the measurement of the molar gas constant R and use of the relation

$$k = \frac{R}{N_A}.$$

284. "A Johnson noise thermometer with traceability to electrical standards," L. Callegaro, V. D'Elia, M. Pisani, and A. Pollarolo, Metrologia **46**(5), 409–415 (2009). (A)
285. "Reduced non-linearities and improved temperature measurements for the NIST Johnson noise thermometer," J. Qu, S. P. Benz, H. Rogalla, and D. R. White, Metrologia **46**(5), 512–524 (2009). (A)
286. "Improvements in the NIST Johnson noise thermometry system," S. P. Benz, J. Qu, H. Rogalla, D. R. White, P. D. Dresselhaus, W. L. Tew, and S. W. Nam, IEEE Trans. Instrum. Meas. **58**(4), 884–890 (2009). (A)
287. "Primary tunnel junction thermometry," J. P. Pekola, T. Holmqvist, and M. Meschke, Phys. Rev. Lett. **101**, 206801 (2008), 4 pp. (A)
288. "Measurement time and statistics for a noise thermometer with a synthetic-noise reference," D. R. White, S. P. Benz, J. R. Labenski, S. W. Nam, J. F. Qu, H. Rogalla, and W. L. Tew, Metrologia **45**(4), 395–405 (2008). (A)
289. "Primary gas thermometry by means of laser-absorption spectroscopy: Determination of the Boltzmann constant," G. Casa, A. Castrillo, G. Galzerano, R. Wehr, A. Merlone, D. Di Serafino, P. Laporta, and L. Gianfrani, Phys. Rev. Lett. **100**, 200801 (2008), 4 pp. (A)
290. "Direct determination of the Boltzmann constant by an optical method," C. Daussy, M. Guinet, A. Amy-Klein, K. Djerroud, Y. Hermier, S. Briaudeau, Ch. J. Bordé, and C. Chardonnet, Phys. Rev. Lett. **98**, 250801 (2007), 4 pp. (A)
291. "Polarizability of helium and gas metrology," J. W. Schmidt, R. M. Gavioso, E. F. May, and M. R. Moldover, Phys. Rev. Lett. **98**, 254504 (2007), 4 pp. (A)
292. "Determination of the Boltzmann constant—Status and prospects," B. Fellmuth, Ch. Gaiser, and J. Fischer, Meas. Sci. Technol. **17**(10), R145–R159 (2006). (A)
293. "Determination of thermodynamic temperature and ^{4}He virial coefficients between 4.2 K and 27.0 K by dielectric constant gas thermometry," H. Luther, K. Grohmann, and B. Fellmuth, Metrologia **33**(4), 341–352 (1996). (A)

H. Josephson constant K_J

Josephson junctions are used as highly precise voltage standards. The Josephson constant K_J is the constant of proportionality that relates the frequency of microwave radiation applied to a superconducting Josephson junction to the induced voltage across the junction. It is related to the elementary charge and the Planck constant by

$$K_J = \frac{2e}{h} = \frac{1}{\Phi_0},$$

where Φ_0 is the magnetic flux quantum.

The universality tests mentioned below provide supporting evidence for the validity of the equation for K_J, which is relevant to the possible redefinition of the SI units.

1. Overviews

294. "Application of the Josephson effect in electrical metrology," B. Jeanneret and S. P. Benz, Eur. Phys. J. Spec. Top. **172**, 181–206 (2009). (I)
295. "Application of the Josephson effect to voltage metrology," S. P. Benz and C. A. Hamilton, Proc. IEEE **92**(10), 1617–1629 (2004). (I)
296. "Josephson voltage standards," J. Kohlmann, R. Behr, and T. Funck, Meas. Sci. Technol. **14**(8), 1216–1228 (2003). (I)
297. R. Behr, F. Müller, and J. Kohlmann, "Josephson junction arrays for voltage standards," in **Studies of Josephson Junction Arrays II, Studies of High Temperature Superconductors: Advances in Research and Applications**, Vol. 40, edited by A. Narlikar (Nova Science, Hauppauge, NY, 2002), pp. 155–184. (A)
298. "Josephson voltage standards," C. A. Hamilton, Rev. Sci. Instrum. **71**(10), 3611–3623 (2000). (I)
299. "Josephson voltage standard—A review," C. A. Hamilton, C. J. Burroughs, and S. P. Benz, IEEE Trans. Appl. Supercond. **7**(2), 3756–3761 (1997). (I)
300. J. C. Gallop, **SQUIDs, the Josephson Effects and Superconducting Electronics** (Adam-Hilger, Bristol, 1991). (I)
301. **Superconducting Quantum Electronics**, edited by V. Kose (Springer, Berlin, 1989). (I)

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

302. K. K. Likharev, **Dynamics of Josephson Junctions and Circuits** (Gordon and Breach, Amsterdam, 1986). (I)
303. A. Barone and G. Paternò, **Physics and Applications of the Josephson Effect** (Wiley, New York, 1982). (I)
304. T. Van Duzer and C. W. Turner, **Principles of Superconductive Devices and Circuits** (Elsevier, New York, 1981). (I)
305. "Implications of solid-state corrections to the Josephson voltage-frequency relation," T. A. Fulton, Phys. Rev. B **7**(3), 981–982 (1973). (I)
306. "Josephson effect in a superconducting ring," F. Bloch, Phys. Rev. B **2**(1) 109–121 (1970). (A)
307. "The Josephson effect and e/h," J. Clarke, Am. J. Phys. **38**(9), 1071–1095 (1970). (I)
308. "Simple interpretation of the Josephson effect," F. Bloch, Phys. Rev. Lett. **21**(17), 1241–1243 (1968). (I)
309. "Josephson currents in superconducting tunneling: The effect of microwaves and other observations," S. Shapiro, Phys. Rev. Lett. **11**(2), 80–82 (1963). (A)
310. "Possible new effects in superconductive tunneling," B. D. Josephson, Phys. Lett. **1**(7), 251–253 (1962). (A)

2. SI determinations

311. "Realization of the kilogram by measuring at 100 kV with the voltage balance ETF," V. Bego, J. Butorac, and D. Ilić, IEEE Trans. Instrum. Meas. **48**(2), 212–215 (1999). (A)
312. "Determination of the volt with the improved PTB voltage balance," T. Funck and V. Sienknecht, IEEE Trans. Instrum. Meas. **40**(2), 158–161 (1991). (A)
313. "A determination of the volt," W. K. Clothier, G. J. Sloggett, H. Bairnsfather, M. F. Currey, and D. J. Benjamin, Metrologia **26**(1), 9–46 (1989). (A)
314. "Absolute determination of the volt using a liquid electrometer," G. J. Sloggett, W. K. Clothier, M. F. Currey, D. J. Benjamin, and H. Bairnsfather, IEEE Trans. Instrum. Meas. **IM-34**(2), 187–191 (1985). (A)

3. Universality tests

315. "A review of Josephson comparison results," B. M. Wood and S. Solve, Metrologia **46**(6), R13–R20 (2009). (I)
316. "Highly precise comparison of $Nb/Al/AlO_x/Al/AlO_x/Al/Nb$ Josephson junction arrays using a SQUID as a null detector," I. Y. Krasnopolin, R. Behr, and J. Niemeyer, Supercond. Sci. Technol. **15**(7), 1034–1036 (2002). (A)
317. "Accurate measurements of quantum voltage steps on arrays of bicrystal Josephson junctions," A. M. Klushin, R. Behr, K. Numssen, M. Siegel, and J. Niemeyer, Appl. Phys. Lett. **80**(11), 1972–1974 (2002). (A)
318. "Precision measurement of the Josephson constant in $YBa_2Cu_3O_{7-\delta}$ superconductor," Yu. V. Tarbeyev, E. D. Koltik, V. I. Krzhimovsky, A. S. Katkov, O. P. Galakhova, S. V. Kozyrev, and S. E. Khabarov, Metrologia **28**(4), 305–307 (1991). (A)
319. "Precision of series-array Josephson voltage standards," R. L. Kautz and F. L. Lloyd, Appl. Phys. Lett. **51**(24), 2043–2045 (1987). (A)
320. "Test for relativistic gravitational effects on charged particles," A. K. Jain, J. E. Lukens, and J.-S. Tsai, Phys. Rev. Lett. **58**(12), 1165–1168 (1987). (A)
321. "High precision measurement of a possible resistive slope of Josephson array voltage steps," J. Niemeyer, L. Grimm, C. A. Hamilton, and R. L. Steiner, IEEE Electron Device Lett. **EDL-7**(1), 44–46 (1986). (A)
322. "Stable Josephson reference voltages between 0.1 and 1.3 V for high-precision voltage standards," J. Niemeyer, L. Grimm, W. Meier, J. H. Hinken, and E. Vollmer, Appl. Phys. Lett. **47**(11), 1222–1223 (1985). (A)
323. "High-precision test of the universality of the Josephson voltage-frequency relation," J.-S. Tsai, A. K. Jain, and J. E. Lukens, Phys. Rev. Lett. **51**(4), 316–319 (1983); Erratum **51**(12), 1109 (1983). (A)
324. "Comparison between the constant-potential steps of two point-contact Josephson junctions," J. C. Macfarlane, Appl. Phys. Lett. **22**(10), 549–550 (1973). (A)
325. "Comparison of microwave-induced constant-voltage steps in Pb and Sn Josephson junctions," T. D. Bracken and W. O. Hamilton, Phys. Rev. B **6**(7), 2603–2609 (1972). (A)
326. "Comments on quantum-electrodynamic corrections to the electron charge in metals," D. N. Langenberg and J. R. Schrieffer, Phys. Rev. B **3**(5), 1776–1778 (1971). (I)
327. "Absence of quantum-electrodynamic corrections to the charge of the electron as measured in Josephson junction experiments," J. B. Hartle, D. J. Scalapino, and R. L. Sugar, Phys. Rev. B **3**(5), 1778–1781 (1971). (A)
328. "Precise comparison of the Josephson frequency-voltage relation at microwave and far-infrared frequencies," T. F. Finnegan, A. Denenstein, D. N. Langenberg, J. C. McMenamin, D. E. Novoseller, and L. Cheng, Phys. Rev. Lett. **23**(5), 229–233 (1969). (A)
329. "Determination of e/h using macroscopic quantum phase coherence in superconductors. I. Experiment," W. H. Parker, D. N. Langenberg, A. Denenstein, and B. N. Taylor, Phys. Rev. **177**(2), 639–664 (1969). (A)
330. "Experimental comparison of the Josephson voltage-frequency relation in different superconductors," J. Clarke, Phys. Rev. Lett. **21**(23), 1566–1569 (1968). (A)

I. von Klitzing constant R_K

The quantum Hall effect provides a highly precise resistance standard. The quotient of the voltage (at a plateau) across a quantum Hall device and the current flowing through it is proportional to the von Klitzing constant R_K, which is given by

$$R_K = \frac{h}{e^2} = \frac{\mu_0 c}{2\alpha}.$$

See also Sec. V A 3.

The universality tests mentioned below provide supporting evidence for the validity of the equation for R_K, which is relevant to the possible redefinition of the SI units.

1. Overviews

331. "Resistance metrology based on the quantum Hall effect," W. Poirier and F. Schopfer, Eur. Phys. J. Spec. Top. **172**, 207–245 (2009). (I)

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

332. "Quantum Hall effect and ohm metrology," W. Poirier and F. Schopfer, Int. J. Mod. Phys. B **23** (12–13), 2779–2789 (2009). (I)
333. "Realizing the farad from two ac quantum Hall resistances," J. Schurr, V. Bürkel, and B. P. Kibble, Metrologia **46**(6), 619–628 (2009). (A)
334. "Experimental observation of the quantum Hall effect and Berry's phase in graphene," Y. Zhang, Y.-W. Tan, H. L. Stormer, and P. Kim, Nature (London) **438**(7065), 201–204 (2005). (A)
335. K. von Klitzing, "25 years of quantum Hall effect (QHE): A personal view on the discovery, physics and applications of this quantum effect," in **The Quantum Hall Effect**, edited by B. Douçot, B. Duplantier, V. Pasquier, and V. Rivasseau (Birkhäuser, Basel, 2005), pp. 1–21. (I)
336. "Developments in the quantum Hall effect," K. von Klitzing, Philos. Trans. R. Soc. London, Ser. A **363**(1834), 2203–2219 (2005). (I)
337. "25 years quantum Hall effect: How it all came about," G. Landwehr, Physica E (Amsterdam) **20**(1–2), 1–13 (2003). (E)
338. "The quantum Hall effect as an electrical resistance standard," B. Jeckelmann and B. Jeanneret, Rep. Prog. Phys. **64**(12), 1603–1655 (2001). (I)
339. "Optimization of QHE-devices for metrological applications," B. Jeckelmann, A. Rüfenacht, B. Jeanneret, F. Overney, K. Pierz, A. von Campenhausen, and G. Hein, IEEE Trans. Instrum. Meas. **50**(2), 218–222 (2001). (A)
340. S. D. Sarma and A. Pinczuk, **Perspectives in Quantum Hall Effects** (Wiley, New York, 1997). (A)
341. "Hall conductance as a topological invariant," G. Watson, Contemp. Phys. **37**(2), 127–143 (1996). (A)
342. T. Chakraborty and P. Pietiläinen, **The Quantum Hall Effects: Integral and Fractional**, 2nd ed. (Springer, Berlin, 1995). (A)
343. M. Janßen, O. Viehweger, U. Fastenrath, and J. Hajdu, **Introduction to the Theory of the Integer Quantum Hall Effect**, edited by J. Hajdu (VCH, Weinheim, Germany, 1994). (I)
344. **Quantum Hall Effect**, edited by M. Stone (World Scientific, Singapore, 1992). (I)
345. **The Integral and Fractional Quantum Hall Effects**, edited by C. T. Van Degrift, M. E. Cage, and S. M. Girvin (American Association of Physics Teachers, College Park, MD, 1991). (I)
346. **Quantum Hall Effect: A Perspective**, edited by A. H. MacDonald (Kluwer Academic, Dordrecht, 1990). (A)
347. **The Quantum Hall Effect**, 2nd ed., edited by R. E. Prange and S. M. Girvin, (Springer-Verlag, New York, 1990). (I)
348. G. Morandi, **Quantum Hall Effect** (Bibliopolis, Naples, Italy, 1988). (A)
349. "Integral quantum Hall effect for nonspecialists," D. R. Yennie, Rev. Mod. Phys. **59**(3), 781–824 (1987). (I)
350. "The quantized Hall effect," K. von Klitzing, Rev. Mod. Phys. **58**(3), 519–531 (1986). (I)
351. "The quantized Hall effect," B. I. Halperin, Sci. Am. **254**(4), 52–60 (1986). (I)
352. "New method for high-accuracy determination of the fine-structure constant based on quantized Hall resistance," K. von Klitzing, G. Dorda, and M. Pepper, Phys. Rev. Lett. **45**(6), 494–497 (1980). (A)

2. SI determinations

See Sec. V A 3.

3. Universality tests

353. "Quantum Hall effect in quantum electrodynamics," A. A. Penin, Phys. Rev. B **79**, 113303 (2009), 4 pp. (A)
354. "Quantum resistance metrology in graphene," A. J. M. Giesbers, G. Rietveld, E. Houtzager, U. Zeitler, R. Yang, K. S. Novoselov, A. K. Geim, and J. C. Maan, Appl. Phys. Lett. **93**, 222109 (2008), 3 pp. (A)
355. "Testing universality of the quantum Hall effect by means of the Wheatstone bridge," F. Schopfer and W. Poirier, J. Appl. Phys. **102**, 054903 (2007), 9 pp. (A)
356. "High-precision measurements of the quantized Hall resistance: Experimental conditions for universality," B. Jeckelmann, B. Jeanneret, and D. Inglis, Phys. Rev. B **55**(19), 13124–13134 (1997). (A)
357. "Direct comparison of the quantized Hall resistance in gallium arsenide and silicon," A. Hartland, K. Jones, J. M. Williams, B. L. Gallagher, and T. Galloway, Phys. Rev. Lett. **66**(8), 969–973 (1991). (A)
358. "Direct comparison of quantized Hall resistances," F. Piquemal, B. Etienne, J.-P. Andre, and J.-N. Patillon, IEEE Trans. Instrum. Meas. **40**(2), 234–236 (1991). (A)
359. "Comparison of the quantized Hall resistance in different $GaAs/Al_xGa_{1-x}As$ heterostructures," B. Jeckelmann, W. Schwitz, H.-J. Bühlmann, R. Houdré, M. Ilegems, D. Jucknischke, and M. A. Py, IEEE Trans. Instrum. Meas. **40**(2), 231–233 (1991). (A)
360. "Universality of quantum Hall effect: Topological invariant and observable," H. Aoki and T. Ando, Phys. Rev. Lett. **57**(24), 3093–3096 (1986). (A)
361. "Precise quantized Hall resistance measurements in $GaAs/Al_xGa_{1-x}As$ and $In_xGa_{1-x}As/InP$ heterostructures," F. Delahaye, D. Dominguez, F. Alexandre, J. P. Andre, J. P. Hirtz, and M. Razeghi, Metrologia **22**(2), 103–110 (1986). (A)
362. "Quantum Hall effect in silicon metal-oxide-semiconductor inversion layers: Experimental conditions for determination of h/e^2," K. Yoshihiro, J. Kinoshita, K. Inagaki, C. Yamanouchi, T. Endo, Y. Murayama, M. Koyanagi, A. Yagi, J. Wakabayashi, and S. Kawaji, Phys. Rev. B **33**, 6874–6896 (1986). (A)
363. "A measurement system for the determination of h/e^2 in terms of the SI ohm and the maintained ohm at the NPL," A. Hartland, G. J. Davies, and D. R. Wood, IEEE Trans. Instrum. Meas. **IM-34**(2), 309–314 (1985). (A)
364. "High precision measurements of the quantized Hall resistance at the PTB," L. Bliek, E. Braun, F. Melchert, P. Warnecke, W. Schlapp, G. Weimann, K. Ploog, G. Ebert, and G. E. Dorda, IEEE Trans. Instrum. Meas. **IM-34**(2), 304–305 (1985). (A)
365. "Temperature dependence of the quantum Hall resistance," M. E. Cage, B. F. Field, R. F. Dziuba, S. M. Girvin, A. C. Gossard, and D. C. Tsui, Phys. Rev. B **30**(4), 2286–2288 (1984). (A)
366. "Dissipation and dynamic nonlinear behavior in the quantum Hall regime," M. E. Cage, R. F. Dziuba, B. F. Field, E. R. Williams, S. M. Girvin, A. C. Gossard, D.

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

C. Tsui, and R. J. Wagner, Phys. Rev. Lett. **51**(15), 1374–1377 (1983). (A)

J. Electron mass m_e

The electron mass is most accurately known in atomic mass units (in which the relative atomic mass of the carbon-12 atom is exactly 12) denoted by $A_r(e)$. Three methods of determining the mass in these units are described in the following three sections. The subsequent section gives an expression for the electron mass in terms of other constants, which provides the best value in kilograms.

1. Bound electron g-factor

The best value of the electron mass in relative atomic mass units is provided by measurements on hydrogenlike ions of the ratio of the electron spin-flip resonance frequency f_s to the cyclotron frequency f_c of the ion in a uniform magnetic field. The electron mass follows from the relation

$$A_r(e) = -\frac{g_e(I)}{2(Z-1)}\frac{f_c}{f_s}A_r(I),$$

where $g_e(I)$ is the g-factor of the electron in the ion I, Z is the nuclear charge number of the ion, and $A_r(I)$ is its relative atomic mass. The atomic masses, from which $A_r(I)$ is readily obtained, are generally known with sufficient accuracy for this purpose (see Sec. V L). The g-factor is calculated from the atomic theory of the bound electron, with the following references. References for the experiments are given next.

367. "Mass measurements and the bound-electron g factor," U. D. Jentschura, A. Czarnecki, K. Pachucki, and V. A. Yerokhin, Int. J. Mass Spectrom. **251**(2–3), 102–108 (2006). (A)

368. "Virtual light-by-light scattering and the g factor of a bound electron," R. N. Lee, A. I. Milstein, I. S. Terekhov, and S. G. Karshenboim, Phys. Rev. A **71**, 052501, 7 pages (2005). (A)

369. "Complete two-loop correction to the bound-electron g factor," K. Pachucki, A. Czarnecki, U. D. Jentschura, and V. A. Yerokhin, Phys. Rev. A **72**, 022108, 11 pages (2005). (A)

370. "Nonrelativistic QED approach to the bound-electron g factor," K. Pachucki, U. D. Jentschura, and V. A. Yerokhin, Phys. Rev. Lett. **93**, 150401 (2004), 4 pp.; Erratum **94**, 229902 (2005), 1 p. (A)

371. "A consistent set of nuclear rms charge radii: Properties of the radius surface $R(N,Z)$," I. Angeli, At. Data. Nucl. Data Tables **87**(2), 185–206 (2004). (A)

372. "The g factor in a light two-body atomic system: A determination of fundamental constants to test QED," S. G. Karshenboim and V. G. Ivanov, Can. J. Phys. **80**(11), 1305–1312 (2002). (A)

373. "Finite nuclear size correction to the bound-electron g factor in a hydrogenlike atom," D. A. Glazov and V. M. Shabaev, Phys. Lett. A **297**(5–6), 408–411 (2002). (A)

374. "Delbrück scattering and the g-factor of a bound electron," S. G. Karshenboim and A. I. Milstein, Phys. Lett. B **549**(3–4), 321–324 (2002). (A)

375. "Self-energy correction to the bound-electron g factor in H-like ions," V. A. Yerokhin, P. Indelicato, and V. M. Shabaev, Phys. Rev. Lett. **89**, 143001 (2002), 4 pp. (A)

376. "Magnetic moment of a two-particle bound state in quantum electrodynamics," A. P. Martynenko and R. N. Faustov, Phys. At. Nucl. **65**(2), 271–276 (2002). (A)

377. "The g factors of bound particles in quantum electrodynamics," A. P. Martynenko and R. N. Faustov, Sov. Phys. JETP **93**(3), 471–476 (2001). (A)

378. "Vacuum polarization in a hydrogen-like relativistic atom: g factor of a bound electron," S. G. Karshenboim, V. G. Ivanov, and V. M. Shabaev, JETP **93**(3), 477–484 (2001). (A)

379. "Some analytic results on the Uehling correction to the g-factor of a bound electron (muon)," S. G. Karshenboim, V. G. Ivanov, and V. M. Shabaev, Can. J. Phys. **79**(1), 81–86 (2001). (A)

380. "Anomalous magnetic moment of a bound electron," A. Czarnecki, K. Melnikov, and A. Yelkhovsky, Phys. Rev. A **63**, 012509 (2000), 4 pp. (A)

381. "The g_j factor of a bound electron and the hyperfine structure splitting in hydrogenlike ions," T. Beier, Phys. Rep. **339**(2–3), 79–213 (2000). (A)

382. "g_j factor of an electron bound in a hydrogenlike ion," T. Beier, I. Lindgren, H. Persson, S. Salomonson, P. Sunnergren, H. Häffner and N. Hermanspahn, Phys. Rev. A **62**, 032510 (2000), 31 pp. (A)

383. "Non-relativistic calculations of the g-factor of a bound electron," S. G. Karshenboim, Phys. Lett. A **266**(4–6), 380–386 (2000). (A)

384. "Gyromagnetic ratios of bound particles," M. I. Eides and H. Grotch, Ann. Phys. (N.Y.) **260**(1), 191–200 (1997). (A)

385. "Relativistic extension of the electromagnetic current for composite systems," F. E. Close and H. Osborn, Phys. Lett. B **34**(5), 400–404 (1971). (A)

386. "Electron g factor in hydrogenic atoms," H. Grotch, Phys. Rev. Lett. **24**(2), 39–42 (1970). (A)

387. "Magnetic moment of the hydrogen atom," R. Faustov, Phys. Lett. B **33**(6), 422–424 (1970). (A)

388. "The magnetic moment of the electron," G. Breit, Nature (London) **122**(3078), 649 (1928). (A)

Measurements have been made on hydrogenlike carbon and oxygen.

389. "Highly charged ions, quantum-electrodynamics, and the electron mass," G. Werth, J. Alonso, T. Beier, K. Blaum, S. Djekic, H. Häffner, N. Hermanspahn, W. Quint, S. Stahl, J. Verdú, T. Valenzuela, and M. Vogel, Int. J. Mass Spectrom. **251**(2–3), 152–158 (2006). (A)

390. "Electronic g factor of hydrogenlike oxygen $^{16}O^{7+}$," J. Verdú, S. Djekić, S. Stahl, T. Valenzuela, M. Vogel, G. Werth, T. Beier, H.-J. Kluge, and W. Quint, Phys. Rev. Lett. **92**, 093002 (2004), 4 pp. (A)

391. "Double Penning trap technique for precise g factor determinations in highly charged ions," H. Häffner, T. Beier, S. Djekić, N. Hermanspahn, H.-J. Kluge, W. Quint, S. Stahl, J. Verdú, T. Valenzuela, and G. Werth, Eur. Phys. J. D **22**(2), 163–182 (2003). (A)

392. "The magnetic moment anomaly of the electron bound in hydrogen-like oxygen $^{16}O^{7+}$," J. Verdú, T. Beier, S. Djekić, H. Häffner, H.-J. Kluge, W. Quint, T. Valenzuela, M. Vogel, and G. Werth, J. Phys. B **36**(3), 655–663 (2003). (A)

393. "Measurement of the g_j factor of a bound electron in

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

hydrogen-like oxygen $^{16}O^{7+}$," J. L. Verdú, T. Beier, S. Djekic, H. Häffner, H.-J. Kluge, W. Quint, T. Valenzuela, and G. Werth, Can. J. Phys. **80**(11), 1233–1240 (2002). (A)

394. "New determination of the electron's mass," T. Beier, H. Häffner, N. Hermanspahn, S. G. Karshenboim, H.-J. Kluge, W. Quint, S. Stahl, J. Verdú, and G. Werth, Phys. Rev. Lett. **88**, 011603 (2002), 4 pp. (A)

2. Transitions in antiprotonic helium

An antiprotonic helium atom is the atomic state of a helium nucleus, an antiproton, and an electron. Transition frequencies between highly excited states of the antiproton in the atom have been measured and compared to the theoretical values, which depend on the mass ratios of the three particles. This comparison provides a determination of the electron mass since the masses of the helium nucleus and the antiproton are more accurately known from other measurements. This assumes that *CPT* (combined charge conjugation, parity reflection, and time reversal) invariance holds, in which case the masses of the antiproton and the proton are the same.

The theory is described in the following references.

395. V. I. Korobov, "Precise spectroscopy of antiprotonic helium and sensitivity of transitions to the antiproton mass," in **Proceedings of the EXA05: International Conference on Exotic Atoms and Related Topics**, edited by A. Hirtl, J. Marton, E. Widmann, and J. Zmeskal (Austrian Academy of Sciences, Vienna, 2005), pp. 391–400. (A)

396. "High-precision Coulomb three-body calculation of antiprotonic helium atoms," Y. Kino, H. Kudo, and M. Kamimura, Mod. Phys. Lett. A **18**(2–6), 388–397 (2003). (A)

397. "Metastable states in the antiprotonic helium atom decaying via Auger transitions," V. I. Korobov, Phys. Rev. A **67**, 062501 (2003), 7 pp. (A)

The experiments use precision laser spectroscopy.

398. "Weighing the antiproton by parts-per-billion-scale laser spectroscopy of antiprotonic helium," R. S. Hayano, Can. J. Phys. **85**(5), 453–460 (2007). (A)

399. "Determination of the antiproton-to-electron mass ratio by precision laser spectroscopy of $\bar{p}He^+$," M. Hori, A. Dax, J. Eades, K. Gomikawa, R. S. Hayano, N. Ono, W. Pirkl, E. Widmann, H. A. Torii, B. Juhász, D. Barna, and D. Horváth, Phys. Rev. Lett. **96**, 243401 (2006), 4 pp. (A)

400. "Direct measurement of transition frequencies in isolated $\bar{p}He^+$ atoms, and new *CPT*-violation limits on the antiproton charge and mass," M. Hori, J. Eades, R. S. Hayano, T. Ishikawa, W. Pirkl, E. Widmann, H. Yamaguchi, H. A. Torii, B. Juhász, D. Horváth, and T. Yamazaki, Phys. Rev. Lett. **91**, 123401 (2003), 4 pp. (A)

401. "Antiprotonic helium," T. Yamazaki, N. Morita, R. S. Hayano, E. Widmann, and J. Eades, Phys. Rep. **366**(4–5), 183–329 (2002). (A)

402. "Sub-ppm laser spectroscopy of antiprotonic helium and a *CPT*-violation limit on the antiprotonic charge and mass," M. Hori, J. Eades, R. S. Hayano, T. Ishikawa, J. Sakaguchi, E. Widmann, H. Yamaguchi, H. A. Torii, B. Juhász, D. Horváth, and T. Yamazaki, Phys. Rev. Lett. **87**, 093401 (2001), 4 pp. (A)

3. Mass measurement in a Penning trap

403. "Determination of the electron's atomic mass and the proton/electron mass ratio via Penning trap mass spectroscopy," D. L. Farnham, R. S. Van Dyck, Jr., and P. B. Schwinberg, Phys. Rev. Lett. **75**(20), 3598–3601 (1995). (A)

4. Electron mass in kilograms

The electron mass in kilograms is evaluated with the relation

$$m_e = \frac{2hR_\infty}{\alpha^2 c}$$

from the definition of the Rydberg constant in Sec. V C using values of the constants obtained as discussed elsewhere in this Resource Letter.

K. Electron-muon mass ratio m_e/m_μ

Muonium is the atomic state of a positive muon and an electron. The ground state has two hyperfine levels, depending on the relative alignment of the spins of the muon and electron. The electron-muon mass ratio is determined by a comparison of the measured and calculated values of the hyperfine splitting, where the mass ratio is one of the parameters in the theory. The expression is

$$\Delta\nu_{\mathrm{Mu}} = \frac{16}{3} cR_\infty \alpha^2 \frac{m_e}{m_\mu}\left(1 + \frac{m_e}{m_\mu}\right)^{-3} \mathcal{F}(\alpha, m_e/m_\mu),$$

where the function $\mathcal{F}$, which is 1 to lowest order in the variables, contains the details of the theory but depends only weakly on the mass ratio.

The value of the fine-structure constant α can also be determined from this relation if an independent value of the mass ratio is used. However, the value for α obtained this way is less accurate than values obtained from other sources.

1. Theory

404. "Three-loop radiative-recoil corrections to hyperfine splitting in muonium," M. I. Eides, H. Grotch, and V. A. Shelyuto, Can. J. Phys. **83**(4), 363–373 (2005).

405. "Three-loop reducible radiative photon contributions to Lamb shift and hyperfine splitting," M. I. Eides and V. A. Shelyuto, Phys. Rev. A **70**, 022506 (2004), 4 pp. (A)

406. "Three-loop radiative-recoil corrections to hyperfine splitting generated by one-loop fermion factors," M. I. Eides, H. Grotch, and V. A. Shelyuto, Phys. Rev. D **70**, 073005 (2004), 7 pp. (A)

407. "Polarization operator contributions to the Lamb shift and hyperfine splitting," M. I. Eides and V. A. Shelyuto, Phys. Rev. A **68**, 042106 (2003), 10 pp. (A)

408. "Three-loop radiative-recoil corrections to hyperfine splitting in muonium," M. I. Eides, H. Grotch, V. A. Shelyuto, Phys. Rev. D **67**, 113003 (2003), 16 pp. (A)

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

409. "The role of pions and kaons in τ decays, $(g_\mu-2)$, the running of α_{QED} and the muonium hyperfine splitting," J. E. Palomar, Nucl. Phys. B (Proc. Suppl.) **121**, 183–186 (2003). (A)

410. "Two-loop polarization contributions to radiative-recoil corrections to hyperfine splitting in muonium," M. I. Eides, H. Grotch, and V. A. Shelyuto, Phys. Rev. D **65**, 013003 (2001), 8 pp. (A)

411. "Hadronic effects in leptonic systems: Muonium hyperfine structure and anomalous magnetic moment of muon," S. I. Eidelman, S. G. Karshenboĭm, and V. A. Shelyuto, Can. J. Phys. **80**(11), 1297–1303 (2002). (A)

412. M. Nio, "Complete result of the $\alpha(Z\alpha)^3$ muonium hyperfine structure," in **Quantum Electrodynamics and Physics of the Vacuum**, edited by G. Cantatore, AIP Conference Proceedings 564 (AIP, Melville, NY, 2001). (A)

413. "One-loop self-energy correction to the 1*s* and 2*s* hyperfine splitting H-like systems," V. A. Yerokhin and V. M. Shabaev, Phys. Rev. A **64**, 012506 (2001), 6 pp. (A)

414. "Polarization of vacuum in a hydrogen-like relativistic atom: Hyperfine structure," S. G. Karshenboĭm, V. G. Ivanov, and V. M. Shabaev, JETP **90**(1), 59–65 (2000). (A)

415. "Analytic results on the VP contribution to the energy in hydrogen-like ions," S. G. Karshenboim, V. G. Ivanov, and V. M. Shabaev, Phys. Scr. **T80**, 491–492 (1999). (A)

416. "All-order binding corrections to muonium hyperfine splitting," S. A. Blundell, K. T. Cheng, and J. Sapirstein, Phys. Rev. Lett. **78**(26), 4914–4917 (1997). (A)

417. "Weak-interaction contributions to hyperfine splitting and Lamb shift," M. I. Eides, Phys. Rev. A **53**(5), 2953–2957 (1996). (A)

418. "One-logarithmic recoil correction in muonium hyperfine splitting," G. Li, M. A. Samuel, and M. I. Eides, Phys. Rev. A **47**(2), 876–878 (1993). (A)

See also Ref. 206.

2. Experiments

419. "High precision measurements of the ground state hyperfine structure interval of muonium and of the muon magnetic moment," W. Liu, M. G. Boshier, S. Dhawan, O. van Dyck, P. Egan, X. Fei, M. G. Perdekamp, V. W. Hughes, M. Janousch, K. Jungmann, D. Kawall, F. G. Mariam, C. Pillai, R. Prigl, G. zu Putlitz, I. Reinhard, W. Schwarz, P. A. Thompson, and K. A. Woodle, Phys. Rev. Lett. **82**(4), 711–714 (1999). (A)

420. "High precision measurement of the hfs interval of muonium and of the muon magnetic moment," F. G. Mariam, W. Beer, P. R. Bolton, P. O. Egan, C. J. Gardner, V. W. Hughes, D. C. Lu, P. A. Souder, H. Orth, J. Vetter, U. Moser, and G. zu Putlitz, Phys. Rev. Lett. **49**(14), 993–996 (1982). (A)

L. Atom, ion, and nucleon masses

Masses of atoms, ions, and nuclei are most accurately known in atomic mass units (in which the mass of the carbon-12 atom is exactly 12). Mass measurements are generally made with ions rather than neutral atoms. For an ion I with n electrons and nucleus N, the mass is

$$m_I = m_N + nm_e - E_b/c^2,$$

where E_b is the total binding energy of the n electrons. For neutral atoms $n=Z$, where Z is the atomic number. The mass of the neutron is mainly determined by measuring the wavelength of γ-rays emitted by the capture of neutrons in hydrogen.

421. "ISOLTRAP: An on-line Penning trap for mass spectrometry on short-lived nuclides," M. Mukherjee, D. Beck, K. Blaum, G. Bollen, J. Dilling, S. George, F. Herfurth, A. Herlert, A. Kellerbauer, H.-J. Kluge, S. Schwarz, L. Schweikhard, and C. Yazidjian, Eur. Phys. J. A **35**(1), 1–29 (2008). (A)

422. "Determination of the proton mass from a measurement of the cyclotron frequencies of D^+ and H_2^+ in a Penning trap," A. Solders, I. Bergström, Sz. Nagy, M. Suhonen, and R. Schuch, Phys. Rev. A **78**, 012514 (2008), 6 pp. (A)

423. "Dipole moment of PH^+ and the atomic masses of ^{28}Si, ^{31}P by comparing cyclotron frequencies of two ions simultaneously trapped in a Penning trap," M. Redshaw, J. McDaniel, and E. G. Myers, Phys. Rev. Lett. **100**, 093002 (2008), 4 pp. (A)

424. "Precise atomic masses for fundamental physics determined at SMILETRAP," R. Schuch, I. Bergström, T. Fritioff, Sz. Nagy, A. Solders, and M. Suhonen, Adv. Quantum Chem. **53**, 67–81 (2008), Chap. 6. (A)

425. "HITRAP: A facility at GSI for highly charged ions," H.-J. Kluge, T. Beier, K. Blaum, L. Dahl, S. Eliseev, F. Herfurth, B. Hofmann, O. Kester, S. Koszudowski, C. Kozhuharov, G. Maero, W. Nörtershäuser, J. Pfister, W. Quint, U. Ratzinger, A. Schempp, R. Schuch, Th. Stöhlker, R. C. Thompson, M. Vogel, G. Vorobjev, D. F. A. Winters, and G. Werth, Adv. Quantum Chem. **53**, 83–98 (2008), Chap. 7. (A)

426. "Precision mass spectrometry and polarizability shifts with one and two ions in a penning trap," M. Redshaw, W. Shi, J. McDaniel, E. Wingfield, and E. G. Myers, Hyperfine Interact. **174**(1–3), 65–69 (2007). (A)

427. "The UW-PTMS: Recent measurements and technological progress," D. B. Pinegar, S. L. Zafonte, and R. S. Van Dyck, Jr., Hyperfine Interact. **174**(1–3), 47–53 (2007). (A)

428. "Atomic weights of the elements 2005 (IUPAC Technical Report)," M. E. Wieser, Pure Appl. Chem. **78**(11), 2051–2066 (2006). (A)

429. "Accurate mass measurements of ^{26}Ne, $^{26-30}Na$, $^{29-33}Mg$ performed with the MISTRAL spectrometer," C. Gaulard, G. Audi, C. Bachelet, D. Lunney, M. de Saint Simon, C. Thibault, and N. Vieira, Nucl. Phys. A **766**, 52–73 (2006). (A)

430. "The history of nuclidic masses and of their evaluation," G. Audi, Int. J. Mass Spectrom. **251**(2–3), 85–94 (2006). (A)

431. "On the Q-value of the tritium β-decay," Sz. Nagy, T. Fritioff, M. Björkhage, I. Bergström, and R. Schuch, Europhys. Lett. **74**(3), 404–410 (2006). (A)

432. "The UW-PTMS: Systematic studies, measurement

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

progress, and future improvements," R. S. Van Dyck Jr., D. B. Pinegar, S. V. Liew, and S. L. Zafonte, Int. J. Mass Spectrom. **251**(2–3), 231–242 (2006). (A)

433. "A direct test of $E=mc^2$," S. Rainville, J. K. Thompson, E. G. Myers, J. M. Brown, M. S. Dewey, E. G. Kessler, Jr., R. D. Deslattes, H. G. Börner, M. Jentschel, P. Mutti, and D. E. Pritchard, Nature (London) **438**(7071), 1096–1097 (2005). (A)

434. "The AME2003 atomic mass evaluation (I). Evaluation of input data, adjustment procedures," A. H. Wapstra, G. Audi, and C. Thibault, Nucl. Phys. A **729**(1), 129–336 (2003). (A)

435. "The AME2003 atomic mass evaluation (II). Tables, graphs and references," G. Audi, A. H. Wapstra, and C. Thibault, Nucl. Phys. A **729**(1), 337–676 (2003). (A)

436. "The deuteron binding energy and the neutron mass," E. G. Kessler, Jr., M. S. Dewey, R. D. Deslattes, A. Henins, H. G. Börner, M. Jentschel, C. Doll, and H. Lehmann, Phys. Lett. A **255**(4–6), 221–229 (1999). (A)

Ionization energies for evaluation of ion masses are in the following references. The binding energy uncertainties are insignificant compared to the ion mass uncertainties.

437. The National Institute of Standards and Technology spectroscopic database, ⟨http://physics.nist.gov/ASD⟩. (I)

438. "Recent trends in the determination of nuclear masses," D. Lunney, J. M. Pearson, and C. Thibault, Rev. Mod. Phys. **75**(3), 1021–1082 (2003); see Appendix A. (A)

439. "Atomic and ionic spectrum lines below 2000 Å: Hydrogen through krypton. Part I (H–Cr)," R. L. Kelly, J. Phys. Chem. Ref. Data **16** (Supplement No. 1), 1–649 (1987). (A)

440. "Neutral-atom electron binding energies from relaxed-orbital relativistic Hartree–Fock–Slater calculations $2 \leq Z \leq 106$," K.-N. Huang, M. Aoyagi, M. H. Chen, B. Crasemann, and H. Mark, At. Data. Nucl. Data Tables **18**(3), 243–291 (1976). (A)

M. Other elementary-particle constants

Constants associated with weak and strong interaction physics, such as the Fermi coupling constant, G_F, not included in the preceding sections, are surveyed by the Particle Data Group in periodic publications. The latest references are the following:

441. Particle Data Group Website, ⟨http://pdg.lbl.gov⟩. (I)

442. "Review of particle physics," Particle Data Group, C. Amsler, *et al.* Phys. Lett. B **667**(1–5), 1–1340 (2008). (A)

VI. DERIVED CONSTANTS

The best values of the constants in this section are currently not obtained from direct measurements but instead are calculated from theoretical identities, relating them to directly measured constants considered in the previous section.

A. Elementary charge e

In the classic Millikan oil-drop experiment, the elementary charge e was measured directly, but the current best value is based on the definition of the fine-structure constant as it appears in the SI system, which yields the relation

$$e = (2\epsilon_0 hc\alpha)^{1/2} = \left(\frac{2\alpha h}{\mu_0 c}\right)^{1/2}.$$

The elementary charge is a universal constant because the charge of any isolated system is an integer multiple of e, as far as it has been tested experimentally. Limits can be placed on possible differences between the magnitude of positive and negative elementary charges from measurements of the neutrality of matter.

443. "The electrical neutrality of atoms and of bulk matter," C. S. Unnikrishnan and G. T. Gillies, Metrologia **41**(5), S125–S135 (2004). (I)

444. "Automated electric charge measurements of fluid microdrops using the Millikan method," E. R. Lee, V. Halyo, I. T. Lee, and M. L. Perl, Metrologia **41**(5), S147–S158 (2004). (A)

445. H. Bachmair, "The elementary charge," in **Units and Fundamental Constants in Physics and Chemistry: Fundamental Constants in Physics and Chemistry**, edited by J. Bortfeldt, and B. Kramer, Subvolume b (Springer, Berlin, 1992), Chap. 3.2.4, pp. 79–82. (I)

446. "The electric neutrality of matter: A summary," M. Marinelli and G. Morpurgo, Phys. Lett. **137B**(5–6), 439–442 (1984). (I)

B. Bohr radius a_0

The Bohr radius, which is also the atomic unit of length, is given by

$$a_0 = \frac{\alpha}{4\pi R_\infty}.$$

C. Reduced Compton wavelength of the electron $\lambda\!\!\!^{-}_C$

The reduced Compton wavelength, *i.e.* the Compton wavelength divided by 2π, is given by

$$\lambda\!\!\!^{-}_C = \frac{\hbar}{m_e c} = \alpha a_0.$$

D. Bohr magneton μ_B

The Bohr magneton follows from

$$\mu_B = \frac{e\hbar}{2m_e} = \left(\frac{c\alpha^5 h}{32\pi^2 \mu_0 R_\infty^2}\right)^{1/2}.$$

E. Nuclear magneton μ_N

The nuclear magneton is given by

$$\mu_N = \frac{A_r(e)}{A_r(p)}\mu_B, \tag{1}$$

where $A_r(e)=m_e/m_u$ and $A_r(p)=m_p/m_u$ are the relative atomic masses of the electron and proton and m_u is the relative atomic mass unit given in Sec. VI G.

F. Electron volt, eV

The electron volt is a unit of energy given by

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

$$1\ \text{eV} = \frac{e}{1\ \text{C}} 1\ \text{J},$$

where C is a coulomb and J is a joule.

G. Relative atomic mass unit, u

The relative atomic mass unit is

$$1\ \text{u} = m_{\text{u}} = \tfrac{1}{12} m(^{12}\text{C}) = 10^{-3}\ \text{kg mol}^{-1}/N_{\text{A}},$$

where $m(^{12}\text{C})$ is the mass of the carbon-12 atom.

VII. POSSIBLE REDEFINITION OF SI UNITS

As methods of measurement have advanced over the years, the International System of Units has also evolved. The latest change in the definition of an SI unit was made in 1983 when the meter was redefined by the statement: "The meter is the length of the path traveled by light in vacuum during a time interval of 1/299 792 458 of a second." This definition implies that the speed of light c is exactly 299 792 458 m/s. (See Ref. 73.)

Currently there is interest in providing similar definitions of other SI units in terms of fundamental constants. The Consultative Committee for Units has recommended to the International Committee for Weights and Measures of the International Bureau of Weights and Measures that the kilogram, ampere, kelvin, and mole be redefined by specifying exact values for the Planck constant h, the elementary unit of charge e, the Boltzmann constant k, and the Avogadro constant N_{A}. The change is likely to take place when it is felt that the experiments needed to determine mass based on the proposed new definition are sufficiently accurate.

447. "The quantum SI: A possible new international system of units," P. J. Mohr, Adv. Quantum Chem. **53**, 27–36 (2008), Chap. 3. (I)

448. "Defining units in the quantum based SI," P. J. Mohr, Metrologia **45**(2), 129–133 (2008). (A)

449. "Note on uncertainties resulting from proposed kilogram redefinitions," B. P. Leonard, Metrologia **44**(2), L4–L6 (2007). (A)

450. "On the role of the Avogadro constant in redefining SI units for mass and amount of substance," B. P. Leonard, Metrologia **44**(1), 82–86 (2007). (A)

451. "The atomic units, the kilogram and the other proposed changes to the SI," B. W. Petley, Metrologia **44**(1), 69–72 (2007). (A)

452. "Preparative steps towards the new definition of the kelvin in terms of the Boltzmann constant," J. Fischer, S. Gerasimov, K. D. Hill, G. Machin, M. R. Moldover, L. Pitre, P. Steur, M. Stock, O. Tamura, H. Ugur, D. R. White, I. Yang, and J. Zhang, Int. J. Thermophys. **28**(6), 1753–1765 (2007). (A)

453. "Considerations on future redefinitions of the kilogram, the mole and of other units," P. Becker, P. De Bièvre, K. Fujii, M. Glaeser, B. Inglis, H. Luebbig, and G. Mana, Metrologia **44**(1), 1–14 (2007). (A)

454. "Scope for replacing the prototype kilogram by an atomic mass unit," S. A. Kononogov and V. V. Khrushchev, Meas. Tech. **49**(10), 953–956 (2006). (A)

455. "Some methods for redefining the unit of mass standard," S. A. Kononogov, M. Yu. Konstantinov, and V. V. Khrushchev, Meas. Tech. **49**(4), 311–316 (2006). (A)

456. "On the redefinition of the kilogram and ampere in terms of fundamental physical constants," S. G. Karshenboim, Phys. Usp. **49**(9), 947–954 (2006). (A)

457. "Primary mass standard based on atomic masses," P. Becker and M. Gläser, Int. J. Mass Spectrom. **251**(2–3), 220–230 (2006). (A)

458. "Redefinition of the kilogram, ampere, kelvin and mole: A proposed approach to implementing CIPM recommendation 1 (CI-2005)," I. M. Mills, P. J. Mohr, T. J. Quinn, B. N. Taylor, and E. R. Williams, Metrologia **43**(3), 227–246 (2006). (A)

459. "Base units of the SI, fundamental constants and modern quantum physics," C. J. Bordé, Philos. Trans. R. Soc. London, Ser. A **363**(1834), 2177–2201 (2005). (A)

460. "Quantum standard for units of length, time, mass and fundamental physical constants," S. A. Kononogov and V. V. Khrushchev, Meas. Tech. **48**(8), 739–745 (2005). (A)

461. "Towards an electronic kilogram: an improved measurement of the Planck constant and electron mass," R. L. Steiner, E. R. Williams, D. B. Newell, and R. Liu, Metrologia **42**(5), 431–441 (2005). (A)

462. "Possible new definitions of the kilogram," R. S. Davis, Philos. Trans. R. Soc. London, Ser. A **363**(1834), 2249–2264 (2005). (A)

463. "The kilogram redefinition—An interim solution," J. L. Flowers and B. W. Petley, Metrologia **42**(5), L31–L34 (2005). (A)

464. "Redefinition of the kilogram: A decision whose time has come," I. M. Mills, P. J. Mohr, T. J. Quinn, B. N. Taylor, and E. R. Williams, Metrologia **42**(2), 71–80 (2005). (A)

465. "Redefining the kilogram," I. Robinson, Phys. World **17**(5), 31–35 (2004). (E)

466. "Wer gewinnt den Wettlauf um das Kilogramm?," E. O. Göbel, Phys. Bl. **57**(1), 35–41 (2004). (A)

467. "Towards a new kilogram definition based on a fundamental constant," W. Schwitz, B. Jeckelmann, and P. Richard, C. R. Phys. **5**(8), 881–892 (2004). (A)

468. "Tracing the atomic mass unit to the kilogram by ion accumulation," M. Gläser, Metrologia **40**(6), 376–386 (2003). (A)

469. "The SI unit of mass," R. Davis, Metrologia **40**(6), 299–305 (2003). (I)

470. "Tracing the definition of the kilogram to the Avogadro constant using a silicon single crystal," P. Becker, Metrologia **40**(6), 366–375 (2003). (A)

471. "Avogadro constant and ion accumulation: Steps towards a redefinition of the SI unit of mass," P. Becker and M. Gläser, Meas. Sci. Technol. **14**(8), 1249–1258 (2003). (A)

472. "Replacing the kilogram," B. P. Kibble and I. A. Robinson, Meas. Sci. Technol. **14**(8), 1243–1248 (2003). (A)

473. "The role of fundamental constants in the international system of units (SI): Present and future," B. N. Taylor and P. J. Mohr, IEEE Trans. Instrum. Meas. **50**(2), 563–567 (2001). (A)

474. "On the redefinition of the kilogram," B. N. Taylor and P. J. Mohr, Metrologia **36**(1), 63–64 (1999). (A)

475. "The silicon path to the kilogram," P. Becker, H. Bettin, L. Koenders, J. Martin, A. Nicolaus, and S. Röttger,

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11

PTB-Mitt. **106**(5), 321–329 (1996). (A)

476. "Early history and future outlook for the x-ray crystal density method," R. D. Deslattes, Metrologia **31**(3), 173–179 (1994). (A)

477. "The third periodic verification of national prototypes of the kilogram (1988–1992)," G. Girard, Metrologia **31**(4), 317–336 (1994). (A)

478. "The stability of the SI unit of mass as determined from electrical measurements," R. S. Davis, Metrologia **26**(1), 75–76 (1989). (A)

479. "The atomic mass unit," B. W. Petley, IEEE Trans. Instrum. Meas. **38**(2), 175–179 (1989). (A)

ACKNOWLEDGMENT

The authors thank their NIST colleague, B. N. Taylor, for his help in maintaining the bibliographic database of the NIST Fundamental Constants Data Center used in the preparation of this Resource Letter.



X Ray Photograph. The photograph shows an instructor at the Kenyon Military Academy using Kenyon College apparatus to make an X ray photograph of his hand. The date is about 1905, and the picture was taken in Ascension Hall on the Kenyon campus. The tube is still in existence and the glass in front of the cathode, through which the X rays passed, has been colored a light purple due to the formation of color centers. The photographic plate is wrapped up in opaque paper and is underneath the instructor's hand. Note the complete absence of shielding from stray radiation. (Photograph and Notes by Thomas B. Greenslade, Jr., Kenyon College)

This article is copyrighted as indicated in the article. Reuse of AAPT content is subject to the terms at: http://scitation.aip.org/termsconditions. Downloaded to IP: 129.6.169.29 On: Wed, 23 Sep 2015 20:29:11